
# ENVIRONMENTAL BIOLOGY OF FISHES AND GEAR PERFORMANCE IN THE PELACIC TUNA LONGLINE FISHERY

Liming Song



# Environmental Biology of Fishes and Gear Performance in the Pelagic Tuna Longline Fishery

Liming Song





#### 图书在版编目 (CIP) 数据

金枪鱼延绳钓渔业中的鱼类环境生物学和渔具性能研究=Environmental Biology of Fishes and Gear Performance in the Pelagic Tuna Longline Fishery: 英文/宋利明著. 一北京: 科学出版社, 2015.10

ISBN 978-7-03-046108-7

I. ①金··· II. ①宋··· III. ①金枪鱼-海洋渔业-环境生物学-研究-英文 ②金枪鱼-海洋渔业-渔具-研究-英文 IV. ①S97

中国版本图书馆 CIP 数据核字 (2015) 第 253499 号

责任编辑:陈露文茜/责任校对:郑金红责任印制:谭宏宇/封面设计:殷靓

#### 斜学出版社出版

北京东黄城根北街 16号 邮政编码: 100717 http://www.sciencep.com

江苏凤凰数码印务有限公司印刷

科学出版社发行 各地新华书店经销

2015年10月第 — 版 开本: 787×1092 1/16 2015年10月第一次印刷 印张: 11 字数: 260 000

定价: 98.00元

(如有印装质量问题, 我社负责调换)

## This book is funded by the following programmes

- 1. National High Technology Research and Development Program of China (2012AA 092302)
- 2. Specialized Research Fund for the Doctoral Program of Higher Education (20113104 110004)
- 3. The Shanghai Municipal Education Commission Innovation Project (12ZZ168)
- 4. The Ministry of Agriculture of the P.R of China under Project of Distant Water Fishery Exploration from 2005 to 2013
- 5. The First-class Discipline Construction Project of Shanghai Higher Education, the Construction of Key Disciplines Project of Shanghai Municipal (S30702)
- 6. The 2013 Shanghai Municipal Professional Graduate Education Reform Pilot Project

## Preface

In recent decades, the offshore fishery resources over the world have been depleted due to climate change, environmental pollution, overfishing and other factors, while there are still increasing demands on the marine living resources. Therefore, the international society has attached great importance to the issue as how to achieve equilibrium between the development of fishery economy and the conservation of fishery resources. After the entry into force of the 1982 UN Convention on the Law of the Sea, the coastal countries have further improved the capability of management and protection of fishery resources within their jurisdiction and strengthened the management of fishing operation from foreign countries in their Exclusive Economic Zones. Subsequently, more than 10 regional fisheries management organizations have been established in the world to manage the high value species on the high seas of the three oceans and in the Antarctic Ocean.

Tuna and tuna like species are the large pelagic fish and belong to the highly migratory species. Although most of the tuna resources have been in full exploitation at the end of the last century, the tuna resources are still the key fishery resources exploited by distant water fishing nations or entities due to its high economic value (the fishing industry called the tuna fisheries as "golden fisheries").

The biological characteristics, the habitat of tunas and the longline fishing gear performance are a part of the hot topics of the regional tuna fishery management organizations of the world. This book illustrates the age and growth of bigeye tuna, the biological characteristics of tunas, the integrated habitat index of yellowfin tuna, catch rate calculation methods, and effects of environmental factors and fishing gear on catch performance. It provides references for the reduction of uncertainties in the tuna resource assessment, the standardization of the tuna abundance index (CPUE), the improvement of stock assessment accuracy, the study of the habitat selection, the bycatch mitigation of sharks, turtles and seabirds, protection to marine ecological environment, the improvement of the catch rate of target species, and the study of the biological characteristics, habitat of tunas and longline fishing gear performance.

This book is intended as a reference for fishery researchers, students, managers, conservation enthusiasts, and fisheries industrials. It can also be used as a textbook for fisheries science and technology at undergraduate and graduate levels.

Shanghai Ocean University has been engaged in the research of the fishery biology, fishery oceanography, and fishing technology for a long time. During the past decade, studies have been conducted on the environment biology of tunas and longline fishing gear performance, which have been funded by the National High Technology Research and Development Program

of China (2012AA092302), Specialized Research Fund for the Doctoral Program of Higher Education (20113104110004), the Shanghai Municipal Education Commission Innovation Project (12ZZ168), the Ministry of Agriculture of the P.R of China under Project of Distant Water Fishery Exploration from 2005 to 2013, the First-class Discipline Construction Project of Shanghai Higher Education, the Construction of Key Disciplines Project of Shanghai Municipal (S30702), the 2013 Shanghai Municipal Professional Graduate Education Reform Pilot Project. There were more than 20 graduate students (Kaikai Lv, Zhenxin Hu, Yaping Wu, Mingming Hui, Jialiang Yang, Weiyun Xu, Jie Li, Dongjing Li, Haiyang Liu, Hao Chen, et al.) having participated in this research field. During the survey, Chinese Overseas Fisheries Association, Guangvuang Fishery Group Ltd of Guangdong province, Liancheng Overseas Fishery (Shenzhen) Co. Ltd., Zheijang Ocean Family Co. Ltd., and China Southern Fishery (Shenzhen) Co. Ltd. have also offered strong support. Thanks go to them for their effort and support. I sincerely thank professors Yong Chen at University of Maine, Pingguo He of the University of Massachusetts, and Senior scientist Keith Bigelow of the Pacific Islands Fisheries Science Center, NOAA Fisheries, et al. for their extraordinary efforts in reviewing some of the parts, as well as to those experts who have evaluated our manuscripts and page proofs and provided valuable comments and suggestions!

I am grateful to my wife, Jie Wang, and Son, Haobo Song, for their strong support. It would not have been possible without their support and understanding.


This book is one of the stage achievements of the author in the tuna longline fisheries research, and is only a portion of China's pelagic tuna longline fisheries research. Finally, the critical feedback for improving our work is always welcome!

Liming Song
Shanghai Ocean University
July 2015

# Contents

	Age and growth of bigeye tuna	
	e and growth of bigeye tuna, using fin spines	
	Introduction	
1.1.2	Materials and methods····	
1.1.3		
	Discussions ····	
	e and growth of bigeye tuna, using vertebra ·····	
1.2.1	Introduction	
1.2.2		
1.2.3		
1.2.4	Discussions ····	17
Chapter 2 1	Biological characteristics of tunas	
2.1 A c	comparison of biological characteristics of yellowfin tuna	22
2.1.1		
2.1.2	Materials and methods	23
2.1.3	Results	25
2.1.4	Discussions ·····	31
2.2 The	e length structure of bigeye tuna and yellowfin tuna catch	36
2.2.1	Introduction	36
2.2.2	Materials and methods·	38
2.2.3	Results	43
2.2.4	Discussions ····	45
Chapter 3	Integrated habitat index	
3.1 Ac	comparison of methods about the prediction of integrated habitat index	49
3.1.1	Introduction	49
3.1.2	Materials and methods	50
3.1.3	Results	57
3.1.4	Discussions	61
3.2 Ac	comparison on calculation methods about an integrated habitat index	67
3.2.1	Introduction	67
3.2.2	Materials and methods	
3.2.3	Results	73
324	Discussions	75

Chapter 4	Tuna catch rate calculation methods
4.1 A c	omparison of two catch rate calculation methods79
4.1.1	Introduction 79
4.1.2	Materials and methods 81
4.1.3	Results 86
4.1.4	Discussions 90
4.2 A c	omparison of catch rate on bigeye tuna of two longline fishing gears94
4.2.1	Introduction 94
4.2.2	Materials and methods 95
4.2.3	Results 99
4.2.4	Discussions 101
Chapter 5	Effects of fishing gear and environmental factors on catch performances
5.1 A c	omparison of catch performance between circle hooks and tuna hooks
5.1.1	Introduction 105
5.1.2	Materials and methods 106
5.1.3	Results 108
5.1.4	Discussions 110
5.2 Eff	ects of environmental factors and fishing gear on catch rates of sharks
5.2.1	Introduction 113
5.2.2	Materials and methods 115
5.2.3	Results 120
5.2.4	Discussions 125
5.3 The	e relationships between the catch rates and the depth and sea surface
tem	perature 134
5.3.1	Introduction 134
5.3.2	Materials and methods 136
5.3.3	Results 139
5.3.4	Discussions 141
5.4 Dev	velopment of a three-dimensional numerical model for pelagic longlines ·······146
5.4.1	Introduction 146
5.4.2	Materials and methods 148
5.4.3	Results 158
5 4 4	Discussions



## 1.1 Age and growth of bigeye tuna, using fin spines

**Abstract** In order to quantitatively study the morphological characteristics, the phenomena of vascularized core of bigeye tuna (Thunnus obesus) fin spines, and the age and growth of bigeye tuna, a total of 1304 bigeye tuna were sampled in the waters near Marshall Islands from November 2009 to January 2010 and their fork length were measured. About 436 dorsal spines have been randomly sampled. We also measured the total length (L) and the width (C) at the base of the fin spines. In the laboratory, the cross sections of the spines were obtained in the position above the condyle base in the distance of half C with an Isomet Low Speed Saw. The cross sections ranging from 400-500 µm thickness were examined with a dissecting microscope with transmitted light. Images of the sections were captured and analysed by the software Image-Pro plus version 3.1. In this study, the following parameters were measured for each spine: the total surface area of the cross section (S, mm<sup>2</sup>), the surface area of the vascularized core (S<sub>0</sub>, mm<sup>2</sup>), and the annual ring surface areas of one to three years old fish including the translucent rings  $(S_1, S_2, S_3, \text{mm}^2)$ . The results showed that: ①The relationships between the L, C and the fork length were the power function; @We could identify the age of one to three years old fish because of the vascularized core; 3The relationships between S and fork length,  $S_0$  and fork length were the power function and the ratio of  $S_0$  to S was about 0.4; ④ From the frequency analysis of the translucent ring surface area, the fork length of bigeye tuna from one to three years old fish was 48 cm, 71 cm, and 90 cm.

Keywords Thunnus obesus, fin spine, age, growth, Marshall Islands

#### 1.1.1 Introduction

The age structure is an important parameter to the growth rate, mortality, productivity of fish stocks, and the stock assessment (Ragonese and Reale, 1992; Campana and Thorrold, 2001). There are many methods to identify the age of fish, such as tagging and releasing, length

frequency and the hard tissue, scales (Yukinawa and Yabuta, 1963), vertebra (Alves *et al.*, 2002), otolith (Farley *et al.*, 2006), fin spine (Gaikov *et al.*, 1980; Antoine *et al.*, 1983; Sun *et al.*, 2001; Stéquert and Conand, 2003).

Lehodey and Leroy (1999) used the tag releasing and otolith to study the age of bigeye tuna (*Thunnus obesus*) in the Western and Central Pacific. Cayre and Diauf (1984) tagged and released 139 bigeye tuna and recapture them in the eastern Atlantic to study its growth. Hallier *et al.* (2005) studied the age and growth of bigeye tuna by tagging and releasing 625 bigeye tuna in the eastern Atlantic.

The structure of rings of fin spine is comprised of euphotic zone and opaque zone. There were many studies about the age and growth of the fish by reading the number of rings of the fin spine cross-section (Gaikov et al., 1980; Antoine et al., 1983; Alves et al., 1998; Sun et al., 2001; Stéquert and Conand, 2003). Alves et al. (1998) used the first dorsal fin spine to study the age and growth of bigeye tuna in the Madeira Archipelago waters and found that the age and growth of different gender bigeye tuna were basically the same and estimated the fork length of the one to eight years old fish. Stéquert and Conand (2003) suggested that the method of using sliced spine was not suitable to study the age because the cross-section of the sliced spines was not clear, each ring could not be accurately identified, and the incorrect number of rings would cause the bias. Gaikov et al. (1980) and Draganik and Pelczarski (1984) used spines to identify the age of bigeye tuna in the Atlantic. Sun et al. (2001) found that the growth ring of dorsal fin spines of bigeye tuna in the Western and Central Pacific was comprised of euphotic zone and opaque zone. They suggested that the bigeye tuna dorsal fin spines were convenient to be collected and conserved, the structure of rings were clear, and could be used to identify the age. But they also found that the area of the vascularized core continued to expand with the fish growth. In order to quantitatively study the morphological characteristics, the phenomenon of the vascularized core of bigeye tuna fin spines, and accurately identify the age of bigeye tuna, the bigeye tuna dorsal fin spines were collected in the waters near Marshall Islands, and the relationship between fork length and morphology size of fin spine were analyzed.

#### 1.1.2 Materials and methods

#### Sampling duration and area

The sampling duration was from November, 2009 to January, 2010. The samples were from the longliners of Liancheng Overseas Fishery (Shenzhen) Corporation fishing in waters near Marshall Islands. The sampling area was defined as 3°30′N-5°00′N, 166°48′E-169°48′E (Fig. 1-1-1).

#### Data collection

We used caliper to measure the fork length of bigeye tuna in Majuro tuna processing factory of Liancheng Overseas Fishery (Shenzhen) Corporation. The measurement accuracy was 0.5 cm.

The fork length of 1304 bigeye tuna was measured, and randomly sampled 436 spines of bigeye tuna. In this study, the fin spine was sampled from the first ray of the first dorsal fin.




Fig. 1-1-1 Sampling area of bigeye tuna in waters near Marshall Islands

#### Spine processing

Spine processing procedures were as follows:

- (1) Removing the muscle at the base of the fin spine with a knife.
- (2) Immersing the spine quickly into the boiling water (the duration depends on the spine size), taking it out and cooling it in the cold water.
  - (3) Removing the remaining connective tissue and black epidermis of fin spine.
  - (4) Putting it into 4% KOH solution to immerse 5 hours.
  - (5) Rinsing with water.
- (6) Measuring the total length (L, mm) and the width at the base of the fin spines (C, mm) after drying, L and C were shown in Fig. 1-1-2.

We assumed that the growth of hard part was proportional to the fish growth. The power regression was used to fit the relationship between the fin spine length L, width at the base C and the fish fork length FL.

Spine cross section was obtained in the position above the condyle base in the distance of half C (Fig. 1-1-2) with an Isomet Low Speed Saw (Buehler, Isomet low-speed saw). The thickness of spine cross section was about 2 mm. It was polished to about 1 mm by the 600 and 2000 grist waterproof and sandpaper. Lastly, we used 0.3  $\mu$ m thickness alumina flannel to polish it to show the clear rings. The cross sections ranging from 400-500  $\mu$ m thickness were examined with a dissecting microscope (Nikon ZOOM645S) with transmitted light.

We took the picture of spine cross section and input it to the computer for the data measurements.




Fig. 1-1-2 Schema of the dorsal spine and the section of the dorsal spine

#### Ring structure interpretation and measurement

Spine cross section under a microscope showed the translucent zone and the opaque zone alternately. We could not estimate the ring formation time because the sampling duration was only two and a half month. Based on the conclusions of other scholars [the translucent zone formed in March (Nose et al., 1967) ], we identify the ring and age of bigeye tuna.

By observing the cross section of fin spines (Fig. 1-1-2), we found the opaque zone with translucent zone formed the annual ring. It was impossible to read each ring accurately because the spine sliced cross-section was not clear or vascularized. The incorrect numbers of ring would influence the accuracy of the result. In this study, we used the ring area method proposed by Stéquert and Conand (2003) to study the ring area of fin spine and the age of bigeye tuna. Analysing the spine cross-section photograph by the software Image-Pro plus version 3.1, the ring area of each annual ring  $(S_1, S_2, S_3, mm^2)$ , the slice surface area  $(S, mm^2)$  and vascular area  $(S_0, \text{ mm}^2)$  (Fig. 1-1-2) were measured. The boundary line between the out edge of the translucent zone and opaque zone was the measuring baseline. The data of the clear ring were only measured. The unclear or incomplete rings were discarded. Comparing the frequency distribution for all measured ring area, the ring area might be the ring area of growth ring if the frequency of ring area was relatively high.

#### 1.1.3 Results

#### Relationships between spine size and fork length

The first dorsal fin spine length L and the base width C were measured for 389 bigeye tuna

(Fig. 1-1-2), and we collected 375 spine length L and 180 complete spine width C. The relationships between the fin spine length L, base width C and fork length FL were shown in equations 1-1-1 and 1-1-2 (Fig. 1-1-3 and Fig. 1-1-4).

$$L = 1.758FL^{0.937}, R^2 = 0.900$$
 (1-1-1)

$$C = 0.278FL^{0.983}, R^2 = 0.884$$
 (1-1-2)




Fig. 1-1-3 The relationship between fork length and the length of dorsal fin spine




Fig. 1-1-4 The relationship between fork length and the width of dorsal fin spine

#### The relationship between fin spine area and fork length

The spine cross section surface area S (mm<sup>2</sup>) of 179 fish was measured. The relationship between fork length FL (cm) and the spine cross section surface area was shown in equation 1-1-3 (Fig. 1-1-5). We found that the greater the cross section of spine was, the greater the area of vascularized core  $S_0$  (mm<sup>2</sup>) was. So, in this study, the area of vascularized core  $S_0$  of the fin spine cross section was measured. The relationship between fork length FL and the area of vascularized core  $S_0$  was indicated in equation 1-1-4 (Fig. 1-1-6).

$$S = 6.220 \times 10^{-4} FL^{2.438}, R^2 = 0.955$$
 (1-1-3)

$$S_0 = 2.607 \times 10^{-4} FL^{2.497}, R^2 = 0.900$$
 (1-1-4)

Based on equation 1-1-3 and equation 1-1-4, the ratio between the surface area of the vascularized core and the area of the dorsal fin spine was 0.41-0.45 while the fork length of bigeye tuna was 45.0-170.0 cm (Fig. 1-1-7). The area of vascularized core expanded with the fork length increasing.




Fig. 1-1-5 The relationship between fork length and the surface area of section of dorsal fin spine




Fig. 1-1-6 The relationship between fork length and the surface area of vascularized core of dorsal fin spine




Fig. 1-1-7 The ratio between the surface area of vascularized core and the area of dorsal fin spine

#### The average growth rate of the dorsal fin spine

The ratio between the surface area of the vascularized core and the area of the dorsal fin spine was 0.41-0.45. The ring in the first round would be into the area of vascularized core when the fork length was greater than 45 cm, but the ring in the first round has been seen. However, as the fork length was greater than 90 cm, the first round of the spine became very illegibility. When the fork length was greater than 100 cm, the ring in the first round would disappear, and it was only to be estimated by the experience. Therefore, this study only identified the rings of fin spine when the fork length was less than 100 cm for age determination. The frequency distribution of the area of fin spine cross section was indicated in Fig. 1-1-8. The optimal area of fin spine cross section was

8 mm<sup>2</sup>, 20 mm<sup>2</sup>, 36 mm<sup>2</sup>, 54 mm<sup>2</sup>, 70 mm<sup>2</sup>, and 82 mm<sup>2</sup>, for the one to six years old fish, and their corresponding fork length was 48 cm, 71 cm, 90 cm, 106 cm, 118 cm, and 126 cm, respectively. We only used the fork length data of one to three years old fish to estimate the growth rate of fish because the fin spine was only suitable to identify the age of bigeye tuna which fork length was below 100 cm. The fork length of one to three years old fish was 48 cm, 71 cm, and 90 cm. Their average growth rate was 48 cm/yr, 23 cm/yr, and 19 cm/yr, respectively.




Fig. 1-1-8 Frequency distribution of surface area of the dorsal fin spine cross section

#### 1.1.4 Discussions

This study suggested that the fin spine was only suitable to identify the age of bigeye tuna whose age was smaller than three years old. If the fish was older than three years, the vascularized core would affect the accuracy of the annual ring data. The numbers of annual ring were only identified by the subjective judgments, and the results would be in bias. For the serious vascularized core of the fin spine section, it was impossible to accurately read each ring of the cross section, especially the first and second ring. In this study, by measuring the area of vascularized core of dorsal fin spine, we found that the area of vascularized core of the dorsal fin spine was about 0.4 of the total area of the fin spine. We calculated the area of vascularized core of the dorsal spine of two years old fish and found that the location of the first ring was near the edge of the vascularized core. When the fish was three years old, the first ring was in the inner of the area of vascularized core, but it could be identified. When the fish was four years old, the first ring was completely disappear.

This study suggested that the relationships between the fin spine length and base width of bigeye tuna and fork length was a power function (equation 1-1-1 and equation 1-1-2), and the fin spine was the suitable material to identify the fish age smaller than three years old. Sun *et al.* (2001) studied the age of bigeye tuna in Western and Central Pacific (FL=70-189 cm) by spine, and found the relationship between the length of fin spine and fork length was a linear function. This was generally consistent with our results. Li (2010) studied the relationship between the base width of fin spine and the fork length of bigeye tuna in the Indian Ocean, and suggested there was no corelation. This might be caused by the small samples of the base width of fin spine.

The ring area method proposed by Stéquert and Conand (2003) avoided the shortcomings of reader's subjectivity for judgment, and the precision was high. Stéquert and Conand (2003) used the method to infer the fork length of one to three years old bigeye tuna, and pointed out that this method wasn't suitable when the samples were over three years old. They did not give specific reasons. In this study, we measured the area of vascularized core of the dorsal spine, and found the first ring was disappeared in the inner area of vascularized core while the age was over three years old. That was the reason why the fin spine was only suitable to identify the age of fish which was three years old or small. The fork length of one to three years old fish in this study was similar to the results obtained by Cayré and Diouf (1984) and Alves *et al.* (2002). There were some differences from the other studies (Shomura and Keala, 1963; Yukinawa and Yabuta, 1963; Kume and Joseph, 1966; Gaikov *et al.*, 1980; Lehodey *et al.*, 1999; Li, 2010) (Tab. 1-1-1). That might be resulted from the different methods and study areas.

Author	Time	Area	1	2	3
This study	2011	Marshall Islands	48	71	90
Shomura & Keala	1963	Pacific	67	101	125
Yukinawa & Yabuta	1963	Pacific	41	73	100
Kume & Joseph	1966	the eastern Pacific	31	80	114
Gaikov et al.	1980	Atlantic	46	79	107
Cayré & Diauf	1984	Atlantic	44	69	92
Lehodey	1999	Western and Central Pacific	63	96	123
Alves	2002	Madeira Archipelago	48	73	94
Li	2010	Indian Ocean	58	79	97

Tab. 1-1-1 The bigeve tuna fork length of ages one to three in various studies

In this study, we did not consider the impact of gender on various parameters. We could not use the method of marginal increment to analysis and verify the formation time of annual ring. The samples were from longliners, and the individuals whose fork length smaller than 85 cm or higher than 160 cm were less. That might cause the samples smaller than one to two years old fish and higher than seven years old fish less.

In the future study, the sampling area, time and rang of individual size should be increased. We should use more suitable methods to identify the age structure by vertebrae, otolith, scale, and tagging, and compare the results to achieve an optimal method to identify the age of bigeye tuna. Combining with the biological data of gonad of bigeye tuna, we can study the growth of bigeye tuna in different life span (such as maturity and ovulation).

#### Acknowledgements

The project is granted by Ministry of Fisheries of the Republic of Marshall Islands. The project is funded by the National High Technology Research and Development Program of China (Project No. 2012AA092302), the Ministry of Agriculture of P.R of China under Project of Distant-water Fishery Exploration in 2009 (Project No. D8006090066), Shanghai Municipal Education