

国外数学名著系列 (续一)

(影印版) 66

Yu. V. Egorov M. A. Shubin (Eds.)

Partial Differential Equations IV
Microlocal Analysis and Hyperbolic Equations

偏微分方程 IV
微局部分分析和双曲型方程

图字: 01-2008-5381

Yu. V. Egorov, M. A. Shubin(Eds.): Partial Differential Equations IV: Microlocal Analysis and Hyperbolic Equations

© Springer-Verlag Berlin Heidelberg 1993

This reprint has been authorized by Springer-Verlag (Berlin/Heidelberg/New York) for sale in the People's Republic of China only and not for export therefrom

本书英文影印版由德国施普林格出版公司授权出版。未经出版者书面许可,不得以任何方式复制或抄袭本书的任何部分。本书仅限在中华人民共和国销售,不得出口。版权所有,翻印必究。

图书在版编目(CIP)数据

偏微分方程 IV: 微局部分分析和双曲型方程=Partial Differential Equations IV: Microlocal Analysis and Hyperbolic Equations/(俄罗斯) 叶戈罗夫(Egorov, Y. V.)等编著. —影印版. —北京: 科学出版社, 2009

(国外数学名著系列; 66)

ISBN 978-7-03-023509-1

I. 偏… II. 叶… III. ①偏微分方程-英文 ②双曲型方程-英文
IV. O175.2

中国版本图书馆 CIP 数据核字(2008) 第 186169 号

责任编辑: 范庆奎/责任印刷: 钱玉芬/封面设计: 黄华斌

科学出版社出版

北京东黄城根北街 16 号

邮政编码: 100717

<http://www.sciencep.com>

双青印刷厂印刷

科学出版社发行 各地新华书店经销

*

2009 年 1 月第 一 版 开本: B5(720 × 1000)

2009 年 1 月第一次印刷 印张: 16

印数: 1—2 500 字数: 304 000

定价: 58.00 元

(如有印装质量问题, 我社负责调换〈双青〉)

《国外数学名著系列》(影印版)专家委员会

(按姓氏笔画排序)

丁伟岳 王 元 文 兰 石钟慈 冯克勤 严加安
李邦河 李大潜 张伟平 张继平 杨 乐 姜伯驹
郭 雷

项目策划

向安全 林 鹏 王春香 吕 虹 范庆奎 王 璐

执行编辑

范庆奎

《国外数学名著系列》(影印版)序

要使我国的数学事业更好地发展起来,需要数学家淡泊名利并付出更艰苦地努力。另一方面,我们也要从客观上为数学家创造更有利的发展数学事业的外部环境,这主要是加强对数学事业的支持与投资力度,使数学家有较好的工作与生活条件,其中也包括改善与加强数学的出版工作。

从出版方面来讲,除了较好较快地出版我们自己的成果外,引进国外的先进出版物无疑也是十分重要与必不可少的。从数学来说,施普林格(Springer)出版社至今仍然是世界上最具权威的出版社。科学出版社影印一批他们出版的好的新书,使我国广大数学家能以较低的价格购买,特别是在边远地区工作的数学家能普遍见到这些书,无疑是对推动我国数学的科研与教学十分有益的事。

这次科学出版社购买了版权,一次影印了 23 本施普林格出版社出版的数学书,就是一件好事,也是值得继续做下去的事情。大体上分一下,这 23 本书中,包括基础数学书 5 本,应用数学书 6 本与计算数学书 12 本,其中有些书也具有交叉性质。这些书都是很新的,2000 年以后出版的占绝大部分,共计 16 本,其余的也是 1990 年以后出版的。这些书可以使读者较快地了解数学某方面的前沿,例如基础数学中的数论、代数与拓扑三本,都是由该领域大数学家编著的“数学百科全书”的分册。对从事这方面研究的数学家了解该领域的前沿与全貌很有帮助。按照学科的特点,基础数学类的书以“经典”为主,应用和计算数学类的书以“前沿”为主。这些书的作者多数是国际知名的大数学家,例如《拓扑学》一书的作者诺维科夫是俄罗斯科学院的院士,曾获“菲尔兹奖”和“沃尔夫数学奖”。这些大数学家的著作无疑将会对我国的科研人员起到非常好的指导作用。

当然,23 本书只能涵盖数学的一部分,所以,这项工作还应该继续做下去。更进一步,有些读者面较广的好书还应该翻译成中文出版,使之有更大的读者群。

总之,我对科学出版社影印施普林格出版社的部分数学著作这一举措表示热烈的支持,并盼望这一工作取得更大的成绩。

王 元

2005 年 12 月 3 日

List of Editors, Authors and Translators

Editor-in-Chief

R.V. Gamkrelidze, Russian Academy of Sciences, Steklov Mathematical Institute, ul. Vavilova 42, 117966 Moscow, Institute for Scientific Information (VINITI), ul. Usievicha 20a, 125219 Moscow, Russia

Consulting Editors

Yu.V. Egorov, Department of Mathematics, Moscow State University, Leninskie Gory, 119899 Moscow, Russia
M.A. Shubin, Department of Mathematics, Moscow State University, Leninskie Gory, 119899 Moscow, Russia

Authors

Yu.V. Egorov, Department of Mathematics, Moscow State University, Leninskie Gory, 119899 Moscow, Russia
V.Ya. Ivrii, Department of Mathematics, Magnitogorsk Mining-Metallurgical Institute, 455000 Magnitogorsk, Russia, and
Department of Mathematics, University of Toronto, 100 St. George Street, Toronto, M5S 1A1, Ontario, Canada

Translator

P.C. Sinha, Jagat Narayan Road, 800003 Patna, India

Contents

I. Microlocal Analysis

Yu.V. Egorov

1

II. Linear Hyperbolic Equations

V.Ya. Ivrii

149

Author Index

237

Subject Index

240

I. Microlocal Analysis

Yu.V. Egorov

Translated from the Russian
by P.C. Sinha

Contents

Preface	7
Chapter 1. Microlocal Properties of Distributions	8
§ 1. Microlocalization	8
§ 2. Wave Front of Distribution. Its Functorial Properties	9
2.1. Definition of the Wave Front	9
2.2. Localization of Wave Front	10
2.3. Wave Front and Singularities of One-Dimensional Distributions	11
2.4. Wave Fronts of Pushforwards and Pullbacks of a Distribution	12
§ 3. Wave Front and Operations on Distributions	12
3.1. The Trace of a Distribution. Product of Distributions	12
3.2. The Wave Front of the Solution of a Differential Equation	13
3.3. Wave Fronts and Integral Operators	13
Chapter 2. Pseudodifferential Operators	14
§ 1. Algebra of Pseudodifferential Operators	14
1.1. Singular Integral Operators	14
1.2. The Symbol	15
1.3. Boundedness of Pseudodifferential Operators	16
1.4. Composition of Pseudodifferential Operators	16
1.5. The Formally Adjoint Operator	17
1.6. Pseudolocality. Microlocality	17
1.7. Elliptic Operators	17
1.8. Gårding's Inequality	18
1.9. Extension of the Class of Pseudodifferential Operators	18

§ 2. Invariance of the Principal Symbol Under Canonical Transformations	18
2.1. Invariance Under the Change of Variables	18
2.2. The Subprincipal Symbol	19
2.3. Canonical Transformations	20
2.4. An Inverse Theorem	20
§ 3. Canonical Forms of the Symbol	21
3.1. Simple Characteristic Points	21
3.2. Double Characteristics	23
3.3. The Complex-Valued Symbol	23
3.4. The Canonical Form of the Symbol in a Neighbourhood of the Boundary	23
§ 4. Various Classes of Pseudodifferential Operators	24
4.1. The $L_{\rho,\delta}^m$ Classes	24
4.2. The $L_{\phi,\phi}^m$ Classes	26
4.3. The Weyl Operators	29
§ 5. Complex Powers of Elliptic Operators	32
5.1. The Definition of Complex Powers	32
5.2. The Construction of the Symbol for the Operator A^z	33
5.3. The Construction of the Kernel of the Operator A^z	35
5.4. The ζ -Function of an Elliptic Operator	36
5.5. The Asymptotics of the Spectral Function and Eigenvalues ...	37
5.6. Complex Powers of an Elliptic Operator with Boundary Conditions	38
§ 6. Pseudodifferential Operators in $\mathbb{R}^n$ and Quantization	39
6.1. The Analogy Between the Microlocal Analysis and the Quantization	39
6.2. Pseudodifferential Operators in $\mathbb{R}^n$	41
Chapter 3. Fourier Integral Operators	43
§ 1. The Parametrix of the Cauchy Problem for Hyperbolic Equations	43
1.1. The Cauchy Problem for the Wave Equation	43
1.2. The Cauchy Problem for the Hyperbolic Equation of an Arbitrary Order	44
1.3. The Method of Stationary Phase	45
§ 2. The Maslov Canonical Operator	46
2.1. The Maslov Index	46
2.2. Pre-canonical Operator	47
2.3. The Canonical Operator	49
2.4. Some Applications	49
§ 3. Fourier Integral Operators	50
3.1. The Oscillatory Integrals	50
3.2. The Local Definition of the Fourier Integral Operator	52

3.3. The Equivalence of Phase Functions	52
3.4. The Connection with the Lagrange Manifold	53
3.5. The Global Definition of the Fourier Distribution	55
3.6. The Global Fourier Integral Operators	56
§ 4. The Calculus of Fourier Integral Operators	58
4.1. The Adjoint Operator	58
4.2. The Composition of Fourier Integral Operators	59
4.3. The Boundedness in L_2	61
§ 5. The Image of the Wave Front Under the Action of a Fourier Integral Operator	62
5.1. The Singularities of Fourier Integrals	62
5.2. The Wave Front of the Fourier Integral	63
5.3. The Action of the Fourier Integral Operator on Wave Fronts	64
§ 6. Fourier Integral Operators with Complex Phase Functions	65
6.1. The Complex Phase	65
6.2. Almost Analytic Continuation	65
6.3. The Formula for Stationary Complex Phase	66
6.4. The Lagrange Manifold	67
6.5. The Equivalence of Phase Functions	68
6.6. The Principal Symbol	69
6.7. Fourier Integral Operators with Complex Phase Functions ...	71
6.8. Some Applications	72
Chapter 4. The Propagation of Singularities	73
§ 1. The Regularity of the Solution at Non-characteristic Points	73
1.1. The Microlocal Smoothness	73
1.2. The Smoothness of Solution at a Non-characteristic Point	73
§ 2. Theorems on Removable Singularities	74
2.1. Removable Singularities in the Right-Hand Sides of Equations	74
2.2. Removable Singularities in Boundary Conditions	75
§ 3. The Propagation of Singularities for Solutions of Equations of Real Principal Type	76
3.1. The Definition and an Example	76
3.2. A Theorem of Hörmander	76
3.3. Local Solvability	77
3.4. Semiglobal Solvability	77
§ 4. The Propagation of Singularities for Principal Type Equations with a Complex Symbol	78
4.1. An Example	78
4.2. The Fixed Singularity	79
4.3. A Special Case	79
4.4. The Propagation of Singularities in the Case of a Complex Symbol of the General Form	80

§ 5. Multiple Characteristics	80
5.1. Non-involutive Double Characteristics	80
5.2. The Levi Condition	81
5.3. Operators Having Characteristics of Constant Multiplicity ...	82
5.4. Operators with Involutive Multiple Characteristics	82
5.5. The Schrödinger Operator	83
Chapter 5. Solvability of (Pseudo)Differential Equations	83
§ 1. Examples	83
1.1. Lewy's Example	83
1.2. Mizohata's Equation	84
1.3. Other Examples	85
§ 2. Necessary Conditions for Local Solvability	85
2.1. Hörmander's Theorem	85
2.2. The Zero of Finite Order	86
2.3. The Zero of Infinite Order	87
2.4. Multiple Characteristics	87
§ 3. Sufficient Conditions for Local Solvability	87
3.1. Operators of Real Principal Type	87
3.2. Operators of Principal Type	88
3.3. Operators with Multiple Characteristics	90
Chapter 6. Smoothness of Solutions of Differential Equations	90
§ 1. Hypoelliptic Operators	90
1.1. Definition and Examples	90
1.2. Hypoelliptic Differential Operators with Constant Coefficients	91
1.3. The Gevrey Classes	91
1.4. Partially Hypoelliptic Operators	92
1.5. Hypoelliptic Equations in Convolutions	93
1.6. Hypoelliptic Operators of Constant Strength	93
1.7. Hypoelliptic Differential Operators with Variable Coefficients	94
1.8. Pseudodifferential Hypoelliptic Operators	95
1.9. Degenerate Elliptic Operators	96
1.10. Partial Hypoellipticity of Degenerate Elliptic Operators	97
1.11. Double Characteristics	98
1.12. Hypoelliptic Operators on the Real Line	99
§ 2. Subelliptic Operators	100
2.1. Definition and Simplest Properties	100
2.2. Estimates for First-Order Differential Operators with Polynomial Coefficients	102
2.3. Algebraic Conditions	103

§3. Hypoelliptic Differential Operators of Second Order	105
3.1. The Sum of the Squares	105
3.2. A Necessary Condition for Hypoellipticity	106
3.3. Operators with a Non-negative Quadratic Form	107
§4. Analytic Hypoellipticity	108
4.1. Elliptic Operators	108
4.2. The Analytic Wave Front	108
4.3. Analytic Pseudodifferential Operators	109
4.4. Necessary Conditions for Analytic Hypoellipticity	110
4.5. Differential Equation of the Second Order	112
4.6. The Gevrey Classes	114
4.7. Generalized Analytic Hypoellipticity	115
Chapter 7. Transformation of Boundary-Value Problems	115
§1. The Transmission Property	115
1.1. Operators in a Half-Space	115
1.2. The Transmission Property	116
1.3. Application to the Study of Lacunae	119
§2. Distributions on a Manifold with Boundary	120
2.1. The Distribution Spaces	120
2.2. Contracted Cotangent Bundle	121
§3. Completely Characteristic Operators	122
3.1. Pseudodifferential Operators and their Kernels	122
3.2. The Transmission Property	123
3.3. Completely Characteristic Operators	124
3.4. The Boundary Wave Front	124
§4. Canonical Boundary Transformation	125
4.1. The Generating Function	125
4.2. The Operator of Principal Type	126
4.3. The Differential Operator of Second Order	126
§5. Fourier Integral Operators	127
5.1. The Generating Function of the Canonical Boundary Transformation	127
5.2. The Fourier Integral Operator	128
Chapter 8. Hyperfunctions	129
§1. Analytic Functionals	129
1.1. Definition and the Basic Properties	129
1.2. Operations on Analytic Functionals	130
§2. The Space of Hyperfunctions	130
2.1. Definition and the Basic Properties	130
2.2. The Analytic Wave Front of a Hyperfunction	131
2.3. Boundary Values of a Hyperfunction	131

§ 3. Solutions of Differential Equations	132
3.1. The Cauchy Problem	132
3.2. The Analytic Wave Front	133
§ 4. Sheaf of Microfunctions	135
4.1. Traces of Holomorphic Functions	135
4.2. The Definition of a Sheaf of Microfunctions	135
4.3. Pseudodifferential Operators	136
4.4. Fourier Integral Operators	138
References	139

Preface

The microlocal analysis is the local analysis in cotangent bundle space. The remarkable progress made in the theory of linear partial differential equations over the past two decades is essentially due to the extensive application of the microlocalization idea. The Hamiltonian systems, canonical transformations, Lagrange manifolds and other concepts, used in theoretical mechanics for examining processes in the phase space, have in recent years become the central objects of the theory of differential equations. For example, the evolution of singularities of solutions of a differential equation is described most naturally in terms of Lagrange manifolds and Hamiltonian systems, the solvability conditions are formulated in terms of the behaviour of integral curves of the Hamiltonian system whose Hamiltonian function serves as the characteristic form, the class of pseudodifferential equations arises in a natural way from that of differential equations under the action of canonical transformations, the class of subelliptic operators is defined by means of the Poisson brackets, etc. The difficulty faced in the microlocal analysis is connected with the principle of uncertainty which does not permit us to localize a function in any neighbourhood of a point of the cotangent space.

This paper presents a survey of the most interesting results, from our point of view, of the microlocal analysis achieved over the recent years. Unfortunately, due to lack of space many significant results could not be included. Also incomplete is the list of the literature cited; more complete lists can be found in Egorov [1984], Shubin [1978], Hörmander [1963, 1983, 1985], Taylor [1981] and Trèves [1982].

The author expresses his thanks to V.Ya. Ivrii for his useful critical comments.

Chapter 1

Microlocal Properties of Distributions

§ 1. Microlocalization

The study of the singularities of solutions constitutes one of the most important problems in the theory of differential equations. In this theory, just as in other mathematical disciplines, one often examines functions modulo smooth ones, so that the points where a given function is infinitely differentiable may be neglected. This approach reflects physical realities: singular points correspond to those phenomena which are most interesting from the point of view of each physical theory.

In investigating physical processes that take place in a bounded space an extensive use is made of the principle of *locality*. Its essence lies in that by knowing the state of the process at a given moment of time in a fixed region Ω of the physical space one may determine, by means of physical laws, the course of the process in a region Ω' , lying strictly inside Ω , for a future time interval. During this time interval the effect of processes taking place outside Ω will have no influence on phenomena in Ω' because the effect is propagated with a finite velocity.

We can introduce a more general principle, the principle of *microlocality*, by examining the phenomenon in a bounded region of the phase space. If we know the state of the process in this region at a certain moment of time, we can describe this process for future close points lying strictly inside the region. In physical terms, this means that the change in the impulse too takes place with a finite velocity because the acting forces are finite.

The above-mentioned principles are reflected in mathematical physics in the investigation of singularities of solutions of differential equations. Namely, local properties of such solutions are those properties which remain unaltered when the solutions are multiplied by smooth functions with a small support. Microlocal properties of a solution refer naturally to those properties which do not change on "multiplication" of the solution by a smooth function having support in a small neighbourhood of the given point in the phase space. However, this operation is much more complicated. In fact, it consists in multiplying by an ordinary smooth cutoff function with a small support, in applying Fourier transformation, in multiplying successively by a smooth cutoff function of dual coordinates, and in applying inverse Fourier transformation. Instead of Fourier transformation we can also use some other decomposition in plane waves; for example, the Radon transformation. In fact, the microlocal analysis is the local analysis on the cotangent bundle space.

A special feature of the microlocal analysis is the fact that localization in the phase space is possible only to a certain extent: the localization of spatial coordinates obstructs that of impulses. In quantum mechanics, this fact is referred to as the Heisenberg uncertainty principle.

The last two decades have seen immensely fruitful applications of the microlocality principle to the theory of partial differential equations. Every function (ordinary or generalized) can be regarded as an aggregate of linear differential equations which this function satisfies. The microlocality principle extends in a natural manner this aggregate to the system of pseudodifferential equations derived from differential equations by transforming the phase space without altering its structure. By applying the microlocality principle we not only obtain a more precise description of singular points of a distribution but we also have a simpler description of the propagation process of these singularities. By this principle we can also extend to distributions the operations defined initially for smooth functions only; for example, the operation of taking trace or the operation of multiplication, etc.

Let us explain the idea of microlocalization with the following simple example. Let $n \geq 2$ be a natural number and let f be a function in $\mathbb{R}^n$ of the form $f(x) = g(\alpha \cdot x)$, where $\alpha \in \mathbb{R}^n \setminus 0$, $\alpha \cdot x = \sum_{j=1}^n \alpha_j \cdot x_j$ and g is a function of a single variable. If $g(t)$ has a singularity, for example, if $g(t)$ is not differentiable at $t = t_0$, then all the points x lying on the plane $\alpha \cdot x = t_0$ are singular points of f . However, f is a smooth function in each direction lying on this plane so that for it singular will be only the direction of the vector α . Radon's theorem enables us to represent each distribution f in $\mathcal{D}'(\mathbb{R}^n)$ as an integral of plane waves:

$$f(x) = \int_{|\alpha|=1} g_\alpha(\alpha \cdot x) d\alpha.$$

Therefore at each point x those directions α will be singular for which the distribution $g_\alpha(t)$ has a singularity at the point $t = x \cdot \alpha$. If, instead of Radon's theorem, we apply Fourier transformation, then f can be represented as an integral of plane waves:

$$f(x) = \int g(\alpha) e^{i\alpha \cdot x} d\alpha,$$

where the integration is performed over the whole $\mathbb{R}^n$. Now those directions of α become singular for f in which $g(t\alpha)$ does not decrease, as $t \rightarrow \infty$, rapidly enough.

As mentioned earlier, in the modern theory of differential equations the microlocality principle is extensively applied to investigate the singularities of the solution. Many important results achieved in recent years by means of this principle in the theory of boundary-value problems, in the spectral theory, in the theory of functions of several complex variables, and in other branches of mathematics point towards great potentialities of the microlocal analysis.

§2. Wave Front of Distribution. Its Functorial Properties

2.1. Definition of the Wave Front. The notion of a singular point of a distribution does not have only one meaning. Depending on the problem under discussion, a singular point may signify a point of discontinuity, or a point where

the function becomes infinite, or a point where the function has an essential singularity in the sense of the theory of complex variables, etc. For the general theory of distributions, the most natural is the following

Definition 2.1. A point x_0 is a non-singular point for a distribution u if there exists a function φ in $C_0^\infty(\mathbb{R}^n)$ such that $\varphi(x_0) \neq 0$ and $\varphi u \in C^\infty(\mathbb{R}^n)$.

It follows from the definition that the singular points of a distribution constitute a closed set. This set is known as the *singular support* of the distribution u and is denoted by $\text{sing supp } u$. We can easily see that this set is invariant under diffeomorphisms of the space; thus the definition of a singular support can be readily extended to distributions on a smooth manifold.

For a smooth manifold X , we denote by $T^*(X)$ the cotangent bundle space and by $T^*(X) \setminus 0$ the same space with the zero section removed (see, for example, Arnol'd [1974], Egorov [1984]). The following definition and examples are due to Hörmander. It should be remarked that in the general theory there are many close concepts that are extensively used, namely, the analytic wave front, Gevrey wave front, oscillation front, etc (see Hörmander [1983, 1985], Trèves [1982], and §4.2 of Chapter 6).

Definition 2.2. A point $(x_0, \xi_0) \in T^*(\mathbb{R}^n) \setminus 0$ does not belong to the *wave front* of a distribution u in $\mathcal{D}'(\mathbb{R}^n)$ if there are a function φ in $C_0^\infty(\mathbb{R}^n)$, with $\varphi(x_0) \neq 0$, and a cone Γ in $\mathbb{R}^n$, with vertex at the origin of coordinates, which contains in its interior the ray $\{\xi; \xi = t\xi_0, t > 0\}$ such that the relations

$$|\tilde{\varphi}u(\xi)| = O((1 + |\xi|)^{-N})$$

holds for all $\xi \in \Gamma$ and all integers N .

The wave front of u will be denoted by $\text{WF}(u)$.

Example 2.1. If the distribution u is a plane wave, that is, if $u(x) = g(\alpha \cdot x)$, where $\alpha \in \mathbb{R}^n \setminus 0$ and $g \in \mathcal{D}'(\mathbb{R})$, then every direction ξ_0 , non-collinear with the vector α , is non-singular for u . That is, $\text{WF}(u)$ may only contain points (x_0, ξ_0) for which $\xi_0 = t\alpha$ with $t \in \mathbb{R} \setminus 0$ and $\alpha \cdot x_0 \in \text{sing supp } g$.

Example 2.2. Suppose that $\eta \in \mathbb{R}^n \setminus 0$, $\tilde{\varphi} \in C_0^\infty(\mathbb{R}^n)$, $\tilde{\varphi}(0) = 1$ and $\tilde{\varphi}(0) > 0$. Then for the function

$$u(x) = \sum_{k=1}^{\infty} \frac{\varphi(kx)}{k^2} e^{ik^2(x, \eta)}$$

which is continuous in $\mathbb{R}^n$, the wave front consists of the ray $\{(0, t\eta), t > 0\}$ (Hörmander [1979a]).

Summing over η , we obtain from this example a function whose wave front coincides with an arbitrary conical closed subset of $T^*(\mathbb{R}^n) \setminus 0$.

2.2. Localization of Wave Front. It is comparatively easy to establish the following properties of the wave front (see Hörmander [1971, 1983, 1985], Egorov [1984]):

1. If $u \in \mathcal{D}'(\mathbb{R}^n)$ and $\varphi \in C_0^\infty(\mathbb{R}^n)$, then

$$\text{WF}(\varphi u) \subset \text{WF}(u).$$

2. If $\pi: T^*(\mathbb{R}^n) \rightarrow \mathbb{R}^n$ is the natural projection, then $\pi \text{WF}(u) = \text{sing supp } u$ for every u in $\mathcal{D}'(\mathbb{R}^n)$.

2.3. Wave Front and Singularities of One-Dimensional Distributions. Let X and Y be smooth manifolds, and let $f: X \rightarrow Y$ be a smooth map. f is called *proper* if the set $f^{-1}(K)$ is compact for every compact set K in Y . For a $\varphi \in C_0^\infty(Y)$ and a proper map f we set

$$f^* \varphi(x) = \varphi(f(x)).$$

Then $f^* \varphi \in C_0^\infty(X)$ and f^* is a continuous map from $\mathcal{D}(Y)$ into $\mathcal{D}(X)$. This enables us to define, by means of duality, the *pushforward* $f_* u$ of every *distribution* $u \in \mathcal{D}'(X)$ by the formula

$$\langle f_* u, \varphi \rangle = \langle u, f^* \varphi \rangle.$$

It is obvious that the same construction remains valid for an arbitrary map f (not necessarily proper) when u has a compact support, that is, when $u \in \mathcal{E}'(X)$.

The *pullback* $f^* u$ of a *distribution* is defined when f is a submersion of X onto Y . This last condition means that for every point $y \in Y$ the set $f^{-1}(y)$ is a smooth submanifold of X and all these submanifolds are diffeomorphic to a fixed k -dimensional smooth manifold. Locally such a map is a projection, and in a suitable local coordinate system it becomes the projection $\mathbb{R}^k \times \mathbb{R}^l \rightarrow \mathbb{R}^k$. If $\varphi \in C_0^\infty(X)$ and the support of φ lies in a coordinate neighbourhood, and if $x = (x', x'')$, where $x' \in \mathbb{R}^k$, $x'' \in \mathbb{R}^l$, then $f_* \varphi(x') = \int \varphi(x', x'') dx''$. For any $u \in \mathcal{D}'(Y)$ we now set

$$\langle f_* u, \varphi \rangle = \langle u, f_* \varphi \rangle.$$

If f is both proper and submersion (for example, if f is a diffeomorphism), then $f^* u$ and $f_* v$ are simultaneously defined for every $u \in \mathcal{D}'(Y)$, $v \in \mathcal{D}'(X)$.

Example 2.3. Let $\pi: \mathbb{R}^n \rightarrow \mathbb{R}^1$ be the projection onto the x_1 -axis. In this case both $\pi^* u$, $u \in \mathcal{D}'(\mathbb{R})$, and $\pi_* v$, $v \in \mathcal{E}'(\mathbb{R}^n)$, are defined, and

$$\pi^* v(x^1) = u(x^1) \otimes 1_{x_2, \dots, x_n};$$

$$\pi_* v(x) = \int v(x_1, x_2, \dots, x_n) dx_2 \dots dx_n.$$

Using these concepts, we can establish the following

Theorem 2.1. Let Ω be a domain in $\mathbb{R}^n$, and let $u \in \mathcal{D}'(\Omega)$. A point (x_0, ξ_0) of $T^*(\Omega) \setminus 0$ does not lie in $\text{WF}(u)$ if and only if there exist a function φ in $C_0^\infty(\Omega)$, with $\varphi(x_0) \neq 0$, and an $\varepsilon > 0$ such that for every smooth function $f: \text{supp } \varphi \rightarrow \mathbb{R}$, with $|\text{grad } f(x_0) - \xi_0| < \varepsilon$, the function $f_*(\varphi u)(t)$ is infinitely differentiable on the real line (Guillemin and Sternberg [1977], Egorov [1985a]).