

Yehuda Pinchover and Jacob Rubinstein

An Introduction to
**Partial Differential
Equations**



CAMBRIDGE

0175.2
P647

AN INTRODUCTION TO PARTIAL
DIFFERENTIAL EQUATIONS

YEHUDA PINCHOVER AND JACOB RUBINSTEIN



CAMBRIDGE
UNIVERSITY PRESS

CAMBRIDGE UNIVERSITY PRESS
Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, São Paulo

CAMBRIDGE UNIVERSITY PRESS
The Edinburgh Building, Cambridge CB2 2RU, UK

www.cambridge.org
information on this title: www.cambridge.org/9780521848865

© Cambridge University Press 2005

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2005

Printed in the United Kingdom at the University Press, Cambridge

A catalog record for this book is available from the British Library

Library of Congress Cataloging in Publication data

ISBN-13 978-0-521-84886-2 hardback

ISBN-10 0-521-84886-5 hardback

ISBN-13 978-0-521-61323-X paperback

ISBN-10 0-521-61323-1 paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this book, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

AN INTRODUCTION TO PARTIAL DIFFERENTIAL EQUATIONS

A complete introduction to partial differential equations, this textbook provides a rigorous yet accessible guide to students in mathematics, physics and engineering. The presentation is lively and up to date, with particular emphasis on developing an appreciation of underlying mathematical theory.

Beginning with basic definitions, properties and derivations of some fundamental equations of mathematical physics from basic principles, the book studies first-order equations, the classification of second-order equations, and the one-dimensional wave equation. Two chapters are devoted to the separation of variables, whilst others concentrate on a wide range of topics including elliptic theory, Green's functions, variational and numerical methods.

A rich collection of worked examples and exercises accompany the text, along with a large number of illustrations and graphs to provide insight into the numerical examples.

Solutions and hints to selected exercises are included for students whilst extended solution sets are available to lecturers from solutions@cambridge.org.

To our parents

*The equation of heaven and earth
remains unsolved.*
(Yehuda Amichai)

המשוואה של שמים וארץ
נשארת קלי פתרון.
(יהודה עמיחי)

Preface

This book presents an introduction to the theory and applications of partial differential equations (PDEs). The book is suitable for all types of basic courses on PDEs, including courses for undergraduate engineering, sciences and mathematics students, and for first-year graduate courses as well.

Having taught courses on PDEs for many years to varied groups of students from engineering, science and mathematics departments, we felt the need for a textbook that is concise, clear, motivated by real examples and mathematically rigorous. We therefore wrote a book that covers the foundations of the theory of PDEs. This theory has been developed over the last 250 years to solve the most fundamental problems in engineering, physics and other sciences. Therefore we think that one should not treat PDEs as an abstract mathematical discipline; rather it is a field that is closely related to real-world problems. For this reason we strongly emphasize throughout the book the relevance of every bit of theory and every practical tool to some specific application. At the same time, we think that the modern engineer or scientist should understand the basics of PDE theory when attempting to solve specific problems that arise in applications. Therefore we took great care to create a balanced exposition of the theoretical and applied facets of PDEs.

The book is flexible enough to serve as a textbook or a self-study book for a large class of readers. The first seven chapters include the core of a typical one-semester course. In fact, they also include advanced material that can be used in a graduate course. Chapters 9 and 11 include additional material that together with the first seven chapters fits into a typical curriculum of a two-semester course. In addition, Chapters 8 and 10 contain advanced material on Green's functions and the calculus of variations. The book covers all the classical subjects, such as the separation of variables technique and Fourier's method (Chapters 5, 6, 7, and 9), the method of characteristics (Chapters 2 and 9), and Green's function methods (Chapter 8). At the same time we introduce the basic theorems that guarantee that the problem at

hand is well defined (Chapters 2–10), and we took care to include modern ideas such as variational methods (Chapter 10) and numerical methods (Chapter 11).

The first eight chapters mainly discuss PDEs in two independent variables. Chapter 9 shows how the methods of the first eight chapters are extended and enhanced to handle PDEs in higher dimensions. Generalized and weak solutions are presented in many parts of the book.

Throughout the book we illustrate the mathematical ideas and techniques by applying them to a large variety of practical problems, including heat conduction, wave propagation, acoustics, optics, solid and fluid mechanics, quantum mechanics, communication, image processing, musical instruments, and traffic flow.

We believe that the best way to grasp a new theory is by considering examples and solving problems. Therefore the book contains hundreds of examples and problems, most of them at least partially solved. Extended solutions to the problems are available for course instructors using the book from solutions@cambridge.org. We also include dozens of drawing and graphs to explain the text better and to demonstrate visually some of the special features of certain solutions.

It is assumed that the reader is familiar with the calculus of functions in several variables, with linear algebra and with the basics of ordinary differential equations. The book is almost entirely self-contained, and in the very few places where we cannot go into details, a reference is provided.

The book is the culmination of a slow evolutionary process. We wrote it during several years, and kept changing and adding material in light of our experience in the classroom. The current text is an expanded version of a book in Hebrew that the authors published in 2001, which has been used successfully at Israeli universities and colleges since then.

Our cumulative expertise of over 30 years of teaching PDEs at several universities, including Stanford University, UCLA, Indiana University and the Technion – Israel Institute of Technology guided us to create a text that enhances not just technical competence but also deep understanding of PDEs. We are grateful to our many students at these universities with whom we had the pleasure of studying this fascinating subject. We hope that the readers will also learn to enjoy it.

We gratefully acknowledge the help we received from a number of individuals. Kristian Jenssen from North Carolina State University, Lydia Peres and Tiferet Saadon from the Technion – Israel Institute of Technology, and Peter Sternberg from Indiana University read portions of the draft and made numerous comments and suggestions for improvement. Raya Rubinstein prepared the drawings, while Yishai Pinchover and Aviad Rubinstein assisted with the graphs. Despite our best efforts, we surely did not discover all the mistakes in the draft. Therefore we encourage observant readers to send us their comments at pincho@techunix.technion.ac.il. We will maintain a webpage with a list of errata at <http://www.math.technion.ac.il/~pincho/PDE.pdf>.

Contents

<i>Preface</i>	<i>page xi</i>
1 Introduction	1
1.1 Preliminaries	1
1.2 Classification	3
1.3 Differential operators and the superposition principle	3
1.4 Differential equations as mathematical models	4
1.5 Associated conditions	17
1.6 Simple examples	20
1.7 Exercises	21
2 First-order equations	23
2.1 Introduction	23
2.2 Quasilinear equations	24
2.3 The method of characteristics	25
2.4 Examples of the characteristics method	30
2.5 The existence and uniqueness theorem	36
2.6 The Lagrange method	39
2.7 Conservation laws and shock waves	41
2.8 The eikonal equation	50
2.9 General nonlinear equations	52
2.10 Exercises	58
3 Second-order linear equations in two independent variables	64
3.1 Introduction	64
3.2 Classification	64
3.3 Canonical form of hyperbolic equations	67
3.4 Canonical form of parabolic equations	69
3.5 Canonical form of elliptic equations	70
3.6 Exercises	73

4	The one-dimensional wave equation	76
4.1	Introduction	76
4.2	Canonical form and general solution	76
4.3	The Cauchy problem and d'Alembert's formula	78
4.4	Domain of dependence and region of influence	82
4.5	The Cauchy problem for the nonhomogeneous wave equation	87
4.6	Exercises	93
5	The method of separation of variables	98
5.1	Introduction	98
5.2	Heat equation: homogeneous boundary condition	99
5.3	Separation of variables for the wave equation	109
5.4	Separation of variables for nonhomogeneous equations	114
5.5	The energy method and uniqueness	116
5.6	Further applications of the heat equation	119
5.7	Exercises	124
6	Sturm–Liouville problems and eigenfunction expansions	130
6.1	Introduction	130
6.2	The Sturm–Liouville problem	133
6.3	Inner product spaces and orthonormal systems	136
6.4	The basic properties of Sturm–Liouville eigenfunctions and eigenvalues	141
6.5	Nonhomogeneous equations	159
6.6	Nonhomogeneous boundary conditions	164
6.7	Exercises	168
7	Elliptic equations	173
7.1	Introduction	173
7.2	Basic properties of elliptic problems	173
7.3	The maximum principle	178
7.4	Applications of the maximum principle	181
7.5	Green's identities	182
7.6	The maximum principle for the heat equation	184
7.7	Separation of variables for elliptic problems	187
7.8	Poisson's formula	201
7.9	Exercises	204
8	Green's functions and integral representations	208
8.1	Introduction	208
8.2	Green's function for Dirichlet problem in the plane	209
8.3	Neumann's function in the plane	219
8.4	The heat kernel	221
8.5	Exercises	223

9	Equations in high dimensions	226
9.1	Introduction	226
9.2	First-order equations	226
9.3	Classification of second-order equations	228
9.4	The wave equation in \mathbb{R}^2 and \mathbb{R}^3	234
9.5	The eigenvalue problem for the Laplace equation	242
9.6	Separation of variables for the heat equation	258
9.7	Separation of variables for the wave equation	259
9.8	Separation of variables for the Laplace equation	261
9.9	Schrödinger equation for the hydrogen atom	263
9.10	Musical instruments	266
9.11	Green's functions in higher dimensions	269
9.12	Heat kernel in higher dimensions	275
9.13	Exercises	279
10	Variational methods	282
10.1	Calculus of variations	282
10.2	Function spaces and weak formulation	296
10.3	Exercises	306
11	Numerical methods	309
11.1	Introduction	309
11.2	Finite differences	311
11.3	The heat equation: explicit and implicit schemes, stability, consistency and convergence	312
11.4	Laplace equation	318
11.5	The wave equation	322
11.6	Numerical solutions of large linear algebraic systems	324
11.7	The finite elements method	329
11.8	Exercises	334
12	Solutions of odd-numbered problems	337
A.1	Trigonometric formulas	361
A.2	Integration formulas	362
A.3	Elementary ODEs	362
A.4	Differential operators in polar coordinates	363
A.5	Differential operators in spherical coordinates	363
	<i>References</i>	364
	<i>Index</i>	366

1

Introduction

1.1 Preliminaries

A partial differential equation (PDE) describes a relation between an unknown function and its partial derivatives. PDEs appear frequently in all areas of physics and engineering. Moreover, in recent years we have seen a dramatic increase in the use of PDEs in areas such as biology, chemistry, computer sciences (particularly in relation to image processing and graphics) and in economics (finance). In fact, in each area where there is an interaction between a number of independent variables, we attempt to define functions in these variables and to model a variety of processes by constructing equations for these functions. When the value of the unknown function(s) at a certain point depends only on what happens in the vicinity of this point, we shall, in general, obtain a PDE. The general form of a PDE for a function $u(x_1, x_2, \dots, x_n)$ is

$$F(x_1, x_2, \dots, x_n, u, u_{x_1}, u_{x_2}, \dots, u_{x_{n1}}, \dots) = 0, \quad (1.1)$$

where x_1, x_2, \dots, x_n are the independent variables, u is the unknown function, and u_{x_i} denotes the partial derivative $\partial u / \partial x_i$. The equation is, in general, supplemented by additional conditions such as initial conditions (as we have often seen in the theory of ordinary differential equations (ODEs)) or boundary conditions.

The analysis of PDEs has many facets. The classical approach that dominated the nineteenth century was to develop methods for finding explicit solutions. Because of the immense importance of PDEs in the different branches of physics, every mathematical development that enabled a solution of a new class of PDEs was accompanied by significant progress in physics. Thus, the method of characteristics invented by Hamilton led to major advances in optics and in analytical mechanics. The Fourier method enabled the solution of heat transfer and wave

propagation, and Green's method was instrumental in the development of the theory of electromagnetism. The most dramatic progress in PDEs has been achieved in the last 50 years with the introduction of numerical methods that allow the use of computers to solve PDEs of virtually every kind, in general geometries and under arbitrary external conditions (at least in theory; in practice there are still a large number of hurdles to be overcome).

The technical advances were followed by theoretical progress aimed at understanding the solution's structure. The goal is to discover some of the solution's properties before actually computing it, and sometimes even without a complete solution. The theoretical analysis of PDEs is not merely of academic interest, but rather has many applications. It should be stressed that there exist very complex equations that cannot be solved even with the aid of supercomputers. All we can do in these cases is to attempt to obtain qualitative information on the solution. In addition, a deep important question relates to the formulation of the equation and its associated side conditions. In general, the equation originates from a model of a physical or engineering problem. It is not automatically obvious that the model is indeed consistent in the sense that it leads to a solvable PDE. Furthermore, it is desired in most cases that the solution will be unique, and that it will be stable under small perturbations of the data. A theoretical understanding of the equation enables us to check whether these conditions are satisfied. As we shall see in what follows, there are many ways to solve PDEs, each way applicable to a certain class of equations. Therefore it is important to have a thorough analysis of the equation before (or during) solving it.

The fundamental theoretical question is whether the problem consisting of the equation and its associated side conditions is well posed. The French mathematician Jacques Hadamard (1865–1963) coined the notion of *well-posedness*. According to his definition, a problem is called well-posed if it satisfies all of the following criteria

1. **Existence** The problem has a solution.
2. **Uniqueness** There is no more than one solution.
3. **Stability** A small change in the equation or in the side conditions gives rise to a small change in the solution.

If one or more of the conditions above does not hold, we say that the problem is *ill-posed*. One can fairly say that the fundamental problems of mathematical physics are all well-posed. However, in certain engineering applications we might tackle problems that are ill-posed. In practice, such problems are unsolvable. Therefore, when we face an ill-posed problem, the first step should be to modify it appropriately in order to render it well-posed.

1.2 Classification

We pointed out in the previous section that PDEs are often classified into different types. In fact, there exist several such classifications. Some of them will be described here. Other important classifications will be described in Chapter 3 and in Chapter 9.

- **The order of an equation**

The first classification is according to the *order* of the equation. The order is defined to be the order of the highest derivative in the equation. If the highest derivative is of order k , then the equation is said to be of order k . Thus, for example, the equation $u_{tt} - u_{xx} = f(x, t)$ is called a second-order equation, while $u_t + u_{xxxx} = 0$ is called a fourth-order equation.

- **Linear equations**

Another classification is into two groups: linear versus nonlinear equations. An equation is called *linear* if in (1.1), F is a linear function of the unknown function u and its derivatives. Thus, for example, the equation $x^7 u_x + e^{xy} u_y + \sin(x^2 + y^2)u = x^3$ is a linear equation, while $u_x^2 + u_y^2 = 1$ is a nonlinear equation. The nonlinear equations are often further classified into subclasses according to the type of the nonlinearity. Generally speaking, the nonlinearity is more pronounced when it appears in a higher derivative. For example, the following two equations are both nonlinear:

$$u_{xx} + u_{yy} = u^3, \quad (1.2)$$

$$u_{xx} + u_{yy} = |\nabla u|^2 u. \quad (1.3)$$

Here $|\nabla u|$ denotes the norm of the gradient of u . While (1.3) is nonlinear, it is still linear as a function of the highest-order derivative. Such a nonlinearity is called *quasilinear*. On the other hand in (1.2) the nonlinearity is only in the unknown function. Such equations are often called *semilinear*.

- **Scalar equations versus systems of equations**

A single PDE with just one unknown function is called a *scalar equation*. In contrast, a set of m equations with l unknown functions is called a *system of m equations*.

1.3 Differential operators and the superposition principle

A function has to be k times differentiable in order to be a solution of an equation of order k . For this purpose we define the set $C^k(D)$ to be the set of all functions that are k times continuously differentiable in D . In particular, we denote the set of continuous functions in D by $C^0(D)$, or $C(D)$. A function in the set C^k that satisfies a PDE of order k , will be called a *classical* (or *strong*) solution of the PDE. It should be stressed that we sometimes also have to deal with solutions that are not classical. Such solutions are called *weak* solutions. The possibility of weak solutions and their physical meaning will be discussed on several occasions later,

see for example Sections 2.7 and 10.2. Note also that, in general, we are required to solve a problem that consists of a PDE and associated conditions. In order for a strong solution of the PDE to also be a strong solution of the full problem, it is required to satisfy the additional conditions in a smooth way.

Mappings between different function sets are called *operators*. The operation of an operator L on a function u will be denoted by $L[u]$. In particular, we shall deal in this book with operators defined by partial derivatives of functions. Such operators, which are in fact mappings between different C^k classes, are called *differential operators*.

An operator that satisfies a relation of the form

$$L[a_1u_1 + a_2u_2] = a_1L[u_1] + a_2L[u_2],$$

where a_1 and a_2 are arbitrary constants, and u_1 and u_2 are arbitrary functions is called a *linear operator*. A linear differential equation naturally defines a linear operator: the equation can be expressed as $L[u] = f$, where L is a linear operator and f is a given function.

A linear differential equation of the form $L[u] = 0$, where L is a linear operator, is called a *homogeneous equation*. For example, define the operator $L = \partial^2/\partial x^2 - \partial^2/\partial y^2$. The equation

$$L[u] = u_{xx} - u_{yy} = 0$$

is a homogeneous equation, while the equation

$$L[u] = u_{xx} - u_{yy} = x^2$$

is an example of a *nonhomogeneous equation*.

Linear operators play a central role in mathematics in general, and in PDE theory in particular. This results from the important property (which follows at once from the definition) that if for $1 \leq i \leq n$, the function u_i satisfies the linear differential equation $L[u_i] = f_i$, then the linear combination $v := \sum_{i=1}^n \alpha_i u_i$ satisfies the equation $L[v] = \sum_{i=1}^n \alpha_i f_i$. In particular, if each of the functions u_1, u_2, \dots, u_n satisfies the homogeneous equation $L[u] = 0$, then every linear combination of them satisfies that equation too. This property is called the *superposition principle*. It allows the construction of complex solutions through combinations of simple solutions. In addition, we shall use the superposition principle to prove uniqueness of solutions to linear PDEs.

1.4 Differential equations as mathematical models

PDEs are woven throughout science and technology. We shall briefly review a number of canonical equations in different areas of application. The fundamental

laws of physics provide a mathematical description of nature's phenomena on a variety of scales of time and space. Thus, for example, very large scale phenomena (astronomical scales) are controlled by the laws of gravity. The theory of electromagnetism controls the scales involved in many daily activities, while quantum mechanics is used to describe phenomena on the atomic scale. It turns out, however, that many important problems involve interaction between a large number of objects, and thus it is difficult to use the basic laws of physics to describe them. For example, we do not fall to the floor when we sit on a chair. Why? The fundamental reason lies in the electric forces between the atoms constituting the chair. These forces endow the chair with high rigidity. It is clear, though, that it is not feasible to solve the equations of electromagnetism (Maxwell's equations) to describe the interaction between such a vast number of objects. As another example, consider the flow of a gas. Each molecule obeys Newton's laws, but we cannot in practice solve for the evolution of an Avogadro number of individual molecules. Therefore, it is necessary in many applications to develop simpler models.

The basic approach towards the derivation of these models is to define new quantities (temperature, pressure, tension, . . .) that describe average macroscopic values of the fundamental microscopic quantities, to assume several fundamental principles, such as conservation of mass, conservation of momentum, conservation of energy, etc., and to apply the new principles to the macroscopic quantities. We shall often need some additional ad-hoc assumptions to connect different macroscopic entities. In the optimal case we would like to start from the fundamental laws and then average them to achieve simpler models. However, it is often very hard to do so, and, instead, we shall sometimes use experimental observations to supplement the basic principles. We shall use x , y , z to denote spatial variables, and t to denote the time variable.

1.4.1 The heat equation

A common way to encourage scientific progress is to confer prizes and awards. Thus, the French Academy used to set up competitions for its prestigious prizes by presenting specific problems in mathematics and physics. In 1811 the Academy chose the problem of heat transfer for its annual prize. The prize was awarded to the French mathematician Jean Baptiste Joseph Fourier (1768–1830) for two important contributions. (It is interesting to mention that he was not an active scientist at that time, but rather the governor of a region in the French Alps – actually a politician!). He developed, as we shall soon see, an appropriate differential equation, and, in addition developed, as we shall see in Chapter 5, a novel method for solving this equation.

The basic idea that guided Fourier was conservation of energy. For simplicity we assume that the material density and the heat capacity are constant in space and time, and we scale them to be 1. We can therefore identify heat energy with temperature. Let D be a fixed spatial domain, and denote its boundary by ∂D . Under these conditions we shall write down the change in the energy stored in D between time t and time $t + \Delta t$:

$$\begin{aligned} & \int_D [u(x, y, z, t + \Delta t) - u(x, y, z, t)] dV \\ &= \int_t^{t+\Delta t} \int_D q(x, y, z, t, u) dV dt - \int_t^{t+\Delta t} \int_{\partial D} \vec{B}(x, y, z, t) \cdot \hat{n} dS dt, \end{aligned} \quad (1.4)$$

where u is the temperature, q is the rate of heat production in D , \vec{B} is the heat flux through the boundary, dV and dS are space and surface integration elements, respectively, and \hat{n} is a unit vector pointing in the direction of the outward normal to ∂D . Notice that the heat production can be negative (a refrigerator, an air conditioner), as can the heat flux.

In general the heat production is determined by external sources that are independent of the temperature. In some cases (such as an air conditioner controlled by a thermostat) it depends on the temperature itself but not on its derivatives. Hence we assume $q = q(x, y, z, t, u)$. To determine the functional form of the heat flux, Fourier used the experimental observation that ‘heat flows from hotter places to colder places’. Recall from calculus that the direction of maximal growth of a function is given by its gradient. Therefore, Fourier postulated

$$\vec{B} = -k(x, y, z) \vec{\nabla} u. \quad (1.5)$$

The formula (1.5) is called *Fourier’s law of heat conduction*. The (positive!) function k is called the *heat conduction (or Fourier) coefficient*. The value(s) of k depend on the medium in which the heat diffuses. In a homogeneous domain k is expected to be constant. The assumptions on the functional dependence of q and \vec{B} on u are called *constitutive laws*.

We substitute our formula for q and \vec{B} into (1.4), approximate the t integrals using the mean value theorem, divide both sides of the equation by Δt , and take the limit $\Delta t \rightarrow 0$. We obtain

$$\int_D u_t dV = \int_D q(x, y, z, t, u) dV + \int_{\partial D} k(x, y, z) \vec{\nabla} u \cdot \hat{n} dS. \quad (1.6)$$

Observe that the integration in the second term on the right hand side is over a different set than in the other terms. Thus we shall use Gauss’ theorem to convert

the surface integral into a volume integral:

$$\int_D [u_t - q - \vec{\nabla} \cdot (k\vec{\nabla}u)]dV = 0, \quad (1.7)$$

where $\vec{\nabla} \cdot$ denotes the divergence operator. The following simple result will be used several times in the book.

Lemma 1.1 *Let $h(x, y, z)$ be a continuous function satisfying $\int_{\Omega} h(x, y, z)dV = 0$ for every domain Ω . Then $h \equiv 0$.*

Proof Let us assume to the contrary that there exists a point $P = (x_0, y_0, z_0)$ where $h(P) \neq 0$. Assume without loss of generality that $h(P) > 0$. Since h is continuous, there exists a domain (maybe very small) D_0 , containing P and $\epsilon > 0$, such that $h > \epsilon > 0$ at each point in D_0 . Therefore $\int_{D_0} h dV > \epsilon \text{Vol}(D_0) > 0$ which contradicts the lemma's assumption. \square

Returning to the energy integral balance (1.7), we notice that it holds for any domain D . Assuming further that all the functions in the integrand are continuous, we obtain the PDE

$$u_t = q + \vec{\nabla} \cdot (k\vec{\nabla}u). \quad (1.8)$$

In the special (but common) case where the diffusion coefficient is constant, and there are no heat sources in D itself, we obtain the classical heat equation

$$u_t = k\Delta u, \quad (1.9)$$

where we use Δu to denote the important operator $u_{xx} + u_{yy} + u_{zz}$. Observe that we have assumed that the solution of the heat equation, and even some of its derivatives are continuous functions, although we have not solved the equation yet. Therefore, in principle we have to reexamine our assumptions a posteriori. We shall see examples later in the book in which solutions of a PDE (or their derivatives) are *not* continuous. We shall then consider ways to provide a meaning for the seemingly absurd process of substituting a discontinuous function into a differential equation. One of the fundamental ways of doing so is to observe that the integral balance equation (1.6) provides a more fundamental model than the PDE (1.8).

1.4.2 Hydrodynamics and acoustics

Hydrodynamics is the physical theory of fluid motion. Since almost any conceivable volume of fluid (whether it is a cup of coffee or the Pacific Ocean) contains a huge number of molecules, it is not feasible to describe the fluid using the law of electromagnetism or quantum mechanics. Hence, since the eighteenth century