

A GUIDE TO CLASSIFICATION IN GEOLOGY

J. W. MURRAY

英 汉 对 照

地 质 学 中 的 分 类 指 南

张 疆 译

A GUIDE TO CLASSIFICATION IN GEOLOGY

J. W. MURRAY, D.I.C., B.Sc., Ph.D., F.G.S.
Professor of Geology
University of Exeter



ELLIS HORWOOD LIMITED
Publishers · Chichester

Halsted Press: a division of
JOHN WILEY & SONS

New York · Chichester · Brisbane · Toronto

1981

照 依 元 英

南 部 英 分 出 中 華 商 報

英 里 德 亞 不 美

新 華 英

英 華 德 英 華 英

英 汉 对 照
地 质 学 中 的 分 类 指 南

[英] J. W. 默里 著

张 疆 译

籍 传 茂、汪熊麟 审校

1982

照 依 元 英

南 部 类 今 世 中 半 部 部

委 里 耀 可 不 〔美〕

社 译 委

外 中 编 前 五 英 部 部

S901

Table of Contents

Author's Preface	9
Chapter 1 – Introduction	11
Chapter 2 – Sediments and Sedimentary Rocks	
2.1 Major groups of sedimentary rocks	13
2.2 Terrigenous sediments and rocks.	14
2.2.1 Rudites – conglomerates and breccias.	14
2.2.2 Sandstones – arenites and wackes.	19
2.2.3 Siltstones.	22
2.2.4 Lutites – mudstone, mudrock, claystone.	22
2.3 Volcanic sedimentary deposits	25
2.3.1 Pyroclastic rocks.	28
2.3.2 Hydroclastic and Hyaloclastic rocks ...	31
2.3.3 Phreatomagmatic rocks.	31
2.3.4 Autoclastic rocks.	31
2.3.5 Alloclastic rocks	32
2.3.6 Terrigenous sediments rich in volcanic fragments	32
2.4 Residual deposits.	33
2.5 Organic deposits	33
2.5.1 Carbonaceous deposits	33
2.5.2 Carbonate sediments and carbonate rocks (limestones).	35
2.5.3 Dolomitic limestones and dolomites ...	44
2.5.4 Siliceous sediments and rocks	45

2.6	Chemical precipitates	46
2.6.1	Evaporites	46
2.6.2	Phosphorites	46
2.6.3	Ironstones	46
2.6.4	Caliche and calcrete	46
2.6.5	Silcrete	47
2.6.6	Chert and flint	47
2.7	Special groups	47
2.7.1	Deep sea sediments	47
2.7.2	Process name rock types	48
2.8	References	49

Chapter 3 – Igneous Rocks

3.1	Plutonic rocks	55
3.1.1	$M < 90\%$	55
3.1.2	$M > 90\%$ – ultramafic rocks	56
3.1.3	Charnockites	65
3.2	Hypabyssal rocks	65
3.3	Volcanic rocks	68
3.3.1	Glassy rocks	72
3.3.2	Igneous rock series	72
3.3.3	Metamorphosed volcanic rocks	73
3.4	References	74

Chapter 4 – Metamorphic Rocks

4.1	Names based on texture	77
4.2	Names based on composition	78
4.3	Gneiss	80
4.4	References	83

Chapter 5 – Mixed Rocks and Rock Associations

5.1	Migmatite	84
5.1.1	Structural types	85
5.1.2	Metatexite and diatexite	86
5.2	Ophiolite	86

5.3 Rodingite	87
5.4 Kimberlite	87
5.5 References	87
Chapter 6 – Stratigraphic Classification	
6.1 Lithostratigraphy	89
6.2 Biostratigraphy	90
6.3 Chronostratigraphy	90
6.4 References	93
Chapter 7 – Engineering Geology	
7.1 Rock materials	97
7.2 Rock masses	100
7.3 Core logging	102
7.4 Aggregates	102
7.5 References	103
Index	105

Table of Contents

87	2.3. Rodenticide
87	2.4. Kimberrite
87	2.5. References
Chapter 6 - Stratigraphic Classification	
87	6.1. Lithostratigraphy
89	6.2. Biostratigraphy
90	6.3. Chronostratigraphy
93	6.4. References
Chapter 7 - Engineering Geology	
94	7.1. Rock masses
97	7.2. Rock masses
100	7.3. Core logging
102	7.4. Aggregates
103	7.5. References
103	Index

Author's Preface

Most professional geologists have specialist knowledge of only part of the subject. When they read papers outside their field of specialisation they may encounter problems with unfamiliar terminology. The problem is even greater for students and amateurs who are in the process of building up their knowledge. The majority of modern textbooks concentrate primarily on processes and mechanisms of producing rocks. My aim has therefore been to provide a precis of the classification systems in common use in geology in the hope that this small book will prove a useful source of reference.

I wish to thank Mr Ellis Horwood for encouraging me to write this book and Mrs G. F. Murray for typing the manuscript. Professor D. T. Donovan, Mr P. Grainger and Dr M. Stone kindly offered comment on various chapters.

Exeter
December 1979

Author's Preface

Most professional geologists have specialist knowledge of only part of the subject. When they read paper outside their field of specialization they may encounter problems with unfamiliar terminology. The problem is even greater for students and amateurs who are in the process of building up their knowledge. The majority of modern textbooks concentrate primarily on processes and mechanisms of geobasic rocks. My aim has therefore been to provide a précis of the classification systems in common use in geology in the hope that this small book will prove a useful source of reference.

I wish to thank Mr Ellis Horwood for encouraging me to write this book and Mrs G. E. Murray for typing the manuscript. Professor D. T. Donovan, Mr P. G. Grist, and Dr M. Stone kindly offered comment on various chapters.

Exeter

December 1979

Chapter 1

Introduction

To classify is defined as 'to arrange or distribute in classes according to a method or system' (*Shorter Oxford English Dictionary*). Accepting this definition, a classification does three basic things:

- it sets out criteria for distinguishing between the items being classified
- it allows grouping of similar items in an hierarchical scheme of classes
- it establishes a scheme of nomenclature.

In geology, classification plays an important role in the organisation of data. Rocks and structures need to be defined by standard schemes of nomenclature in order to assist communication between geologists. There have been many attempts to standardise nomenclature, but not all classification schemes are universally accepted. This proliferation of systems presents no problems so long as authors state clearly which classification scheme they are using. Moreover, it is doubtful whether it is desirable or necessary to employ a totally standard classification system, as new research inevitably reveals alternative ways of approaching the same problem.

Some geologists believe that the ideal classification is one that is purely descriptive. However, many geological

classifications are genetic. For instance, rocks are classified, according to the mode of formation, into igneous, sedimentary, and metamorphic classes. In many cases individual rock names not only describe the rock but also define their chemical composition, texture and grain size, and the environment of formation, for example, quartz arenite, a sedimentary rock composed of detrital grains 0.0625 - 2 mm in diameter, more than 95% of which are quartz.

With so many variables involved it is not surprising that different authors place greatest emphasis on different variables, which in turn leads to disagreement. Thus, the term 'greywacke' fell into disrepute because it meant different things to different people: to some it was a muddy sandstone whereas to others the product of a turbidity current. However, not all muddy sandstones are the product of turbidity currents, and not all turbidity currents deposit muddy sand. The problem can be solved by using 'greywacke' for a muddy sandstone and, for those laid down by turbidity currents, by using the term 'turbiditic greywacke.'

Some classifications, although founded on sound logic, have not been accepted, often because they involve cumbersome terminology. The ideal classification is simple to use, provides a workable system of nomenclature, and is acceptable to those using it. Chapters 2 - 7 represent a compilation of those classifications proposed by international working parties (for example, igneous rocks) and others commonly used. No attempt has been made to include all available classification schemes.

Chapter 2

Sediments and sedimentary rocks

Sediment: an unconsolidated deposit of either detritus derived from pre-existing rock or new minerals formed by chemical or biochemical processes or organically formed materials.

Sedimentary rock: a lithified sediment, including those sediments which have undergone total mineralogical reorganisation under conditions of low temperature and low pressure (but excluding true metamorphism).

2.1 MAJOR GROUPS OF SEDIMENTARY ROCKS

A commonly used subdivision is as follows:

Terrigenous deposits - composed of detrital transported material.

Volcanic sedimentary deposits - composed of fragmented volcanic material extruded penecontemporaneously with sediment deposition. Sometimes this group is termed 'pyroclastic' but this type is only one of several volcanic sedimentary deposits.

Residual deposits - left after chemical weathering of rock, for example laterite.

Organic deposits - composed of the remains of plants or animals (all limestones are conveniently included here even though some contain only a low organic component).

Chemical precipitates - notably evaporite mineral deposits whether primary or secondary (diagenetic) in origin.

An additional category has been added here to cater for process name rock types and deep sea sediments.

2.2 TERRIGENOUS SEDIMENTS AND ROCKS

Terrigenous sediments consists of the detritus of pre-existing rocks which has been produced by weathering (physical and chemical), has been transported by wind, water or ice, has suffered varying degrees of abrasion, and is finally deposited.

The processes of lithification include compaction, cementation, and the chemical and mineralogical alteration of unstable minerals, for example feldspars being altered to clays.

The classification of terrigenous sediments and rocks is based on their composition and grain size.

The principal components are fragments (clasts) of rock, quartz and feldspar, together with clay minerals.

Grain size varies from boulder to clay (Figs. 2.1 and 2.2)

2.2.1 Rudites - conglomerates and breccias

These are made up of clasts > 2 mm in diameter.

Gravels and conglomerates

In the majority of cases the clasts are derived from a variety of source rocks separate from the site of deposition (extraformational). However, in some instances penecontemporaneous reworking of a sediment during deposition produces cohesive fragments which are redeposited (intraformational).

ϕ	diameter mm	Particle	Size Terms (rock group name)		Common types of terrigenous sediment
- 8	256	Boulder	Gravel	Rudaceous (rudite)	Conglomerate or breccia
- 6	64	Cobble			
- 2	4	Pebble			
- 1	2	Granule			
0	1	Very coarse sand	Arenaceous (arenite)	Psammitic (psammite)	Sandstone
1	0.5	Coarse sand			
2	0.25	Medium sand			
3	0.125	Fine sand			
4	0.0625	Very fine sand			
8	0.0039	Silt	Argillaceous or Lutaceous (lutite)	Pelitic (pelite)	Siltstone
		Clay			Mudstone

Fig. 2.1 - Sedimentary size grades and terms. The ϕ (phi) scale is a logarithmic transformation of the Wentworth scale: $\phi = -\log_2 d$, where d is the diameter in mm (Krumbein, 1934, 1936). Particle data after Wentworth (1922).

Rock types		
Rudite	Conglomerates	Normal, 4 – 64 mm Fine, <4 mm Sandy, >20% sand Clayey, >20% clay
Arenite	Sandstones	Conglomeratic, >20% pebble Pebbly, 10 – 20% pebbles Very coarse, 1 – 2 mm Coarse, 0.5 – 1 mm Normal Medium, 0.25 – 0.5 mm Fine, 0.125 – 0.25 mm Very fine, 0.063 – 0.125 mm Silty, >20% silt Clayey, >20% clay
	Fine clastics	Sandy siltstones, >20% sand Siltstone Silty shale Shale
Lutite		

Fig. 2.2 - Classification of terrigenous rocks based on grade size (after Krynine, 1948).