

NONLINEAR
PHYSICAL
SCIENCE

Sergey P. Kuznetsov

Hyperbolic Chaos

A Physicist's View

双曲混沌

一个物理学家的观点



高等教育出版社
HIGHER EDUCATION PRESS

Hyperbolic Chaos

A Physicist's View

双曲混沌

一个物理学家的观点

Shuangqu Hundun

With 214 figures



高等教育出版社·北京
HIGHER EDUCATION PRESS BEIJING

Author

Sergey P. Kuznetsov

Kotel'nikov's Institute of Radio-Engineering and

Electronics of RAS, Saratov Branch

410019, Zelenaya 38, Saratov, Russian Federation

Email: spkuz@rambler.ru

© 2011 Higher Education Press, 4 Dewai Dajie, 100120, Beijing, P. R. China

图书在版编目 (CIP) 数据

双曲混沌: 一个物理学家的观点 = Hyperbolic Chaos : A Physicist's View :

英文 / (俄罗斯) 库兹涅佐夫著 . — 北京 : 高等教育出版社, 2011.9

(非线性物理科学 / 罗朝俊, (瑞典) 伊布拉基莫夫, (墨) 阿弗莱
诺维奇主编)

ISBN 978-7-04-031964-4

I. ①双… II. ①库… III. ①混沌理论 - 研究 - 英文

IV. ① O415.5

中国版本图书馆 CIP 数据核字 (2011) 第 156888 号

策划编辑 王丽萍

责任编辑 李华英

封面设计 杨立新

责任校对 胡晓琪

责任印制 朱学忠

出版发行 高等教育出版社

社 址 北京市西城区德外大街 4 号

邮政编码 100120

印 刷 涿州市星河印刷有限公司

开 本 787 × 1092 1/16

印 张 21

字 数 400 000

购书热线 010-58581118

咨询电话 400-810-0598

网 址 <http://www.hep.edu.cn>

<http://www.hep.com.cn>

网上订购 <http://www.landaco.com>

<http://www.landaco.com.cn>

版 次 2011 年 9 月第 1 版

印 次 2011 年 9 月第 1 次印刷

定 价 69.00 元

本书如有缺页、倒页、脱页等质量问题, 请到所购图书销售部门联系调换

版权所有 侵权必究

物 料 号 31964-00

Sales only inside the mainland of China

{仅限中国大陆地区销售}

本书海外版由 Springer 负责在中国大陆地区以外区域销售, ISBN 为 978-3-642-23665-5

NONLINEAR PHYSICAL SCIENCE

NONLINEAR PHYSICAL SCIENCE

Nonlinear Physical Science focuses on recent advances of fundamental theories and principles, analytical and symbolic approaches, as well as computational techniques in nonlinear physical science and nonlinear mathematics with engineering applications.

Topics of interest in *Nonlinear Physical Science* include but are not limited to:

- New findings and discoveries in nonlinear physics and mathematics
- Nonlinearity, complexity and mathematical structures in nonlinear physics
- Nonlinear phenomena and observations in nature and engineering
- Computational methods and theories in complex systems
- Lie group analysis, new theories and principles in mathematical modeling
- Stability, bifurcation, chaos and fractals in physical science and engineering
- Nonlinear chemical and biological physics
- Discontinuity, synchronization and natural complexity in the physical sciences

SERIES EDITORS

Albert C.J. Luo

Department of Mechanical and Industrial
Engineering
Southern Illinois University Edwardsville
Edwardsville, IL 62026-1805, USA
Email: aluo@siue.edu

Nail H. Ibragimov

Department of Mathematics and Science
Blekinge Institute of Technology
S-371 79 Karlskrona, Sweden
Email: nib@bth.se

Valentin Afraimovich

San Luis Potosi University
IICO-UASLP, Av. Karakorum 1470
Lomas 4a Seccion, San Luis Potosi
SLP 78210, Mexico
Email: valentin@cactus.iico.uaslp.mx

INTERNATIONAL ADVISORY BOARD

Ping Ao

Eugene Benilov

Maurice Courbage

James A. Glazier

Jose Antonio Tenreiro Machado

Josep J. Masdemont

Sergey Prants

Jian Qiao Sun

Pei Yu

Jan Awrejcewicz

Eshel Ben-Jacob

Marian Gidea

Shijun Liao

Nikolai A. Magnitskii

Dmitry E. Pelinovsky

Victor I. Shrira

Abdul-Majid Wazwaz

Preface

One of intensively developing research directions is exploration of complex dynamics and chaos in nonlinear systems. We regard an object as a *dynamical system*, if its state at any given time may be obtained from the initial state according to a certain rule defined for the given system. It is remarkable that such definition, although representing ideal of deterministic vision, does not exclude a possibility of *chaotic behavior* of the object, with dependence of the state on time looking like a random process. Chaos occurs in systems of different nature, e.g. relating to mechanics, hydrodynamics, electronics, laser physics and nonlinear optics, chemical kinetics, and biomedical disciplines. The main attribute of the dynamical chaos is *sensitivity to small perturbations of initial conditions*, which makes it impossible to predict the future states actually for times longer than some characteristic scale, which usually depends logarithmically on the inaccuracy of the initial conditions (“horizon of predictability”).

For dissipative dynamical systems, chaos is associated with presence in the state space of a curious object called *the strange attractor*. At present, the collection of models with strange attractors is very rich, including artificial mathematical examples, models of physical, chemical, and biological systems.

A classic example of chaotic attractor is *Lorenz attractor*. It occurs in a set of three first-order differential equations modeling fluid convection or dynamics of a single-mode laser. The Lorenz model was a subject of extensive and careful studies in recent decades. Accurate mathematical foundation of chaotic nature of dynamics of the Lorenz model was found to be a subtle and delicate problem. It was announced as the 14-th Smale problem (from the list of difficult mathematical problems suggested by Steven Smale in 1998, as challenging mathematicians of the 21st century, similar to the list of Hilbert problems advanced for the 20th century). The solution was given by W. Tucker in 2002 by combination of technique of computer-assisted proof and careful analytic considerations.

Maybe, there was a lost opportunity for the nonlinear science to find out an alternative, a less painful way to discover physically meaningful and mathematically validated examples of chaotic behavior.

About 40 years ago, in mathematical studies a special type of chaotic attractors was introduced, namely, the *uniformly hyperbolic attractors*. They relate to systems belonging to the so-called *axiom A class* and are considered in the *hyperbolic theory* associated with the names of Anosov, Alekseev, Smale, Williams, Sinai, Plykin, Ruelle, Pesin, Newhouse and others. Chaotic nature of dynamics on the uniformly hyperbolic attractors is proved rigorously. They possess a property of *structural stability*, i.e., the phase space arrangement, character of dynamics, and its statistical characteristics are insensitive to variation of parameters and functions in the governing equations.

Originally, it was expected that the uniformly hyperbolic attractors will be relevant to many physical situations of occurrence of dynamical chaos. However, as time passed and many examples of chaotic systems of different nature had been suggested and studied, it became clear that these examples do not fit the narrow frames of the early hyperbolic theory. Therefore, people started to think of the uniformly hyperbolic attractors rather as of refined abstract image of chaos that has no relation to real systems. Efforts of mathematicians were redirected at developing generalizations applicable to broader classes of systems. For example, notions of singular-hyperbolic (or quasi-hyperbolic) attractors, non-uniformly hyperbolic attractors, partially hyperbolic attractors, and quasi-attractors were introduced, and certain progress in their exploration was achieved.

Abandoned for a long time and not clarified until recently is a question on a possibility of occurrence of the dynamical behavior associated with uniformly hyperbolic attractors in real world systems or, at least, in specially designed systems in physics and technology.

In textbooks on and reviews of nonlinear dynamics, the uniformly hyperbolic attractors are represented usually by artificial discrete time models based on geometric constructions, often explained qualitatively, in words, or by graphic images. Of course, for a physicist this is only a starting point for a research.

First of all, in addition to the geometric constructions it is desirable to have examples in a form of explicitly written equations, which allow application of computer methods for analyzing the dynamics and calculation of quantitative characteristics interesting to possible applications.

For some physical systems description in discrete time is very natural, and it would be worth examining possibilities of occurrence of the hyperbolic attractors in maps relating to such systems.

Next, it is important to turn to continuous time systems, which are of the first place importance, e.g., in physics and technology.

It is desirable to have a clear vision of how to implement the dynamics on uniformly hyperbolic attractors using combinations of structural elements known in the context of the theory of oscillations and in applications (oscillators, coupled systems, and feedback loops).

Finally, the proposed models should be implemented as real operating devices, for example, in electronics, mechanics, nonlinear optics, and technical applications of such devices should be indicated with explanation of advantages over alternative possible solutions.

In the theory of oscillations, since classic works of Andronov and his school, rough or structurally stable systems are regarded as those subjected to priority research, and as the most important for practice. It seems natural that the same should relate to systems with structurally stable uniformly hyperbolic attractors. The lack of physical examples in this regard looks as an evident dissonance. From a methodological point of view, the situation is similar to that in the early 20th century concerning the limit cycles, before their role as mathematical images for self-oscillations was established. In a similar way, the uniformly hyperbolic chaotic attractors should find their place as images for phenomena in real systems. It will help to link the abstract hyperbolic theory developed by mathematicians with description of real systems and attribute this theory to physical content.

This book is devoted to review the modern state of the outlined research program. The material is presented in four parts.

The introductory Part I consists of two chapters. Chapter 1 is devoted to the necessary basic concepts, including the notion of dynamical system, attractor, Poincaré map, and hyperbolicity. We introduce and discuss classic examples of uniformly hyperbolic attractors: Smale-Williams solenoid, DA-attractor of Smale, and attractors of Plykin type. An overview is given to the content of the hyperbolic theory (the cone criterion, structural stability, Markov partitions and symbolic dynamics, measures of Sinai-Ruelle-Bowen, etc.). Chapter 2 is a review of the literature indicating various possible situations of occurrence of uniformly hyperbolic attractors in dynamical systems.

Part II is the basic one. Here, we introduce a number of examples of systems, which allow physical implementation, and possess uniformly hyperbolic attractors with one-dimensional unstable manifolds (one positive Lyapunov exponent). In Chap. 3 we consider models, which admit a natural stroboscopic discrete time description. These are systems operating under periodic pulse kicks and systems whose dynamics is composed of a periodic sequence of stages, each of which is governed by a particular form of the differential equations. Chapter 4 discusses models designed on a base of two self-oscillators which are excited alternately because of forced external parameter modulation and transmit the excitation to each other in such way that its phase transforms in accordance with expanding circle map. Chapter 5 is devoted to autonomous systems operating according to the same principle. In Chap. 6 we consider schemes of parametric generators of chaos with hyperbolic attractors. Chapter 7 is devoted to methods of computer verification of the hyperbolic nature of attractors illustrated with the examples suggested in the previous chapters. Particularly, we consider technique of visualization of mutual location of stable and unstable manifolds, statistics of angles of intersections between them, visualization of the natural invariant measure distributions on the attractors, and verification of the cone criterion.

Part III is devoted to model systems, for which mathematical justification of the hyperbolic nature of the attractors is more problematic. Hypothetically, however, the hyperbolicity apparently takes place. This assertion is based on the fact that from a physical point of view, those models are built following the same principles as the systems considered in Chaps. 4 and 5, for which the hyperbolicity is justified at

the level of computations. In Chap. 8 we consider non-autonomous systems of four alternately excited oscillators. Among them, there is a model, in which the transformation of phases during a period of parameter modulation follows the Anosov map on the torus, and a model associated with a map on the torus with hyperchaotic dynamics (more than one positive Lyapunov exponent). Also a system is considered constructed of two coupled elements, with each possessing a hyperbolic attractor. For this case, some details are studied for the transition corresponding to complete synchronization of chaos. In Chap. 9, an autonomous system is considered functioning due to dynamics nearby a heteroclinic cycle in the amplitude equations. On this basis, we implement an attractor of Smale-Williams type in the Poincaré map, an attractor with dynamics of cyclic (phase) variables governed by the Anosov map, and a hyperchaotic attractor. Chapter 10 is devoted to systems with delayed feedback, in which a chaotic map for the phases of successively generated oscillation trains is arranged due to the transfer of excitation from the previous stages of activity of a single oscillatory element (self-oscillator) to the next stages. In Chap. 11 we consider a high-dimensional system built on two ensembles of self-oscillators with global coupling within each of them. Due to the turning the coupling on and off alternately in two ensembles, they undergo periodic Kuramoto transitions from synchronization to desynchronization and back; the interaction between the ensembles is organized in such way that the phase of the oscillatory excitation for the mean field transferred to each other, follows the chaotic map.

Part IV is devoted to examples of hyperbolic (or hypothetically hyperbolic) attractors in electronic experiments available to date. In Chap. 12 we consider experiments with a non-autonomous system of two alternately excited self-oscillators manifesting dynamics on the attractor of Smale-Williams type. In Chap. 13 two variants of non-autonomous time-delay systems with periodic parameter modulation are discussed. One possesses presumably an attractor of Smale-Williams type, embedded in an infinite-dimensional state space of the respective stroboscopic Poincaré map, and the other corresponds to a partially hyperbolic attractor.

The Appendix address some matters, which are essential to the whole presentation and illustrations, but dropping out of the basic structure of the book. We consider algorithm for computing Lyapunov exponents based on the Gram-Schmidt orthogonalization, and derivation of the Hénon and Ikeda maps from physical considerations for systems with impulse kicks. Further, the construction of the Smale horseshoe is presented, which gives a nontrivial example of a non-attractive complicated invariant set. The Kaplan-Yorke formula is derived and explained connecting approximately the Lyapunov exponents and dimensions of chaotic attractors. Formal definition of the model of Hunt is reproduced, which implements suspension of attractor of Plykin type. A simple model of hyperbolic dynamics on a compact surface of negative curvature is considered. The final Appendix is devoted to the shadowing property of hyperbolic chaotic attractors, which is illustrated in computations for a model from Chap. 4 with added noise.

The book may be useful to physicists and engineers interested in the practical application of the theory of deterministic chaos, particularly in obtaining robust chaos insensitive to parameters and characteristics of components, fluctuations, interfer-

ences, etc. This may relate to various disciplines—mechanics, hydrodynamics, electronics, laser physics, and nonlinear optics.

Contents of the book can probably help to think about possibilities of occurrence of structurally stable chaos in systems of different physical nature. Intriguing area for contemplations may be the significance of such chaos in neurodynamics.

On the other hand, the book can be useful for mathematicians interested in concrete applications of the hyperbolic theory. For them, it may be of interest to see how the mathematical theories are refracted from the perspective of the applied disciplines.

The author tried to present the material in a style available to graduate and post graduate students of non-mathematical specialties and to make it, as far as possible, self-consistent to allow a study without recourse to other sources. I tried to avoid formal definitions and formulations with abusing mathematical symbolism, replacing them with rather intuitive qualitative arguments. Perhaps, a part of mathematically oriented readers will find it insufficient; they are referred to rich literature on mathematical theory of dynamical systems. Also, I must warn that the consideration of general content of nonlinear dynamics is restricted here by the minimum needed for understanding the substantive material of the book. So, it can not be regarded as a substitute for a systematic study of integral general courses of nonlinear dynamics, to which the interested reader is referred for this purpose.

I am grateful, for useful discussions, assistance, and constructive criticism, to V.S. Anishchenko, V.S. Afraimovich, B.P. Bezruchko, V.N. Belykh, A.Yu. Jalnine, A.Yu. Zhironov, O.B. Isaeva, A.P. Kuznetsov, P.V. Kuptsov, J. Kurths, A.Yu. Loskutov, R. MacKay, L.A. Mel'nikov, A. Pikovsky, V.I. Ponomarenko, A.G. Rozhnev, N.M. Ryskin, I.R. Sataev, E.P. Seleznev, W. Tucker, D.V. Treshchev, D.I. Trubetskoy, L.V. Turukina.

Also, I thank all colleagues and administrative staff of Saratov Branch of Kotelnikov's Institute of Russian Academy of Sciences for many years of friendly collaboration, support, and creative atmosphere stimulating the scientific work.

The researches laying the foundation of this book were supported, in part, by RFBR grants 06-02-16619, 09-02-00426, 09-02-00707, and DFG-RFBR grants 04-02-04011, 08-02-91963. Additionally, the author acknowledges a support from program of Development of Scientific Potential of Higher Education of Ministry of Education of Ministry of Education and Science of Russian Federation (grant 2.1.1/1738).

Sergey P. Kuznetsov

Saratov, February 2011

Contents

Part I Basic Notions and Review

1	Dynamical Systems and Hyperbolicity	3
1.1	Dynamical systems: basic notions	3
1.1.1	Systems with continuous and discrete time, and their mutual relation	3
1.1.2	Dynamics in terms of phase fluid: Conservative and dissipative systems and attractors	6
1.1.3	Rough systems and structural stability	8
1.1.4	Lyapunov exponents and their computation	10
1.2	Model examples of chaotic attractors	12
1.2.1	Chaos in terms of phase fluid and baker's map	12
1.2.2	Smale-Williams solenoid	15
1.2.3	DA-attractor	16
1.2.4	Plykin type attractors	17
1.3	Notion of hyperbolicity	19
1.4	Content and conclusions of the hyperbolic theory	22
1.4.1	Cone criterion	24
1.4.2	Instability	25
1.4.3	Transversal Cantor structure and Kaplan-Yorke dimension	25
1.4.4	Markov partition and symbolic dynamics	26
1.4.5	Enumerating of orbits and topological entropy	27
1.4.6	Structural stability	28
1.4.7	Invariant measure of Sinai-Ruelle-Bowen	29
1.4.8	Shadowing and effect of noise	30
1.4.9	Ergodicity and mixing	30
1.4.10	Kolmogorov-Sinai entropy	31
	References	31

2	Possible Occurrence of Hyperbolic Attractors	35
2.1	The Newhouse-Ruelle-Takens theorem and its relation to the uniformly hyperbolic attractors	35
2.2	Lorenz model and its modifications	37
2.3	Some maps with uniformly hyperbolic attractors	40
2.4	From DA to the Plykin type attractor	43
2.5	Hunt's example: Suspending the Plykin type attractor	46
2.6	The triple linkage: A mechanical system with hyperbolic dynamics	49
2.7	A possible occurrence of a Plykin type attractor in Hindmarsh-Rose neuron model	51
2.8	Blue sky catastrophe and birth of the Smale-Williams attractor	52
2.9	Taffy-pulling machine	53
	References	54

Part II Low-Dimensional Models

3	Kicked Mechanical Models and Differential Equations with Periodic Switch	59
3.1	Smale-Williams solenoid in mechanical model: Motion of a particle on a plane under periodic kicks	60
3.2	A set of switching differential equations with attractor of Smale-Williams type	65
3.3	Explicit dynamical system with attractor of Plykin type	68
3.3.1	Plykin type attractor on a sphere	68
3.3.2	Plykin type attractor on the plane	73
3.4	Plykin-like attractor in smooth non-autonomous system	76
	References	79
4	Non-Autonomous Systems of Coupled Self-Oscillators	81
4.1	Van der Pol oscillator	81
4.2	Smale-Williams attractor in a non-autonomous system of alternately excited van der Pol oscillators	84
4.3	System of alternately excited van der Pol oscillators in terms of slow complex amplitudes	93
4.4	Non-resonance excitation transfer	94
4.5	Plykin-like attractor in non-autonomous coupled oscillators	95
4.5.1	Representation of states on a sphere and equations of the model	95
4.5.2	Numerical results for the coupled oscillators	98
	References	101
5	Autonomous Low-dimensional Systems with Uniformly Hyperbolic Attractors in the Poincaré Maps	103
5.1	Autonomous system of two coupled oscillators with self-regulating alternating excitation	103

5.2	System constructed on a base of the predator-prey model	107
5.3	Example of blue sky catastrophe accompanied by a birth of Smale-Williams attractor	112
	References	117
6	Parametric Generators of Hyperbolic Chaos	119
6.1	Parametric excitation of coupled oscillators. Three-frequency parametric generator and its operation	120
6.2	Hyperbolic chaos in parametric oscillator with Q-switch and pump modulation	123
6.2.1	Dynamical equations	123
6.2.2	Qualitative explanation of the operation	126
6.2.3	Numerical results	127
6.2.4	Numerical results in the frame of method of slow complex amplitudes	129
6.3	Parametric generator of hyperbolic chaos based on four coupled oscillators with pump modulation	131
6.3.1	Model, operation principle and basic equations	132
6.3.2	Chaotic dynamics: results of computer simulation	134
	References	139
7	Recognizing the Hyperbolicity: Cone Criterion and Other Approaches	141
7.1	Verification of transversality for manifolds	141
7.1.1	Visualization of the manifolds	142
7.1.2	Distributions of angles of the manifold intersections	144
7.2	Visualization of invariant measures	150
7.3	Cone criterion and examples of its application	155
7.3.1	Procedure of verification of the cone criterion	155
7.3.2	Examples of application of the cone criterion	161
	References	169
Part III Higher-Dimensional Systems and Phenomena		
8	Systems of Four Alternately Excited Non-autonomous Oscillators ...	173
8.1	Arnold's cat map dynamics in a system of coupled non-autonomous van der Pol oscillators	173
8.2	Dynamics corresponding to hyperchaotic maps	180
8.2.1	System implementing toral hyperchaotic map	180
8.2.2	Model with cascade transfer of excitation upward the frequency spectrum	182
8.3	Hyperchaos and synchronous chaos in a system of coupled non-autonomous oscillators	187
8.3.1	Equations and basic modes of operation	188
8.3.2	Equations for slow complex amplitudes	193
	References	199

9	Autonomous Systems Based on Dynamics Close to Heteroclinic Cycle	201
9.1	Heteroclinic connection: an example of Guckenheimer and Holmes	201
9.2	Attractor of Smale-Williams type in a system of three coupled self-oscillators	203
9.3	Attractor with dynamics governed by the Arnold cat map	207
9.4	Model with hyperchaos	210
9.5	An autonomous system with attractor of Smale-Williams type with resonance transfer of excitation in a ring array of van der Pol oscillators	213
	References	217
10	Systems with Time-delay Feedback	219
10.1	Some notions concerning differential equations with deviating argument	220
10.2	Van der Pol oscillator with delayed feedback, parameter modulation and auxiliary signal	223
10.2.1	Attractor of Smale-Williams type in the time-delayed system	224
10.2.2	Hyperchaotic attractors	227
10.3	Van der Pol oscillator with two delayed feedback loops and parameter modulation	231
10.4	Autonomous time-delay system	237
	References	240
11	Chaos in Co-operative Dynamics of Alternately Synchronized Ensembles of Globally Coupled Self-oscillators	243
11.1	Kuramoto transition in ensemble of globally coupled oscillators	243
11.2	Model of two alternately synchronized ensembles of oscillators	247
11.2.1	Collective chaos in ensemble of van der Pol oscillators	248
11.2.2	Slow-amplitude approach	251
11.2.3	Description of the dynamics in terms of ensembles of phase oscillators	254
	References	256
Part IV Experimental Studies		
12	Electronic Device with Attractor of Smale-Williams Type	259
12.1	Scheme of the device and the principle of operation	259
12.2	Experimental observation of the Smale-Williams attractor	260
	References	263

13 Delay-time Electronic Devices Generating Trains of Oscillations with Phases Governed by Chaotic Maps	265
13.1 Van der Pol oscillator with delayed feedback, parameter modulation and auxiliary signal	265
13.2 Van der Pol oscillator with two delayed feedback loops and parameter modulation	269
References	272
14 Conclusion	273
References	275
Appendix A Computation of Lyapunov Exponents: The Benettin Algorithm	277
References	279
Appendix B Hénon and Ikeda Maps	281
References	287
Appendix C Smale's Horseshoe and Homoclinic Tangle	289
References	292
Appendix D Fractal Dimensions and Kaplan-Yorke Formula	293
References	297
Appendix E Hunt's Model: Formal Definition	299
References	303
Appendix F Geodesics on a Compact Surface of Negative Curvature	305
References	309
Appendix G Effect of Noise in a System with a Hyperbolic Attractor	311
References	317
Index	319

Part I
Basic Notions and Review

