

# Graduate Texts in Mathematics

Alan F. Beardon

## The Geometry of Discrete Groups

离散群几何

Springer

世界图书出版公司  
[www.wpcbj.com.cn](http://www.wpcbj.com.cn)

Alan F. Beardon

# The Geometry of Discrete Groups

With 93 Illustrations



Springer

## 图书在版编目(CIP)数据

离散群几何 = The Geometry of Discrete Groups:  
英文/(英)比尔登(Beardon, A. F.)著. 一影印  
本. 一北京:世界图书出版公司北京公司, 2011. 7  
ISBN 978-7-5100-3755-9

I. ①离… II. ①比… III. ①离散数学—英文  
②几何学—英文 IV. ①O158 ②O18

中国版本图书馆 CIP 数据核字(2011)第 139062 号

---

书 名: The Geometry of Discrete Groups

作 者: Alan F. Beardon

中译名: 离散群几何

责任编辑: 高蓉 刘慧

---

出 版 者: 世界图书出版公司北京公司

印 刷 者: 三河市国英印务有限公司

发 行 者: 世界图书出版公司北京公司(北京朝内大街 137 号 100010)

联系电话: 010-64021602, 010-64015659

电子信箱: kjb@wpchj.com.cn

---

开 本: 24 开

印 张: 15

版 次: 2011 年 07 月

版权登记: 图字:01-2011-2558

---

书 号: 978-7-5100-3755-9/0·890

定 价: 45.00 元

---

Alan F. Beardon  
University of Cambridge  
Department of Pure Mathematics  
and Mathematical Statistics  
16 Mill Lane  
Cambridge CB2 1SB  
England

*Editorial Board*

S. Axler  
Mathematics Department  
San Francisco State University  
San Francisco, CA 94132  
USA

F.W. Gehring  
Mathematics Department  
East Hall  
University of Michigan  
Ann Arbor, MI 48109  
USA

K.A. Ribet  
Department of  
Mathematics  
University of California  
at Berkeley  
Berkeley, CA 94720  
USA

---

Mathematics Subject Classifications (1991): 30-01, 30 CXX, 20F32, 30 FXX, 51 M10, 20 HXX

---

Library of Congress Cataloging in Publication Data  
Beardon, Alan F.

The geometry of discrete groups.

(Graduate texts in mathematics; 91)

Includes bibliographical references and index.

1. Discrete groups. 2. Isometries (Mathematics)  
3. Möbius transformations. 4. Geometry, Hyperbolic.  
I. Title. II. Series.

QA171.B364 1983 512'.2 82-19268

© 1983 by Springer-Verlag New York Inc.

All rights reserved. No part of this book may be translated or reproduced in any form without written permission from Springer-Verlag, 175 Fifth Avenue, New York, New York 10010, U.S.A.

This reprint has been authorized by Springer-Verlag (Berlin/Heidelberg/New York) for sale in the Mainland China only and not for export therefrom.

ISBN 0-387-90788-2 Springer-Verlag New York Berlin Heidelberg  
ISBN 3-540-90788-2 Springer-Verlag Berlin Heidelberg New York SPIN 10743260

# Graduate Texts in Mathematics 91

*Editorial Board*

S. Axler F.W. Gehring K.A. Ribet

**Springer**

*New York*

*Berlin*

*Heidelberg*

*Barcelona*

*Hong Kong*

*London*

*Milan*

*Paris*

*Singapore*

*Tokyo*

*To Toni*

# Preface

This text is intended to serve as an introduction to the geometry of the action of discrete groups of Möbius transformations. The subject matter has now been studied with changing points of emphasis for over a hundred years, the most recent developments being connected with the theory of 3-manifolds: see, for example, the papers of Poincaré [77] and Thurston [101]. About 1940, the now well-known (but virtually unobtainable) Fenchel–Nielsen manuscript appeared. Sadly, the manuscript never appeared in print, and this more modest text attempts to display at least some of the beautiful geometrical ideas to be found in that manuscript, as well as some more recent material.

The text has been written with the conviction that geometrical explanations are essential for a full understanding of the material and that however simple a matrix proof might seem, a geometric proof is almost certainly more profitable. Further, wherever possible, results should be stated in a form that is invariant under conjugation, thus making the intrinsic nature of the result more apparent. Despite the fact that the subject matter is concerned with groups of isometries of hyperbolic geometry, many publications rely on Euclidean estimates and geometry. However, the recent developments have again emphasized the need for hyperbolic geometry, and I have included a comprehensive chapter on analytical (not axiomatic) hyperbolic geometry. It is hoped that this chapter will serve as a “dictionary” of formulae in plane hyperbolic geometry and as such will be of interest and use in its own right. Because of this, the format is different from the other chapters: here, there is a larger number of shorter sections, each devoted to a particular result or theme.

The text is intended to be of an introductory nature, and I make no apologies for giving detailed (and sometimes elementary) proofs. Indeed,

many geometric errors occur in the literature and this is perhaps due, to some extent, to an omission of the details. I have kept the prerequisites to a minimum and, where it seems worthwhile, I have considered the same topic from different points of view. In part, this is in recognition of the fact that readers do not always read the pages sequentially. The list of references is not comprehensive and I have not always given the original source of a result. For ease of reference, Theorems, Definitions, etc., are numbered collectively in each section (2.4.1, 2.4.2, ...).

I owe much to many colleagues and friends with whom I have discussed the subject matter over the years. Special mention should be made, however, of P. J. Nicholls and P. Waterman who read an earlier version of the manuscript, Professor F. W. Gehring who encouraged me to write the text and conducted a series of seminars on parts of the manuscript, and the notes and lectures of L. V. Ahlfors. The errors that remain are mine.

*Cambridge, 1982*

ALAN F. BEARDON



# Contents

## CHAPTER 1

<b>Preliminary Material</b>	<b>1</b>
1.1. Notation	1
1.2. Inequalities	2
1.3. Algebra	3
1.4. Topology	3
1.5. Topological Groups	5
1.6. Analysis	6

## CHAPTER 2

<b>Matrices</b>	<b>9</b>
2.1. Non-singular Matrices	9
2.2. The Metric Structure	11
2.3. Discrete Groups	14
2.4. Quaternions	16
2.5. Unitary Matrices	17

## CHAPTER 3

<b>Möbius Transformations on <math>\mathbb{R}^n</math></b>	<b>20</b>
3.1. The Möbius Group on $\mathbb{R}^n$	20
3.2. Properties of Möbius Transformations	28
3.3. The Poincaré Extension	33
3.4. Self-mappings of the Unit Ball	36
3.5. The General Form of a Möbius Transformation	40
3.6. Distortion Theorems	42
3.7. The Topological Group Structure	45
3.8. Notes	54

## CHAPTER 4

Complex Möbius Transformations	56
4.1. Representations by Quaternions	56
4.2. Representation by Matrices	60
4.3. Fixed Points and Conjugacy Classes	64
4.4. Cross Ratios	75
4.5. The Topology on $\mathcal{M}$	78
4.6. Notes	82

## CHAPTER 5

Discontinuous Groups	83
5.1. The Elementary Groups	83
5.2. Groups with an Invariant Disc	92
5.3. Discontinuous Groups	94
5.4. Jørgensen's Inequality	104
5.5. Notes	115

## CHAPTER 6

Riemann Surfaces	116
6.1. Riemann Surfaces	116
6.2. Quotient Spaces	117
6.3. Stable Sets	122

## CHAPTER 7

Hyperbolic Geometry	126
---------------------	-----

*Fundamental Concepts*

7.1. The Hyperbolic Plane	126
7.2. The Hyperbolic Metric	129
7.3. The Geodesics	134
7.4. The Isometries	136
7.5. Convex Sets	138
7.6. Angles	141

*Hyperbolic Trigonometry*

7.7. Triangles	142
7.8. Notation	144
7.9. The Angle of Parallelism	145
7.10. Triangles with a Vertex at Infinity	146
7.11. Right-angled Triangles	146
7.12. The Sine and Cosine Rules	148
7.13. The Area of a Triangle	150
7.14. The Inscribed Circle	151

*Polygons*

7.15. The Area of a Polygon	153
7.16. Convex Polygons	154
7.17. Quadrilaterals	156
7.18. Pentagons	159
7.19. Hexagons	160

*The Geometry of Geodesics*

7.20. The Distance of a Point from a Line	162
7.21. The Perpendicular Bisector of a Segment	164
7.22. The Common Orthogonal of Disjoint Geodesics	165
7.23. The Distance Between Disjoint Geodesics	166
7.24. The Angle Between Intersecting Geodesics	166
7.25. The Bisector of Two Geodesics	166
7.26. Transversals	167

*Pencils of Geodesics*

7.27. The General Theory of Pencils	168
7.28. Parabolic Pencils	169
7.29. Elliptic Pencils	170
7.30. Hyperbolic Pencils	170

*The Geometry of Isometries*

7.31. The Classification of Isometries	171
7.32. Parabolic Isometries	172
7.33. Elliptic Isometries	172
7.34. Hyperbolic Isometries	173
7.35. The Displacement Function	174
7.36. Isometric Circles	176
7.37. Canonical Regions	177
7.38. The Geometry of Products of Isometries	179
7.39. The Geometry of Commutators	184
7.40. Notes	187

## CHAPTER 8

## Fuchsian Groups

8.1. Fuchsian Groups	188
8.2. Purely Hyperbolic Groups	190
8.3. Groups Without Elliptic Elements	198
8.4. Criteria for Discreteness	200
8.5. The Nielsen Region	202
8.6. Notes	203

## CHAPTER 9

## Fundamental Domains

9.1. Fundamental Domains	204
9.2. Locally Finite Fundamental Domains	206
9.3. Convex Fundamental Polygons	217
9.4. The Dirichlet Polygon	226
9.5. Generalized Dirichlet Polygons	234
9.6. Fundamental Domains for Coset Decompositions	238
9.7. Side-Pairing Transformations	240
9.8. Poincaré's Theorem	242
9.9. Notes	252

<b>CHAPTER 10</b>	
<b>Finitely Generated Groups</b>	<b>253</b>
10.1. Finite Sided Fundamental Polygons	253
10.2. Points of Approximation	258
10.3. Conjugacy Classes	263
10.4. The Signature of a Fuchsian Group	268
10.5. The Number of Sides of a Fundamental Polygon	274
10.6. Triangle Groups	276
10.7. Notes	286
<b>CHAPTER 11</b>	
<b>Universal Constraints on Fuchsian Groups</b>	<b>287</b>
11.1. Uniformity of Discreteness	287
11.2. Universal Inequalities for Cycles of Vertices	288
11.3. Hecke Groups	293
11.4. Trace Inequalities	295
11.5. Three Elliptic Elements of Order Two	301
11.6. Universal Bounds on the Displacement Function	308
11.7. Canonical Regions and Quotient Surfaces	324
11.8. Notes	327
<b>References</b>	<b>329</b>
<b>Index</b>	<b>335</b>

## CHAPTER 1

# Preliminary Material

### §1.1. Notation

We use the following notation. First,  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}$  and  $\mathbb{C}$  denote the integers, the rationals, the real and complex numbers respectively;  $\mathbb{H}$  denotes the set of quaternions (Section 2.4).

As usual,  $\mathbb{R}^n$  denotes Euclidean  $n$ -space, a typical point in this being  $x = (x_1, \dots, x_n)$  with

$$|x| = (x_1^2 + \dots + x_n^2)^{1/2}.$$

Note that if  $y > 0$ , then  $y^{1/2}$  denotes the positive square root of  $y$ . The standard basis of  $\mathbb{R}^n$  is  $e_1, \dots, e_n$  where, for example,  $e_1 = (1, 0, \dots, 0)$ . Certain subsets of  $\mathbb{R}^n$  warrant special mention, namely

$$B^n = \{x \in \mathbb{R}^n : |x| < 1\},$$

$$H^n = \{x \in \mathbb{R}^n : x_n > 0\},$$

and

$$S^{n-1} = \{x \in \mathbb{R}^n : |x| = 1\}.$$

In the case of  $\mathbb{C}$  (identified with  $\mathbb{R}^2$ ) we shall use  $\Delta$  and  $\partial\Delta$  for the unit disc and unit circle respectively.

The notation  $x \mapsto x^2$  (for example) denotes the function mapping  $x$  to  $x^2$ : the domain will be clear from the context. Functions (maps or transformations) act on the *left*: for brevity, the image  $f(x)$  is often written as  $fx$  (omitting brackets). The composition of functions is written as  $fg$ : this is the map  $x \mapsto f(g(x))$ .

Two sets  $A$  and  $B$  *meet* (or  $A$  *meets*  $B$ ) if  $A \cap B \neq \emptyset$ . Finally, a property  $P(n)$  holds *for almost all*  $n$  (or *all sufficiently large*  $n$ ) if it fails to hold for only a finite set of  $n$ .

## §1.2. Inequalities

All the inequalities that we need are derivable from Jensen's inequality: for a proof of this, see [90], Chapter 3.

**Jensen's Inequality.** Let  $\mu$  be a positive measure on a set  $X$  with  $\mu(X) = 1$ , let  $f: X \rightarrow (a, b)$  be  $\mu$ -integrable and let  $\phi: (a, b) \rightarrow \mathbb{R}$  be any convex function. Then

$$\phi\left(\int_X f d\mu\right) \leq \int_X (\phi f) d\mu. \quad (1.2.1)$$

Jensen's inequality includes Hölder's inequality

$$\int_X fg d\mu \leq \left(\int_X f^2 d\mu\right)^{1/2} \left(\int_X g^2 d\mu\right)^{1/2}$$

as a special case: the discrete form of this is the Cauchy-Schwarz inequality

$$|\sum a_i b_i| \leq (\sum |a_i|^2)^{1/2} (\sum |b_i|^2)^{1/2}$$

for real  $a_i$  and  $b_i$ . The complex case follows from the real case and this can, of course, be proved by elementary means.

Taking  $X = \{x_1, \dots, x_n\}$  and  $\phi(x) = e^x$ , we find that (1.2.1) yields the general Arithmetic-Geometric mean inequality

$$y_1^{\mu_1} \cdots y_n^{\mu_n} \leq \mu_1 y_1 + \cdots + \mu_n y_n,$$

where  $\mu$  has mass  $\mu_j$  at  $x_j$  and  $y_j = \phi f(x_j)$ .

In order to apply (1.2.1) we need a supply of convex functions: a sufficient condition for  $\phi$  to be convex is that  $\phi^{(2)} \geq 0$  on  $(a, b)$ . Thus, for example, the functions  $\cot$ ,  $\tan$  and  $\cot^2$  are all convex on  $(0, \pi/2)$ . This shows, for instance, that if  $\theta_1, \dots, \theta_n$  are all in  $(0, \pi/2)$  then

$$\cot\left(\frac{\theta_1 + \cdots + \theta_n}{n}\right) \leq \frac{\cot \theta_1 + \cdots + \cot \theta_n}{n}. \quad (1.2.2)$$

As another application, we prove that if  $x$  and  $y$  are in  $(0, \pi/2)$  and  $x + y < \pi/2$  then

$$\tan x \tan y \leq \tan^2\left(\frac{x + y}{2}\right). \quad (1.2.3)$$

Writing  $w = (x + y)/2$ , we have

$$\begin{aligned}\frac{\tan x + \tan y}{1 - \tan x \tan y} &= \tan(x + y) \\ &= \frac{2 \tan w}{1 - \tan^2 w}.\end{aligned}$$

As  $\tan$  is convex, (1.2.1) yields

$$\tan x + \tan y \geq 2 \tan w$$

and the desired inequality follows immediately (noting that  $\tan^2 w < 1$  so both denominators are positive).

## §1.3. Algebra

We shall assume familiarity with the basic ideas concerning groups and (to a lesser extent) vector spaces. For example, we shall use elementary facts about the group  $S_n$  of permutations of  $\{1, 2, \dots, n\}$ : in particular,  $S_n$  is generated by transpositions. As another example, we mention that if  $\theta: G \rightarrow H$  is a homomorphism of the group  $G$  onto the group  $H$ , then the kernel  $K$  of  $\theta$  is a normal subgroup of  $G$  and the quotient group  $G/K$  is isomorphic to  $H$ .

Let  $g$  be an element in the group  $G$ . The elements *conjugate* to  $g$  are the elements  $ghg^{-1}$  in  $G$  ( $h \in G$ ) and the conjugacy classes  $\{ghg^{-1}: h \in G\}$  partition  $G$ . In passing, we mention that the maps  $x \mapsto xgx^{-1}$  and  $x \mapsto gxxg^{-1}$  (both of  $G$  onto itself) play a special role in the later work. The *commutator* of  $g$  and  $h$  is

$$[g, h] = ghg^{-1}h^{-1};$$

for our purposes this should be viewed as the composition of  $g$  and a conjugate of  $g^{-1}$ .

Let  $G$  be a group with subgroups  $G_i$  ( $i$  belonging to some indexing set). We assume that the union of the  $G_i$  generate  $G$  and that different  $G_i$  have only the identity in common. Then  $G$  is the *free product* of the  $G_i$  if and only if each  $g$  in  $G$  has a unique expression as  $g_1 \cdots g_n$  where no two consecutive  $g_i$  belong to the same  $G_j$ . Examples of this will occur later in the text.

## §1.4. Topology

We shall assume a knowledge of topology sufficient, for example, to discuss Hausdorff spaces, connected spaces, compact spaces, product spaces and homeomorphisms. In particular, if  $f$  is a 1-1 continuous map of a compact

space  $X$  onto a Hausdorff space  $Y$ , then  $f$  is a homeomorphism. As special examples of topologies we mention the discrete topology (in which every subset is open) and the topology derived from a metric  $\rho$  on a set  $X$ . An isometry  $f$  of one metric space  $(X, \rho)$  onto another, say  $(Y, \sigma)$ , satisfies

$$\sigma(fx, fy) = \rho(x, y)$$

and is necessarily a homeomorphism.

Briefly, we discuss the construction of the quotient topology induced by a given function. Let  $X$  be any topological space, let  $Y$  be any non-empty set and let  $f: X \rightarrow Y$  be any function. A subset  $V$  of  $Y$  is *open* if and only if  $f^{-1}(V)$  is an open subset of  $X$ : the class of open subsets of  $Y$  is indeed a topology  $\mathcal{T}_f$  on  $Y$  and is called the *quotient topology induced by  $f$* . With this topology,  $f$  is automatically continuous. The following two results on the quotient topology are useful.

**Proposition 1.4.1.** *Let  $X$  be a topological space and suppose that  $f$  maps  $X$  onto  $Y$ . Let  $\mathcal{T}$  be any topology on  $Y$  and let  $\mathcal{T}_f$  be the quotient topology on  $Y$  induced by  $f$ .*

- (1) *If  $f: X \rightarrow (Y, \mathcal{T})$  is continuous, then  $\mathcal{T} \subset \mathcal{T}_f$ .*
- (2) *If  $f: X \rightarrow (Y, \mathcal{T})$  is continuous and open, then  $\mathcal{T} = \mathcal{T}_f$ .*

**PROOF.** Suppose that  $f: X \rightarrow (Y, \mathcal{T})$  is continuous. If  $V$  is in  $\mathcal{T}$ , then  $f^{-1}(V)$  is open in  $X$  and so  $V$  is in  $\mathcal{T}_f$ . If, in addition,  $f: X \rightarrow (Y, \mathcal{T})$  is an open map then  $V$  in  $\mathcal{T}_f$  implies that  $f^{-1}(V)$  is open in  $X$  and so  $f(f^{-1}V)$  is in  $\mathcal{T}$ . As  $f$  is surjective,  $f(f^{-1}V) = V$  so  $\mathcal{T}_f \subset \mathcal{T}$ .  $\square$

**Proposition 1.4.2.** *Suppose that  $f$  maps  $X$  into  $Y$  where  $X$  and  $Y$  are topological spaces,  $Y$  having the quotient topology  $\mathcal{T}_f$ . For each map  $g: Y \rightarrow Z$  define  $g_1: X \rightarrow Z$  by  $g_1 = gf$ . Then  $g$  is continuous if and only if  $g_1$  is continuous.*

**PROOF.** As  $f$  is continuous, the continuity of  $g$  implies that of  $g_1$ . Now suppose that  $g_1$  is continuous. For an open subset  $V$  of  $Z$  (we assume, of course, that  $Z$  is a topological space) we have

$$(g_1)^{-1}(V) = f^{-1}(g^{-1}V)$$

and this is open in  $X$ . By the definition of the quotient topology,  $g^{-1}(V)$  is open in  $Y$  so  $g$  is continuous.  $\square$

An alternative approach to the quotient topology is by equivalence relations. If  $X$  carries an equivalence relation  $R$  with equivalence classes  $[x]$ , then  $X/R$  (the space of equivalence classes) inherits the quotient topology induced by the map  $x \mapsto [x]$ . Equally, any surjective function  $f: X \rightarrow Y$  induces an equivalence relation  $R$  on  $X$  by  $xRy$  if and only if  $f(x) = f(y)$  and  $Y$  can be identified with  $X/R$ . As an example, let  $G$  be a group of homeomorphisms of a topological space  $X$  onto itself and let  $f$  map each  $x$  in  $X$



to its  $G$ -orbit  $[x]$  in  $X/G$ . If  $X/G$  is given the induced quotient topology, then  $f: X \rightarrow X/G$  is continuous. In this case,  $f$  is also an open map because if  $V$  is open in  $X$  then so is

$$f^{-1}(fV) = \bigcup_{g \in G} g(V).$$

Finally, the reader will benefit from an understanding of covering spaces and Riemann surfaces although most of the material in this book can be read independently of these ideas. Some of this is discussed briefly in Chapter 6: for further information, the reader is referred to (for example) [4], [6], [28], [50], [63] and [100].

## §1.5. Topological Groups

A *topological group*  $G$  is both a group and a topological space, the two structures being related by the requirement that the maps  $x \mapsto x^{-1}$  (of  $G$  onto  $G$ ) and  $(x, y) \mapsto xy$  (of  $G \times G$  onto  $G$ ) are continuous: obviously,  $G \times G$  is given the product topology. Two topological groups are *isomorphic* when there is a bijection of one onto the other which is both a group isomorphism and a homeomorphism: this is the natural identification of topological groups.

For any  $y$  in  $G$ , the space  $G \times \{y\}$  has a natural topology with open sets  $A \times \{y\}$  where  $A$  is open in  $G$ . The map  $x \mapsto (x, y)$  is a homeomorphism of  $G$  onto  $G \times \{y\}$  and the map  $(x, y) \mapsto xy$  is a continuous map of  $G \times \{y\}$  onto  $G$ . It follows that  $x \mapsto xy$  is a continuous map of  $G$  onto itself with continuous inverse  $x \mapsto xy^{-1}$  and so we have the following elementary but useful result.

**Proposition 1.5.1.** *For each  $y$  in  $G$ , the map  $x \mapsto xy$  is a homeomorphism of  $G$  onto itself: the same is true of the map  $x \mapsto yx$ .*

A topological group  $G$  is *discrete* if the topology on  $G$  is the discrete topology: thus we have the following Corollary of Proposition 1.5.1.

**Corollary 1.5.2.** *Let  $G$  be a topological group such that for some  $g$  in  $G$ , the set  $\{g\}$  is open. Then each set  $\{y\}$  ( $y \in G$ ) is open and  $G$  is discrete.*

Given a topological group  $G$ , define the maps

$$\phi(x) = xax^{-1}$$

and

$$\psi(x) = xax^{-1}a^{-1} = [x, a],$$