

American
DRUG INDEX

1975

American DRUG INDEX

1075



Y073270

Charles O. Wilson, Ph.D.

Dean and Professor of Pharmaceutical Chemistry,
School of Pharmacy, Oregon State University

and

Tony E. Jones, Ph.D.

Professor of Pharmaceutical Chemistry, School of Pharmacy
University of Colorado; Director of
Pharmaceutical Research, Carbisulphoil Company



J. B. LIPPINCOTT COMPANY
Philadelphia • Toronto

AMERICAN DRUG INDEX

1975

Charles O. Wilson, Ph.D.
Dean and Professor of Pharmaceutical Chemistry,
School of Pharmacy, Oregon State University

Copyright © 1975, by J. B. Lippincott Company

Copyright © 1956, 1957, 1958, 1959, 1960, 1961, 1962, 1963, 1964,
1965, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974,
by J. B. Lippincott Company

Tony E. Jones, Ph.D.
Professor of Pharmaceutical Chemistry, School of Pharmacy,
University of Colorado, Director of
Pharmaceutical Research, Cordispharm Company

This book is fully protected by copyright and, with the exception
of brief excerpts for review, no part of it may be reproduced
in any form by print, photoprint, microfilm or any other means,
without the written permission of the publishers.

ISBN 0-397-50343-1

Library of Congress Catalog Card Number 55-6286

Printed in the United States of America

Preface

The *American Drug Index* has been prepared for the identification and correlation of the many pharmaceuticals available to the medical and allied professions. The need for this index has become acute as the number of drugs and drug products in all their tremendous variety has multiplied.

The organization of the index is fundamentally alphabetical with extensive cross-indexing. Names listed are generic (also called non-proprietary, public name or common name); brand (also called trade-mark, proprietary or specialty), chemical, U.S.P. and N.F. Synonyms that are in general use also are included. All names used for a pharmaceutical occur in alphabetical order with the pertinent data given under the brand name and the name by which it is made available. Data included are generic names, chemical names, manufacturer, pharmaceutical forms, size, dosage and use. Generic names occur in alphabetical order and here the recognition of the drug is indicated by U.S.P. (United States Pharmacopeia), N.F. (National Formulary) and N.D. (New Drugs). The information is in accord with the U.S.P. XVIII, and the N.F. XIII.

Because of the multiplicity of specialty names (brand names) used for the same therapeutic agent or the same combination of therapeutic agents, it was apparent that some correlation could be done. As an example of this, please turn to tetracycline HCl. Here under the generic name are given the brand names. Following are combinations of tetracycline HCl organized in a manner to point out relationships among the many products. Reference then is made to the specialty name or names having the indicated composition. Under the specialty name are given manufacturer, composition, available forms, sizes, dosage and use.

The multiplicity of generic names for the same therapeutic agent has complicated the nomenclature of these agents. Examples of multiple generic names for the same chemical substance are (1) parabromdylamine, brompheniramine; (2) acetaminophen, p-hydroxy acetanilid, N-acetyl-p-aminophenol; (3) d-isoeephedrine, d-pseudoephedrine; (4) methapyrilene HCl, thenylpyramine HCl; and (5) pyrilamine, pyranisamine, pyranilamine, pyraminyl and anisopyradamine.

Generic names recognized by the United States Pharmacopeia Committee, and the National Formulary Committee are preceded by an asterisk. United States Adopted Names (USAN's) (generic names) are also identified in the text by an asterisk.

The *American Drug Index* should be most useful to physicians, pharmacists, dentists, nurses, and to sales personnel, students and teachers in the fields incorporating pharmaceuticals. The cross-indexing feature permits the finding of drugs or drug combinations when only one major ingredient is known. For example, a combination of aluminum hydroxide gel and magnesium trisilicate is available. This combination can be found by looking under the name of either of the two ingredients, and in each case the specialty names are given.

CHARLES O. WILSON · TONY E. JONES

A = 1 LAXATIVE. (Western Research) Aloin $\frac{1}{4}$ gr., pow. ext. stramonium $\frac{1}{16}$ gr., pow. ext. cascara sagrada $\frac{1}{2}$ gr., pow. ext. rhubarb $\frac{1}{2}$ gr., ginger $\frac{1}{32}$ gr./Tab. Bot. 1000s. Paks 28.
Use: Laxative.

A.A. ANTACID TABLETS. (Western Research) Balanced combination of calcium carbonate ppt. 583.2 mg., magnesium carbonate light, 194.4 mg., magnesium trisilicate and aluminum hydroxide dried gel 64.8 mg./Chewable Tab. Strip-pak 28s, 1000s.
Use: Antacid.

A.A.A. OINTMENT. (Jenkins) Ammoniated mercury 2%, acid salicylic 1.25%, acid boric 1%, zinc oxide 15%. Jar 1 oz., 1 lb.
Use: Impetigo, nonspecific eczema & minor skin irritations.

A-A PLUS. (Galen Labs.) Al hydroxide gel 400 mg., Mg oxide 175 mg., Na lauryl sulfate 25 mg., Me cellulose 100 mg., hyoscyamine sulfate 0.1037 mg., atropine sulfate 0.0194 mg., hyoscine HBr 0.0065 mg./Tab. Bot. 100s, 1000s. Susp. 1 pt.
Use: Bronchial asthma inhibitor.

AA RANE. (Syntex) Cromolyn sodium 20 mg./Cap. Bot. 60s.
Use: Bronchial asthma inhibitor.

ABAR-APC. (Tmco) Isobutylallyl barbituric acid 0.75 gr., aspirin 3 gr., phenacetin 2 gr., caffeine 0.66 gr./Tab. Bot. 1000s.
Use: Analgesic, sedative.

ABCDG VITAMIN CAPSULES. (Lannett) Vit. A 5000 u., D 400 u., B-1 1 mg., B-2 2 mg., C 30 mg./Cap. Bot. 1000s.
Use: Vitamin supplement.

ABDEC DROPS. (Parke, Davis) Vit. A 5000 u., D 400 u., B-1 1 mg., B-2 1.2 mg., B-6 1 mg., sodium pantothenate 5 mg., nicotinamide 10 mg., C 50 mg./0.6 cc. Bot. $\frac{1}{2}$ oz., $\frac{1}{3}$ oz. W/dropper.
Use: Vitamin deficiencies.

ABDEC BABY VITAMIN DROPS W/FLUORIDE. (Parke, Davis) Vit. A. palmitate 5000 u., calciferol 400 u., B-1 1 mg., B-2 1.2 mg., B-6 1 mg., sodium d-pantothenate 5.0 mg., niacinamide 10 mg., Vit. C 50 mg., sodium fluoride 0.5 mg./0.6 ml. Bot. 50 ml. W/dropper.
Use: Dental caries preventive.

ABDEC KAPSEALS. (Parke, Davis) Vit. A 10,000 u., D 400 u., B-1 5 mg., B-2 3 mg., B-6 1.5 mg., B-12 2 mcg., dl-panthanol 10 mg., nicotinamide 25 mg., Vit. C 75 mg., mixed tocopherols 5 I.U./Cap. Bot. 100s, 250s., unit dose pkg 10s.
Use: Multivitamin therapy.

ABDOL W/Minerals. (Parke, Davis) Vit. C 50 mg., B-2 2.5 mg., B-12 1 mcg., B-6 0.5 mg., B-1 2.5 mg., nicotinamide 20 mg., Vit. A 5000 u., D 400 u., Ca pantothenate 2.5 mg., folic acid 0.1 mg., I 0.15 mg., Mn 1 mg., K 5.0 mg., Fe 15 mg., Cu 1 mg., Zn 0.5 mg., Mg 1 mg., Ca 44 mg., P 34 mg./Cap. Bot. 100s, 250s, 5000s.
Use: Vitamin-mineral deficiency states.

ABDOL W/VIT. C. (Parke, Davis) Vit. A 5000 u., D 400 u., B-1 2.5 mg., B-2 2.5 mg., B-6 0.5 mg., B-12 1 mcg., nicotinamide 20 mg., Vit. C 50 mg., pantothenic acid (as Cal. salt) 5 mg./Cap. Bot. 100s, 5000s.
Use: Vitamin deficiency.

ABEREL. (McNeil) Retinoic acid (Vit. A acid) 0.05%. Bot. 60 cc. Swabs 15s.
Use: Acne.

ABSINOL. (Lemmon) Acetaminophen 150 mg., salicylamide 150 mg., phenyltoloxamine dihydrogen citrate 25 mg., phenylephrine HCl 2.5 mg./Tab. Bot. 1000s.

ABSORBABLE CELLULOSE COTTON OR GAUZE.

See: Oxidized Cellulose (Various Mfr.)
• **ABSORBABLE DUSTING POWDER, U.S.P. XVII.**

ABSORBABLE GELATIN FILM. Sterile nonantigenic, absorbable, water-insoluble, gelatin film.
See: Gelfilm (Upjohn)

• **ABSORBABLE GELATIN SPONGE, U.S.P. XVIII.** Gelatin sponge.

See: Gelfoam, Preps. (Upjohn)

• **ABSORBABLE SURGICAL SUTURE, U.S.P. XVIII.** (Various Mfr.) Surgical Gut, Surgical Catgut, Catgut suture.

ABSORBENT BASE. (Upsher-Smith) Oint. Base: Cetyl alcohol, stearyl alcohol, lauryl sulfate, methyl p-hydroxybenzoate & hydrocarbons. Jar 1 lb.
Use: Ointment base.

ABSORBENT GAUZE, U.S.P. XVIII.

Use: Surgical aid.

ABSORBINE ARTHRITIC PAIN LOTION.

(Young) Bot. 2 oz., 4 oz.

ABSORBINE FOOT POWDER. (Young) Plastic bot. 2 oz. w/shaker top.

Use: Athlete's foot.

ABSORBINE, JR. (Young) Wormwood, thymol, YG-7 (chloroxylenol), menthol, acetone. Liq. Bot. 1 oz., 2 oz., 4 oz., 12 oz. w/applicator.

Use: Muscle soreness, stiffness.

ABSORBINE MEDICATED SOAP. (Young). Bar 3 $\frac{1}{2}$ oz.

ABUSCREEN. (Roche Diagnostics) An immunological and radiochemical assay for morphine and morphine glucuronide in nanogram levels. Utilizes ^3H labeled morphine assayed with liquid scintillation counters using conventional beta counting procedures and ^{125}I labeled morphine requiring gamma scintillation equipment. 100 tests.

A.C.A. CAPS. (Scrip) Aspirin 3 $\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr., acetophenetidin 2 $\frac{1}{2}$ gr./Cap. Bot. 100s.

A.C.A. No. 2. (Scrip) Aspirin 4 gr., caffeine $\frac{1}{4}$ gr., acetophenetidin 2 gr./Tab. Bot. 100s.

A.C.A. No. 5. (Scrip) Phenacetin 2 gr., salicylamide 2 gr., atropine sulfate 1/600 gr., tr. gelsemium 3 min., camphor monobromate $\frac{1}{2}$ gr./Tab. Bot. 100s.

A-CAINE. (A.V.P.) Benzocaine 2%, cod liver oil, zinc oxide, balsam peru, bismuth subgallate, ephedrine sulfate and lanolin in a special base. Tube w/applicator, 1 $\frac{1}{2}$ oz.

- Use: Temporary relief of hemorrhoids, minor rectal irritation and as an aid in rectal examination.
- A CAPSULES.** (Stanlabs) Natural Vit. A 5,000 u., 10,000 u., 25,000 u., 50,000 u./Cap. Bot. 100s. Synthetic or soluble, 10,000 u., 25,000 u., 50,000 u./Cap. Bot. 100s.
- **ACACIA**, U.S.P. XVIII. (Penick) Acacia senegal, gum arabic. Mucilage N.F. XII; pow., $\frac{1}{4}$ lb., tears, $\frac{1}{4}$ -1 lb., syr. N.F. XIII. whole, $\frac{1}{4}$ -1 lb. Use: Demulcent, emulsifier.
- A.C.A. NASAL DROPS.** (Emerson) Phenylephrine HCl in a saline solution. Dropper bot. 1 oz. Use: Relief of nasal blockage in infants and children.
- A-CAP.** (A.V.P.) Vit. A 50,000 u./Cap. Bot. 100s.
- A-CAPS.** (Drug Industries) Vit. A 50,000 u./Cap. Bot. 100s, 500s. Use: Vitamin A therapy.
- ACCELERASE.** (Organon) Pancrelipase 165 mg., bile salts 65 mg., cellulase 2 mg., calcium carbonate 20 mg./Cap. Bot. 60s, 500s. Use: Functional digestive disorders.
- ACCELERASE-PB.** (Organon) Same as Accelerase, with belladonna alkaloids (87.5% hyoscyamine and 12.5% atropine sulf.) 0.2 mg., phenobarbital 16 mg./Cap. Bot. 60s, 500s. Use: Digestive enzyme.
- ACCOPAIN.** (Vitarine) Codeine phosphate $\frac{1}{4}$ gr., buffered aspirin $\frac{3}{2}$ gr., phenacetin $2\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./Cap. Bot. 24s. Use: Analgesic.
- A.C.D. CAPSULES.** (Philips Roxane) Pwd. opium 1.62 mg., phenacetin 97.5 mg., aspirin 97.5 mg., caffeine 4 mg., camphor monobromated 16.2 mg., atropine sulfate 0.065 mg./Cap. Bot. 1000s.
- A-C-D SOLUTION.** U.S.P. Sod. citrate, citric acid & dextrose in sterile pyrogen-free solution. Abbott-(Abbo-Vac) 250, 450, 500 cc. (non-Vac) 500 ml. (Pliapak) w/Blood collection set 500 cc. Baxter-500 cc. bot. with 70, 120 or 300 cc. of sol.; 1000 cc. with 500 cc. of sol. Cutter-500 cc. bot. with 75 or 120 cc. of sol.; 650 cc. bot. with 80 or 130 cc. of sol. Diamond-(Abbo-Vac) 250 cc., 500 cc. Use: Anticoagulant for preparation of plasma or whole blood.
- A-C-D SOLUTION MODIFIED.** (Squibb) Acid citrate dextrose anticoagulant solution modified. Use: Diagnostic agent for cancer.
- **ACECLIDINE.** (Kingshill) 3-Quinuclidinol acetate (ester). Glauco-stat. Under study. Use: Parasympathomimetic.
- **ACEDAPSONE.** (Parke, Davis) 4',4"-Sulfonylbis (acetanilide). Under study. Use: Antimalarial; antibacterial (leprostatic).
- ACEDOVAL.** (Vale) Dover's powder $\frac{1}{4}$ gr., phenacetin $2\frac{1}{2}$ gr., caffeine anhydrous $\frac{1}{8}$ gr., camphor $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s, 5000s.
- ACEDYNE.** (Vale) Phenacetin $\frac{1}{2}$ gr., aspirin $\frac{1}{2}$ gr., caffeine anhydrous $\frac{1}{2}$ gr., atropine sulfate $\frac{1}{2}$ gr., gr., magnesium trisilicate 1 gr./Tab. Bot. 100s, 1000s, 5000s.
- ACELAX.** (Marcen) Danthron 100 mg., prune concentrate 64 mg., chondrus drispus 64 mg./Tab. Bot. 100s, 1000s. Use: Laxative.
- ACENOCOUMARIN.** See: Sintrom, Tab. (Geigy)
- **ACENOCOUMAROL.** N.D. 1967. 3-(Alpha-acetyl-4-nitrobenzyl)-4-hydroxycoumarin. See: Sintrom, Tab. (Geigy).
- **ACEPROMAZINE.** 10(3-Dimethylaminopropyl)-3-acetylphenothiazine maleate. Plegicil Use: Tranquillizer.
- ACEROLA C.** (Kirkman) Vit. C 100 mg./Wafer. Bot. 100s.
- ACETA.** (Century) Acetaminophen. Elix. (120 mg./5 cc) Bot. 4 oz., pint, gal. Tabs. (5 gr.) Bot. 100s, 1000s, 3000s, 5000s. Supp. 300 mg., 600 mg. 12s. Use: Analgesic.
- ACETA W/ CODEINE.** (Century) Acetaminophen 325 mg., codeine phosphate $\frac{1}{2}$ gr./Tab. Bot. 100s. Use: Analgesic.
- ACETABAR.** (Philips Roxane) Phenobarbital $\frac{1}{4}$ gr., acetophenetidin $2\frac{1}{2}$ gr., acetylsalicylic acid $\frac{3}{2}$ gr., caffeine $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s. Use: Analgesic-sedative.
- ACETAGESIC.** (Kay) Acetaminophen 120 mg./5 cc. Elix. Bot. pts., gals. Use: Analgesic and antipyretic, in pain and fever syndrome.
- p-ACETAMIDOBENZONIC ACID SALT** of 2-dimethylaminoethanol. See: Deanan, Tab. (Riker).
- **ACETAMINOPHEN**, N.F. XIII. Elix., Tabs: N.F. XIII. N-Acetyl-p-amino-phenol. See: Aceta, Tab. Elix., Supp. (Century) Acetagesic (Kay) Anapap, Tab. (Fellows) Anapol, Liq. (Kenyon) Anelix, Elix. (Kirkman) Apadon, Tab. (Elder) Apamide, Tab. (Denab) Apap, Cap. (TPR Pharmaceuticals) Canital, Tab. (Carnrick) Ce. Apap, Liq. (Central) Chlo. 1-Tyl. (Jenkins) Conace. J. Elix., Tab. (C.M.C.) Dapa, Tab. (Ferndale) Dimindol, Tab., Elix. (Blue Line) Dularin, Tab. Elix. (Donner) Duo-Darth, Tab. (Lemmon) Elixodyne, Liq. (Lemmon) Febridol, Elix., Tab. (Amid) G-1 (Palmedico) Janupap, Tab. (Tutag) Lestemp, Elix. (Reid-Provident) Lyteca, Syr. (Westerfield) Med-Apap, Elix. (Med. Chem.) Medigesic (Medics) Menalgisia, Tab. (Otis Clapp) Metalid, Tab. (Philips Roxane) Nebs, Tab., Liq. (Eaton)

• Generic name: approved by F.D.A., or recognized as USAN or by U.S.P., N.F. or N.D.

Neopap, Supp. (Webcon)
 Nilprin 7½, Tab. (AVP)
 Pirin, Tab. (Scrip)
 Proval, Cap., Elix., Drop., Tab. (Reid-Provident)
 Pyrapap, Tab., Susp. (Savage)
 Rystal, Syr. (Grail)
 SK-Apap, Tab., Elix. (SmithKline)
 Taper, Elix., Tab. (Parke, Davis)
 Temetan, Elix., Tab. (Nevin)
 Temptra, Drop, Syr., Tab. (Mead Johnson)
 Tenlap, Elix. (Dow)
 Tussapap, Tab. (Suttliff & Case)
 Tylaprin, Elix. (Cenci)
 Tylenol, Elix., Tab. (McNeil)
 Valadol, Tab., Liq. (Squibb)
 Valgesic, Elix. (Kirk)
 Valorin Tab. (Otis Clapp)

ACETAMINOPHEN W/COMBINATIONS

Absinol, Tab. (Lemmon)
 Aceta w/Codeine, Tab. (Century)
 Algecol, Tab. (Winston)
 Algoson (McNeil)
 Allylgesic, Tab. (Elder)
 Allylgesic W/Ergotamine, Cap. (Elder)
 Allylvon, Cap. (Elder)
 Alumadrine, Tab. (Fleming)
 Amino-Bar, Tab. (Bowman)
 Aminodyne, Elix. (Bowman)
 Amodynos Forte, Tab. (Buffington)
 Apap w/Codeine, Tab. (Central)
 Apapalone, Tab. (Amfre-Grant)
 Arthol Tabs. (Towne)
 Arthralgen, Tab. (Robins)
 Bancaps (Westerfield)
 Bancaps-C (Westerfield)
 Ban-O-Pain, Cap. (Daniels)
 Banesin-Forte, Tab. (Westerfield)
 Bexahist, Cap. (Bexar)
 Biophens-S, Tab. (La Crosse)
 Bromo Quinine Cold Tabs (Vantage)
 Butigetec, Tab. (McNeil)
 Calm Aids, Cap. (Daniels, Robert)
 Capital w/Codeine, Tab. (Carnrick)
 Ceroze Comp., Cap. (Ives)
 Chexit, Tab. (Dorsey)
 Coastaldyne, Tab. (Coastal)
 Coastalgesic, Tab. (Coastal)
 Codalan, Tab. (Lannett)
 Codap (Tutag)
 Codrene W/Codeine, Syr. (Daniels, Robert)
 Codrene-D S.F., Syr. (Daniels, Robert)
 Codimal, Tab. (Central)
 Coldene, Syr. (Pharmcraft)
 Colrex Compound, Elix. (Rowell)
 Conar-A, Tab., Susp. (Beecham-Massengill)
 Corzans, Tab. (Moore Kirk)
 Co-Tylenol, Tab. (McNeil)
 Dapase, Tab. (Ferndale)
 Darth, Tab. (Denab)
 Darvocet-N, Tab. (Lilly)
 Darvocet-N 100, Tab. (Lilly)
 Decorex, Elix. (Donley-Evans)

Desa-Hist AT, Tab. (Pharmics)
 Dialog, Prods. (Ciba)
 Dolopar, Liq. (O'Neal)
 Drinacet Cap. (Philips Roxane)
 Drinophen, Cap. (Lannett)
 D-Sinus, Cap. (Laser)
 Dularin-TH, Tab. (Dooner)
 Duo-Gesic, Tab. (Trout)
 Duoprin, Tab. (Dunhall)
 Duramid, Tab. (O'Neal)
 Dysonil, Tab. (Circle Pharm.)
 Elpanal, Tab. (Lemmon)
 Enpayne, Tab. (Burgin-Arden)
 Escotussin, Cap. (Esco)
 Esemgesic, Tab. (Spencer-Mead)
 Esgic, Tab. (Gilbert)
 Excedrin and P.M., Tab. (Bristol-Myers)
 Fever-Tabs. (Jaymass)
 Fevopain, Susp., Tab. (Crest)
 G-2, Tab. (Palmedico)
 G-3, Cap. (Palmedico)
 Gaysal, Tab. (Geriatric)
 G-Lixir (Palmedico)
 Guaamine, Cap. (Suttliff & Case)
 Hasacode, Tab. (Arnar-Stone)
 Hasamal, Tab. (Arnar-Stone)
 Histogesic, Tab. (Century)
 Histosal #2, Tab. (Ferndale)
 Hi-Temp, Tab. (Saron)
 Hycomine Comp., Tab. (Endo)
 Indogesic, Tab. (Century)
 Isomel, Cap. (Tutag)
 Kleer, Syr. (Scrip)
 Koly-Tabs. (Scrip)
 Koryza Tab. (Fellows-Testagar)
 Lique-C, Liq., Tab. (Elder)
 Medigesic, Tab. (Medics)
 Mense, Tab. (Daniels, Robert)
 Midran, Tab. (Columbia)
 Midrin, Cap. (Carnrick)
 Minotal, Tab. (Carnrick)
 Mycatussin, Cap., Syr. (Myers-Carter)
 Myolate, Tab. (Panray)
 Naldetuss, Syr. (Bristol-Myers)
 N-D Gesic, Tab. (Hyrex-Key)
 Neo-Pyranistan, Tab. (Standex Labs.)
 Neo-Vadrin Sinus Headache Formula, Tab.
 (First Texas)
 Nokane, Tab. (Wren)
 Novahistine Singlet, Tab. (Dow)
 Nylorac PB, Tab. (Beard)
 Opacedrin, Cap. (Elder)
 Ornex, Cap. (SmithKline)
 Pallacin, Tab. (Denab)
 Panitol H.M.B., Tab. (Wesley)
 Panritis, Tab. (Pan Amer.)
 Parafon Forte, Tab. (McNeil)
 Pavadon Elix. (Coastal)
 Pedittuss, Liq. (Suttliff & Case)
 Pedric, Elix., Wafers (Vale)
 Percogesic, Tab. (Endo)
 Phenahist, Elix. (Amid)

Phrenilin, Tab. (Carnrick)
 Prodolor, Tab. (Panray)
 Prdolor, Tab. (Amfre-Grant)
 Pyradyne, Tab. (Lemmon)
 Pyradyne Comp., Tab. (Lemmon)
 Pyrhist Cold Cap. (Daniels, Robert)
 Renpap, Tab. (Wren)
 Repan, Tab. (Everett)
 Rhinex, Tab. (Lemmon)
 Rhinidrin, Tab. (Central)
 Rhinogestic, Tab. (Vale)
 Rhinspec, Tab. (Lemmon)
 S.A.C. Sinus, Tabs. (Towne)
 Salaphen, Tab. (Ferndale)
 Salatin w/Codeine 1/2 Gr., Tab. (Ferndale)
 Salimeph, Preps. (Kremers Urban)
 Salphenyl, Cap., Liq. (Mallard)
 Sanspen, Tab. (Spencer-Mead)
 Santussin, Prep. (Sandia)
 Scotgesic, Cap., Elix. (Scott/Cord)
 Scotuss Pediatric Cough Syr. (Scott/Cord)
 Sedalgestic, Tab. (Table Rock)
 Sedapap, Elix. (Mayrand)
 Sedragesic, Tab. (Lannett)
 Sinarest, Tab. (Pharmacraft)
 Sinubid, Tab. (Warner Chilcott)
 Sinudan, Tab. (Daniels, Robert)
 Sinuwes (Western Research)
 Sinulin, Tab. (Carnrick)
 Sinumal, Tab., Liq. (Cook)
 Sinus Tab., Tab. (Kenyon)
 Sinutab Preps. (Warner-Chilcott)
 Spantuss, Liq. (Arco)
 Spendrisin, Tab. (Spencer-Mead)
 Stopain, Tab. (Columbia)
 Strascogestic, Tab. (Pennwalt)
 Symptomax, Liq. (Reid-Provident)
 T-Caps, Cap. (Burlington)
 Tega-Code, Cap. (Ortega)
 Tegapap, Liq., Tab. (Ortega)
 Temco, Liq. (Cook)
 Teragen, Tab. (O'Neal)
 Tercol, Liq. (Commerce)
 T-Gesic, Cap. (Tennessee)
 Triaminicin, Tab. (Dorsey)
 Triaprin, Cap. (Dunhall)
 Trind, Syr. (Mead Johnson)
 Trind-DM, Syr. (Mead Johnson)
 Tussagesic, Tab., Susp. (Dorsey)
 Two-Dyne, Tab. (Hyrex-Key)
 Valhist, Cap. (Clapp)
 Vannor, Cap. (North Amer. Pharm.)
 Windolor, Tab. (Winston)

ACETAMINOPHENOL

See: Acetaminophen.

- **ACETANILID.** (Various Mfr.) (Acetylaminobenzene, acetylaniline, antifebrin) N-phenylacetamide cry.

Use: Antipyretic, analgesic.

Powder Compound, N.F. (Various Mfr.)

ACETARSOL.

See: Acetarsone, Tabs.

- **ACETARSONE.** (City Chemical) N.F. XI., Tablets, N.F. XI. 3-Acetamido-4-hydroxy-phenylarsonic acid (Acetarsol, Acetphenarsine, Amarsan, Dynarsan, Ehrlich 594, Limarsol, Orarsan, Osarsal, Osvarsan, Paroxyl, Stovarsol.)

ACETARSONE SALT OF ARECOLINE.

See: Drocarbil, N.F.

ACETASEM. (Phillips Roxane) Phenacetin 162.5

mg., aspirin 227.5 mg., caffeine 8.1 mg., gelsemium
 pwd. extract 4.62 mg./Tab. Bot. 1000s.

Use: Analgesic.

- **ACETAZOLAMIDE**, U.S.P. XVIII. N-(5-Sulfamoyl-1,3,4-thiadiazol-2-yl) acetamide. Tab., U.S.P. XVIII.

See: Diamox, Tab. (Lederle)

Use: Carbonic anhydrase inhibitor.

- **ACETAZOLAMIDE SODIUM**, U.S.P. XVIII. Sterile, U.S.P. XVIII.

See: Diamox, Tabs. (Lederle)

Diamox Sequels, Caps. (Lederle)

ACET-DIA-MER-SULFONAMIDE, N.F. Sulfacetamide, sulfadiazine & sulfamerazine.

See: Buffonamide, Susp., Tab. (Tutag)

Buf-Sul, Tabs. & Susp. (Sheryl)

Cetazine, Tab. (Bowman)

Coco-Diazine, Susp. (Lilly)

Incorposul, Susp. & Tabs. (Blue Line)

Sulfadiazine, sulfamerazine & sulfacetamide,

Susp., N.F.; Tab. N.F.

Trizyl, Susp. (Central)

W/Sulfamethazine.

See: Delomets, Tab. (Durst)

Katrasul, Tabs., Susp. (Conal)

- **ACETEST.** (Ames) Sod. nitroprusside, disodium phosphate, aminoacetic acid, lactose. Tab. Bot. 100s, 250s.

Use: Test for acetone or diacetic acid in urine.

- **ACETIC ACID**, U.S.P. XVIII, Glacial, U.S.P. XVIII, Diluted, N.F. XIII.

ACETICYL.

See: Acetylsalicylic Acid.

ACETILUM ACIDULATUM.

See: Acetylsalicylic Acid. (Various Mfr.)

- **ACETOHEXAMIDE**, N.F. XIII. N.D. 1967. (Lilly.)

N-(p-acetylphenylsulfonyl)-N'-cyclo-hexylurea.

1-((p-acetylphenyl)sulfonyl)-3-cyclohexylurea.

See: Dymelor, Tab. (Lilly)

Use: Blood sugar lowering compound.

- **ACETOJEN, JR.** (Jenkins) Acetylsalicylic acid 1 1/2 gr., phenacetin 1 gr., hyoscyamus ext. 0.0125 gr., (total alkaloids 0.0002 gr.) gelsemium ext. 0.0125 gr./Tab. Bot. 1000s.

ACETOL.

See: Acetylsalicylic Acid. (Various Mfr.)

- **ACETOMEROCTOL.** Acetato [2-hydroxy-5-(1,1,3,3-tetramethylbutyl) phenyl] mercury. Under study.

Use: Topical antiseptic.

ACETONE, N.F. XIII.

• Generic name: approved by F.D.A., or recognized as USAN or by U.S.P., N.F. or N.D.

ACETONE or DIACETIC ACID TEST.

See: Acetest, Tab. (Ames)

ACETONYL. (Upjohn) Granular efferv. salt. Aspirin as sod. salt $7\frac{1}{2}$ gr. w/pot. & sod. citrates & tartrates having an alkalinity equiv. to $33\frac{1}{2}$ gr. Na bicarbonate, calcium lactate, sod. phosphate & magnesium sulfate/60 gr. Bot. 4 oz.

Use: Analgesic & alkaliizer.

ACETOPHEN CAPSULE. (Jenkins) Aspirin $3\frac{1}{2}$ gr., phenacetin $2\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./Cap. Bot. 1000s

ACETOPHEN C.T. GREEN. (Jenkins) Aspirin $3\frac{1}{2}$ gr., phenacetin $2\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./Tab. Bot. 1000s.

•ACETOPHENAZINE MALEATE, N.F. XIII, N.D. 1967. 2-Acetyl-10-(3-[4-(β -hydroxyethyl)-piperazinyl] propyl) phenothiazine dimaleate. 10-(3-[4-(2-Hydroxyethyl)-1-piperazinyl] propyl)-phenothiazin-2-yl methyl ketone maleate (1:2). See: Tindal Maleate, Tab. (Schering)
Use: Tranquilizer.

ACETOPHENETIDIN, U.S.P. XVI. Tablets. U.S.P. XVI. (Various Mfr.) Phenacetin, Ethoxycetanilide. Pow. $\frac{1}{4}$ lb., 1 lb., 5 lb.
Use: Antipyretic, analgesic.

ACETOSAL.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOSALIC ACID.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOSALIN.

See: Acetylsalicylic Acid. (Various Mfr.)

•ACETOSULFONE. 4,4'-Diaminodiphenylsulfone-2-N-acetylsulfonamide sodium. (N'-acetyl-6-sulfanilylmetanilamido) sodium.
See: Promacatin, Tab. (Parke, Davis)

ACETOXYPHENYLMERCURY, N.F. XIII.

See: Phenylmercuric acetate.

ACETPHENARSINE.

See: Acetarsone, Tab. (Abbott)

ACETPHENOLISATIN.

See: Isalax, Tab. (Vale)

Oxyphenisatin Acetate Prep.

W/Combinations.

See: Hydrocil Fortified, Tab. (Fuller)

W/D-Calcium pantothenate, sodium sulfosuccinate, acetphenolisatin.

See: Peri-Pantyl, Tab. (McGregor)

W/Dioctyl sod. sulfosuccinate.

See: Nuvac, Cap., Supp. (La Crosse)

W/Dioctyl sodium sulfosuccinate, guar gum.

See: Guarsol, Tabs (Western Research)

W/Dioctyl sodium sulfosuccinate, sod. carboxymethylcellulose.

See: Scrip-Lax, Tab. (Scrip)

•ACETRIZOATE SODIUM. Injection, U.S.P. XVI
3-Acetamido-2,4,6-triiodobenzoate sodium
See: Cystokon Dilution, Sol. (Mallinckrodt)

W/Polyvinylpyrrolidone

See: Salpax, Vial (Ortho)

ACETRIZOIC ACID, U.S.P. XV 3-Acetylaminio-2,4,6-triiodobenzoic acid

See: Acetrizolate, Sod.

•ACETRYPTIN HYDROCHLORIDE. (Warner-Chilcott) 5-Acetyltryptamine hydrochloride.
Use: Antihypertensive agent.

ACET-THEOCIN SODIUM.

See: Theophylline Sod. acetate (Various Mfr.)

ACETYCOL. (O'Neal) Colchicine salicylated 0.25 mg., aspirin 325.0 mg., p-aminobenzoic acid 162.0 mg., Vit. C 20.0 mg., B-1 5.0 mg., niacin 15.0 mg./Tab. Bot. 100s
Use: Analgesic.

ACETYL ADALIN.

See: Acetylcarbromal. (Various Mfr.)

p-ACETYLAMINOBENZALDEHYDE THIO-SEMICARBAZONE. (Amithiozone, Antib, Berculon A, Benzothiozon, Conteben, Myuizone, Neustab, Tethethion, Thiomicid, Thioparamizone, Thiacetazone).
Use: With streptomycin in the treatment of tuberculosis.

ACETYLAMINO BENZENE.

See: Acetanilid. (Various Mfr.)

N-ACETYL-p-AMINOPHENOL, N.F. XII.
See: Acetaminophen

ACETYLANILINE.

See: Acetanilid. (Various Mfr.)

ACETYL-BETA-METHYLCHOLINE CHLORIDE, U.S.P.
See: Mecholyl Chloride, Amp. (Baker, J.T.)

ACETYL-BROMO-DIETHYLACETYL-CARBAMIDE.
See: Acetylcarbromal. (Various Mfr.)

•ACETYLCARBOMAL. (Meyer) Acetyl adalin, acetylbromodiethylacetyl-carbamide. Pow. for mfg.
See: Carbased, Tab. (Mallard)

Paxarel, Tab. (Circle)

Sedamyl. (Riker)

Use: Sedative.

W/Bromisovalum, acetaminophen.

See: Isomel, Cap. (Tutag)

Medigesic, Tab. (Medics)

Vannor, Cap. (North Amer. Pharm.)

W/Bromisovalum, scopolamine aminoxide HBr.

See: Tranquinal, Tab. (Barnes-Hind)

ACETYLCYCLIC CHLORIDE.

Use: S.C., I.M., I.V. Parasympathomimetic agent & vasodilator. Paroxysmal tachycardia.

See: Miochol Ophthalmic (Smith, Miller & Patch)

ACETYLCYCLIC-LIKE THERAPEUTIC AGENTS.

See: Cholinergic agents.

•ACETYLCYSTEINE. F.D.A. (Mead Johnson)

N-Acetyl-L-cysteine

Use: Mucolytic agent.

See: Mucomyst, Sol. (Mead Johnson)

•ACETYLDIGITOXIN, N.F. XIII Tab. N.F. XIII

α -Digitoxin monoacetate: The natural-occurring cardiac glycoside defined as the alpha acetyl ester of digitoxin

See: Acylanid, Tab. (Sandoz)

ACETYLIN.

See: Acetylsalicylic Acid. (Various Mfr.)

•ACETYLMETHADOL. Methadyl acetate

ACETYL MIDICEL.

See: Midicel Acetyl Susp. (Parke, Davis)

ACETYLPHENYLISATIN.

See: Oxyphenisatin Acetate.

1-((p-ACETYLPHENYL)SULFONYL)-3-CYCLO-HEXYLUREA, N.F. XIII.

See: Acetohexamide.

ACETYLRESORCINOL.

See: Resorcinol Monoacetate. (Various Mfr.)

ACETYSAL.

See: Acetylsalicylic Acid

ACETYSALICYLATE ALUMINUM.

See: Aluminum Aspirin

ACETYSALICYLIC ACID, U.S.P. XVII. Cap-

sules, N.F.; Tablets, U.S.P. XVII. (Various Mfr.)

Aspirin, U.S.P. XVII. Acidum acetylsalicylicum, 2-acetoxybenzoic acid, Acetillum Acidulatum, Acetophen, Acetol, Acetosol, Acetosalic Acid, Acetosalin, Aceticyl, Acetylsal, Acetylpyrin, Aspro, Helicon, Rhodine, Salacetin, Salcetogen, Saletin.

See: Aspirin Preps. (Various Mfr.)

ACETYSALICYLIC ACID, ACETOPHENE-TIDIN AND CAFFEINE, N.F.

See: A.P.C., Prep. (Various Mfr.)

***ACETYL SULFAMETHOXYPYRIDAZINE.**

N.D. 1967. 3-(N-Acetylsulfanilamido)-6-methoxy-pyridazine.

4'-(ACETYLSULFAMOYL)PHTHALANILIC ACID, N.F. XIII.

See: Phthalylsulfacetamide

N'-ACETYSULFANILAMIDE. (Albucid; p-Aminobenzenesulfonacetamide; Sulfacet; Sulfacetamide, N-Sulfanilylacetamide.)

Use: Sulfonamide therapy.

***ACETYL SULFISOXAZOLE, N.F. XIII. Oral**

Susp. N.F. XIII. N'-Acetyl-N'-(3,4-dimethyl-5-isoxazolyl)-sulfanilamide: N-(3,4-dimethyl-5-isoxazolyl)-N-sulfanilylacetamide.

See: Gantrisin Acetyl, Syr., Susp. (Roche)

Lipo-Gantrisin Acetyl, Liq. (Roche)

***ACETYL TANNIC ACID. Tannic acid acetate.**

Use: Antiperistaltic.

5-ACETYLTRYPTAMINE HYDROCHLORIDE.

(Warner Chilcott) Potent, rapid acting oral antihypertensive agent; pending release.

ACHLORHYDRIA DETERMINATION.

See: Diagnex Blue, Prep. (Squibb)

ACHLORHYDRIA THERAPY.

See: Acidol. (Various Mfr.)

Acidol-Pepsin, Cap. (Winthrop)

Glyco-HCl, Cap. (Burnham)

Glutamic Acid HCl (Various Mfr.)

Murispin, Tab. (Norgine)

Normacid, Tab. (Stuart)

ACHOL. (Enzyme Process) Vit. A 4000 u., keto-

cholic acids 62 mg./Tab. Bot. 100s, 250s.

Use: Vitamin A supplement and therapy.

ACHOLEST. (Fougera) Sebatic acid bromthynol blue. Kit w/30 tests.

Use: Plasma cholinesterase test kit.

ACHROMYCIN. N.D. 1966. (Lederle) Tetra-

cycline HCl.

Intramuscular: (100 mg., 250 mg.) Vial. Box 1s,

100s.

Intravenous: (100 mg., 250 mg., 500 mg. w/

Vit. C.) Vial.

Oint.: (3%) Tube ½ & 1 oz.

Oint.: Eye & ear 1% ½ oz. tube.

Oint. (Ophth.) 1% Tubes ½ oz.

Ophth. Oil Susp.: 1%, Bot. w/dropper 4 cc.

ACHROMYCIN-V. (Lederle) Tetracycline HCl

100 mg./Cap. Bot. 100s; 250 mg. Bot. 16s, 100s,

1000s; Unit dose 100s; 500 mg. Bot. 100s, 1000s;

Unit dose 100s.

Syr.: 125 mg./5 cc. Bot. 2 oz., 1 pt.

Pediatric Drops: 100 mg./cc. Dropper Bot. 10 cc.

Oral Susp.: 250 mg./5 cc. Bot. 1 oz.

Use: Treatment of infections caused by both gram-negative and gram-positive bacteria.

ACHROSTATIN-V. (Lederle) Tetracycline HCl

250 mg., nystatin 250M u./Cap. Bot. 16s, 100s.

Oral Susp. (1.5 Gm. & nystatin 1,500,000 u.) Bot.

2 oz.

Use: Broad-spectrum antibiotic w/antifungal therapy.

ACICHEK. (Valentine) Dihydroxy aluminum amino-

acetate 250 mg., Sod. carboxymethylcellulose 150

mg., glycine 30 mg./Tab. Bot. 100s, 1000s.

Use: Gastric hyperacidity in peptic ulcer.

ACID ACRIFLAVINE.

See: Acriflavine Hydrochloride (Various Mfr.)

ACID CITRATE DEXTROSE SOLUTION.

See: A.C.D. Sol. (Various Mfr.)

ACID CITRATE DEXTROSE ANTICOAGU-

LANT SOLUTION MODIFIED.

See: A-C-D Solution Modified (Squibb)

ACI-DERMA. (Princer) An aqueous emulsion of

glyceryl monostearate, cetyl alcohol, sodium lauryl sulfate, propyl and benzyl p-hydroxy benzoates and propylene glycol, buffered to pH 2-3. Bot. 4 oz., 1 pt.

Use: Restore the acid mantle of the skin.

ACIDEX. (Smith, Miller & Patch) Ca carbonate 3½

gr., Mg carbonate 2 gr., bismuth subnitrate ½ gr./

Tab. Bot. 60s.

Use: Hyperacidity and peptic ulcer.

ACID HISTAMINE PHOSPHATE.

See: Histamine Phosphate. (Various Mfr.)

ACID MANTLE CREME. (Dome) Aluminum ace-

tate in specially prep. water-soluble hydrophilic cream at pH 4.2. Tube 1 oz. Jar 4 oz., 1 lb.

Use: Skin prophylactic agent in restoring normal skin acidity.

ACID MANTLE LOTION. (Dome) Aluminum ace-

tate lotion. pH 4.5 Bot. 4 oz., squeeze, 1 pt.

Use: Dermatitis.

ACIDOL.

See: Betaine Hydrochloride (Various Mfr.)

ACIDOL-PEPSIN. (Winthrop) Betaine HCl 4 gr.,

pepsin 3.5 gr./Cap. Bot. 50s.

Use: Achlorhydria.

*Generic name: approved by F.D.A., or recognized as USAN or by U.S.P., N.F. or N.D.

ACID TRYPFLAVINE.

See: Acriflavine Hydrochloride. (Various Mfr.)

ACIDULIN. (Lilly) Glutamic acid HCl 340 mg./

Pulvule. Bot. 100s, 1000s.

Use: Achlorrhya.

ACIGLUMIN.

See: Glutamic Acid Hydrochloride. (Various Mfr.)

ACI-JEL. (Ortho) Acetic acid 0.92%, ricinoleic acid 0.70%, oxyquinoline sulfate 0.025%, boric acid 3.0%, and glycerine 5% in a vegetable gum base.

Jelly, Tube with applicator, 85 Gm.

Use: Vaginal antiseptic.

ACILLIN. (ICN Pharm.) Ampicillin trihydrate equiv.

to ampicillin 250 mg., 500 mg./Cap. Bot. 100s, 1000s.

Use: Penicillin therapy.

ACIQUEL. Potassium gluconate.**ACLETIN.** (Associated Concentrates) Lecithin,

soya 8 gr. with vitamins/Cap. Bot. 100s, 250s.

Use: Fat metabolism.

A.C.N. (Person & Covey) Vitamin A palmitate

25M u., ascorbic acid 250 mg., niacinamide 25 mg.,

hesperidin 50 mg./Tab. Bot. 50s, 100s, 1000s.

Use: Acne vulgaris.

ACNAVEEN. (Cooper) Aveeno colloidal oatmeal

8%, sulfur 2%, salicylic acid 2%, in a soap-free

cream base. Plastic tube 4 oz. Bar 3.8 oz.

Use: Antipruritic cleanser and shampoo.

ACNA-VITE. (Cenci) Vit. A 10,000 u., C 250 mg.,

hesperidin 50 mg., niacinamide 25 mg./Cap. Bot. 75s.

Use: Acne vulgaris with Vit. A & C deficiency.

ACNE-AID CREAM. (Stiefel) Sulfur 2.5%, resorcinol

1.25%, parachlorometaxylene 0.375% w. Micel

(brand of microporous cellulose). Tube, 1½ oz.

Use: Oily acne.

ACNE-AID DETERGENT SOAP. (Stiefel) Natural

soap with nonirritating detergent. Bar 4.1 oz.

Use: Cleanse oily skin.

ACNE-AID LOTION. (Stiefel) Sulfur 10%, alcohol

10% in a washable base. Clear or Tinted Bot. plastic 1 oz. or 2 oz.

Use: Treatment of acne.

ACNE CORT DOME. (Dome) Hydrocortisone

acetate ¼%, sulfur 4%, hexachlorophene 1%,

resorcinol monoacetate 3%. Cream. Tube 1 oz.

Lotion. Squeeze Bot. 1 oz.

Use: Treatment of acne.

ACNE CREAM. (De Pree) Sulfur 8%, resorcinol 2%,

hexachlorophene 1%, allantoin, zinc oxide, alcohol

9.25%. Tube 1½ oz.

Use: Dermatologic.

ACNEDERM. (Lannett) Lotion: Dispersable

sulfur 5%, zinc sulfate 3%. Soap: Zinc oxide 2%,

zinc sulfate 1%, colloidal sulfur 5%. Cake 3.2 oz.

Use: Skin conditions

ACNE-DOME CREAM & LOTION. (Dome)

Regular: sulfur 4%, resorcinol monoacetate 3%.

Cream: Tube 1 oz. Lotion: Bot. 2 oz.

Use: Treatment of acne.

ACNE-DOME MEDICATED CLEANSER.

(Dome) Sulfur 2%, salicylic acid 2% in a colloidal

soybean complex. Medicated w/sponge 4 oz.

Use: Medicated skin cleanser.

ACNESARB LOTION. (C & M Pharmcal) Salicylic

acid 3%, boric acid 2%, hyamine 10X 0.08%, isopropyl alcohol. Bot. 4 oz., pt.

Use: Anti-acne lotion.

ACNESOL. (C & M Pharmcal) Salicylic acid 3%,

boric acid 2%, isopropyl alcohol 63%, hexachlorophene 0.5%. Bot. 4 oz., pt.

Use: Dermatologic.

ACNESTROL LOTION. (Dermik) Diethylstilbestrol

0.0875% (equiv. to 0.70 mg. diethylstilbestrol),

29% isopropyl alcohol. Bot. 2 oz.

Use: Acne vulgaris.

ACNETONE-ACNE LOTION. (Whorton) Lot. 4 oz.,

16 oz.

ACNIPLEX. (G. S. Herbert) Sulfur, resorcinol, hexa-

chlorophene, aluminum chlorhydroxy allantoinate, alcohol 3.7%. Cream ¼ oz.

Use: Antiseptic w/sulfur.

ACNO. (Cummins) Micronized sulfur 3%, isopropanol

25%, salicylic acid 2%. Bot. 8 oz.

Use: Acne & related skin conditions.

ACNOMEAD. (Spencer-Mead) Colloidal sulfur 2%,

salicylic acid 1% w/soapless cleansers, wetting agents, and polyethylene granules. Tube 1½ oz., jar 1 lb.

Use: Acne, scrub cream.

ACNOMEL. (SmithKline) Resorcinol 2%, sulfur 8%,

hexachlorophene 0.25% w/bentonite, propylene glycol, alcohol & titanium dioxide.

Cream. Tube 1 oz. Cake. Sulfur 4%, resorcinol 1%,

hexachlorophene 0.25%. 1-oz. containers.

Use: Acne therapy.

ACNOPHILL. (Torch) Sulfur 5%, Pot. and zinc sul-

fides and polysulfides 5%, zinc oxide 10% in hydrophilic vehicle. Plain or tinted. Jar 1 lb. and 2 oz.

Use: Acne.

ACNOTEX. (C & M Pharmcal) Sulfur 8%, salicylic

acid 2.25%, hyamine 10X 0.08%, propylene glycol, powder base, acetone, isopropyl alcohol. Bot. 2 oz., pt.

Use: Dermatologic lotion.

ACNYCIN CREAM. (Columbia) Hexachlorophene

0.5%, sulfur 5%, resorcinol 2% and zinc oxide.

Tubes, 1 oz.

Use: Acne.

ACOGESIC. (Pennwalt) Phenyltoloxamine dihydro-

gen citrate 25 mg., acetyl-p-aminophenol 300 mg., salicylamide 200 mg./Cap. Bot. 50s.

Use: Analgesic.

ACON. (Endo) Water-soluble vitamin A. Cap. (25,000

& 50,000 u./Cap.) Bot. 100s.

Use: Vitamin A therapy.

ACONITINE. Aconitine U.S.P.X.**ACOTUS.** (Whorton) Phenylephrine HCl 5 mg.,

guaiaacal glyceryl ether 100 mg., menthol 1 mg.,

alcohol, by vol. 10% 5 cc. Bot. 4 oz., gal.

Use: Colds, coughs.

ACRIFLAVINE. N.F. X. (Euflavine, Gonacrine,

Neutroflavine.) Acriflavine neutral; a mixture

of 2, 8-diamino-10-methylacridinium chloride

& 2,8-diaminoacridine. (Lilly) Tab. 1½ gr.

Bot. 100s.

Use: External antiseptic.

ACRIFLAVINE HYDROCHLORIDE, N.F. X, (Various Mfr.) Hydrochloride form of acriflavine, N. F. Acid acriflavine, acid tryptaflavine, flavine, tryptaflavine.

National Aniline—Pow., Bot. (1, 5, 10, 25, & 50 Gm.). Tab. (1½ gr.) Bot. 50s & 100s.

Use: External bacteriostatic.

ACRISAN EAR DROPS, (Recsei) Aminoacridine propionate 0.3%, alkyl-parahydroxy-benzoate 1%, acetic acid 2%, chlorobutanol 3%, benzyl alcohol 1.5%, hydrocortisone 0.1% in propylene glycol. Ear drops, ½ oz. also available w/Hydrocortisone. Bot. ½ oz.

Use: Anti-infective.

• **ACRISORCIN**, N.D. 1967. 9-Aminoacridinium 4-hexylresorcinolate.

See: Akrinol Cream (Schering)

Use: Topical, tinea versicolor.

• **ACRONINE**, (Lilly) 3,12-Dihydro-6-methoxy-3,12-trimethyl-7H-pyranol[2,3-c]acridin-7-one. Under study.

Use: Antineoplastic.

ACTABS, (Blair) Aspirin w/ascorbic acid. Bot. 100s, 1000s.

ACTABS BUFFERED, (Blair) Aspirin w/ascorbic acid and cal. carbonate. Bot. 100s, 1000s.

ACTAMER, (Monsanto Chem.)

See: Bithionol, U.S.P.

ACTAMIN, (Burlington) Acetaminophen 325 mg./Tab. Dispens-A-Kit: 500s.

Use: Pain reliever.

ACTH-ACTEST GEL, (Fellows) Repository corticotropin 40 or 80 u./cc. Vial 5 cc.

• **ACTH**, N.D. 1966. (Adrenocorticotrophic hormone.

Adrenocorticotrop(h)in).

See: Corticotropin, U.S.P.

Arden Pharm. (40 or 80 u./ml., 5 ml.)

National Drug—20 u./ml., 2 ml., 10 ml.

Parke, Davis—25 units/Vial; 40 units/Vial.

Testagar—40 units/ml., vial 5 ml.

Wesley Pharm. (40 or 80 u./ml., 5 ml.)

ACTH GEL, PURIFIED, Arcum—40 or 80 units/ml., vial 5 ml. Conal—40 or 80 units/ml., vial 5 ml. Hart Labs.—40 units/ml., vial 5 ml. Bowman—Adrenocorticotrophic hormone 40 u., aqueous gelatin 16%, phenol 0.5%/cc. Vial 5 cc.

Use: Repository corticotropin.

Fellows-Testagar—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.

Intra Prods.—40 units/ml., dosette 5 ml.; 80 units/ml., dosette 5 ml.

King Pharm.—40 units/ml., vial 1 ml.

Medical Chem.—40 or 80 units/ml., vial 5 ml.

Merrell-National—20 u./ml. 2 ml., 10 ml., 40 units/ml., vial 1 ml., 5 ml.; 80 units/ml., vial 5 ml.

Philips Roxane—Repository corticotropin inj., U.S.P.

40 units in 16% aq. gelatin. Vial 5 ml.

Rocky Mtn. Pharmacal—40 units/ml., 80 units/ml. Vial 5 ml.

Wesley Pharm.—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.

ACTHAR, N.D. 1966. (Armour) Corticotropin, U.S.P. Vial 25, 40 u. (Lyophilized).

ACTHAR GEL HIGHLY PURIFIED, (Armour) H. P. acthar gel, a new, highly purified, repository ACTH preparation that may be given subcutaneously or intramuscularly. 40 & 80 u. per cc., 1 cc., 5 cc. multiple dose vials.

See: Corticotropin, U.S.P.

Use: Stimulant of adrenal cortex, anti-inflammatory, antiallergic & useful in collagen diseases.

ACTICORT LOTION 100, (Cummins) Hydrocortisone alcohol 1% in a specially formulated vehicle. Bot. 4 oz.

Use: Anti-inflammatory & antipruritic.

ACTIDL, (Burroughs Wellcome) Triprolidine HCl Tab. (2.5 mg.) Bot. 100s. Syr. (1.25 mg./5 cc.) Bot. 1 pt.

Use: Antihistamine.

ACTI-DIONE, (Upjohn) Cycloheximide. (3-[2-(3,5-Dimethyl-2-oxocyclohexyl)-2-hydroxyethyl]-glutamide) 85-100% w/w.

Not for medicinal use. For professional use only.

Bot. 4 Gm.

ACTIFED, (Burroughs Wellcome) Triprolidine HCl. 2.5 mg., pseudoephedrine HCl. 60 mg./Tab. Bot. 100s, 1000s. Triprolidine HCl 1.25 mg., pseudoephedrine HCl 30 mg./5 cc. Syr. Bot. 4 oz., 1 pt.

Use: Common Cold.

ACTIFED-C EXPECTORANT, (Burroughs Wellcome) Codeine phosphate 10 mg., triprolidine HCl 2 mg., pseudoephedrine HCl 30 mg., glyceryl guaiacolate 100 mg./5 cc. Bot. pt.

Use: Treatment of the common cold.

ACTINOMYCIN, (Approved name in Great Britain) Antimicrobial substances W/antitumor activity produced by Streptomyces antibioticus and Streptomyces chrysomallus.

ACTINOMYCIN C. Name previously used for Cactinomycin.

• **ACTINOMYCIN D**. Dactinomycin, N.D. 1966.

See: Cosmegen (Merck Sharp & Dohme)

ACTINOSPECTOCIN. Name previously used for Spectinomycin.

ACTIVATED ATTAPULGITE.

W/Aluminum hydroxide, magnesium carbonate coprecipitate, compressed gel.

See: Hykasil, Cream (Philips Roxane)

W/Polysorbate 80, colloidal sulfur, salicylic acid, propylene glycol.

See: Sebaborb Lotion (Summers Labs.)

• **ACTIVATED 7-DEHYDROCHOLESTEROL**, U.S.P. XVI.

See: Vitamin D-3 (Various Mfr.)

ACTIVATED ERGOSTEROL.

See: Calciferol, U.S.P.

ACTIVOL, (Carbulsulphoil) Liquid soap. Bot. 1 gal, 55 gal. drum.

Use: Surgical Scrub-ups.

• Generic name: approved by F.D.A., or recognized as USAN or by U.S.P., N.F. or N.D.

ACTOL EXPECTORANT. (Beecham-Massengill)

Noscopine 30 mg., glyceryl guaiacolate 200 mg./5 cc. or Tab. Syr. Bot. pt., gal. Tab. Bot. 100s.

Use: Treatment of symptoms associated with common cold.

ACTOSPAR. (Sandoz) Sparteine sulfate.***ACTOQUINOL SODIUM.** 8-Ethoxy-5-quinoline-sulfonic acid sodium salt.

Use: Ultraviolet screen.

ACTYLATE. (Kinney) Ammonium salicylate 80 mg.,

pot. salicylate 80 mg., strontium salicylate 80 mg., pot. para-aminobenzoate 0.32 Gm., ascorbic acid 20 mg./Tab. Bot. 100s.

Use: Rheumatoid arthritis.

ACUDERM. (Nortech) Zinc oxide, starch, talc, castor oil, petrolatum. Oint. Tube 1 oz.**ACUSONE OINTMENT.** (Nortech) Hydrocortisone 0.5%, zinc oxide, starch, talc, castor oil, petrolatum.

Oint.: Tube 1/2 oz., 1 oz.

ACUTRAN. (Mallinckrodt) Amphencloral.**ACUTUSS.** (Philips Roxane) Clophedianol HCl 7.5

mg., diphenylpyraline HCl 0.5 mg., phenylephrine HCl 2.5 mg., glyceryl guaiacolate 25.0 mg., acetaminophen 162.5 mg./Tab. Bot. 100s.

Use: Relief of cold symptoms.

ACUTUSS EXPECTORANT W/CODEINE.

(Philips Roxane) Codeine phosphate 10 mg., phenylpropanolamine HCl 5 mg., phenylephrine HCl 5 mg., chlorpheniramine maleate 2 mg., glyceryl guaiacolate 100 mg., chloroform 13.5 mg. alcohol 5%/Tsp.

(5 cc.) Bot. 8 oz., gal.

Use: For acute cough.

ACUTUSS EXPECTORANT. (Philips Roxane)

Clophedianol HCl 15 mg., diphenylpyraline HCl 1 mg., phenylephrine HCl 5 mg., glyceryl guaiacolate 100 mg., alcohol 6%/5 cc. Bot. 12 fl. oz., gal.

Use: Cough suppressant, expectorant, decongestant.

ACYLANID. (Sandoz) Acetyldigitoxin. Tab. 0.1 mg.

(pink) 100s.; 0.2 mg., (white) 100s.

Use: Digitoxin therapy.

ACYLPYRIN.

See: Aspirin

A AND D CREAM. (Schering) Vits. A & D, lanolin in

cream. Tube 2 oz.

Use: Dry skin, sunburn, irritations.

A AND D HEMORRHOIDAL SUPPOSITORIES.

(Schering) Vitamins A 1500 u. & D 200 u., Peruvian balsam, bismuth subgallate, & zinc oxide in an emollient base of theobroma oil. Box 12s.

Use: Internal hemorrhoids.

A AND D HEMORRHOIDAL SUPPOSITORIES

WITH HC. (Schering) Hydrocortisone acetate 10 mg., Vit. A 1500 u., Vit. D-3 200 u., Peruvian balsam, bismuth subgallate, zinc oxide/Supp. Box 12s.

Use: Relief of internal hemorrhoids.

A AND D OINTMENT. (Schering) Vitamin A & D in

a petrolatum-lanolin base. Tube 1 1/2, 4 oz.; Jar 10 oz., 1 lb., Hospital packs, Tubes 1/2 oz., 1 1/2 oz., 4 oz.

W/Prednisolone 0.5%, Vit. A 1500 u., D 200 u./Gm. Tube 10 Gm., 25 Gm.

A AND D VITAMIN CAPSULES. (Lannett) Vit. A

5000 u., D 400 u./Cap. Bot. 500s, 1000s.

Use: Vit. supplement.

A AND D VITAMIN CREAM. (DePree) Bot. 1 1/2 oz., 4 oz.**A AND D VITAMIN OINTMENT.** (Lannett) In a petrolatum-lanolin base. Jar 1, 5 lb.**A AND D VITAMIN OINTMENT.** (Ulmer) Jar 1 lb.**A-D DRESSING.** (LaCrosse) Vit. A 6000 u., D 600 u.,

allantoin 2%. Tubes 1 oz., Jars 1 lb., 5 lb.

Use: Multi-purpose dressing.

A-D DRESSING W/HEXACHLOROPHENE. (La

Crosse) Vit. A 6000 u., D 600 u., allantoin 2%/oz.

Tube 1 oz. Jar 1 lb., 5 lb.

Use: Antiseptic-antibacterial ointment.

A-D-A. (LaCrosse) Amobarbital 60 mg., dextroamphet-

amine sulfate 15 mg./Cap. Bot. 100s, 1000s.

Use: Obesity.

ADABEE & ADABEE with MINERALS. (Robins)

Vit. A 10,000 u., C 250 mg., B-1 15 mg., B-2 10 mg.,

B-6 5 mg., niacinamide 50 mg./Tab. Bot. 100s, 500s.

Adabee with Minerals. In addition to above formula, iron

15 mg., calcium 103 mg., phosphorus 80 mg./Tab.

Bot. 100s, 500s.

Use: Vitamin supplement.

ADALAN LANATABS. (Lannett) Amobarbital 50

mg., homatropine methylbromide 7.5 mg., metham-

phetamine HCl 10 mg./T.D. Tab. Bot. 100s, 1000s.

ADAMANTANAMINE HCl. (Endo) Amantadine

HCl. Anti-flu tablet. This drug is thought to protect

cells against entry of the flu virus without actually destroying the virus.

See: Symmetrel, Cap., Syr. (Endo)

ADAPIN. (Pennwalt) Doxepin HCl 10 mg., 25 mg./

Cap. Bot. 100s, 50 mg. Bot. 50s. Tab. Bot. 100s.

Use: Antianxiety.

ADAPT. (Burton, Parsons) Povidone, EDTA 0.1%,

thimerosal 0.004%. Wetting sol. Bot. 15 cc.

Use: Contact lens wetting sol.

A.D.C. DROPS. (Parke, Davis) Vit. A 5000 u., D

400 u., C 50 mg., B-6 1 mg./0.6 cc. Dropper bot. 15 cc., 45 cc. w/fluoride 0.5 mg., Dropper bot. 30 cc.

Use: Vitamin deficiencies in infants and children.

A.D.C. DROPS. (Consol. Midland) Vit. A 3000 u.,

D 400 u., C 60 mg./0.6 cc. Bot. 60 cc.

AD-CEBRIN DROPS. (Lilly) Vitamin A 4,000 u.,

D 400 u., C 60 mg./0.6 cc. Dropper Bot. 30 ml.

Use: Dietary supplement.

AD-CEBRIN with FLUORIDE DROPS. (Lilly)

Vitamin A 4,000 u., D 400 u., C 60 mg., fluoride 0.5 mg./0.6 cc. Dropper Bot. 30 ml.

Use: Dietary supplement & prevention of dental caries.

ADECON. (Consol. Midland) Vit. A 5000 u., D 1000

u./Cap. Bot. 1000s.

Use: Supplement.

ADEFLOR B DROPS. (Upjohn) Vitamin A 0.6 mg.,

D 10 mcg., B-1 1 mg., B-2 1 mg., C 50 mg., nicotinamide 10 mg., B-6 1 mg., d-pantothenyl alcohol 3

mg., fluoride (from sodium fluoride) 0.5 mg./0.6 cc. Calibrated drop. Bot. 50 cc.

Use: Aid in the prevention of dental caries w/vitamin supplement.

ADEFLOR CHEWABLE. (Upjohn) Fluoride 1 mg. &

0.5 mg., Vitamin A 1.2 mg., D 10 mcg., C 75 mg., thiamine monitrate (B-1) 2 mg., B-2 2 mg., nico-

tinamide 18 mg., B-6 1 mg., calcium pantothenate 5 mg., B-12 2 mcg./0.5 or 1 mg. Chewable Tab. Bot. 100s, 500s.

Use: Prevention of dental caries.

ADEFLOR DROPS. (Upjohn) Vit. A 0.6 mg., D 10 mcg., C 50 mg., B-6 1 mg., fluoride (as sodium fluoride) 0.5 mg./0.6 cc. 30 or 50 cc. Bot. w/dropper. Use: Prevention of dental caries.

ADEFLOR M TABLETS. (Upjohn) Vit. A 1.8 mg., D 10 mcg., B-1 1.5 mg., B-2 2.5 mg., C 100 mg., nicotinamide 20 mg., Vit. B-6 10 mg., calcium pantothenate 10 mg., Vit. B-12 2 mcg., folic acid 0.1 mg., fluoride (as sodium fluoride) 1 mg., calcium (as calcium carbonate) 250 mg., iron 30 mg./Tab. Bot. 100s, 500s.

Use: Dietary supplement.

ADENIC. (Kenyon) Adenosine-5-monophosphate 25 mg., B-12 75 mcg., niacin 20 mg./cc. Bot. 10 cc.

Use: Anti-inflammatory.

ADENIC FORTE GEL. (Myers-Carter) Adenosine-5-monophosphate 25 mg., B-12 300 mcg., niacin 10 mg., methionine 75 mg./cc. Vial 10 cc.

Use: Anti-inflammatory.

ADENINE. (Various Mfr.) 6-Aminopurine.

ADENO-B. (Blaine) Adenosine 50 mg., Vit. B-12 120 mcg./cc. Vial 10 cc.

ADENO TWELVE GEL INJECTION. (Fellows) Adenosine-5-monophosphate 100 mg., Vit. B-12 100 mcg./cc. Vial 10 cc.

ADENOCREST. (Nutrition) Adenosine-5-monophosphate sod. 25 mg., Vit. B-12 100 mcg., nicotinic acid as sod. salt. 25 mg., gelatin 100 mg./cc. Vials 10 cc.

ADENOLIN FORTE. (Lincoln) Adenosine 5-monophosphate, 25 mg., Vit. B-12 300 mcg., methionine 25 mg., niacin 10 mg./cc. Vial 15 cc.

Use: Arthritis, bursitis, tendinitis & other degenerative diseases.

• **ADENOSINE.** F.D.A. 6-Amino-9-beta-D-Ribofuranosyl-9H-purine.

ADENOSINE B-12. (Vale) Adenosine-5-phosphoric acid 25 mg., cyanocobalamin 100 mcg./ml. Vial 10 ml.

ADENOSINE B-12 GEL. (Medical Chem.) Adenosine-5-monophosphate 100 mg., Vit. B-12 100 mcg./cc. Vial 10 cc.

ADENOSINE-B-12-NICOTINIC. (Med. Chem.) Adenosine-5-monophosphate 25 mg., B-12 75 mg., nicotinic acid 20 mg./cc. Vial 10 cc.

ADENOSINE COMBINATION SOLUTION. (Ries) Adenosine-5-monophosphoric acid 25 mg., Vit. B-12 100 mcg. Vial 10 cc.

ADENOSINE & VIT. B-12 IN GELATINE. (Fellows) Forte: Adenosine 5-monophosphate 50 mg., Vit. B-12 100 mcg./cc. Super: Adenosine 5-monophosphate 100 mg., Vit. B-12 100 mcg./cc.

ADENOSINE W/VITAMIN B-12. (Med. Chem.) Adenosine-5-monophosphoric acid 25 mg., Vit. B-12 60 mcg., water q. s./cc. Vials 10 cc.

• Generic name: approved by F.D.A., or recognized as USAN or by U.S.P., N.F. or N.D.

ADENOSINE-5-MONOPHOSPHATE AS THE SODIUM SALT.

See: Adenic, Prods. (Myers-Carter)

Adenocrest, Amp. (Nutrition)

W/Vit. B-12

See: Adenosine B-12 Gel. (Medical Chem.)

ADENOSINE-5-MONOPHOSPHORIC ACID W/CYANOCOBALAMINE.

See: Adenosine B-12. (Var. Mfr.)

Adenovite-12 (Bowman)

Myo-B, Inj. (Sig)

Vitasine-Gel, Inj. (Burgin Arden)

•ADENOSINE PHOSPHATE. 5-Adenylic acid.

See: My-B-Den (Dome)

ADENOTINATE. (Medics) Adenosine-5-monophosphate 25 mg., Vit. B-12 75 mcg., nicotinic acid 20 mg., benzyl alcohol/cc. Vials 10 cc.

Use: Neuritis, bursitis, etc.

• **ADENOVIRUS VACCINE.** Aqueous preparation of Adenoviruses, types 3, 4, and 7 in approximately equal proportions grown in monkey kidney tissue cultures and inactivated with formaldehyde. Parke, Davis (types 3, 6, 7) Vial 5 cc.

ADEPSINE OIL.

See: Petrolatum Liquid (Various Mfr.)

ADIPHENINE HYDROCHLORIDE. (Various Mfr.) (2-Diethylaminoethyl diphenylacetate hydrochloride.)

Pow. 1 lb. for mfg. use.

See: Trasentine, Tab. (Ciba)

Use: Antispasmodic.

W/Phenobarbital.

See: Trasentine-Phenobarbital, Tab. (Ciba)

ADIRON. (Modern) Organic iron. Tab. Bot. 160s.

ADLEAF. (Panray) Vit. A 5000 u., D 850 u./Cap. Bot. 100s.

ADONIDINE. (City Chem.) Bot. 1 Gm.

Use: Cardiac stimulant.

ADOPHYLLIN ELIXIR. (Bowman) Theophylline, anhydrous 80 mg., alcohol 20%/15 cc. Bot. 3 oz., pt. gal.

Use: Asthma.

ADPHEN TABLETS. (Ferndale) Phendimetrazine tartrate 35 mg./Tab. Bot. 100s.

Use: Anorexiant.

A-D. (Lincoln) dl-Methionine 250 mg., corn starch lubricant 42 mg./Cap. Bot. 100s, 500s.

Use: An. to acid supplement.

ADRENAL CORTEX EXTRACTS. (Various Mfr.) Purified Extract of Adrenal Glands. Influences metabolism of proteins & carbohydrates. Also promotes formation of carbohydrates from proteins & fats. Metabolism of electrolytes (K & Na ions) & water is affected. No international unit standard for the hormones of the adrenal cortex (0.1 mg. 17-hydroxycorticosterone - 1 glycogen unit - 2.5 rat units - 50 dog units) 100 mcg./cc., 10 cc., 200 mcg./cc., 10 cc.

W/Methylandrostenediol, Vit. B-12.

See: Geri-Ace, Inj. (Brown)

ADRENAL CORTEX HORMONES.

See: Gluco-Corticoids

Mineral-Corticoids

ADRENAL CORTEX INJECTION, N.F. XII.

(Aqueous). (Various Mfr.) Mixed natural adrenocortical hormones.

See: Eschatin, Vial. (Parke, Davis)

ADRENAL CORTEX PREPARATIONS.

See: Eschatin, Vial (Parke, Davis)

Lipo-Adrenal Cortex, in oil, Vial (Upjohn)

W/Vitamin C.

See: Cortalex, Tab. (Upjohn)

ADRENALIN(E).

See: Epinephrine. (Various Mfr.)

ADRENALIN IN OIL. (Parke, Davis) Epinephrine 2 mg., Ampuls (1:500) 1 cc. 10s.**ADRENALIN CHLORIDE SOLUTION.** (Parke, Davis) Epinephrine HCl. Principle of the medullary portion of suprarenal glands. Amp. (1:1000) 1 cc., 10s. Bot. (1:1000) 1 oz., (1:100) ¼ oz. Steril-Vials (1:1000) 30 cc.

Use: Sympathomimetic agent.

W/Procaine.

See: Adrocaïne, Sol. (Parke, Davis)

ADRENALINE HYDROCHLORIDE.

See: Epinephrine Hydrochloride. (Various Mfr.)

ADRENALONE. 3',4'-Dihydroxy-2-(methylamino)-acetophenone.

Use: Adrenergic (ophthalmic).

ADRENAMINE.

See: Epinephrine (Various Mfr.)

ADRENERGIC AGENTS.

See: Sympathomimetic agents

ADRENERGIC-BLOCKING AGENTS.

See: Sympatholytic agents

ADRENINE.

See: Epinephrine. (Various Mfr.)

ADRENOCHROME.

See: Carbazochrome Salicylate.

ADRENOCHROMAZONE.

See: Carbazochrome Salicylate.

ADRENOCHROME MONOSEMICARBAZONE.

See: Carbazochrome Salicylate

ADRENOCORTICOTROPIC HORMONE. ACTH

acts by stimulating the endogenous production of cortisone and the final result is similar whichever substance is administered. The nature of action of each is unknown. Both are necessary in the metabolism of carbohydrates, protein & fat & exert profound effects on neuromuscular metabolism.

See: ACTH gel, Inj. (Bowman)

Corticotropin, U.S.P.

Cortigel (Savage)

Cortropic, Gel, Inj. (Tutag)

ADRENO-MIST. (SmithKline) Epinephrine HCl

1:100, chlorobutanol base in glycerin, Sod. bisulfite. Sol. Bot. ½, 1 oz.

Use: Inhalation, paroxysms & symptoms of asthma.

ADRENOSEM SALICYLATE. (Beecham-Massengill)

Carbazochrome salicylate, benzyl alcohol 2% by volume. Amp. (5 mg./cc.) 1 cc., 5s, 100s; (10 mg./2 cc.) 2 cc., 5s, 100s. Syr. (2.5 mg./5 cc.) Bot. 4 oz., Tab. (2.5 mg.) Bot. 50s, 1000s.

Use: Orally, I.M., to increase capillary resistance and suppress bleeding from capillary bed.

ADRENUCLEO. (Enzyme Process) Vit. C 250 mg., d-cal. pantothenate 12.5 mg., bioflavonoids 62.5 mg./Tab. Bot. 100s, 250s.

Use: Vitamin-C supplement therapy.

ADRIAMYCIN. Under study.

Use: Treatment of hemoblastoses.

ADRICOT. (Kay) Cortisone acetate 50 mg., polysorbate-80 4 mg., sodium chloride 9 mg., sodium carboxymethylcellulose 5 mg./cc. Vial 10 cc.**ADROCAINE SOLUTION.** (Parke, Davis) Adrenalin 1:100,000, procaine HCl 1½%, chloretone 0.5%. Vial 50 cc. 10s.

Use: Local anesthesia.

ADRON. (Sheryl) Iron (elemental) 100 mg., ascorbic acid 200 mg./Tab. 60s.

Use: Treatment of simple iron deficiency anemia.

ADROYD. N.D. 1967. (Parke, Davis) Oxymetholone 5 mg. & 10 mg./Tab. Bot. 30s.

Use: Anabolic steroid.

ADSORBONAC OPHTH. SOLUTION. (Burton-Parsons) Sodium chloride 2% or 5% in vehicle of polyvinylpyrrolidone and BP water soluble polymers. Dropper Vial 15 ml.

Use: Reduction of corneal edema and clearing of the cornea for diagnostic examination.

ADSOBTEAR. (Burton, Parsons) Hydroxyethylcellulose 0.44%, w/polyvinylpyrrolidone and water-soluble polymers. Bot. dropper 15 ml.

Use: Artificial tears.

ADSORMONE. (Herring) Complete set of reagents required. Pow. 100 Gm.

Use: Pregnancy test.

ADVANCE. (Ross) Milk substitute for infants. Cans 13 oz.

Use: Pediatric supplement.

ADVENT. (Winston) Vit. A 4000 u., D 400 u., C 200 mg., B-1 2 mg., B-2 1.5 mg., B-6 0.5 mg., B-12 1 mcg., niacinamide 10 mg., Ca. lactate 400 mg., ferrous fumarate 200 mg., I 0.01 mg., Pot 0.835 mg., Cu 0.15 mg., Mn 0.05 mg., Zn 0.085 mg./Tab. Bot. 100s, 1000s.

Use: Pregnancy and lactation.

A.E.A. (Stan Lab.) Aminophylline 120 mg., ephedrine HCl 25 mg., amobarbital 25 mg./Tab. Bot. 100s, 1000s.**A.E.P.** (Reaco Products) Aminophylline 1½ gr., ephedrine sulfate ¼ gr., phenobarbital ¼ gr./Tab. Bot. 100s, 1000s.

Use: Symptomatic relief of asthma.

AEROAID. (Aeroceuticals) Thimerosal 0.033%, alcohol 26.6%. Spray Bot. Applications, 1 oz., 4 oz., 6 oz.

Use: Metered first aid antiseptic spray.

AEROCAINE. (Aeroceuticals) Benzyl alcohol 22.7%, benzocaine 13.6%, benzethonium chloride 0.5%. Spray Bot. ¾ oz., 5 oz.

Use: Topical anesthetic and antiseptic.

AEROCELL. (Aeroceuticals), Exfoliative cytology fixative spray. Bot. 4 oz.**AERODINE.** (Aeroceuticals) Povidone-iodine aerosol N.F. Bot. 4.5 oz.

Use: Broad range antiseptic germicide.

AEROFOAM CLEANER. (Ulmer) o-Phenylphenol 0.10%, 4-chloro-2-cyclopentylphenol 0.08%, diethanolamine-lauric acid amide 0.20%, triethanolamine dodecyl benzene sulfonate 0.30%. Bot. 20 oz. Use: Germicidal foam.

AEROFREEZE. (Aeroceuticals) Trichloromonofluoromethane & dichlorodifluoromethane. Aerosol containers 8 oz. (12s). Use: Topical anesthetic.

AEROHALOR. (Abbott) Aerohalor Cartridges to introduce dry solid aerosols into the upper respiratory tract and lungs. Individual boxes.

AEROLATE SR. & JR. (Fleming) Caps. Theophylline 1 gr., 4 gr. for Sr. and 2 gr. for Jr./Cap. Bot. 100s, 1000s. Elix. 160 mg./15 ml., Bot. pt., gal. Use: Bronchodilation.

AEROLONE COMPOUND. (Lilly) Clopane (clo-pentamine) HCl 0.5 Gm., isoproterenol HCl 0.25 Gm., in propylene glycol, ascorbic acid/100 cc. Sol. Bot. 1 fl. oz. Use: Asthma, status asthmaticus.

AEROMINT. (Aeroceuticals) 200 metered sprays. Aerosol 0.5 oz. Use: Breath freshener.

AEROPURE. (Aeroceuticals) Isopropanol 7.8%, triethylene glycol 3.9%, essential oils 3%, methyl-dodecyl benzyl trimethyl ammonium chloride 0.12%, methyl-dodecylxylene bis (trimethyl ammonium chloride) 0.03%, inert ingredients, 85.15%. Bot. 1 1/4 oz., 6 oz. Use: Concentrated air sanitizer and deodorizer.

AEROSEB-D. (G.S. Herbert Labs) Dexamethasone 10 mg./90 Gm. aerosol. Use: Dermatoses.

AEROSEB-HC. (G.S. Herbert Labs) Hydrocortisone 0.53 mg./1 sec. of spray. Aerosol pkg. 225 mg./90 Gm. Use: Seborrheic dermatitis.

AEROSEPT. (Dalin) Lidocaine, cetyltrimethyl-ammonium bromide, hexachlorophene. Aerosol Pkg. 6 oz. Use: Antiseptic, anesthetic.

AEROSIL. (Aeroceuticals) Dimethylpolysiloxane, trichloromonofluoromethane, dichlorodifluoromethane. Bot. 12 oz.

Use: Silicone lubricant and protective spray.

AEROSOL MERTHIOLATE. (Aeroceuticals) Merthiolate (thimerosal N.F., Lilly) 0.033%, alcohol 22%. Spray Bot. 4 oz. Use: First aid antiseptic spray.

AEROSOL OT.

See: Dioctyl Sodium Sulfosuccinate, U.S.P.

AEROSPORIN OTIC SOLUTION. (Burroughs Wellcome) Polymyxin B sulfate 0.1% (10,000 units/cc.) in acidified propylene glycol. Bot. w/dropper 10 cc.

Use: Topically, infections of the ear.

AEROSPORIN SULFATE. (Burroughs Wellcome) Polymyxin B sulfate 50 mg., sterile pow. (500,000 u.)

Vial (20 cc.): pow., non-sterile, for Rx compounding (100,000,000 u.). Otic Sol., sterile. Bot. 10 cc.

See: Polymyxin B sulfate, U.S.P.

Use: Orally for bacillary dysentery, parenteral form for hospital use only.

W/Bacitracin.

See: Polysporin, Oint. (Burroughs Wellcome)

W/Bacitracin & neomycin sulfate.

See: Neosporin, oint. & Ophth. Oint. (Burroughs Wellcome)

W/Bacitracin, neomycin, hydrocortisone.

See: Cortisporin, Prep. (Burroughs Wellcome)

AEROTHERM. (Aeroceuticals) Benzyl alcohol 22.7%, benzocaine 13.6%. Spray Bot. 5 oz. Use: First aid spray for burns.

AEROZOIN. (Aeroceuticals) Comp. tr. of benzoin 22%, hexachlorophene 0.1%, isopropyl alcohol 33%. Spray Bot. 6 oz., 15 oz. Use: Skin conditioner for tape and cast dressing.

AETHYLIS CHLORIDUM.

See: Ethyl Chloride (Various Mfr.)

A.F.F. (Invenex) Ferrous Fumarate 195 mg., ascorbic acid 100 mg./Tab. Bot. 100s.

Use: Treatment of anemias.

A-FIL. (Texas Pharm.) Menthyl anthranilate 5%, titanium dioxide 5% in vanishing cream. Tubes 1.5 oz. neutral or dark. Use: Sun screen.

AFRIKOL. (Citroleum) Suntan lotion w/no cocoa butter or lanolin. Bot. 4 oz.

AFRIN. (Schering) N.D. 1967. Oxymetazoline 0.5 mg., glycine 3.8 mg., sorbitol 40 mg., phenylmercuric acetate 0.02 mg., benzalkonium Cl 0.2 mg./cc. nasal spray or soln. Spray: Bot., Plastic, 15 cc. Bot. Soln: 30 cc. dpr. bot. Industrial pkg. 3 cc. Bot.

Use: Nasal decongestant.

•AGAR, U.S.P. XVIII. (Various Mfr.)

Agar-agar, Bengal Gelatin, Ceylon.

Use: Bulk laxative.

W/Mineral oil.

See: Agoral, Emulsion (Warner-Chilcott) Petrogalar (Wyeth)

AGGREGATED ALBUMIN.

See: Albumotope-LS (Squibb)

AGOBYL. (C.M.C.) Magnesium sulfate 37.0 Gm., succinic acid 2.0 Gm., peptone (bacteriological) 9.0 Gm., citric acid 4.0 Gm., sod. bicarbonate 6.0 Gm., sucrose (mint flavored) 42.0 Gm./100 Gm. Pkg. 125 Gm.

AGOFOLLIN.

See: Estradiol. (Various Mfr.)

AGORAL. W/Phenolphthalein (Warner Chilcott)

Phenolphthalein & mineral oil in an emulsion containing agar-gel, tragacanth, egg-albumen, acacia, glycerin and water. Marshmallow flavor. Bot. 8, 16 fl. oz., Raspberry flavored Bot. pint.

Use: Laxative.

AGORAL, PLAIN. (Warner Chilcott) Emulsion of mineral oil and an agar-gel. Bot. 16 oz.

Use: Laxative.

•Generic name: approved by F.D.A., or recognized as USAN or by U.S.P., N.F. or N.D.

A/G-PRO. (Miller Pharmacal) Protein hydrolysate 8.3 gr. w/essential and nonessential amino acids 45%, l-lysine 50 mg., methionine 12.5 mg. and Fe, Cu, I, Mn, K, Zn, Mg./6 Tab. Bot. 180s.
Use: Tissue repair.

A.G. SULFA. (Mericon) Sulfisoxazole 500 mg., phenylazopyridine HCl 50 mg./Tab. Bot. 100s, 1000s.

AGURIN.

See: Theobromine Sod. Acetate (Various Mfr.)

AHAN. (Eric, Kirk and Gary) Vitamin A acetate 25,000 u./Cap. Bot. 100s. 50,000 or 100,000 u./Cap. Bot. 100s.

AIDANT. (Noyes) Acetophenetidin 2 gr., aspirin 3 gr., caffeine $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s.
Also available w/Dover's pow., or gelsemium.
Use: Analgesic.

AIRET. (Baylor Labs.) Dihydroxypropyl theophylline 100 mg., alcohol 10%/15 ml. Bot. 1 pt.
Use: Treatment of bronchial asthma.

AIRET G.G. (Baylor Labs.) Dihydroxypropyl theophylline 200 mg., glyceryl guaiacolate 200 mg./Cap. Bot. 100s. Elixir: dihydroxypropyl theophylline 100 mg., glyceryl guaiacolate 100 mg., alcohol 17%/15 ml. Bot. 1 pt.
Use: Treatment of bronchial asthma.

AIRET L.A. (Baylor Labs.) Dihydroxypropyl theophylline 400 mg./Tab. Bot. 100s.
Use: Treatment of bronchial asthma.

AIRET R. (Baylor Labs.) Dihydroxypropyl theophylline 200 mg., ephedrine HCl 24 mg., butabarbital 24 mg./Tab. Bot. 100s.
Use: Treatment of bronchial asthma w/sedative.

AIRET Y. (Baylor Labs) Theophylline 200 mg., ephedrine HCl 15 mg., butabarbital 15 mg./Tab. Bot. 100s. Theophylline 100 mg., ephedrine HCl 15 mg., butabarbital 15 mg., alcohol 10%/15 cc. Bot. pt.
Use: Relief of bronchopulmonary insufficiency.

AKINETON. N.D. 1967. (Knoll) Biperiden HCl 2 mg./Tab. Bot. 100s, 1000s.
Use: Anticholinergic and myospasmodic for Parkinson's disease and drug-induced parkinsonism.

AKINETON LACTATE. N.D. 1967. (Knoll) Biperiden lactate 5 mg. in aqueous 1.4% sodium lactate solution/cc. Amp 1 cc. Box 10s.
Use: Relief of Parkinson-like dystonic crises.

AKNE DRYING LOTION. (Alto) Zinc oxide 12%, urea 10%, sulfur 6%, salicylic acid 2%, benzalkonium chloride 0.2%, isopropyl alcohol 70%, in a base containing menthol, silicon dioxide, iron oxide and perfume. Bot. 2 $\frac{1}{4}$ oz.
Use: Acne vulgaris, acne rosacea, seborrhea.

AKNE ORAL KAPSULETS. (Alto) Theionized sulfur 325 mg./Capsule-shaped tablet. Bot. 20s.
Use: Oral sulfur therapy.

AKNE SCRUB. (Alto) Povidone iodine with polyethylene granules. Bot. $\frac{3}{4}$ oz.
Use: Acne cleanser, antiseptic, microbicidal

AKRINOL CREAM. N.D. 1967. (Schering) Acrisorcin, the hexylresorcinol salt of 9-amino-acridine, 2 mg./Gm. Tube 50 Gm.
Use: Topical use tinea versicolor.

ALA-CORT. (Del-Ray Lab.) Hydrocortisone 1%.

Cream: Tube 1 oz.

Use: Glucocorticoid.

ALADRINE. (Merit) Ephedrine sulfate $\frac{1}{8}$ gr., secobarbital sod. $\frac{1}{4}$ gr./Tab. Bot. 100s, 1000s. Ephedrine sulfate $\frac{1}{8}$ gr., secobarbital sodium $\frac{1}{4}$ gr./5 cc. Bot. pt., gal.

Use: Antiasthmatic.

ALAMAG. (Barre Drug) Magnesium-aluminum hydroxide gel. Susp. Bot. pt.

W/Belladonna alkaloid, Susp. Bot. 8 oz.

ALAMECIN. Antibiotic produced by *Trichoderma viride* Pers. ex fries. (Unspecific action).

ALAMINE-C. (North Amer. Pharm.) Codeine phosphate 5 mg., phenylephrine HCl 10 mg., chlorpheniramine maleate 2 mg., chloroform 13.5 mg., 1-menthol 1 mg., alcohol 5%/5 cc. Bot. pt.

ALAMINE EXPECTORANT. (North Amer. Pharm.) Codeine phosphate 5 mg., glyceryl guaiacolate 100 mg., phenylephrine HCl 10 mg., chlorpheniramine maleate 2 mg., chloroform 13.5 mg., 1-menthol 1 mg., alcohol 5%/5 cc. Bot. pt.

ALAMINE LIQUID. (North Amer. Pharm.) Phenylephrine HCl 5 mg., chlorpheniramine maleate 1 mg., chloroform 13.5 mg., 1-menthol 1 mg., alcohol 5%/5 cc. Bot. pt.

ALAMINE TABLETS. (North Amer. Pharm.) Propenpyridamine maleate 12.5 mg., phenylephrine HCl 5 mg./Tab. Bot. 100s.

ALASED. (Norgine) Homatropine methylbromide $\frac{1}{16}$ gr., dried aluminum hydroxide $7\frac{1}{2}$ gr., phenobarbital $\frac{1}{4}$ gr., magnesium trisilicate $2\frac{1}{2}$ gr./Tab. Bot. 100s, 250s.
Use: Antispasmodic with antacid.

AL-AY. (Bowman) Phenylephrine HCl 5 mg., chlorpheniramine maleate 2 mg., acetylsalicylic acid 162 mg., caffeine 15 mg., aminoacetic acid 162 mg./Tab. Bot. 100s, 500s, 1000s.
Use: Nasal decongestant.

ALAZANINE TRICHLORPHATE. 3-Ethyl-2-[3-(3-ethyl-2-benzothiazolylidene) propenyl]-benzothiazolium 2,4,5-trichlorophenoxide salt with two formula weights of 2,4,5(trichlorophenol).
Use: Anthelmintic.

ALBALON. (Allergan) Naphazoline HCl 0.1%, polyvinyl alcohol 1.4%, benzalkonium Cl, disodium edetate, citric acid, sodium citrate, sodium Cl, sodium hydroxide. Bot. 15 cc. Unit dose 1 cc.
Use: Vasoconstrictor.

ALBAMYCIN. N.D. 1966. (Upjohn) Novobiocin sod. 250 mg./Cap. Bot. 16s, 100s.
Use: Wide-range antibiotic therapy.

ALBAMYCIN CALCIUM. N.D. 1966.
See: Albamycin, Syr. (Upjohn)

ALBAMYCIN MIX-O-VIAL. (Upjohn) In upper compartment, N,N-Dimethyl-acetamide 0.47 cc. in sterile water for injection; lower compartment, novobiocin sodium (Albamycin), 500 mg.; nicotinamide 175 mg./Vial 5 cc. Box 1s, 25s.

ALBAMYCIN SYR. (Upjohn) Novobiocin cal. 125 mg./5 cc. Bot. 2 oz.