


Cox Rings



**IVAN ARZHANTSEV
ULRICH DERENTHAL
JÜRGEN HAUSEN
ANTONIO LAFACE**



Cox Rings

IVAN ARZHANTSEV
Moscow State University

ULRICH DERENTHAL
Leibniz Universität Hannover

JÜRGEN HAUSEN
Eberhard Karls Universität Tübingen

ANTONIO LAFACE
Universidad de Concepción



CAMBRIDGE
UNIVERSITY PRESS

CAMBRIDGE
UNIVERSITY PRESS

32 Avenue of the Americas, New York, NY 10013-2473, USA

Cambridge University Press is part of the University of Cambridge.

It furthers the University's mission by disseminating knowledge in the pursuit of education, learning, and research at the highest international levels of excellence.

www.cambridge.org

Information on this title: www.cambridge.org/9781107024625

© Ivan Arzhantsev, Ulrich Derenthal, Jürgen Hausen, and Antonio Laface 2015

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2015

Printed in the United States of America

A catalog record for this publication is available from the British Library.

Library of Congress Cataloging in Publication Data

Arzhantsev, I. V. (Ivan Vladimirovich), 1972–

[Kol'tsa Koksa. English]

Cox rings / Ivan Arzhantsev, Department of Algebra, Faculty of Mechanics and Mathematics, Moscow [and three others].

pages cm. – (Cambridge studies in advanced mathematics)

Includes bibliographical references and index.

ISBN 978-1-107-02462-5 (hardback)

1. Algebraic varieties. 2. Rings (Algebra) I. Title.

QA564.A7913 2015

516.3'53–dc23 2014005540

ISBN 978-1-107-02462-5 Hardback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party Internet Web sites referred to in this publication and does not guarantee that any content on such Web sites is, or will remain, accurate or appropriate.

Editorial Board

B. BOLLOBÁS, W. FULTON, A. KATOK, F. KIRWAN,
P. SARNAK, B. SIMON, B. TOTARO

COX RINGS

Cox rings are significant global invariants of algebraic varieties, naturally generalizing homogeneous coordinate rings of projective spaces. This book provides a largely self-contained introduction to Cox rings, with a particular focus on concrete aspects of the theory. Besides the rigorous presentation of the basic concepts, other central topics include the case of finitely generated Cox rings and its relation to toric geometry; various classes of varieties with group actions; the surface case; and applications in arithmetic problems, in particular Manin's conjecture. The introductory chapters require only basic knowledge of algebraic geometry. The more advanced chapters also touch on algebraic groups, surface theory, and arithmetic geometry.

Each chapter ends with exercises and problems. These comprise mini-tutorials and examples complementing the text, guided exercises for topics not discussed in the text, and, finally, several open problems of varying difficulty.

Ivan Arzhantsev received his doctoral degree in 1998 from Lomonosov Moscow State University and is a professor in its department of higher algebra. His research areas are algebraic geometry, algebraic groups, and invariant theory.

Ulrich Derenthal received his doctoral degree in 2006 from Universität Göttingen. He is a professor of mathematics at Leibniz Universität Hannover. His research interests include arithmetic geometry and number theory.

Jürgen Hausen received his doctoral degree in 1995 from Universität Konstanz. He is a professor of mathematics at Eberhard Karls Universität Tübingen. His field of research is algebraic geometry, in particular algebraic transformation groups, torus actions, geometric invariant theory, and combinatorial methods.

Antonio Laface received his doctoral degree in 2000 from Università degli Studi di Milano. He is an associate professor of mathematics at Universidad de Concepción. His field of research is algebraic geometry, more precisely linear systems, algebraic surfaces, and their Cox rings.

CAMBRIDGE STUDIES IN ADVANCED MATHEMATICS

EDITORIAL BOARD

B. Bollobás, W. Fulton, A. Katok, F. Kirwan, P. Samak, B. Simon, B. Totaro

All the titles listed below can be obtained from good booksellers or from Cambridge University Press. For a complete series listing visit: www.cambridge.org/mathematics.

Already published

- 104 A. Ambrosetti & A. Malchiodi *Nonlinear analysis and semilinear elliptic problems*
- 105 T. Tao & V. H. Vu *Additive combinatorics*
- 106 E. B. Davies *Linear operators and their spectra*
- 107 K. Kodaira *Complex analysis*
- 108 T. Ceccherini-Silberstein, F. Scarabotti, & F. Tolli *Harmonic analysis on finite groups*
- 109 H. Geiges *An introduction to contact topology*
- 110 J. Faraut *Analysis on Lie groups: An introduction*
- 111 E. Park *Complex topological K-theory*
- 112 D. W. Stroock *Partial differential equations for probabilists*
- 113 A. Kirillov, Jr *An introduction to Lie groups and Lie algebras*
- 114 F. Gesztesy et al. *Soliton equations and their algebro-geometric solutions, II*
- 115 E. de Faria & W. de Melo *Mathematical tools for one-dimensional dynamics*
- 116 D. Applebaum *Lévy processes and stochastic calculus (2nd Edition)*
- 117 T. Szamuel *Galois groups and fundamental groups*
- 118 G. W. Anderson, A. Guionnet, & O. Zeitouni *An introduction to random matrices*
- 119 C. Perez-Garcia & W. H. Schikhof *Locally convex spaces over non-Archimedean valued fields*
- 120 P. K. Friz & N. B. Victoir *Multidimensional stochastic processes as rough paths*
- 121 T. Ceccherini-Silberstein, F. Scarabotti, & F. Tolli *Representation theory of the symmetric groups*
- 122 S. Kalikow & R. McCutcheon *An outline of ergodic theory*
- 123 G. F. Lawler & V. Limic *Random walk: A modern introduction*
- 124 K. Lux & H. Pahlings *Representations of groups*
- 125 K. S. Kedlaya *p -adic differential equations*
- 126 R. Beals & R. Wong *Special functions*
- 127 E. de Faria & W. de Melo *Mathematical aspects of quantum field theory*
- 128 A. Terras *Zeta functions of graphs*
- 129 D. Goldfeld & J. Hundley *Automorphic representations and L-functions for the general linear group, I*
- 130 D. Goldfeld & J. Hundley *Automorphic representations and L-functions for the general linear group, II*
- 131 D. A. Craven *The theory of fusion systems*
- 132 J. Väänänen *Models and games*
- 133 G. Malle & D. Testerman *Linear algebraic groups and finite groups of Lie type*
- 134 P. Li *Geometric analysis*
- 135 F. Maggi *Sets of finite perimeter and geometric variational problems*
- 136 M. Brodmann & R. Y. Sharp *Local cohomology (2nd Edition)*
- 137 C. Muscalu & W. Schlag *Classical and multilinear harmonic analysis, I*
- 138 C. Muscalu & W. Schlag *Classical and multilinear harmonic analysis, II*
- 139 B. Helffer *Spectral theory and its applications*
- 140 R. Pemantle & M. C. Wilson *Analytic combinatorics in several variables*
- 141 B. Branner & N. Fagella *Quasiconformal surgery in holomorphic dynamics*
- 142 R. M. Dudley *Uniform central limit theorems (2nd Edition)*
- 143 T. Leinster *Basic category theory*

Contents

Introduction	<i>page</i> 1
1 Basic concepts	6
1.1 Graded algebras	7
1.1.1 Monoid graded algebras	7
1.1.2 Veronese subalgebras	10
1.2 Gradings and quasitorus actions	13
1.2.1 Quasitori	13
1.2.2 Affine quasitorus actions	16
1.2.3 Good quotients	20
1.3 Divisorial algebras	24
1.3.1 Sheaves of divisorial algebras	24
1.3.2 The relative spectrum	27
1.3.3 Unique factorization in the global ring	30
1.3.4 Geometry of the relative spectrum	32
1.4 Cox sheaves and Cox rings	35
1.4.1 Free divisor class group	35
1.4.2 Torsion in the divisor class group	39
1.4.3 Well-definedness	41
1.4.4 Examples	44
1.5 Algebraic properties of the Cox ring	47
1.5.1 Integrity and normality	47
1.5.2 Localization and units	50
1.5.3 Divisibility properties	52
1.6 Geometric realization of the Cox sheaf	56
1.6.1 Characteristic spaces	56
1.6.2 Divisor classes and isotropy groups	59
1.6.3 Total coordinate space and irrelevant ideal	63
1.6.4 Characteristic spaces via GIT	65
Exercises to Chapter 1	69

2 Toric varieties and Gale duality	75
2.1 Toric varieties	75
2.1.1 Toric varieties and fans	75
2.1.2 Some toric geometry	78
2.1.3 The Cox ring of a toric variety	82
2.1.4 Geometry of Cox's construction	86
2.2 Linear Gale duality	89
2.2.1 Fans and bunches of cones	89
2.2.2 The GKZ decomposition	93
2.2.3 Proof of Theorem 2.2.1.14	96
2.2.4 Proof of Theorems 2.2.2.2, 2.2.2.3, and 2.2.2.6	100
2.3 Good toric quotients	103
2.3.1 Characterization of good toric quotients	103
2.3.2 Combinatorics of good toric quotients	107
2.4 Toric varieties and bunches of cones	110
2.4.1 Toric varieties and lattice bunches	110
2.4.2 Toric geometry via bunches	113
Exercises to Chapter 2	117
3 Cox rings and combinatorics	123
3.1 GIT for affine quasitorus actions	124
3.1.1 Orbit cones	124
3.1.2 Semistable quotients	128
3.1.3 A_2 -quotients	133
3.1.4 Quotients of H -factorial affine varieties	137
3.2 Bunched rings	142
3.2.1 Bunched rings and their varieties	142
3.2.2 Proofs to Section 3.2.1	147
3.2.3 Example: Flag varieties	151
3.2.4 Example: Quotients of quadrics	155
3.2.5 The canonical toric embedding	161
3.3 Geometry via defining data	167
3.3.1 Stratification and local properties	167
3.3.2 Base loci and cones of divisors	172
3.3.3 Complete intersections	178
3.3.4 Mori dream spaces	181
3.4 Varieties with a torus action of complexity 1	186
3.4.1 Detecting factorial gradings	186
3.4.2 Factorially graded rings of complexity 1	191
3.4.3 T -varieties of complexity 1 via bunched rings	197
3.4.4 Geometry of T -varieties of complexity 1	202
Exercises to Chapter 3	207

4 Selected topics	215
4.1 Toric ambient modifications	216
4.1.1 The Cox ring of an embedded variety	216
4.1.2 Algebraic modification	220
4.1.3 Toric ambient modifications	226
4.1.4 Computing examples	232
4.2 Lifting automorphisms	236
4.2.1 Quotient presentations	236
4.2.2 Linearization of line bundles	241
4.2.3 Lifting group actions	246
4.2.4 Automorphisms of Mori dream spaces	251
4.3 Finite generation	257
4.3.1 General criteria	257
4.3.2 Finite generation via multiplication map	262
4.3.3 Finite generation after Hu and Keel	268
4.3.4 Cox–Nagata rings	274
4.4 Varieties with torus action	279
4.4.1 The Cox ring of a variety with torus action	279
4.4.2 H -factorial quasiaffine varieties	284
4.4.3 Proof of Theorems 4.4.1.2, 4.4.1.3, and 4.4.1.6	291
4.5 Almost homogeneous varieties	297
4.5.1 Homogeneous spaces	297
4.5.2 Small embeddings	303
4.5.3 Examples of small embeddings	308
4.5.4 Spherical varieties	314
4.5.5 Wonderful varieties and algebraic monoids	319
Exercises to Chapter 4	326
5 Surfaces	334
5.1 Mori dream surfaces	334
5.1.1 Basic surface geometry	334
5.1.2 Nef and semiample cones	339
5.1.3 Rational surfaces	345
5.1.4 Extremal rational elliptic surfaces	350
5.1.5 K3 surfaces	356
5.1.6 Enriques surfaces	361
5.2 Smooth del Pezzo surfaces	366
5.2.1 Preliminaries	366
5.2.2 Generators of the Cox ring	369
5.2.3 The ideal of relations	374
5.2.4 Del Pezzo surfaces and flag varieties	381

5.3	K3 surfaces	385
5.3.1	Abelian coverings	385
5.3.2	Picard numbers 1 and 2	390
5.3.3	Nonsymplectic involutions	397
5.3.4	Cox rings of K3 surfaces	400
5.4	Rational \mathbb{K}^*-surfaces	405
5.4.1	Defining data and their surfaces	405
5.4.2	Intersection numbers	411
5.4.3	Resolution of singularities	417
5.4.4	Gorenstein log del Pezzo \mathbb{K}^* -surfaces	422
	Exercises to Chapter 5	431
6	Arithmetic applications	437
6.1	Universal torsors and Cox rings	437
6.1.1	Quasitori and principal homogeneous spaces	437
6.1.2	Universal torsors	442
6.1.3	Cox rings and characteristic spaces	449
6.2	Existence of rational points	454
6.2.1	The Hasse principle and weak approximation	454
6.2.2	Brauer–Manin obstructions	457
6.2.3	Descent and universal torsors	459
6.2.4	Results	461
6.3	Distribution of rational points	464
6.3.1	Heights and Manin’s conjecture	464
6.3.2	Parameterization by universal torsors and Cox rings	467
6.4	Toward Manin’s conjecture for del Pezzo surfaces	469
6.4.1	Classification and results	469
6.4.2	Strategy	474
6.4.3	Parameterization via Cox rings	475
6.4.4	Counting integral points on universal torsors	478
6.4.5	Interpretation of the integral	485
6.5	Manin’s conjecture for a singular cubic surface	489
6.5.1	Statement of the result	489
6.5.2	Geometry and Cox ring	490
6.5.3	Parameterization via Cox rings	491
6.5.4	Counting integral points on universal torsors	493
6.5.5	Interpretation of the integral	497
	Exercises to Chapter 6	498
	<i>Bibliography</i>	501
	<i>Index</i>	517

Introduction

The basic principles regarding Cox rings become visible already in the classical example of the projective space \mathbb{P}^n over a field \mathbb{K} , which we assume to be algebraically closed and of characteristic zero. The elements of \mathbb{P}^n are the lines $\ell \subseteq \mathbb{K}^{n+1}$ through the origin $0 \in \mathbb{K}^{n+1}$. Such a line ℓ is concretely specified by its *homogeneous coordinates* $[z_0, \dots, z_n]$, where (z_0, \dots, z_n) is *any* point on ℓ , different from the origin. Hence, this description comes with an intrinsic ambiguity. More formally speaking, that means that we should regard the projective space as a quotient by a group action

$$\begin{array}{ccc} \mathbb{K}^{n+1} \setminus \{0\} & \subseteq & \mathbb{K}^{n+1} \\ z \mapsto [z] & \downarrow & / \mathbb{K}^* \\ & \mathbb{P}^n & \end{array}$$

where \mathbb{K}^* acts on \mathbb{K}^{n+1} via scalar multiplication. This presentation of the projective space \mathbb{P}^n as the quotient of its *characteristic space* $\mathbb{K}^{n+1} \setminus \{0\}$ by the action of the *characteristic torus* \mathbb{K}^* is the geometric way of thinking of Cox rings. In algebraic terms, the action of \mathbb{K}^* on \mathbb{K}^{n+1} is encoded by the associated decomposition of the polynomial ring into homogeneous parts

$$\mathbb{K}[T_0, \dots, T_n] = \bigoplus_{k \geq 0} \mathbb{K}[T_0, \dots, T_n]_k,$$

where $\mathbb{K}[T_0, \dots, T_n]_k$ is the vector space of homogeneous polynomials f of degree k , which means that $f(tz) = t^k f(z)$ holds for all t and z . The polynomial ring together with this classical grading is the *Cox ring* of the projective space. Note that to construct \mathbb{P}^n as a \mathbb{K}^* -quotient, we have to remove the origin, which is the vanishing locus of the *irrelevant ideal*, from the *total coordinate space* \mathbb{K}^{n+1} . The Cox ring can be seen in terms of algebraic geometry intrinsically

from the projective space as

$$\mathbb{K}[T_0, \dots, T_n] = \bigoplus_{k \geq 0} \mathbb{K}[T_0, \dots, T_n]_k \cong \bigoplus_{k \in \mathbb{Z}} \Gamma(\mathbb{P}^n, \mathcal{O}(kD)),$$

where the class of the hyperplane $D := V(T_0) \subseteq \mathbb{P}^n$ freely generates the *divisor class group* $\text{Cl}(\mathbb{P}^n) \cong \mathbb{Z}$ and the sections $1, T_1/T_0, \dots, T_n/T_0$ of degree 1 on the right-hand side correspond to the homogeneous generators T_0, \dots, T_n on the left-hand side. The brief discussion of this simple example shows us that Cox rings are located in the intersection of three fields: graded algebras, group actions and quotients, and divisors and their section rings.

Let us also take a brief look at the arithmetic aspects. The rational points $[z_0, \dots, z_n]$ in the projective space \mathbb{P}^n over \mathbb{Q} are parameterized uniquely up to sign by primitive vectors, that is, tuples of coprime integers z_0, \dots, z_n . This description of rational points is related to Cox rings via the diagram

$$\begin{array}{ccc} \mathbb{Z}_{\text{prim}}^{n+1} \setminus \{0\} & \subseteq & \mathbb{Z}^{n+1} \\ z \mapsto [z] \downarrow / \mathbb{Z}^* & & \\ \mathbb{P}^n(\mathbb{Q}) & & \end{array}$$

A typical problem is to estimate the number of rational points with bounded height $H([z_0, \dots, z_n]) := \max\{|z_0|, \dots, |z_n|\}$, which in our example essentially amounts to estimating the number of lattice points in an $(n+1)$ -dimensional box. This is an instance of Manin's conjecture on the number of rational points of bounded height on Fano varieties.

The current interest in Cox rings has several sources. A first one dates back to the 1970s when Colliot-Thélène and Sansuc [96, 98] introduced universal torsors as a tool in arithmetic geometry in particular to investigate the existence of rational points on varieties. In the last few years, Salberger's approach [263] to study the distribution of rational points via universal torsors and their explicit representations in terms of Cox rings caused a considerable surge of research. Another source is the occurrence of characteristic spaces and Cox rings in toric geometry in the mid-1990s in work of Audin [23], Cox [104] and others, which had a tremendous impact on this field. Five years later, Hu and Keel [176] observed the fundamental connection between Mori theory and geometric invariant theory via Cox rings; one of the main insights is that, roughly speaking, finite generation of the Cox ring is equivalent to an optimal behavior with respect to the minimal model program. This put the toric case into a much more general framework and established Cox rings as an active field of research in algebraic geometry. For example, the explicit presentation of the Cox ring of a given variety in terms of generators and relations is a

central question. The research in this direction was initiated by the work of Batyrev/Popov [33] and Hassett/Tschinkel [162] on (weak) del Pezzo surfaces.

The intention of this book is to provide an elementary access to Cox rings and their applications in algebraic and arithmetic geometry with a particular focus on the new, concrete aspects that Cox rings bring into these fields. The introductory part, consisting of the first three chapters, requires basic knowledge in algebraic geometry, and, in addition, some familiarity with toric varieties is helpful. The subsequent three chapters consider also more advanced topics such as algebraic groups, surface theory, and arithmetic questions.

Chapter 1 provides the mathematical framework for the ideas occurring in the preceding example discussion. We present the basics on graded algebras, quasitorus actions and their quotients, and divisors and sheaves of divisorial algebras. Building on this, we define an essentially unique Cox ring for any irreducible, normal variety with only constant invertible functions and a finitely generated divisor class group:

$$\mathcal{R}(X) := \bigoplus_{[D] \in \text{Cl}(X)} \Gamma(X, \mathcal{O}(D)).$$

First results concern the algebraic properties of the Cox ring, in particular the divisibility properties: Cox rings are factorially graded rings in the sense that we have unique factorization for homogeneous elements. The further main results of the chapter elaborate the relations between the Cox ring and its geometric counterpart, the presentation of the underlying variety as a quotient of the characteristic space by the action of the characteristic quasitorus. For smooth varieties, this quotient presentation equals the universal torsor and in general it dominates the universal torsor.

Chapter 2 discusses the concepts provided in Chapter 1 for the example class of toric varieties; these come with an action of an algebraic torus having a dense open orbit. The basic feature of toric varieties is their complete description in terms of combinatorial data, so-called lattice fans. Approaching toric varieties via quotient presentations turns out to be combinatorial as well, and the describing data, which we call lattice bunches, correspond to lattice fans via linear Gale duality. Besides being illustrative examples, toric varieties are important in subsequent chapters as adapted ambient varieties.

Chapter 3 is devoted to varieties with a finitely generated Cox ring. In this setting, the varieties sharing the same Cox ring all occur as quotients of open subsets of their common total coordinate space, the spectrum of the Cox ring. Geometric invariant theory gives us a concrete combinatorial description of the possible characteristic spaces, which finally leads to the encoding of our varieties by “bunched rings.” The resulting picture shares many combinatorial features with toric geometry. In fact, the varieties inherit many properties from a canonical toric ambient variety. We take a look from the combinatorial

aspect to invariants such as Picard groups; local divisor class groups; and the cones of effective, movable, semiample, and ample divisors. Moreover, we treat singularities, intersection numbers, and the Mori chamber structure of the effective cone. A particularly interesting class are the rational varieties with a torus action of complexity 1, for example, \mathbb{K}^* -surfaces. Here, the bunched ring description leads to a very efficient approach to the geometry; for example, one obtains a concrete combinatorial resolution of singularities.

Chapter 4 begins with a study of Cox rings of embedded varieties and the effect of modifications, for example, blow-ups on the Cox ring. Then we investigate the various quotient presentations of a variety and show that they are all dominated by the characteristic space, playing here a similar role as the universal covering in topology. The problem of lifting group actions to quotient presentations and the automorphism group are investigated. We provide various criteria for finite generation of the Cox ring, for example, Knop's criterion for unirational varieties with a group action of complexity 1, a characterization via the multiplication map, and the characterization in terms of Mori theory due to Hu and Keel. Moreover, we relate Cox rings of blow-ups of the projective space to invariant rings of unipotent group actions following Nagata. For varieties coming with a torus action, we describe the Cox ring in terms of isotropy groups and a certain quotient; this generalizes the toric case and gives the foundation for the bunched ring approach to the more general case of complexity 1. Finally, we take a look at almost homogeneous varieties. After describing the Cox ring of a homogeneous space, we discuss embeddings with a small boundary in terms of bunched rings and then turn to Brion's description of Cox rings of spherical varieties and wonderful compactifications.

In Chapter 5, we take a close look at Cox rings of complex algebraic surfaces. A first general part is devoted to the classification of smooth Mori dream surfaces. We present a complete picture for surfaces with nonnegative anticanonical Iitaka dimension, and study in detail the cases of elliptic rational surfaces, K3 surfaces, and Enriques surfaces. Then we turn to the explicit description of Cox rings by generators and relations. For del Pezzo surfaces, we show that the Cox ring is generated in anticanonical degree 1 and that the ideal of relations is generated by quadrics. A discussion of the relations between Cox rings of del Pezzo surfaces and flag varieties ends this part. Then we return to K3 surfaces. Here we provide a detailed study in the case of Picard number 2 and complete results are obtained for double covers of del Pezzo surfaces and of blow-ups of Hirzebruch surfaces in at most three points. Finally, we develop the theory of rational \mathbb{K}^* -surfaces. Here we allow singularities and show how their minimal resolution is encoded in the Cox ring. As an example class, we present the Gorenstein log del Pezzo \mathbb{K}^* -surfaces in terms of their Cox rings.

The aim of Chapter 6 is to indicate how Cox rings and universal torsors can be applied to arithmetic questions regarding rational points on varieties. We

begin by discussing Colliot-Thélène's and Sansuc's theory of universal torsors over not necessarily algebraically closed fields, and explore the connection to characteristic spaces and Cox rings. Then we enter the problem of the existence of rational points on varieties over number fields. We discuss the Hasse principle and weak approximation. The failure of these principles is often explained by Brauer–Manin obstructions, and we indicate how they can be approached via universal torsors and give an overview of the existing results. Then we turn to Manin's conjecture. For del Pezzo surfaces, it is known in many cases, and a general strategy emerges. We discuss this strategy in detail and show how it can be applied to prove Manin's conjecture for a singular cubic surface.

Each chapter is followed by a choice of exercises and problems. The collections comprise small general background tutorials and examples complementing the text as well as guided exercises to topics going beyond the text including references and, finally, we pose several open problems (*) of varying presumed difficulty.

We are grateful to all people supporting the work on this text. In particular, we thank Carolina Araujo, Michela Artebani, Hendrik Bäker, Victor Batyrev, Benjamin Bechtold, Cinzia Casagrande, Jean-Louis Colliot-Thélène, Christopher Frei, Giuliano Gagliardi, Fritz Hörmann, Johannes Hofscheier, Elaine Huggenberger, Simon Keicher, Alvaro Liendo, Taras Panov, Marta Pieropan, Yuri Prokhorov, Fred Rohrer, Alexei Skorobogatov, Damiano Testa, Dmitri Timashev, Andrea Tironi, and Luca Ugaglia for helpful remarks and discussions.

1

Basic concepts

In this chapter we introduce the Cox ring and, more generally, the *Cox sheaf* and its geometric counterpart, the *characteristic space*. In addition, algebraic and geometric aspects are discussed. Section 1.1 is devoted to commutative algebras graded by monoids. In Section 1.2, we recall the correspondence between actions of quasitori (also called diagonalizable groups) on affine varieties and affine algebras graded by abelian groups and provide the necessary background on good quotients. Section 1.3 is a first step toward constructing Cox rings. Given an irreducible, normal variety X and a finitely generated subgroup $K \subseteq \text{WDiv}(X)$ of the group of Weil divisors, we consider the associated *sheaf of divisorial algebras*

$$\mathcal{S} = \bigoplus_{D \in K} \mathcal{O}_X(D).$$

We present criteria for local finite generation and consider the relative spectrum. A first result says that $\Gamma(X, \mathcal{S})$ is a unique factorization domain if K generates the divisor class group $\text{Cl}(X)$. Moreover, we characterize divisibility in the ring $\Gamma(X, \mathcal{S})$ in terms of divisors on X . In Section 1.4, the Cox sheaf of an irreducible, normal variety X with finitely generated divisor class group $\text{Cl}(X)$ is introduced; roughly speaking it is given as

$$\mathcal{R} = \bigoplus_{[D] \in \text{Cl}(X)} \mathcal{O}_X(D).$$

The Cox ring then is the corresponding ring of global sections. In the case of a free divisor class group well-definedness is straightforward. The case of torsion needs some effort; the precise way to define \mathcal{R} then is to take the quotient of an appropriate sheaf of divisorial algebras with respect to a certain ideal sheaf. Basic algebraic properties and divisibility theory of the Cox ring are investigated in Section 1.5. Finally, in Section 1.6, we study the characteristic space, that is, the relative spectrum $\hat{X} = \text{Spec}_X \mathcal{R}$ of the Cox sheaf. It comes

with an action of the characteristic quasitorus $H = \text{Spec } \mathbb{K}[\text{Cl}(X)]$ and a good quotient $\widehat{X} \rightarrow X$. We relate geometric properties of X to properties of this action and describe the characteristic space in terms of geometric invariant theory.

1.1 Graded algebras

1.1.1 Monoid graded algebras

We recall basic notions on algebras graded by abelian monoids. In this subsection, R denotes a commutative ring with a unit element.

Definition 1.1.1.1 Let K be an abelian monoid. A K -graded R -algebra is an associative, commutative R -algebra A with a unit and a direct sum decomposition

$$A = \bigoplus_{w \in K} A_w$$

into R -submodules $A_w \subseteq A$ such that $A_w \cdot A_{w'} \subseteq A_{w+w'}$ holds for any two elements $w, w' \in K$. The R -submodules $A_w \subseteq A$ are the (K -)homogeneous components of A . An element $f \in A$ is (K -)homogeneous if $f \in A_w$ holds for some $w \in K$, and in this case w is called the *degree* of f . We write $A_\times \subseteq A$ for the multiplicative monoid of homogeneous elements.

We also speak of a K -graded R -algebra as a monoid graded algebra or just as a graded algebra. To compare R -algebras A and A' that are graded by different abelian monoids K and K' , we work with the following notion of a morphism.

Definition 1.1.1.2 A *morphism* from a K -graded algebra A to a K' -graded algebra A' is a pair $(\psi, \tilde{\psi})$, where $\psi: A \rightarrow A'$ is a homomorphism of R -algebras, $\tilde{\psi}: K \rightarrow K'$ is a homomorphism of abelian monoids, and

$$\psi(A_w) \subseteq A'_{\tilde{\psi}(w)}$$

holds for every $w \in K$. In the case $K = K'$ and $\tilde{\psi} = \text{id}_K$, we denote a morphism of graded algebras just by $\psi: A \rightarrow A'$ and also refer to it as a (K -)graded homomorphism.

Example 1.1.1.3 Given an abelian monoid K and $w_1, \dots, w_r \in K$, the polynomial ring $R[T_1, \dots, T_r]$ can be turned into a K -graded R -algebra by setting

$$R[T_1, \dots, T_r]_w := \left\{ \sum_{v \in \mathbb{Z}_{\geq 0}^r} a_v T^v; a_v \in R, v_1 w_1 + \dots + v_r w_r = w \right\}.$$

This K -grading is determined by $\deg(T_i) = w_i$ for $1 \leq i \leq r$. Moreover, $R[T_1, \dots, T_r]$ comes with the natural $\mathbb{Z}_{\geq 0}^r$ -grading given by

$$R[T_1, \dots, T_r]_{\nu} := R \cdot T^{\nu},$$

and we have a canonical morphism $(\psi, \tilde{\psi})$ from $R[T_1, \dots, T_r]$ to itself, where $\psi = \text{id}$ and $\tilde{\psi}: \mathbb{Z}_{\geq 0}^r \rightarrow K$ sends ν to $\nu_1 w_1 + \dots + \nu_r w_r$.

For any abelian monoid K , we denote by K^{\pm} the associated group of differences and by $K_{\mathbb{Q}} := K^{\pm} \otimes_{\mathbb{Z}} \mathbb{Q}$ the associated rational vector space. Note that we have canonical maps $K \rightarrow K^{\pm} \rightarrow K_{\mathbb{Q}}$, where the first one is injective if and only if K admits cancellation and the second one is injective if and only if K^{\pm} is torsion free. Given $w \in K$, we allow ourselves to write $w \in K^{\pm}$ and $w \in K_{\mathbb{Q}}$ for the respective images.

Definition 1.1.14 Let A be a K -graded R -algebra. The *weight monoid* of A is the submonoid $S(A) \subseteq K$ generated by all $w \in K$ with $A_w \neq 0$. The *weight group* of A is the subgroup $K(A) \subseteq K^{\pm}$ generated by $S(A) \subseteq K$. The *weight cone* of A is the convex cone $\omega(A) \subseteq K_{\mathbb{Q}}$ generated by $S(A) \subseteq K$.

By an *integral R -algebra*, we mean an R -algebra $A \neq 0$ without zero divisors. Note that for an integral R -algebra A graded by an abelian monoid K , the weight monoid of A is given as

$$S(A) = \{w \in K; A_w \neq 0\} \subseteq K.$$

We recall the construction of the algebra associated with an abelian monoid; it defines a covariant functor from the category of abelian monoids to the category of monoid graded algebras.

Construction 1.1.15 Let K be an abelian monoid. As an R -module, the associated *monoid algebra* over R is given by

$$R[K] := \bigoplus_{w \in K} R \cdot \chi^w$$

and its multiplication is defined by $\chi^w \cdot \chi^{w'} := \chi^{w+w'}$. If K' is a further abelian monoid and $\tilde{\psi}: K \rightarrow K'$ is a homomorphism, then we have a homomorphism

$$\psi := R[\tilde{\psi}]: R[K] \rightarrow R[K'], \quad \chi^w \mapsto \chi^{\tilde{\psi}(w)}.$$

The pair $(\psi, \tilde{\psi})$ is a morphism from the K -graded algebra $R[K]$ to the K' -graded algebra $R[K']$, and this assignment is functorial.

Note that the monoid algebra $R[K]$ has K as its weight monoid, and $R[K]$ is finitely generated over R if and only if the monoid K is finitely generated. In general, if a K -graded algebra A is finitely generated over R , then its weight monoid is finitely generated and its weight cone is *polyhedral*, that is, the set of nonnegative linear combinations over a given finite collection of vectors.