

Biotechnology

A Comprehensive Treatise in 8 Volumes
edited by H.-J. Rehm and G. Reed

Volume 5

Volume Editor: G. Reed



Verlag Chemie · Weinheim · Deerfield Beach, Florida · Basel

Biotechnology

*A Comprehensive Treatise
in 8 Volumes*

edited by H.-J. Rehm and G. Reed

Volume 5

*Food and Feed Production
with Microorganisms*

Volume Editor: G. Reed



Weinheim · Deerfield Beach, Florida · Basel

Prof. Dr. H.-J. Rehm
Institut für Mikrobiologie
der Universität
Tibustr. 7-15
D-4400 Münster
Federal Republic of Germany

Dr. G. Reed
Universal Foods Corp.
Technical Center
6143 N 60th Street
Milwaukee, Wisconsin 53218
U.S.A.

Publisher's Editor: Dr. Hans F. Ebel
Copy Editor: Christa Maria Schultz
Production Manager: Peter J. Biel

This book contains 135 figures and 201 tables

Deutsche Bibliothek Cataloguing-in-Publication Data

Biotechnology: a comprehensive treatise in 8 vol. /
ed. by H.-J. Rehm and G. Reed. - [Ausg. in 8 Bd.] -
Weinheim; Deerfield Beach, Florida; Basel: Verlag Chemie

NE: Rehm, Hans-Jürgen [Hrsg.]

Vol. 5. → Food and feed production with microorganisms

Food and feed production with microorganisms / vol. ed.: G. Reed. -
Weinheim; Deerfield Beach, Florida; Basel: Verlag Chemie, 1983.

(Biotechnology; Vol. 5)

ISBN 3-527-25767-5 (Weinheim)

ISBN 0-89573-045-6 (Deerfield Beach, Florida)

NE: Reed, Gerald [Hrsg.]

© Verlag Chemie GmbH, D-6940 Weinheim, 1983.

All rights reserved (including those of translation into foreign languages). No part of this book may be reproduced in any form - by photoprint, microfilm, or any other means - nor transmitted or translated into a machine language without written permission from the publishers.

Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.

Composition and Printer: Zehnersche Buchdruckerei, D-6720 Speyer

Bookbinder: Klambt-Druck GmbH, D-6720 Speyer

Printed in the Federal Republic of Germany

Biotechnology

Volume 5

A Comprehensive Treatise
in 8 Volumes
edited by H.-J. Rehm and G. Reed

Volume 1
Microbial Fundamentals

Volume 2
Fundamentals of Biotechnical Engineering

Volume 3
Biomass, Microorganisms for Special Applications,
Microbial Products I, Energy from Renewable Resources

Volume 4
Microbial Products II

Volume 5
Food and Feed Production with Microorganisms

Volume 6a
Microbial Transformations

Volume 6b
Special Microbial Processes

Volume 7
Enzymes in Biotechnology

Volume 8
Microbial Degradations



Deerfield Beach, Florida · Basel



Biotechnology

*A Comprehensive Treatise
in 8 Volumes*

edited by H.-J. Rehm and G. Reed

Volume 1

Microbial Fundamentals

Volume 2

Fundamentals of Biochemical Engineering

Volume 3

Biomass, Microorganisms for Special Applications,
Microbial Products I, Energy from Renewable Resources

Volume 4

Microbial Products II

Volume 5

Food and Feed Production with Microorganisms

Volume 6a

Microbial Transformations

Volume 6b

Special Microbial Processes

Volume 7

Enzymes in Biotechnology

Volume 8

Microbial Degradations



Weinheim · Deerfield Beach, Florida · Basel

Preface

This volume combines a description of all food fermentations. The purpose of these fermentations is to produce or preserve foods or to make them more palatable, more nutritious, more digestible, or a combination of these properties. In addition food fermentations can be distinguished from all other fermentations by the following, important characteristics.

The origin of these fermentations can usually be traced to antiquity but no specific time period or place of origin can be assigned. All food fermentations are spontaneous or "natural" fermentations, or they began as "natural" fermentations; and all of them show a development from home or communal production to industrial production, at least in the developed countries.

Food fermentations are not pure culture fermentations. The substrates are sometimes pasteurized but never sterilized. The cleanest substrates are probably ethanol for vinegar production and boiled brewer's worts. All food fermentations are mixed culture fermentations comprising not only different strains of microorganisms but different species and genera of yeasts, fungi, and bacteria. Spontaneous fermentations usually show a succession of organisms. For instance, the fermentation of soda cracker sponges begins as a yeast fermentation and ends as a lactic acid fermentation, or the wine fermentation which starts with a succession of yeasts (from less ethanol tolerant to more ethanol tolerant yeasts) and ends often with a malo-lactic fermentation. In many instances the fermentation is carried out entirely by the natural microflora

of the substrates or by inoculation of subsequent batches of a fermented food with a retained portion of the preceding batch. This is particularly true of the indigenous fermented foods treated in Chapter 11. On the other hand, several foods are fermented by massive inoculation with a specific organism that has been produced separately for this purpose. Examples are the production of white bread with baker's yeast or the production of wine in which the native microbial population has been suppressed by the addition of sulfur dioxide followed by inoculation with a desirable wine yeast.

In the brewing industry it is possible to prepare a brewer's wort that is almost sterile and to inoculate it with pure culture yeasts. However, the fermentation cannot be carried out under the stringent pure culture conditions that prevail, for instance, in the production of antibiotics. In industrialized countries the trend is towards massive inoculation with desirable strains of microorganisms, so-called starter cultures, produced in separate facilities and often by separate companies.

There are several food related fermentations which have been included in Volume 3 of "Biotechnology", "Biomass, Microorganisms for Special Applications, Microbial Products". These are the production of edible microbial biomass, the production of baker's yeast, of edible mushrooms, of starter cultures for milk and meat processing, of starter cultures for other purposes, and the production of ethanol, acetic acid, lactic acid, citric acid, amino acids, and of extracellular polysaccharides. The produc-

tion of baker's yeast and of lactic acid starter cultures has the most direct impact on the production of fermented foods and the reader should consult the appropriate chapters.

It is difficult to understand the role of microorganisms without some knowledge of the technology of fermented foods. Therefore, the chapters in this volume treat both the technological and the microbiological aspects of fermented foods. It is hoped that a proper balance between processing information and the microbiological aspects of food fermentations has been achieved.

It is foreseeable that in the near future the novel techniques of genetic engineering will permit the creation of microorganisms with properties especially desirable for food and feed production. Some important results through genetic manipulation have been achieved recently in the production of certain pharmaceuticals and selected pure compounds as referred to in Volume 6a of "Biotechnology", but little progress has

been reported so far from the food and feed industries. Hence, the present volume is rather "classical" in this respect. Some novel achievements such as the application, in the sugar industry, of "synthetic" varieties of *E. coli* or of α -galactosidase isolated therefrom will be reported later (Volumes 6b and 7).

The authors of this volume have worked diligently to make their wide knowledge and expertise available to the reader. The value of this comprehensive volume rests basically on their contributions. The volume editor also wishes to thank Prof. H.-J. Rehm for his invaluable editorship and Dr. H. F. Ebel and Christa Schultz of Verlag Chemie for their skill and dedication in getting this volume into print. The members of the Editorial Advisory Board of "Biotechnology" contributed much useful advice.

Milwaukee, November 1983

Gerald Reed

Editorial Advisory Board

of "Biotechnology"

Prof. Dr. S. Aiba

Osaka University
Osaka, Japan

Prof. Dr. N. Blakebrough

National College of
Food Technology
Weybridge, United Kingdom

Prof. Dr. H. Brauer

Technische Universität Berlin
Berlin, Federal Republic of Germany

Prof. Dr. H. Dellweg

Institut für Gärungsgewerbe
und Biotechnologie
Berlin, Federal Republic of Germany

Prof. Dr. A. L. Demain

Massachusetts Institute of Technology
Cambridge, U.S.A.

Prof. Dr. G. Durand

Institut National des Sciences Appliquées
Toulouse, France

Prof. Dr. K. Esser

Ruhr-Universität Bochum
Bochum, Federal Republic of Germany

Prof. Dr. A. Fiechter

ETH Zürich
Zurich, Switzerland

Prof. Dr. S. Fukui

Kyoto University
Kyoto, Japan

Prof. Dr. Vera Johanides

University of Zagreb
Zagreb, Yugoslavia

Prof. Dr. K. Kieslich

Gesellschaft für Biotechnologische Forschung
Braunschweig-Stöckheim, Federal Republic of
Germany

Prof. Dr. H. Klaushofer

Universität für Bodenkultur
Vienna, Austria

Prof. Dr. A. Kocková-Kratochvilová

Slovak Academy of Sciences
Bratislava, Czechoslovakia

Prof. Dr. R. M. Lafferty

Technische Universität Graz
Graz, Austria

Prof. Dr. M. D. Lilly

University College London
London, United Kingdom

Prof. Dr. R. I. Mateles

Stauffer Chemical Company
Westport, U.S.A.

Prof. Dr. K. Mosbach

University of Lund
Lund, Sweden

Prof. Dr. G. Shelef

Technion - Israel Institute of Technology
Haifa, Israel

Prof. Dr. G. K. Skryabin

USSR Academy of Sciences
Moscow, USSR

Prof. Dr. H. Suomalainen

The Finnish State Alcohol Monopoly
Helsinki, Finland

Prof. Dr. G. Terui

Osaka University
Osaka, Japan

List of Contributors

of Volume 5

Prof. Dr. Larry R. Beuchat

Department of Food Science
University of Georgia
Agricultural Experiment Station
Experiment, Georgia 30212
U.S.A.

Dr. Heinrich Ebner

A-4020 Linz/Donau
Austria

Dr. Leslie Bluhm

Jos. E. Seagrams and Sons
Louisville, Kentucky 10677
U.S.A.

Prof. Dr. Matías J. Fernández Díez

Instituto de la Grasa y sus Derivados
Sevilla 12
Spain

Prof. Dr. Vittorio Bottazzi

Istituto di Microbiologia
Facolta di Agraria
Universita Cattolica del Sacre Cuore
I-29100 Piacenza
Italy

Dr. Heinrich Follmann

Heinrich Frings GmbH & Co KG
D-5300 Bonn
Federal Republic of Germany

Dr. Johan M. Castelein

Rega School
B-3000 Louvain
Belgium

Dr. William A. Hardwick Jr.

Anheuser-Busch Inc.
Technical Center
St. Louis, Missouri 63118
U.S.A.

***Prof. Dr. Suzanne Lafon-
Lafourcade***

Institut d'Oenologie
Université de Bordeaux II et INRA
F-33405 Talence
France

Dr. Gottfried Spicher

Bundesforschungsanstalt für Getreide-
und Kartoffelverarbeitung
D-4930 Detmold
Federal Republic of Germany

Dr. Douglas W. Lehrian

Hershey Foods Corp.
Technical Center
Hershey, Pennsylvania 17033
U.S.A.

Prof. Dr. John R. Stamer

Cornell University
New York State
Agricultural Experiment Station
Geneva, New York 14456
U.S.A.

Dr. Hans-Ulrich Liepe

Rudolf Müller & Co
D-6300 Giessen
Federal Republic of Germany

Prof. Dr. Ebenezer R. Vedamuthu

Microlife Technics
Sarasota, Florida 33578
U.S.A.

Gordon R. Patterson

Hershey Foods Corp.
Corporate Administrative Center
Hershey, Pennsylvania 17033
U.S.A.

Prof. Dr. Hubert Verachtert

Laboratory of Industrial Microbiology
University of Leuven
Heverlee-Louvain
Belgium

Dr. Henry J. Peppler

Universal Foods Corp.
Technical Center
Milwaukee, Wisconsin 53218
U.S.A.

Clinton Washam

Mallinckrodt Inc.
St. Louis, Missouri 63134
U.S.A.

Dr. Gary W. Sanderson

Universal Foods Corp.
Technical Center
Milwaukee, Wisconsin 53218
U.S.A.

Contents

Chapter 1

Baked Goods 1

by *Gottfried Spicher*

Chapter 2

Wine and Brandy 81

by *Suzanne Lafon-Lafourcade*

Chapter 3

Beer 165

by *William A. Hardwick*

Chapter 4

Cheese 231

by *Ebenezer R. Vedamuthu*
and *Clinton Washam*

Chapter 5

Other Fermented Dairy Products 315

by *Vittorio Bottazzi*

Chapter 6

Lactic Acid Fermentation of Cabbage and Cucumbers 365

by *John R. Stamer*

Chapter 7

Olives 379

by *Matías J. Fernández Díez*

Chapter 8

Starter Cultures in Meat Production 399

by *Hans-Ulrich Liepe*

XII Contents

Chapter 9

Vinegar 425

by *Heinrich Ebner*
and *Heinrich Follmann*

Chapter 10

Distilled Beverages 447

by *Leslie Bluhm*

Chapter 11

Indigenous Fermented Foods 477

by *Larry R. Beuchat*

Chapter 12

Cocoa Fermentation 529

by *Douglas W. Lehrian*
and *Gordon R. Patterson*

Chapter 13

Tea Manufacture 577

by *Gary W. Sanderson*

Chapter 14

Coffee Fermentation 587

by *Johan M. Castelein*
and *Hubert Verachtert*

Chapter 15

Fermented Feeds and Feed Supplements 599

by *Henry J. Peppler*

Index 617

Chapter 1

Baked Goods

Gottfried Spicher

Bundesforschungsanstalt für Getreide- und Kartoffelverarbeitung
Detmold, Federal Republic of Germany

- I. Introduction
- II. The Nutritional Contribution of Bread
- III. Outline of the Technology of Baked Goods
- IV. Choice and Preparation of Raw Materials
 - A. Flour
 - B. Water
 - C. Salt
 - D. Other Ingredients
 - E. Leavening Agents
 1. Biological leavening agents
 - a) Yeast
 - b) Sour dough starter cultures
 2. Chemical leavening agents
 3. Physical leavening agents
 - F. Additives Affecting the Processing Characteristics of Flour and Doughs (Dough Conditioners)
 1. Regulation of water absorption in doughs
 2. Improvement of the properties of doughs and baked goods
 3. Effect of dough fermentation
 4. Dough acidification
 - G. Formulated Additives
- V. Formation of the Dough
 - A. Preparation of Raw Materials
 - B. Kneading of Doughs
 1. Mixers
 2. The mixing process
 3. Dough formation
- VI. Leavening of Doughs
 - A. Processing of Wheat Doughs
 1. Straight dough process

2. Sponge dough process
3. Liquid pre-ferment processes
4. Continuous mix process
5. No time doughs
- B. Rye Dough (Sour Dough) Processes
 1. Seed
 2. Sour dough processes with several stages
 3. Two stage sour dough process
 4. Single stage sour doughs
 5. Salt sour processes
 6. Freeze-dried sour dough
 7. Effect of processing conditions on the sour dough fermentation
 8. Effect of processing conditions on the degradation of carboxylic acids
 9. Effect of processing conditions on the hydrolysis of proteins

VII. Processing of Fermented Doughs

- A. Fermentation
- B. Dough Make-up
- C. Proofing

VIII. Baking

- A. Baking Ovens
- B. Processing Conditions During Baking
- C. Physico-chemical Changes During Baking
- D. Formation of Aroma Substances

IX. Bread Quality and its Preservation

- A. Bread Defects
- B. Bread Staling
- C. Microbial Spoilage
- D. Retention of Bread Quality
 1. Storage
 2. Packaging
 3. Preservation
 - a) Physical processes
 - b) Chemical processes

X. References

I. Introduction

Cereals and bread have been the basic food for individuals and population groups for many centuries. In various countries bread accounts for 18–80% of all nutrients.

The history of bread can be traced back about 6 millennia. It is probable that it developed from a gruel. During the earlier stone age the preparation of a gruel made from rubbed or ground grain and water or milk was known by all civilizations. The flat bread developed from this gruel which was either air-dried or baked on hot stones. The first primitive baking was done by placing formed doughs into hot ashes.

Towards the end of the stone age the flat bread had assumed the shape of a disc. These breads were generally eaten warm, or they could be dried and stored. All types of grain could be used for the preparation of the gruel, but the choice of grains for the preparation of baked flat breads is narrower. Therefore, flat breads were introduced only slowly and only in some localities. Barley and other grains (hardly known today) were used. With the change to the loaf bread the choice of suitable grains is still narrower. Today wheat is grown worldwide for this purpose (Fig. 1, Table 1). Beyond this rye is grown in Central and Northern Europe. Wheat and rye are also called "bread" grains because of their suitability for the production of baked goods.

The production of a loaf bread presupposes the development of the baking oven and the discovery of the dough fermentation. The earliest baking oven has been found in the area of the early Babylonians. The Egyptians developed a baking oven approaching the design of a modern oven about 2700 B.C. Since 1750 B.C. there were professional bakers in Egypt. They were familiar with the raising of wheat doughs with brewer's yeast and even with chemical leavening. In 450 B.C. the Egyptians were familiar with the souring of doughs. But the form of this sour dough bread was still similar to that of a flat bread. Discoveries in




Figure 1. Production of various cereals worldwide (calculated from data of the FAO Production Year Book by ROTHE, 1980).

tombs have shown that a sour dough bread was known in rye growing areas north of the Alps since the end of the bronze age (800 B.C.). In 100 B.C. the sour dough bread was generally known throughout the world (VON STOKAR, 1951). The Greeks developed the art of baking further, and especially the design of baking ovens. Greek bakers brought the art of baking bread and cake to Rome. The Romans in turn developed the earlier techniques of other populations to a high degree of technical and organizational perfection. They also developed the form of the baking oven which remained the model until the 19th century. In Central Europe the preparation of bread in its proper sense stems from the middle of the first millennium A.C. The further development was very fast, and at the beginning of the 8th century bakers are mentioned in the "lex alamanorum" (ROHRLICH, 1976).

The introduction of the stone oven permitted an increase in the heating surface and an increase in production. The oldest dough kneading machines date from the second half of the 18th century. Finally, the industrial production of baker's yeast in the last century was decisive for the technology of baking. Only this development permitted the production of the loaf bread as we know it today. The concentration of the

Table 1. Wheat and Rye Production of Some Countries in 1977
(in 1000 tons)

Country	Wheat	Rye	Wheat/Rye Ratio
Italy	6330	31	204/1
Bulgaria	3010	20	150/1
Great Britain	5230	38	138/1
USA	55130	432	128/1
Canada	19650	392	50/1
France	17450	376	46/1
Hungary	5310	147	36/1
Argentina	5300	170	31/1
USSR	92040	8470	10.9/1
Czechoslovakia	5240	870	6.0/1
Sweden	1560	364	4.3/1
Austria	1070	351	3.0/1
West Germany	7180	2538	2.8/1
East Germany	3100	1500	2.1/1
Denmark	605	320	1.9/1
Poland	5310	6200	0.9/1
Totals	386600	23770	16 /1

(ROTHE, 1980)

population in towns at the beginning of the 20th century led to technical advances and the rationalization of production. A lack of workers, undesirable working hours, the desire to make the work easier, and to lower production costs contributed to this development. Mainly since 1950 automation has advanced in a broad front and has led to the establishment of some fully automated bakeries. The typical properties of bread, its form, appearance, texture and flavor, were originally produced manually and in conformity with the quality of available grains. Now the baker has to produce bread in the same form and with the same quality by means of rational, technical processes (WASSERMANN, 1981). This change from the manual to the industrial method of producing bread and other baked goods can be observed in almost all industrialized countries. The conditions for this change are particularly favorable in countries in which the assortment of types of breads and baked goods is not very large. Therefore, the development started earlier in countries producing bread almost exclusively from wheat, as in the United States. During the past years the industrial production of rye

breads and mixed grain breads has progressed considerably. The production of sweet goods and shelf stable baked goods occurs today already largely in industrial facilities (SEIBEL, 1970).

The development of cereal foods has proceeded through several stages: from roasted grain to gruels to flat breads and finally to leavened bread loaves. But the latter stages have not replaced the earlier ones. Today all stages are still practiced. About 60% of the world population, mainly in Central America, parts of South America, Africa, the Middle East and the Far East, eat gruels made from grain crops and flat breads.

II. The Nutritional Contribution of Bread

The individual grain consists of a starchy endosperm whose outer layers are rich in protein, vitamins, and minerals; a largely

indigestible fruit coat (or pericarp); and a germ or embryo. The higher the degree of extraction of the flour, that is, the greater the yield of flour for a given weight of grain, the greater the amount of seed coat and of the outer layer in the flour. Such flours are darker. Short extraction flour (or very white flour) contains almost exclusively portions of the endosperm. Wheat flours generally contain more protein and less fiber than rye flours. The ash content of rye flours is almost twice as high as that of comparable wheat flours.

Grains and products made from milled grains have a favorable combination of nutrients. For 4 billion people they provide more than 50% of the required calories, and in some regions more than 90%. Beyond this they supply considerable percentages of the nutritionally important proteins, B vitamins, and minerals and trace minerals. Bread is a staple food in Germany (as it is in other industrialized countries) and supplies the following percentages of the required nutrients: 20% of the calories, 45–55% of the carbohydrates, 22–25% of the proteins in the form of plant protein. Some specialty breads also contain animal proteins. In addition bread contributes 25–33% of the required vitamins of the B complex (thiamin, riboflavin), 2 grams daily of minerals (among them calcium and iron), as well as a considerable part of the requirement for sodium chloride. If bread is sup-

plemented with foods rich in nutrients such as animal protein, vitamins A, C and D, and some minerals, it can be a suitable basis for a nutritionally optimal diet (CREMER et al., 1969) (Table 2).

Carbohydrates occur mainly in the form of starch (on the average about 45–50% of the weight of fresh bread) (Table 3). Mono- and disaccharides are not very important since they account for less than 5% of the carbohydrates. The carbohydrates of bread serve mainly as carbon and energy sources. In the form of gelatinized starch they are preferable to excessive amounts of carbohydrates in the form of sugar because of the lesser risk of contributing to arteriosclerosis and diabetes (YUDKIN and RODDY, 1964). In addition carbohydrates counteract the tendency toward an excessive intake of fats in industrialized countries. They also aid in the digestion and complete oxidation of fats in humans because of large inner surface and porosity which distributes the fat finely in the stomach and intestinal tract.

Bread, and particularly bread from high extraction flours and whole grain, contains bulk materials. These are the indigestible, organic substances present in the structural molecules and the cell wall of plants. They consist mainly of cellulose, hemicellulose, pectin, and lignin. These substances increase the mass of food in the intestines and increase the feeling of satiety. They regulate the passage of food through the intestinal

Table 2. Contribution of Some Foods to Caloric Supply in West Germany

	Percent of Total Caloric Content		
	1950	1960	1974
1. Bread flour (75% wheat, 25% rye)	32	26	20
2. Pork	3	5	8
3. Beef	2	3	4
4. Sugar	9	11	13
5. Alcohol	2	5	7
Sum of 2 to 5	16	24	32

(MENDEN et al., 1975)