

M. J. D. Powell

Approximation Theory and Methods

逼近理论和方法


CAMBRIDGE

世界图书出版公司
www.wpcbj.com.cn

M. J. D. POWELL

*John Humphrey Plummer Professor of Applied Numerical Analysis
University of Cambridge*

*Approximation theory
and methods*



CAMBRIDGE
UNIVERSITY PRESS

Approximation Theory and Methods (978-0-521-29514-7) by M. J. D. Powell, first published by Cambridge University Press 2001
All rights reserved.

This reprint edition for the People's Republic of China is published by arrangement with the Press Syndicate of the University of Cambridge, Cambridge, United Kingdom.

© Cambridge University Press & Beijing World Publishing Corporation 2014 This book is in copyright. No reproduction of any part may take place without the written permission of Cambridge University Press or Beijing World Publishing Corporation.

This edition is for sale in the mainland of China only, excluding Hong Kong SAR, Macao SAR and Taiwan, and may not be bought for export therefrom.

此版本仅限中华人民共和国境内销售，不包括香港、澳门特别行政区及中国台湾。不得出口。

图书在版编目 (CIP) 数据

逼近理论和方法 = Approximation theory and methods: 英文/(英)鲍威尔
(Powell, M. J. D.) 著. —影印本. —北京: 世界图书出版公司北京公司, 2014. 9
ISBN 978 - 7 - 5100 - 8625 - 0

I. ①逼… II. ①鲍… III. ①逼近论—教材—英文 IV. ① O174. 41

中国版本图书馆 CIP 数据核字 (2014) 第 211071 号

书 名: Approximation Theory and Methods

作 者: M. J. D. Powell

中译名: 逼近理论和方法

责任编辑: 高蓉 刘慧

出 版 者: 世界图书出版公司北京公司

印 刷 者: 三河市国英印务有限公司

发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010 - 64021602, 010 - 64015659

电子信箱: kjb@wpcbj. com. cn

开 本: 24 开

印 张: 15

版 次: 2015 年 1 月

版权登记: 图字: 01 - 2013 - 8236

书 号: 978 - 7 - 5100 - 8625 - 0

定 价: 49.00 元

Approximation theory and methods

PREFACE

There are several reasons for studying approximation theory and methods, ranging from a need to represent functions in computer calculations to an interest in the mathematics of the subject. Although approximation algorithms are used throughout the sciences and in many industrial and commercial fields, some of the theory has become highly specialized and abstract. Work in numerical analysis and in mathematical software is one of the main links between these two extremes, for its purpose is to provide computer users with efficient programs for general approximation calculations, in order that useful advances in the subject can be applied. This book presents the view of a numerical analyst, who enjoys the theory, and who is keenly interested in its importance to practical computer calculations. It is based on a course of twenty-four lectures, given to third-year mathematics undergraduates at the University of Cambridge. There is really far too much material for such a course, but it is possible to speak coherently on each chapter for about one hour, and to include proofs of most of the main theorems. The prerequisites are an introduction to linear spaces and operators and an intermediate course on analysis, but complex variable theory is not required.

Spline functions have transformed approximation techniques and theory during the last fifteen years. Not only are they convenient and suitable for computer calculations, but also they provide optimal theoretical solutions to the estimation of functions from limited data. Therefore seven chapters are given to spline approximations. The classical theory of best approximations from linear spaces with respect to the minimax, least squares and L_1 -norms is also studied, and algorithms are described and analysed for the calculation of these approximations. Interpolation is considered also, and the accuracy of interpolation and

other linear operators is related to the accuracy of optimal algorithms. Special attention is given to polynomial functions, and there is one chapter on rational functions, but, due to the constraints of twenty-four lectures, the approximation of functions of several variables is not included. Also there are no computer listings, and little attention is given to the consequences of the rounding errors of computer arithmetic. All theorems are proved, and the reader will find that the subject provides a wide range of techniques of proof. Some material is included in order to demonstrate these techniques, for example the analysis of the convergence of the exchange algorithm for calculating the best minimax approximation to a continuous function. Several of the proofs are new. In particular, the uniform boundedness theorem is established in a way that does not require any ideas that are more advanced than Cauchy sequences and completeness. Less functional analysis is used than in other books on approximation theory, and normally functions are assumed to be continuous, in order to simplify the presentation. Exercises are included with each chapter which support and extend the text. All references to related work are given in an appendix.

It is a pleasure to acknowledge the excellent opportunities I have received for research and study in the Department of Applied Mathematics and Theoretical Physics at the University of Cambridge since 1976, and before that at the Atomic Energy Research Establishment, Harwell. My interest in approximation theory began at Harwell, stimulated by the enthusiasm of Alan Curtis, and strengthened by Pat Gaffney, who developed some of the theory that is reported in Chapter 24. I began to write this book in the summer of 1978 at the University of Victoria, Canada, and I am grateful for the facilities of their Department of Mathematics, for the encouragement of Ian Barrodale and Frank Roberts, and for financial support from grants A5251 and A7143 of the National Research Council of Canada. At Cambridge David Carter of King's College kindly studied drafts of the chapters and offered helpful comments. The manuscript was typed most expertly by Judy Roberts, Hazel Felton, Margaret Harrison and Paula Lister. I wish to express special thanks to Hazel for her assistance and patience when I was redrafting the text. My wife, Caroline, not only showed sympathetic understanding at home during the time when I worked long hours to complete the manuscript, but also she assisted with the figures. This work is dedicated to Caroline.

Pembroke College, Cambridge
January 1980

M. J. D. POWELL

CONTENTS

Preface	ix
1 The approximation problem and existence of best approximations	1
1.1 Examples of approximation problems	1
1.2 Approximation in a metric space	3
1.3 Approximation in a normed linear space	5
1.4 The L_p -norms	6
1.5 A geometric view of best approximations	9
2 The uniqueness of best approximations	13
2.1 Convexity conditions	13
2.2 Conditions for the uniqueness of the best approximation	14
2.3 The continuity of best approximation operators	16
2.4 The 1-, 2- and ∞ -norms	17
3 Approximation operators and some approximating functions	22
3.1 Approximation operators	22
3.2 Lebesgue constants	24
3.3 Polynomial approximations to differentiable functions	25
3.4 Piecewise polynomial approximations	28
4 Polynomial interpolation	33
4.1 The Lagrange interpolation formula	33
4.2 The error in polynomial interpolation	35
4.3 The Chebyshev interpolation points	37
4.4 The norm of the Lagrange interpolation operator	41
5 Divided differences	46
5.1 Basic properties of divided differences	46
5.2 Newton's interpolation method	48

5.3	The recurrence relation for divided differences	49
5.4	Discussion of formulae for polynomial interpolation	51
5.5	Hermite interpolation	53
6	The uniform convergence of polynomial approximations	61
6.1	The Weierstrass theorem	61
6.2	Monotone operators	62
6.3	The Bernstein operator	65
6.4	The derivatives of the Bernstein approximations	67
7	The theory of minimax approximation	72
7.1	Introduction to minimax approximation	72
7.2	The reduction of the error of a trial approximation	74
7.3	The characterization theorem and the Haar condition	76
7.4	Uniqueness and bounds on the minimax error	79
8	The exchange algorithm	85
8.1	Summary of the exchange algorithm	85
8.2	Adjustment of the reference	87
8.3	An example of the iterations of the exchange algorithm	88
8.4	Applications of Chebyshev polynomials to minimax approximation	90
8.5	Minimax approximation on a discrete point set	92
9	The convergence of the exchange algorithm	97
9.1	The increase in the levelled reference error	97
9.2	Proof of convergence	99
9.3	Properties of the point that is brought into reference	102
9.4	Second-order convergence	105
10	Rational approximation by the exchange algorithm	111
10.1	Best minimax rational approximation	111
10.2	The best approximation on a reference	113
10.3	Some convergence properties of the exchange algorithm	116
10.4	Methods based on linear programming	118
11	Least squares approximation	123
11.1	The general form of a linear least squares calculation	123
11.2	The least squares characterization theorem	125
11.3	Methods of calculation	126
11.4	The recurrence relation for orthogonal polynomials	131

12	Properties of orthogonal polynomials	136
12.1	Elementary properties	136
12.2	Gaussian quadrature	138
12.3	The characterization of orthogonal polynomials	141
12.4	The operator R_n	143
13	Approximation to periodic functions	150
13.1	Trigonometric polynomials	150
13.2	The Fourier series operator S_n	152
13.3	The discrete Fourier series operator	156
13.4	Fast Fourier transforms	158
14	The theory of best L_1 approximation	164
14.1	Introduction to best L_1 approximation	164
14.2	The characterization theorem	165
14.3	Consequences of the Haar condition	169
14.4	The L_1 interpolation points for algebraic polynomials	172
15	An example of L_1 approximation and the discrete case	177
15.1	A useful example of L_1 approximation	177
15.2	Jackson's first theorem	179
15.3	Discrete L_1 approximation	181
15.4	Linear programming methods	183
16	The order of convergence of polynomial approximations	189
16.1	Approximations to non-differentiable functions	189
16.2	The Dini-Lipschitz theorem	192
16.3	Some bounds that depend on higher derivatives	194
16.4	Extensions to algebraic polynomials	195
17	The uniform boundedness theorem	200
17.1	Preliminary results	200
17.2	Tests for uniform convergence	202
17.3	Application to trigonometric polynomials	204
17.4	Application to algebraic polynomials	208
18	Interpolation by piecewise polynomials	212
18.1	Local interpolation methods	212
18.2	Cubic spline interpolation	215
18.3	End conditions for cubic spline interpolation	219
18.4	Interpolating splines of other degrees	221

19	<i>B</i>-splines	227
19.1	The parameters of a spline function	227
19.2	The form of <i>B</i> -splines	229
19.3	<i>B</i> -splines as basis functions	231
19.4	A recurrence relation for <i>B</i> -splines	234
19.5	The Schoenberg-Whitney theorem	236
20	Convergence properties of spline approximations	241
20.1	Uniform convergence	241
20.2	The order of convergence when f is differentiable	243
20.3	Local spline interpolation	246
20.4	Cubic splines with constant knot spacing	248
21	Knot positions and the calculation of spline approximations	254
21.1	The distribution of knots at a singularity	254
21.2	Interpolation for general knots	257
21.3	The approximation of functions to prescribed accuracy	261
22	The Peano kernel theorem	268
22.1	The error of a formula for the solution of differential equations	268
22.2	The Peano kernel theorem	270
22.3	Application to divided differences and to polynomial interpolation	274
22.4	Application to cubic spline interpolation	277
23	Natural and perfect splines	283
23.1	A variational problem	283
23.2	Properties of natural splines	285
23.3	Perfect splines	290
24	Optimal interpolation	298
24.1	The optimal interpolation problem	298
24.2	L_1 approximation by <i>B</i> -splines	301
24.3	Properties of optimal interpolation	307
	Appendix A The Haar condition	313
	Appendix B Related work and references	317
	Index	333

The approximation problem and existence of best approximations

1.1 Examples of approximation problems

A simple example of an approximation problem is to draw a straight line that fits the curve shown in Figure 1.1. Alternatively we may require a straight line fit to the data shown in Figure 1.2. Three possible fits to the discrete data are shown in Figure 1.3, and it seems that lines *B* and *C* are better than line *A*. Whether *B* or *C* is preferable depends on our confidence in the highest data point, and to choose between the two straight lines we require a measure of the quality of the trial approximations. These examples show the three main ingredients of an approximation calculation, which are as follows: (1) A function, or some data, or

Figure 1.1. A function to be approximated.




Figure 1.2. Some data to be approximated.





Figure 1.3. Three straight-line fits to the data of Figure 1.2.



more generally a member of a set, that is to be approximated. We call it f . (2) A set, $\mathcal{A}$ say, of approximations, which in the case of the given examples is the set of all straight lines. (3) A means of selecting an approximation from $\mathcal{A}$.

Approximation problems of this type arise frequently. For instance we may estimate the solution of a differential equation by a function of a certain simple form that depends on adjustable parameters, where the measure of goodness of the approximation is a scalar quantity that is derived from the residual that occurs when the approximating function is substituted into the differential equation. Another example comes from the choice of components in electrical circuits. The function f may be the required response from the circuit, and the range of available components gives a set $\mathcal{A}$ of attainable responses. We have to approximate f by a member of $\mathcal{A}$, and we require a criterion that selects suitable components. Moreover, in computer calculations of mathematical functions, the mathematical function is usually approximated by one that is easy to compute.

Many closely related questions are of interest also. Given f and $\mathcal{A}$, we may wish to know whether any member of $\mathcal{A}$ satisfies a fixed tolerance condition, and, if suitable approximations exist, we may be willing to accept any one. It is often useful to develop methods for selecting a member of $\mathcal{A}$ such that the error of the chosen approximation is always within a certain factor of the least error that can be achieved. It may be possible to increase the size of $\mathcal{A}$ if necessary, for example $\mathcal{A}$ may be a linear space of polynomials of any fixed degree, and we may wish to predict the improvement in the best approximation that comes from enlarging $\mathcal{A}$ by increasing the degree. At the planning stage of a numerical method we may know only that f will be a member of a set $\mathcal{B}$, in which case it is relevant to discover how well any member of $\mathcal{B}$ can be approximated from $\mathcal{A}$. Further, given $\mathcal{B}$, it may be valuable to compare the suitability of two different sets of approximating functions, $\mathcal{A}_0$ and $\mathcal{A}_1$. Numerical methods for the calculation of approximating functions are required. This book presents much of the basic theory and algorithms that are relevant to these questions, and the material is selected and described in a way that is intended to help the reader to develop suitable techniques for himself.

1.2 Approximation in a metric space

The framework of metric spaces provides a general way of measuring the goodness of an approximation, because one of the basic

properties of a metric space is that it has a distance function. Specifically, the distance function $d(x, y)$ of a metric space $\mathcal{B}$ is a real-valued function, that is defined for all pairs of points (x, y) in $\mathcal{B}$, and that has the following properties. If $x \neq y$, then $d(x, y)$ is positive and is equal to $d(y, x)$. If $x = y$, then the value of $d(x, y)$ is zero. The triangle inequality

$$d(x, y) \leq d(x, z) + d(z, y) \quad (1.1)$$

must hold, where x, y and z are any three points in $\mathcal{B}$.

In most approximation problems there exists a suitable metric space that contains both f and the set of approximations $\mathcal{A}$. Then it is natural to decide that $a_0 \in \mathcal{A}$ is a better approximation than $a_1 \in \mathcal{A}$ if the inequality

$$d(a_0, f) < d(a_1, f) \quad (1.2)$$

is satisfied. We define $a^* \in \mathcal{A}$ to be a best approximation if the condition

$$d(a^*, f) \leq d(a, f) \quad (1.3)$$

holds for all $a \in \mathcal{A}$.

The metric space should be chosen so that it provides a measure of the error of each trial approximation. For example, in the problem of fitting the data of Figure 1.2 by a straight line, we approximate a set of points $\{(x_i, y_i); i = 1, 2, 3, 4, 5\}$ by a function of the form

$$p(x) = c_0 + c_1 x, \quad (1.4)$$

where c_0 and c_1 are scalar coefficients. Because we are interested in only five values of x , the most convenient space is $\mathcal{R}^5$. The fact that $p(x)$ depends on two parameters is not relevant to the choice of metric space. We measure the goodness of the approximation (1.4) as the distance, according to the metric we have chosen, from the vector of function values $\{p(x_i); i = 1, 2, 3, 4, 5\}$ to the data values $\{y_i; i = 1, 2, 3, 4, 5\}$.

It may be important to know whether or not a best approximation exists. One reason is that many methods of calculation are derived from properties that are obtained by a best approximation. The following theorem shows existence in the case when $\mathcal{A}$ is compact.

Theorem 1.1

If $\mathcal{A}$ is a compact set in a metric space $\mathcal{B}$, then, for every f in $\mathcal{B}$, there exists an element $a^* \in \mathcal{A}$, such that condition (1.3) holds for all $a \in \mathcal{A}$.

Proof. Let d^* be the quantity

$$d^* = \inf_{a \in \mathcal{A}} d(a, f). \quad (1.5)$$

If there exists a^* in $\mathcal{A}$ such that this bound on the distance is achieved, then there is nothing to prove. Otherwise there is a sequence $\{a_i; i = 1, 2, \dots\}$ of points in $\mathcal{A}$ which gives the limit

$$\lim_{i \rightarrow \infty} d(a_i, f) = d^*. \quad (1.6)$$

By compactness the sequence has at least one limit point in $\mathcal{A}$, a^+ say. Expression (1.6) and the definition of a^+ imply that, for any $\varepsilon > 0$, there exists an integer k such that the inequalities

$$d(a_k, f) < d^* + \frac{1}{2}\varepsilon \quad (1.7)$$

and

$$d(a_k, a^+) < \frac{1}{2}\varepsilon \quad (1.8)$$

are obtained. Hence the triangle inequality (1.1) provides the bound

$$\begin{aligned} d(a^+, f) &\leq d(a^+, a_k) + d(a_k, f) \\ &< d^* + \varepsilon. \end{aligned} \quad (1.9)$$

Because ε can be arbitrarily small, the distance $d(a^+, f)$ is not greater than d^* . Therefore a^+ is a best approximation. $\square$

When $\mathcal{A}$ is not compact it is easy to find examples to show that best approximations may not exist. For instance, let $\mathcal{B}$ be the Euclidean space $\mathcal{R}^2$ and let $\mathcal{A}$ be the set of points that are strictly inside the unit circle. There is no best approximation to any point of $\mathcal{B}$ that is outside or on the unit circle.

1.3 Approximation in a normed linear space

The properties of metric spaces are not sufficiently strong for most of our work, so it is assumed that $\mathcal{A}$ and f are contained in a normed linear space, which we call $\mathcal{B}$ also when we want to refer to it. The norm is a real-valued function $\|x\|$ that is defined for all $x \in \mathcal{B}$. Its properties are such that the function

$$d(x, y) = \|x - y\| \quad (1.10)$$

is suitable as a distance function. Therefore, by letting z be zero in expression (1.1) and by reversing the sign of y , we may deduce the triangle inequality

$$\|x + y\| \leq \|x\| + \|y\|. \quad (1.11)$$

Moreover, the norm must satisfy the homogeneity condition

$$\|\lambda x\| = |\lambda| \|x\| \quad (1.12)$$

for all $x \in \mathcal{B}$ and for all scalars λ .