




Protein Phosphorylation in Parasites

Novel Targets for Antiparasitic Intervention

Edited by Christian Doerig, Gerald Späth, and Martin Wiese



Edited by Christian Doerig, Gerald Späth, and Martin Wiese

Protein Phosphorylation in Parasites

Novel Targets for Antiparasitic Intervention



WILEY Blackwell

The Editors

Volume Editors:

Prof. Dr. Christian Doerig

Monash University
Department of Microbiology
Wellington Road Clayton
Building 76
Victoria 3800
Australia
christian.doerig@monash.edu

Dr. Gerald Spath

Institut Pasteur Department of Parasitology and Mycology Rue du Dr. Roux 25 75015 Paris France gerald.spaeth@pasteur.fr

Dr. Martin Wiese

University of Strathclyde Institute of Pharmacy and Biomedicine Cathedral Street 161 G4 0RE Glasgow United Kingdom martin.wiese@strath.ac.uk

Series Editor:

Prof. Dr. Paul M. Selzer

MSD Animal Health Innovation GmbH Zur Propstei 55270 Schwabenheim Germany paul.selzer@msd.de

Cover legend

See page XIII

Limit of Liability/Disclaimer of Warranty. While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty can be created or extended by sales representatives or written sales materials. The Advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor authors shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

Library of Congress Card No.: applied for

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library.

Bibliographic information published by the Deutsche Nationalbibliothek

The Deutsche Nationalbibliothek lists this publication in the Deutsche Nationalbibliografie; detailed bibliographic data are available on the Internet at http://dnb.d-nb.de>.

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Boschstr. 12, 69469 Weinheim, Germany

Wiley-Blackwell is an imprint of John Wiley & Sons, formed by the merger of Wiley's global Scientific, Technical, and Medical business with Blackwell Publishing,

All rights reserved (including those of translation into other languages). No part of this book may be reproduced in any form – by photoprinting, microfilm, or any other means – nor transmitted or translated into a machine language without written permission from the publishers. Registered names, trademarks, etc. used in this book, even when not specifically marked as such, are not to be considered unprotected by law.

 Print ISBN:
 978-3-527-33235-9

 ePDF ISBN:
 978-3-527-67539-5

 ePub ISBN:
 978-3-527-67537-1

 Mobi ISBN:
 978-3-527-67538-8

oBook ISBN: 978-3-527-67540-1

Cover Design Adam-Design, Weinheim, Germany

Typesetting Thomson Digital, Noida, India

Printing and Binding Markono Print Media Pte Ltd, Singapore

Printed on acid-free paper

Edited by Christian Doerig, Gerald Späth, and Martin Wiese

Protein Phosphorylation in Parasites

Titles of the Series "Drug Discovery in Infectious Diseases"

Selzer, P.M. (ed.)

Antiparasitic and Antibacterial Drug Discovery

From Molecular Targets to Drug Candidates

2009

Print ISBN: 978-3-527-32327-2, also available in Adobe PDF format ISBN: 978-3-527-62682-3

Becker, K. (ed.)

Apicomplexan Parasites Molecular Approaches toward Targeted

Drug Development

Print ISBN: 978-3-527-32731-7, also available as digital format

Caffrey, C.R. (ed.)

Parasitic Helminths

Targets, Screens, Drugs and Vaccines

2012

Print ISBN: 978-3-527-33059-1, also available as digital format

Jäger, T., Koch, O., Flohé, L. (eds.)

Trypanosomatid Diseases Molecular Routes to Drug Discovery

Print ISBN: 978-3-527-33255-7, also available as digital format

Forthcoming Topics of the Series

Gottfried Unden, Eckard Thines, Anja Schüffler (eds.) Antiinfectives Christian Doerig, Gordon Langsley, Pietro Alano (eds.) Malaria Signaling

Related Titles

Li, R., Stafford, J.A. (eds.)

Kinase Inhibitor Drugs

Print ISBN: 978-0-470-27829-1, also available as digital format

Klebl, B., Müller, G., Hamacher, M. (eds.)

Protein Kinases as Drug **Targets**

Print ISBN: 978-3-527-31811-7, also available as digital format

Ghosh, A.K. (ed.)

Aspartic Acid Proteases as Therapeutic Targets

Print ISBN: 978-3-527-31811-7, also available as digital format

Smit, M.J., Lira, S.A., Leurs, R. (eds.)

Chemokine Receptors as Drug **Targets**

Print ISBN: 978-3-527-32118-6, also available as digital format

Foreword: Protein Kinases in Parasites

Today, despite the fact that this is not obvious to the vast majority of the people leaving in industrialized countries, a large part of the world is still massively suffering and dying from parasitic diseases as a result of the lack of efficacious and/or affordable treatments. Each year 30,000 people pass away due to human African Trypanosomiasis (HAT or African sleeping sickness), a disease caused by the parasite Trypanosoma brucei spp. Available treatments for this disease are poor, with unacceptable efficacy and safety profiles, particularly in the late phase of the infection when the parasite has invaded the central nervous system. In South and Central America, Trypanosoma cruzi is the infectious agent of Chagas' disease (American Trypanosomiasis) which represents the most important parasitic infection in this part of the world. It is affecting more than 10 million people, with about 100 million people at risk. Leishmaniasis is due to the infection by protozoa of the genus Leishmania and is affecting more than 10 million people worldwide. These parasites live in the alimentary tract of blood-sucking sand flies, and as nonflagellate intracellular forms mostly within the macrophages of mammalian hosts. The severity of the disease is ranging from cutaneous and/or mucosal to visceral infection. Malaria occurs following infection by Plasmodium spp. and is the most prevalent parasitic disease, affecting more than 250 million of people per year and still responsible for almost a million deaths, the vast majority of which impacting children below 5 years. Not only unicellular parasites bear a huge impact on global public health: parasitic helminths (worms), such as Schistosoma ssp, also represent a serious public health problem, mostly in the developing world. In view of such a dramatic situation, more than ever, it is crucial that the entire scientific community in basic research and industries develops all possible strategies leading to an arsenal of therapeutic weapons that will efficiently treat patients and eradicate these diseases. Among possible drug targets, enzymes that modulate the level of phosphorylation of parasite and host proteins such as protein kinases (PKs) and protein phosphatases are interesting candidates.

First, the kinomes of parasites like kinetoplastids and apicomplexans could reveal promising taxon-specific drug targets. Indeed, signalling pathways are well known to allow any organism to adapt to its environment by coordinating intracellular processes. Bioinformatics approaches revealed a total of 176 PKs in *T. brucei*, 190 in

T. cruzi and 199 in L. major. Compared to trypanosomatids, the human kinome contains 3 times more protein kinases while the size of the *Plasmodium* kinome is only about half that of trypansomatids. Trypanosomatids and Plasmodium do not contain receptor-linked tyrosine kinases, but possess divergent kinases with no orthologues in the mammalian kinome (Ward P, Equinet L, Packer J, Doerig C. 2004. BMC Genomics; Parsons, M, Worthey E, Ward P, Mottram J. 2005. BMC Genomics). The fact that trypanosomatids exhibit a large set of PKs, covering approximately 2% of each genome, suggests that phosphorylation may play a key role in the biology of most parasites.

Despite differences in kinome sizes and composition from one parasite to another, major signalling pathways and functions are conserved. Motility, for instance, is an essential attribute that allows some parasites finding their target cells in human hosts and/or arthropod vectors. In apicomplexans, this key driving force depends on a unique component whereby adhesins contained in the micronemes are released onto the parasite apical extremity and translocated to the posterior end of the cell, thus propelling the parasite forward. In Toxoplasma gondii, Calcium-dependent protein kinase 1 (TgCDPK1) is an essential regulator of calciumdependent exocytosis and this could well be the case in most of the opportunistic human parasites. Recently, the phosphorylation of the alpha subunit of eukaryotic translation initiation factor 2 (eIF2α) was described elevated in dormant forms of apicomplexan parasites such as Plasmodium spp. and Toxoplasma gondii. (Zhang M. et al., Eukaryotic Cell, 2013).

Kinases have been shown to be essential for survival of parasites in their mammalian hosts. Nevertheless, a parasite kinase-specific small molecule inhibitor still awaits to be identified and the question whether drugging the parasite kinome is more a dream or a reality begs for an answer. In complex parasite life cycles such as that of *Plasmodium*, most of the key developmental forms of the parasite such as sexual and liver stages rely on protein kinase-mediated regulations as highlighted by C. Doerig (Nat. Chem. Biol.) commenting the work of Kato et al. who has demonstrated that PfCDPK1 plays a key role in asexual blood stage egress. Considering the 3-dimensional structure of protein kinases, there are increasing evidences that the ATP-binding pocket represents a druggable site. Specific kinomes like the one of Plasmodium display sufficient specificity, compared to the human one, to represent a potentially fertile source of novel targets. Interestingly, the counteracting biochemical reactions driven by Plasmodium phosphatases are similarly specific enough to envisage drug discovery programs targeting molecular events that are modulated by these enzymes (Wilkes and Doerig BMC genomics 2008).

Since helminths are Metazoan and have therefore a kinome that is very similar to that of their hosts, it is unlikely that highly selective targets will be identified. However, the kinome still remains an attractive target in this case to: precisely because of similarities between the helminthic and human kinomes, "piggy-back" approaches exploiting the wealth of resources devoted to targeting human kinases in the context of diseases such as cancer and neurodegenerative diseases is an attractive option as a strategy to combat diseases caused by worms.

Last but not least, the host-parasite interface might well be a target of choice to avoid induction of drug resistance and spreading. Plasmodium infection of host cells takes advantage of the plasticity of this parasite and the different forms produced along its complex life cycle. For instance during the infection of human hepatocytes (liver stage of Plasmodium's life cycle), not only part of the parasite kinome is solicited but some human protein kinases in liver cells such as MET, PRKWNK1, SGK2, STK35 and PKC\(\zeta\) seem to be crucial to Plasmodium sporozoite invasion mechanism and differentiation/growth (Prudêncio M. et al., 2008, PLoS Pathogens). Evidence is emerging that even in the erythrocyte, host signalling pathways are activated and required for parasite survival (Sicard et al 2011). Host protein kinases such as MEKs and downstream MAPKs may play a key role in the host immune response to *Plasmodium*. Indeed, these protein kinases have been shown to regulate the production of pro-inflammatory cytokines produced in response to specific markers of various infectious agents that may modulate the specificity and effectiveness of adaptive immunity. Thus, small molecules could be used as immunomodulatory tools to control pathogen infections and resulting diseases by regulating specific host protein kinases. (Zhu J. et al., 2009, J. Biol. Chem.).

In the present book, the bioinformatics approach leading to the study of parasite kinomes and phosphatomes will be described and followed by chapters addressing the functional analysis of some of the key enzymes. The potential roles of host cell kinome and phosphatome will be discussed. Finally, opportunities for drug discovery programs targeting parasite protein kinases and phosphatases will be explored in protozoan and helminthic parasites alike. There is no doubt that the holistic view described in this book will contribute to the future success of new efficacious and affordable therapeutics to treat the world population severely impacted by parasitic diseases.

Geneva, July 2013

Didier Leroy

Preface

Diseases caused by eukaryotic pathogens have been a scourge of human populations ever since the emergence of our species. Many of the major lineages of eukaryotes, from Excavata (Giardia), through Discicristata (Trypanosoma, Leishmania), Amoebozoa (Entamoeba) and Alveolata (Plasmodium, Toxoplasma, Eimeria), to Opisthokonta (metazoans, fungi), include species that have adapted to a parasitic lifestyle and have co-evolved with their hosts in the lineage that led to Homo sapiens. The burden imposed by parasitic diseases is disproportionally large in the poorest nations. While there has been immense progress in controlling some of these diseases in the second half of the XXth century, notably through the use of specific drugs, the global picture remains very gloomy: first, pathogens have responded to novel treatments by developing resistance against the drugs; thus, for example, that wonder antimalarial drug, chloroquine, has now become ineffective in a majority of malaria-affected countries. Even the latest generation of antimalarial drugs, based on artemisinin, shows signs of losing efficacy in some parts of the world. Second, many of these diseases have remained grossly neglected in terms of investment in research and development of novel control agents, largely because of the poor marketing prospects such agents would offer. Clearly, a renewed effort is urgently needed to address this global issue. Fortunately, awareness has increased in the last decade, which has led to an increase of funding from public institutions such as the European Commission and the research councils and agencies of many governments, as well as private bodies such as the Bill & Melinda Gates foundation. Furthermore, new organisational tools now exist to fund such research; for example, the Medicines for Malaria Venture (MMV, www.mmv.org), a Public-Private Partnership based in Geneva, and the Drugs for Neglected Diseases initiative (DNDi, www. dndi.org), have already had a tangible impact in this area. To eventually bring parasitic diseases under effective control, it is crucial that existing funding for fundamental research on eukaryotic pathogens is maintained and expanded, so as to prime the drug development pipeline. A high priority on the agenda is to develop control agents with un-tapped mechanisms of action.

Protein phosphorylation is an enormously important phenomenon in the biology of eukaryotic cells, where it regulates essentially all complex processes. This fundamental role has singled out protein kinases as potential targets for anticancer

agents, and indeed, a number of protein kinase inhibitors have reached the market in this context.

Could protein kinases represent targets for the treatment of parasitic diseases as well? A group of about 25 researchers interested in this idea convened in Paris in 2001, at the first EU-COST-funded meeting on "Protein kinases of eukaryotic parasites". This forum has reconvened in Glasgow in 2005, and in Lausanne in 2010. By then, the attending community had grown to 80 people, and significant progress had been achieved in (i) our fundamental understanding of the complement of parasite protein kinases and protein phosphatases (kinome and phosphatome) and the function of these enzymes in the biology of the parasites, and (ii) the identification of specific kinase targets in many eukaryotic parasites, and, in a few cases, of parasite kinase inhibitors. It was also emerging that the phosphorylation machinery in the host was playing a crucial role in parasite survival and development, suggesting that kinase inhibitors developed against cancer might be repositioned for the treatment of parasitic diseases.

The present book is an outcome of the 2010 meeting in Lausanne, and offers a written and updated version of some of the highlights that were presented there. It covers bio-informatics analyses of the kinomes and phosphatomes of selected eukaryotic parasites, recent advances in our fundamental understanding of the biology of selected kinases and phosphatases (inclusive of host signalling elements), and finally the state-of-the-art with respect to anti-parasitic drug discovery efforts targeting protein kinases.

We consider protein kinases offer huge potential for the development of urgently needed control agents against devastating diseases caused by eukaryotic parasites. This will happen only if the research community embraces the idea and constitutes compelling supporting data, so that policymakers and industrial partners can be convinced that there would be a significant return on investment in terms of impact on global public health. The purpose of this book is therefore to stimulate interest of established researchers and students in this topic, which offers a combination of both fascinating biology and potential tangible impact.

The Editors are aware of the significant additional commitment that engaging into the writing of a chapter represents in the busy life of research scientists, and are therefore very grateful to all authors for their timely contributions. We are indebted to the series editor, Dr. Paul M. Selzer of MSD Animal Health Innovation GmbH, for his constant encouragements and active involvement in the preparation of this volume, and to Anne du Guerny, Project Editor at Wiley Blackwell, for her patience and excellent support throughout the publication process.

Melbourne, Paris, and Glasgow September 2013 Christian Doerig, Gerald Späth, and Martin Wiese

Cover Legend

The cover is composed of several illustrations coming from or being related to the articles in this volume. The underlying phylogenetic tree illustrates the evolutionary relationships among eukaryotic species, including model organisms and protozoan pathogens, selected across all eukaryotic supergroups (courtesy of D. Miranda-Saavedra, see chapter 1 for details). The protein structure shows the homology model of EtCRK2 a CDK2-like protein of Eimeria tenella with ATP docked into the ATP binding pocket. The protein is shown as ribbons, while ATP is depicted in ball-and-stick representation with atoms colored according to the CPK model (courtesy of R. J. Marhöfer, see chapter 15 for details). The black matrix panel shows fluorescence microscopy images of different parasites. The top row of the panel shows an intra-erythrocytic Plasmodium falciparum schizont. The mitotic regulator Aurora kinase 3 is labeled in green, the Plasmodium homologue of centrosome protein Centrin-3 is labeled in red and the parasite DNA is stained in blue (courtesy of T. Carvalho, see chapter 1 and 13 for details). The middle row of the panel shows immunofluorescent staining of Trypanosoma brucei bloodstream forms. PKA-like kinase substrates are labeled in red, the paraflagellar rod protein in green as reference for the flagellum, and nuclear and kinetoplast DNA are stained blue with DAPI (courtesy of S. Bachmaier and M. Boshart, see chapter 5 for details). The lower row, from left to right, shows in the 1st image, eggs of Schistosoma mansoni purified form livers of infected hosts. Due to tyrosine-rich eggshell precursor proteins, which are fused via quinone tanning during eggshell synthesis, green and red auto-fluorescence is observed by fluorescence microscopy (courtesy of C. G. Grevelding, see chapter 16 for details). The 2nd image shows several human fibroblast cells with large blue nuclei massively infected with a transgenic strain of T. gondii tachyzoites visualized by small blue nuclei expressing GFP in its single mitochondrion. Cellular lipid bodies are stained red with Oil red O (courtesy of F. Seeber, Robert Koch Institute, Berlin, Germany). The 3rd image shows a section of a Schistosoma mansoni male worm labeled with anti-S. mansoni Insulin Receptor 1 antibodies. The antibody was localized at the basal membrane of the tegument, in muscles and in intestinal epithelium of worms (courtesy of C. Dissous, see chapter 16 for details). The 4th image shows an Echinococcus multilocularis protoscolex with DAPI/phalloidin staining (courtesy of K. Brehm, see chapter 17 for details).

List of Contributors

Roman Affentranger

Douglas Connect Baermeggenweg 14 4314 Zeiningen Switzerland

Merhnaz Amani

Structural Genomics Consortium MaRS South Tower, 7th Floor 101 College Street Toronto, Ontario Canada

Alexandra V. Andreeva*

University of Illinois at Chicago Department of Pharmacology 909 S. Wolcott Ave Chicago, IL 60612 USA alexandravandreeva@gmail.com

Gustavo Arrizabalaga

Indiana University School of Medicine Departments of Pharmacology & Toxicology, Microbiology & Immunology 635 Barnhill Drive, MS A-503 Indianapolis, IN 46202 USA

*Corresponding Author

Jennifer D. Artz

Structural Genomics Consortium MaRS South Tower, 7th Floor 101 College Street Toronto, Ontario Canada

Mrigya Babuta

Jawaharlal Nehru University School of Life Sciences New Delhi, 110067 India

Sabine Bachmaier

Ludwig-Maximilians-Universität München Fakultät für Biologie, Genetik Biozentrum Grosshadernerstr. 2-4 82152 Planegg-Martinsried Germany

Svenja Beckmann

Justus-Liebig-University Institute for Parasitology Rudolf-Buchheim-Str. 2 35392 Giessen Germany

Corinna Benz

University of South Bohemia Institute of Parasitology Biology Centre and Faculty of Sciences Česk Budějovice Czech Republic

Alok Bhattacharya*

Jawaharlal Nehru University School of Life Sciences New Delhi, 110067 India alok.bhattacharya@gmail.com; alok0200@mail.jnu.ac.in

Sudha Bhattacharya

Jawaharlal Nehru University School of Environmental Sciences New Delhi, 110067 India

Ira J. Blader*

University at Buffalo Department of Microbiology and Immunology 138 Farber Hall Buffalo, NY 14214 USA iblader@buffalo.edu

Michael Boshart*

Ludwig-Maximilians-Universität München Fakultät für Biologie, Genetik Biozentrum Grosshadernerstr. 2-4 82152 Planegg-Martinsried Germany boshart@lmu.de

Klaus Brehm*

University of Würzburg Institute of Hygiene and Microbiology Josef-Schneider-Strasse 2 97080 Würzburg Germany kbrehm@hygiene.uni-wuerzburg.de

James R. Brown

GlaxoSmithKline Computational Biology Quantitative Sciences, R&D 1250 South Collegeville Road, UP1230 Collegeville, PA 19426-0989 USA

Sharon D. Bryant

Inte:Ligand Software Development & Consulting GmbH Mariahilferstrasse 74B/11 1070 Vienna Austria

Christin Buro

Justus-Liebig-University Institute for Parasitology Rudolf-Buchheim-Str. 2 35392 Giessen Germany

Alessandro Contini

Università degli Studi di Milano Dipartimento di Scienze Farmaceutiche - Sezione di Chimica Organica "A. Marchesini" Via Venezian 21 20133 Milan Italy

Hugo Gutierrez de Teran

Uppsala University Department of Cell and Molecular Biology **BMC** 754 29 Uppsala Sweden

Colette Dissous*

Inserm U1019 CNRS UMR 8204 Center for Infection and Immunity of Lille (CIIL) Institut Pasteur de Lille 1, rue du Prof. Calmette 59019 Lille France Colette.dissous@pasteur-lille.fr

Dirk Dobbelaere

University of Bern Molecular Pathobiology Vetsuisse Faculty 3012 Bern Switzerland

Christian Doerig

Monash University Department of Microbiology Wellington Road Building 76 Clayton, Victoria 3800 Australia

David Drewry

GlaxoSmithKline Department of Chemical Biology 20 T. W. Alexander Drive Research Triangle Park Durham, NC 27709 USA

Francisco-Javier Gamo

GlaxoSmithKline Tres Cantos Medicines Development Campus Severo Ochoa 2 28760 Tres Cantos Spain

Jose F. Garcia-Bustos*

Monash University Department of Microbiology Clayton, Victoria 3800 Australia jose.garcia-bustos@monash.edu

Daniel E. Goldberg*

Washington University in St Louis Departments of Medicine and Molecular Microbiology Howard Hughes Medical Institute 660 S. Euclid Ave St Louis, MO 63110 USA goldberg@borcim.wustl.edu

Nadege Gouignard

Institut Pasteur de Lille Center for Infection and Immunity of Lille (CIIL) Inserm U1019 CNRS UMR 8204 59019 Lille France

Christoph G. Grevelding

Justus-Liebig-University Institute for Parasitology Rudolf-Buchheim-Str. 2 35392 Giessen Germany

Tansy C. Hammarton*

University of Glasgow College of Medical, Veterinary and Life Sciences Institute of Infection, Immunity & Inflammation 120 University Place Glasgow G12 8QQ UK Tansy. Hammarton@glasgow.ac.uk

Barry Hardy*

Douglas Connect Baermeggenweg 14 4314 Zeiningen Switzerland Barry.Hardy@douglasconnect.com

Tanya Hills

Structural Genomics Consortium MaRS South Tower, 7th Floor 101 College Street Toronto, Ontario Canada

Raymond Hui*

Structural Genomics Consortium MaRS South Tower. 7th Floor 101 College Street Toronto, Ontario Canada Raymond.hui@utoronto.ca

Natarajan Kannan

University of Georgia Institute of Bioinformatics Davison Life Sciences Bldg 120 Green Street Athens, GA 30602 USA

and

University of Georgia Department of Biochemistry and Molecular Biology B122 Life Sciences Bldg Athens, GA 30602 USA

Mikhail A. Kutuzov*

University of Illinois at Chicago Department of Pharmacology 909 S. Wolcott Ave Chicago, IL 60612 USA m.kutuzov@usa.net

Didier Leroy*

Director Drug Discovery Medicines for Malaria Venture Rte de Pre Bois 20, 1215 Geneva leroyd@mmv.org

Silke Leutner

Justus-Liebig-University Institute for Parasitology Rudolf-Buchheim-Str. 2 35392 Giessen Germany

Linda, Y. Lin

Structural Genomics Consortium MaRS South Tower, 7th Floor 101 College Street Toronto, Ontario Canada

Isabelle Lucet

Monash University Department of Microbiology Wellington Road Building 76 Clayton, Victoria 3800 Australia

Dustin J. Maly

University of Washington Department of Chemistry Chemistry Building Seattle, WA 98195 USA

Richard J. Marhöfer

MSD Animal Health Innovation **GmbH** Zur Propstei 55270 Schwabenheim Germany

Keith R. Matthews

University of Edinburgh School of Biological Sciences Institute of Immunology and Infection Research Centre for Immunity, Infection and Evolution King's Building, West Mains Road Edinburgh EH9 3JT UK

Ethan A. Merritt

University of Washington Department of Biochemistry 1959 NE Pacific Street Health Sciences Building Seattle, WA 98195 USA

Diego Miranda-Saavedra*

Institute of Cellular Medicine Newcastle University Medical School Framlington Place Newcastle upon, Tyne NE2 4HH UK diego@ifrec.osaka-u.ac.jp

Jeremy C. Mottram

University of Glasgow College of Medical, Veterinary and Life Sciences Institute of Infection, Immunity and Inflammation Wellcome Trust Centre for Molecular Parasitology 120 University Place Glasgow G12 8TA UK

Mirela Neculai

Structural Genomics Consortium MaRS South Tower, 7th Floor 101 College Street Toronto, Ontario Canada

Sandra Nelson

Chief Technology Officer PHD Diagnostics, LLC 632 Russell Street Covington, Kentucky 41011 USA

Victor Nussenzweig*

NYU Langone Medical Center Department of Pathology 550 1st Ave New York, NY 10016 USA Victor.Nussenzweig@nyumc.org

Kayode K. Ojo

University of Washington Department of Medicine Division of Allergy and Infectious Diseases 750 Republican Street Seattle, WA 98109 USA

Mahesh Kumar Padwal

National Centre for Cell Science Ganeshkhind Pune 411007 India

Ruben Papoian

University of Cincinnati Drug Discovery Center 2180 East Galbraith Road Cincinnati, OH 45237 USA

Bhaskar Saha*

National Centre for Cell Science Ganeshkhind Pune 411007 India sahab@nccs.res.in

Uddipan Sarma

National Centre for Cell Science Ganeshkhind Pune 411007 India

William L. Seibel

University of Cincinnati Drug Discovery Center Compound Library and Cheminformatics 2180 E. Galbraith Road Cincinnati, OH 45237 USA

Paul M. Selzer*

MSD Animal Health Innovation GmbH Zur Propstei 55270 Schwabenheim Germany paul.selzer@msd.de

and

University of Glasgow College of Medical, Veterinary and Life Sciences Institute of Infection, Immunity and Inflammation Wellcome Trust Centre for Molecular Parasitology 120 University Place Glasgow G12 8TA UK

and

University of Tübingen Interfaculty Institute of Biochemistry Hoppe-Seyler-Str. 4 72076 Tübingen Germany

Somlata

Jawaharlal Nehru University School of Life Sciences New Delhi, 110067 India

Jeff Spitzner

Amperand, Ltd. 311 Kendall PL Columbus, OH 43205 USA

Raki Sudan

National Centre for Cell Science Ganeshkhind Pune 411007 India