

Graduate Texts in Mathematics **1**

Wang Kunyang and Li Luoqing

Harmonic Analysis and Approximation on the Unit Sphere

(球面上的调和分析与逼近)



Science Press

Wang Kunyang and Li Luoqing

Harmonic Analysis and Approximation on the Unit Sphere

(球面上的调和分析与逼近)

Project supported by the National Natural Science Foundation

Book published by the support of the China Talent Fund



SCIENCE PRESS
Beijing/New York

Responsible Editor : Lü Hong

Copyright ©2000 by Science Press

Published by Science Press

16 Donghuangchenggen North Street

Beijing 100717, China

Printed in Beijing

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the copyright owner.

ISBN 7-03-008366-0 / O • 1216 (Beijing)

ISBN 1-880132-64-8 (New York)

Graduate Texts in Mathematics **1**

To our supervisor

Professor Sun Yongsheng

Preface

In recent years the research in Fourier Analysis and Approximation Theory has been extended from the classical setting, i.e., from the investigation on \mathbb{R}^n and \mathbb{T}^n , respectively, to the investigation on manifolds. Nikol'skii has published a series of papers in this respect. The unit sphere

$$\Omega_n := \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1^2 + \dots + x_n^2 = 1\}, \quad n \geq 2,$$

is a typical manifold in \mathbb{R}^n . Fourier analysis on Ω_n is also called Fourier-Laplace analysis since the Laplace-Beltrami operator takes an important role in almost all problems. When $n = 2$, we have $\Omega_2 = \mathbb{T}$ and get back to the classical case.

The investigation of Fourier-Laplace analysis has already a long history. The earliest research papers may have been published at the beginning of the 20th century. A basic lecture note is the one by C. Müller. There were also important papers we would like to point out. In 1968 Berens, Butzer and Pawelke studied basic approximation and saturation problems on the sphere. In 1973 Bonami and Clerc established important theorems on the convergence of Cesàro means of Fourier-Laplace series. Since 1980 the approximation of functions on the sphere has become an even more active field with the publication of the papers by Nikol'skii, Lizorkin, Kamzolov and others.

Why do analysts pay more attention to Ω_n recently? Here are two reasons. Firstly, although classical Fourier analysis on Ω_2 is getting more and more complete, quite a few problems on Ω_n for $n \geq 3$ are still left open. Secondly, the research on the sphere is in demand by practical problems in physics, geography, seismology, etc.

Since 1990 Professor Sun Yongsheng has urged his group to do research on the sphere, and under his guidance and encouragement his students started the research on Fourier-Laplace analysis and related problems on approximation. Since then the author's research has been supported by the NSF of China for the three periods, 1992~1994 (No. 19171008), 1995~1997 (19471007), and 1998~2000 (No.19771009).

The monograph is a summary of our research until 1998.

In order to make the book self-contained we wrote a preliminary first chapter. The second chapter provides basic knowledge on Fourier-Laplace series and some early research results.

In the third chapter, we present a kind of operators which are equiconvergent with Cesàro means with the same orders. These operators are simply convolutions with Jacobi polynomials as kernels. So it is convenient to investigate convergence problems with help of these operators. Also these operators can be applied to investigate general linear summability and strong summability of Fourier-Laplace series.

In the fourth chapter, we discuss how to describe the constructive properties of functions defined on the sphere. Then in Chapter 5 we give a detailed proof of Jackson inequality. This inequality says that for any function defined on the sphere, the best approximation by polynomials is dominated by its modulus of continuity. In L^p metric this inequality was established in 1987, but for L^1 and uniform metric this problem has been keeping open until 1994 the first author of the book and Riemenschneider found a constructive proof jointly.

In the last chapter, we discuss the problem of approximation by linear means such as Riesz means, Cesàro means and de la Vallée Poussin means.

The first five chapters are written originally by the first author and have been served as the material for his seminars. The last chapter (Chapter 6) is written by the second author.

We are sincerely grateful to our supervisor Professor Sun Yong Sheng. We are also grateful to Professor H.Berens, Professor G.Brown, Professor S.Riemenschneider and Professor Z.Ditzian. With them we have been collaborating since long and our research gets their kind concern and encouragement.

We thank the National Natural Science Foundation of China very much for the financial support during 1992~2000. Also we thank the China Talent Fund very much for the support to publish this book.

Contents

Chapter 1 Preliminaries	1
1.1 Basic concepts	1
1.1.1 Definition of μ_k^n , \mathcal{A}_k^n and \mathcal{H}_k^n	1
1.1.2 $L^2(\Omega_n)(n \geq 2)$	4
1.1.3 The case $n = 2$	6
1.1.4 Zonal harmonics	7
1.1.5 Representation for spherical harmonics	14
1.1.6 Laplace-Beltrami Operator	16
1.1.7 The convolution for functions on sphere	17
1.2 Gegenbauer and Jacobi polynomials	21
1.2.1 Rodrigues' formula	21
1.2.2 Funk-Hecke formula	24
1.2.3 Laplace representation	26
1.2.4 Generating formulas	27
1.2.5 The leading coefficient of P_k^n	31
1.2.6 Differential equations for P_k^n	31
1.2.7 Jacobi Polynomials	32
1.3 Jacobi polynomials with complex indices	34
Chapter 2 Fourier-Laplace Series	43
2.1 Introduction	43
2.2 Convergence, Lebesgue constant	45

2.3	Cesàro means(Early results)	48
2.4	Translation operator and mean operator	56
2.5	Maximal translation operator	63
2.5.1	The proof of Theorem 2.5.1	64
2.5.2	Proof of Theorem 2.5.2	71
2.6	Projection operators	75
Chapter 3	Equiconvergent Operators of Cesàro Means	85
3.1	Definition	85
3.2	Localization	95
3.2.1	The case $\delta \geq n - 2$	95
3.2.2	The necessity of “antipole conditions” when $-1 < \delta < n - 2$	97
3.2.3	Antipole conditions when $\frac{n-2}{2} - 1 < \delta < n - 2$...	100
3.2.4	Corollary of Theorems 3.2.5 and 3.2.6	105
3.3	Pointwise convergence	106
3.3.1	Equivalent conditions for convergence	106
3.3.2	Tests for convergence	108
3.3.3	A test of Salem type	113
3.4	Maximal operator E_* and a. e. convergence	116
3.5	Application to linear summability	129
3.5.1	Introduction	129
3.5.2	Auxiliary lemmas	131
3.5.3	Convergence everywhere	142
3.5.4	Convergence at Lebesgue points	150
Chapter 4	Constructive Properties of Spherical Functions	161
4.1	Best approximation operator	161
4.2	Pointwise Derivatives	162
4.2.1	Preliminary	163
4.2.2	Estimate for the tangent gradients	165
4.2.3	Estimate for the normal gradient of harmonic polynomials	168
4.3	Fractional derivative and integral	170
4.4	Fractional integrals of variable order	172

4.4.1	Definitions	172
4.4.2	Properties of Poisson integrals on the sphere	174
4.4.3	Proof of Theorem 4.4.1	180
4.5	Modulus of continuity	182
4.6	Derivatives and finite differences	188
Chapter 5	Jackson Type Theorems	193
5.1	Jackson inequality and K -functional	193
5.1.1	Estimates for ultraspherical polynomials	194
5.1.2	Estimates for the best approximation	212
5.1.3	Estimate for derivative of polynomials	214
5.1.4	Proof of Theorems 5.1.1 and 5.1.2	215
5.2	Difference $\ast \Delta_t^k$ and space H^r	216
Chapter 6	Approximation by Linear Means	229
6.1	Almost everywhere approximation	229
6.1.1	Introduction	229
6.1.2	Approximation by Riesz means on sets of full measure	231
6.1.3	Approximation by partial sums on sets of full measure	236
6.1.4	Strong approximation by Cesàro Means	241
6.2	Approximation in norm	254
6.2.1	Riesz means and Peetre K -moduli	254
6.2.2	Riesz means and the best approximation	257
6.2.3	Riesz means with critical index	259
6.2.4	Riesz means and Cesàro means	263
6.3	The de la Vallée Poussin Means	266
6.3.1	Convergence and approximation in norm	268
6.3.2	Pointwise convergence and approximation	270
6.3.3	Weak type inequalities for the best approximation	273
6.3.4	Characterization through a classical modulus of smoothness in C	275
6.3.5	Approximation for zonal functions	280
References	285
Index	299

Chapter 1

Preliminaries

The materials in the first two sections of this chapter are mostly taken from the books [SW], [M] and [Sz]. As for the concept of convolution of functions defined on the sphere we follow the argument in [D]. In section 1.2 we give a very basic introduction on Jacobi polynomials and particularly on Gegenbauer polynomials. We also introduce the normalized Gegenbauer polynomials from the point of view of the harmonic analysis on the sphere. This follows the argument in [M]. For a complete discussion on Jacobi polynomials we refer the reader to Szegő's famous book [Sz].

In the section 1.3 we will extend the Jacobi polynomials to the case of complex indices and give for such polynomials some asymptotic estimates. The results of this section have been given in [BW].

1.1 Basic concepts

1.1.1 Definition of \mathcal{P}_k^n , \mathcal{A}_k^n and \mathcal{H}_k^n

Let \mathbb{N} denote the set of natural numbers and let \mathbb{R}^n ($n \in \mathbb{N}$) be the n -dimensional Euclidean space with norm $|x| := \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}$ for $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. What we will treat with is complex valued functions defined on \mathbb{R}^n . As usual we denote by $\Delta = \Delta_n$ the Laplace operator, i.e.,

$$\Delta \equiv \Delta_n := \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \cdots + \frac{\partial^2}{\partial x_n^2}.$$

Suppose f is a function defined on \mathbb{R}^n . If f satisfies the Laplace equation, i.e., $\Delta_n f = 0$, we say that f is harmonic. If for all $\alpha \in \mathbb{C}$ and all $x \in \mathbb{R}^n$, $f(\alpha x) = \alpha^k f(x)$ with $k \in \mathbb{Z}_+ = \{0\} \cup \mathbb{N}$, then we say that f is homogeneous of degree k .

Definition 1.1.1 The set of all homogeneous polynomials of n variables of degree k is written as \mathcal{P}_k^n . The subset of all harmonic functions in \mathcal{P}_k^n is denoted by \mathcal{A}_k^n . An element of \mathcal{A}_k^n is called *solid spherical harmonic* (see [SW, p.141]).

When $n \geq 2$ we denote by Ω_n the unit spherical surface of \mathbb{R}^n , i.e.,

$$\Omega_n := \{\xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n : \xi_1^2 + \dots + \xi_n^2 = 1\}.$$

Definition 1.1.2 Suppose $f \in \mathcal{A}_k^n$ ($n \geq 2$). The restriction of f on Ω_n is called spherical harmonic of n variables of degree k . The set of all such functions is denoted by \mathcal{H}_k^n .

It is obvious that \mathcal{A}_k^n and \mathcal{H}_k^n are all linear spaces over complex scalar field \mathbb{C} and they have the same dimension which is denoted by a_k^n . In order to find the value of a_k^n we first consider the linear space \mathcal{P}_k^n . We denote the dimension of \mathcal{P}_k^n by d_k^n . Let $P \in \mathcal{P}_k^n$. Then P has the following representation:

$$P(x) = \sum_{\langle \alpha \rangle = k} c_\alpha x^\alpha, \quad \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_+^n, \quad c_\alpha \in \mathbb{C},$$

where $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$ and $\langle \alpha \rangle = \alpha_1 + \dots + \alpha_n$. From this we can deduce by induction on the dimensions that

$$d_k^n = C_{n-1+k}^{n-1} = \frac{(n+k-1)!}{k!(n-1)!}. \quad (1.1.1)$$

We now prove the following lemma from which the relation of d_k^n and a_j^n will be derived.

Lemma 1.1.1 For any $P \in \mathcal{P}_k^n$ ($n \geq 2$) there is a unique decomposition:

$$P(x) = \sum_{j=0}^{\ell} |x|^{2j} P_{k-2j}(x), \quad \ell = \left\lfloor \frac{k}{2} \right\rfloor, \quad (1.1.2)$$

where $P_{k-2j} \in \mathcal{A}_{k-2j}^n$ ($j = 0, 1, \dots, \ell$), and

$$d_k^n = \sum_{j=0}^{\ell} a_{k-2j}^n. \quad (1.1.3)$$

Proof We introduce an inner product on \mathcal{P}_k^n by defining

$$\langle P, Q \rangle_k := P(D) \overline{Q}, \quad \forall P, Q \in \mathcal{P}_k^n,$$

where $P(D)$ denotes the differential operator determined by P , while D^α is defined for any $\alpha \in \mathbb{Z}_+^n$ by

$$D^\alpha := \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}}.$$

It is easy to verify that $(\mathcal{P}_k^n, \langle \cdot, \cdot \rangle_k)$ is a complete inner product space. And the convergence in $(\mathcal{P}_k^n, \langle \cdot, \cdot \rangle_k)$ is just equivalent to the convergence of the coefficients of polynomials.

The cases $k = 0$ and $k = 1$ are trivial. We now assume $k \geq 2$ and define for $j \geq 2$

$$\mathcal{B}_j^n := |x|^2 \mathcal{P}_{j-2}^n = \{P(x) = |x|^2 Q(x) : Q \in \mathcal{P}_{j-2}^n\}.$$

Obviously, \mathcal{B}_j^n is a closed subspace of \mathcal{P}_j^n with inner product $\langle \cdot, \cdot \rangle_j$. We write \mathcal{B}_j^n as M temporarily. We are going to prove \mathcal{A}_j^n is just the orthogonal complement of M in \mathcal{P}_j^n .

Suppose $P \in \mathcal{P}_j^n$ and $R(x) = |x|^2 Q(x) \in M$. If $\langle P, R \rangle_j = 0$ then

$$\langle R, P \rangle_j = \Delta Q(D) \overline{P} = Q(D) \overline{\Delta P} = \langle Q, \Delta P \rangle_{j-2} = 0,$$

where $\Delta = \Delta_n$. If $\langle Q, \Delta P \rangle_{j-2} = 0$ holds for all $Q \in \mathcal{P}_{j-2}^n$ ($j \geq 2$) then $\Delta P = 0$. Hence we know $P \in \mathcal{A}_j^n$. This means $M^\perp \subset \mathcal{A}_j^n$.

Now let $P \in \mathcal{A}_j^n$ and $R(x) = |x|^2 Q(x) \in M$. Then

$$\langle P, R \rangle_j = \langle Q, \Delta P \rangle_{j-2} = 0.$$

So $M^\perp \supset \mathcal{A}_j^n$.

We have proved $M^\perp = \mathcal{A}_j^n$. Therefore by the orthogonal decomposition theorem we deduce that for any $P \in \mathcal{P}_k^n$ ($k \geq 2$) there are unique $P_0 \in \mathcal{A}_k^n$ and unique $Q \in \mathcal{P}_{k-2}^n$ such that

$$P(x) = P_0(x) + |x|^2 Q(x) \quad (1.1.4)$$

and obviously (for $k \geq 2$)

$$d_k^n = \dim \mathcal{A}_k^n + \dim \mathcal{B}_j^n = a_k^n + d_{k-2}^n. \quad (1.1.5)$$

By using (1.1.4) and (1.1.5) repeatedly for ℓ times we get (1.1.2) and (1.1.3) and complete the proof. \square

Observing that

$$d_0^n = a_0^n = 1 \quad \text{and} \quad d_1^n = a_1^n = n$$

and noticing (1.1.1) we get the following

Corollary 1.1.2 Let $n \geq 2$. Then

$$a_k^n = \begin{cases} 1, & \text{if } k = 0, \\ (n + 2k - 2) \frac{(n + k - 3)!}{k!(n - 2)!}, & \text{if } k \in \mathbb{N}. \end{cases} \quad (1.1.6)$$

Also we deduce from Lemma 1.1.1 the following useful

Corollary 1.1.3 The restriction on Ω_n of any polynomial is just a sum of some spherical harmonics.

1.1.2 $L^2(\Omega_n)$ ($n \geq 2$)

The space $L^2(\Omega_n)$ has naturally an inner product defined by

$$\langle f, g \rangle := \int_{\Omega_n} f(\xi) \overline{g(\xi)} d\omega_n(\xi), \quad (1.1.7)$$

where $d\omega_n$ denotes the surface element of Ω_n and the letter ξ in connection with $d\omega_n$ means that the integration is carried out with