



# Ions *in* Solution *and their* Solvation

YIZHAK MARCUS

WILEY

# IONS IN SOLUTION AND THEIR SOLVATION

---

**Yizhak Marcus**

*Hebrew University of Jerusalem*



**WILEY**

Copyright © 2015 by John Wiley & Sons, Inc. All rights reserved

Published by John Wiley & Sons, Inc., Hoboken, New Jersey

Published simultaneously in Canada

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at [www.copyright.com](http://www.copyright.com). Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permissions>.

**Limit of Liability/Disclaimer of Warranty:** While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at [www.wiley.com](http://www.wiley.com).

***Library of Congress Cataloging-in-Publication Data:***

Marcus, Y.

Ions in solution and their solvation / Yizhak Marcus.

pages cm

Includes indexes.

ISBN 978-1-118-88914-5 (cloth)

1. Ionic solutions. 2. Ions. 3. Solution (Chemistry) 4. Solvation. I. Title.

QD561 .M3365

541'.372-dc23

2015020563

Cover credits: Art courtesy of the author & Background Image: Roman Okopny/iStockphoto

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

1 2015

## **IONS IN SOLUTION AND THEIR SOLVATION**



# PREFACE

Thirty years ago, my book *Ion Solvation* was published, which became very popular and has been since then extensively cited, in particular for the numerical data concerning the properties of ions. However, much has happened during these years regarding experimental and computer simulation results pertinent to this subject as well as regarding physicochemical insights of the interactions that occur between ions and their surrounding solvents and other solutes too. It is therefore high time for an updated and revised edition of the early book to be available to scientists and engineers who deal with ions in solution. It is expected that the present book fulfills its mission to provide the needed up-to-date information.

It is appropriate to quote here some passages from the preface of my earlier book. “My main purpose ... on the one hand is to look back on what has been accomplished in a given field, organize it in an orderly fashion, and present it in a comprehensible and unified manner. On the other hand, I use this opportunity to locate gaps in our knowledge, and either fill these with new research while the book is being written, or to do this to the best of my ability in the course of time.” The latter purposes I still pursue in spite of a rather advanced age.

A further quotation from the preface of my earlier book is as true at the present time as, if not even more so than, at the time my earlier book was written. “University research nowadays ... is not conducive to the carrying out of extensive and systematic sets of measurements of high accuracy on the properties of systems. Both from the stand point of student interest and from that of the necessary financial support, work on systems that are not of immediate practical importance ... is not encouraged.” Still, impressive amounts of work of high quality are produced in universities too, and where relevant have found their way into the present book.

Numerical tables were an essential part of my earlier book on *ion solvation* and are also necessarily so in the present book on ions in solution and their solvation. These are interspersed in the places in the text where the quantities are being discussed, rather than being relegated to appendices, although the tables may disrupt the smooth flow of the text. Their provision, it is hoped, should make the present book as useful as a source book of data as was its predecessor. However, it must be stressed that there are some newer and corrected numerical data, for example, the CODATA thermochemical compilations or data examined by the European Commission on Atomic Energy that could not find their way into the preset book. Readers are also encouraged to look up recent volumes of the *Journal of Physical and Chemical Reference Data* for the latest critically examined values.

The significance of ions in solution and of their solvation is dealt with in the introductory chapter, which also describes the organization of the material in the chapters, as is also reflected in the list of contents.

It remains for me to express my appreciation of my university, of which I am now an emeritus professor for 15 years, which has provided me with the facilities required for doing nonlaboratory research and for writing reviews and books. The academic environment and discussions with colleagues here as well as at conferences that I attend, being supported by my university, are invaluable for doing what I have undertaken in this book.

Jerusalem, August 2014

# CONTENTS

<b>Preface</b>	<b>ix</b>
<b>1 Introduction</b>	<b>1</b>
1.1 The Significance and Phenomenology of Ions in Solution,	1
1.2 List of Symbols and Abbreviations,	5
<b>2 Ions and Their Properties</b>	<b>10</b>
2.1 Ions as Isolated Particles,	10
2.1.1 Bare Ions,	11
2.1.2 Ions in Clusters,	26
2.2 Sizes of Ions,	30
2.3 Ions in Solution,	35
2.3.1 Thermodynamics of Ions in Aqueous Solutions,	38
2.3.1.1 <i>Heat Capacities of Aqueous Ions,</i>	38
2.3.1.2 <i>Entropies of Aqueous Ions,</i>	39
2.3.1.3 <i>Enthalpies of Formation of Aqueous Ions,</i>	43
2.3.1.4 <i>Gibbs Energies of Formation of Aqueous Ions,</i>	44
2.3.1.5 <i>Ionic Molar Volumes in Aqueous Solutions,</i>	44
2.3.2 Other Properties of Aqueous Ions,	49
2.3.2.1 <i>Ionic Conductivities in Aqueous Solutions,</i>	49
2.3.2.2 <i>Ionic Self-Diffusion in Aqueous Solutions,</i>	50
2.3.2.3 <i>Ionic Effects on the Viscosity,</i>	51
2.3.2.4 <i>Ionic Effects on the Relaxation of NMR Signals,</i>	55
2.3.2.5 <i>Ionic Dielectric Decrements,</i>	55
2.3.2.6 <i>Ionic Effects on the Surface Tension,</i>	56
References,	58



<b>3</b>	<b>Solvents for Ions</b>	<b>63</b>
3.1	Solvent Properties that Suit Ion Dissolution,	63
3.2	Physical Properties of Solvents,	64
3.2.1	Volumetric Properties,	64
3.2.2	Thermodynamic Properties,	69
3.2.3	Electrical, Optical, and Magnetic Properties,	70
3.2.4	Transport Properties,	75
3.3	Chemical Properties of Solvents,	77
3.3.1	Structuredness,	77
3.3.2	Solvent Properties Related to Their Ion Solvating Ability,	80
3.3.2.1	<i>Polarity,</i>	81
3.3.2.2	<i>Electron Pair Donicity and Ability to Accept a Hydrogen Bond,</i>	83
3.3.2.3	<i>Hydrogen Bond Donicity and Electron Pair Acceptance,</i>	84
3.3.2.4	<i>Softness,</i>	85
3.3.3	Solvents as Acids and Bases,	86
3.3.4	Miscibility with and Solubility in Water,	88
3.3.5	Spectroscopic and Electrochemical Windows,	90
3.4	Properties of Binary Aqueous Cosolvent Mixtures,	90
3.4.1	Physical Properties of Binary Aqueous Mixtures with Cosolvents,	90
3.4.1.1	<i>Thermodynamic Properties of the Mixtures,</i>	92
3.4.1.2	<i>Some Electrical, Optical, and Transport Properties of the Mixtures,</i>	98
3.4.2	Chemical Properties of Binary Aqueous Mixtures with Cosolvents,	98
3.4.2.1	<i>Structuredness,</i>	98
3.4.2.2	<i>Properties Related to the Ion Solvating Ability,</i>	101
	References,	104
<b>4</b>	<b>Ion Solvation in Neat Solvents</b>	<b>107</b>
4.1	The Solvation Process,	107
4.2	Thermodynamics of Ion Hydration,	109
4.2.1	Gibbs Energies of Ion Hydration,	109
4.2.1.1	<i>Accommodation of the Ion in a Cavity,</i>	110
4.2.1.2	<i>Electrostatic Interactions,</i>	110
4.2.2	Entropies of Ion Hydration,	116
4.2.3	Enthalpies of Ion Hydration,	116
4.3	Transfer Thermodynamics into Nonaqueous Solvents,	117
4.3.1	Selection of an Extra-Thermodynamic Assumption,	117
4.3.2	Thermodynamics of Transfer of Ions into Nonaqueous Solvents,	118
4.3.2.1	<i>Gibbs Energies of Transfer,</i>	118
4.3.2.2	<i>Enthalpies of Transfer,</i>	126
4.3.2.3	<i>Entropies of Transfer,</i>	130

4.3.2.4	<i>Ionic Heat Capacities in Nonaqueous Solvents,</i>	130
4.3.2.5	<i>Ionic Volumes in Nonaqueous Solvents,</i>	133
4.4	The Structure of Solvated Ions,	135
4.4.1	Hydration Numbers from Diffraction Studies,	138
4.4.2	Hydration Numbers from Computer Simulations,	139
4.4.3	Hydration Numbers from Bulk Properties,	141
4.4.4	Solvation Numbers in Nonaqueous Solvents,	147
4.5	The Dynamics of Solvated Ions,	147
4.5.1	The Mobility of Ions in Solution,	147
4.5.2	Rate of Solvent Exchange Near Ions,	150
4.6	Acid/Base Properties of Ions in Solution,	151
	References,	153
<b>5</b>	<b>Mutual Effects of Ions and Solvents</b>	<b>156</b>
5.1	Ion Effects on the Structure of Solvents,	156
5.1.1	Experimental Studies of Ion Effects on the Structure of Solvents,	156
5.1.1.1	<i>Self-diffusion of Water Molecules,</i>	156
5.1.1.2	<i>Viscosity B-Coefficients,</i>	157
5.1.1.3	<i>NMR Signal Relaxation,</i>	159
5.1.1.4	<i>Dielectric Relaxation,</i>	159
5.1.1.5	<i>Vibrational Spectroscopy,</i>	160
5.1.1.6	<i>X-Ray Absorption and Scattering,</i>	162
5.1.1.7	<i>Structural Entropy,</i>	163
5.1.1.8	<i>Transfer from Light to Heavy Water,</i>	165
5.1.1.9	<i>Internal Pressure,</i>	168
5.1.1.10	<i>Some Other Experimental Results,</i>	170
5.1.2	Computer Simulations of Ion Effects on the Structure of Solvents,	170
5.2	Ion Effects on the Dynamics of the Solvent,	171
5.2.1	Mean Residence Times of Solvent Molecules Near Ions,	171
5.2.2	Experimental Studies of Ion Effects on the Solvent Orientation Dynamics,	174
5.2.2.1	<i>Ultrafast Infrared Spectroscopy,</i>	174
5.2.2.2	<i>High-frequency Dielectric Relaxation Spectroscopy,</i>	176
5.2.2.3	<i>NMR Relaxation Times,</i>	178
5.2.3	Computer Simulations of Reorientation Times,	180
5.3	Solvent Effects on the Properties of Ions in Solution,	180
5.3.1	Bulk Properties,	180
5.3.2	Molecular Properties,	186
	References,	187
<b>6</b>	<b>Ions in Mixed Solvents</b>	<b>193</b>
6.1	Ion Transfer into Solvent Mixtures,	194
6.2	Properties of Ions in Solvent Mixtures,	199
6.2.1	Thermodynamic Properties of Ions in Mixed Solvents,	199
6.2.2	Transport Properties of Ions in Mixed Solvents,	203

6.3	Preferential Solvation of Ions, 205	
6.3.1	Spectroscopic Studies, 207	
6.3.2	Results from Thermodynamic Data, 210	
6.3.2.1	<i>The QLQC Method, 211</i>	
6.3.2.2	<i>The IKBI Method, 213</i>	
6.3.2.3	<i>Treatments Based on Stepwise Solvent Replacements, 215</i>	
	References, 216	
<b>7</b>	<b>Interactions of Ions with Other Solutes</b>	<b>219</b>
7.1	Ion–Ion Interactions, 219	
7.1.1	Activity Coefficients of Electrolyte Solutions, 220	
7.1.2	Ion Hydration Related to Ion–Ion Interactions, 223	
7.2	Ion Association, 227	
7.2.1	Electrostatic Theory of Ion Association, 230	
7.2.1.1	<i>Activity Coefficients of Neutral Ion Pairs, 231</i>	
7.2.2	Methods for Studying Ion Association, 232	
7.2.3	Thermodynamic Quantities Pertaining to Ion Association, 234	
7.2.4	Aggregation of Ions in Solutions, 237	
7.3	Salting-in and Salting-out, 239	
7.3.1	Empirical Setschenow Constant Data, 240	
7.3.2	Interpretation of Salting Phenomena, 240	
	References, 244	
<b>8</b>	<b>Applications of Solutions of Ions</b>	<b>247</b>
8.1	Applications in Electrochemistry, 248	
8.1.1	Batteries and Supercapacitors, 248	
8.1.2	Solvent-Independent pH and Electrode Potential Scales, 251	
8.2	Applications in Hydrometallurgy, 257	
8.3	Applications in Separation Chemistry, 259	
8.3.1	Solvent Extraction of Alkali Metal Cations, 259	
8.3.2	Solvation of Ionizable Drug Molecules, 262	
8.4	Applications to Chemical Reaction Rates, 264	
8.5	Solvated Ions in Biophysical Chemistry, 269	
8.5.1	The Hofmeister Series, 270	
8.5.1.1	<i>The Anion Hofmeister Series, 270</i>	
8.5.1.2	<i>The Cation Hofmeister Series, 271</i>	
8.5.1.3	<i>Interpretation of the Hofmeister Series, 272</i>	
8.5.2	Water Structure Effects of Ions, 275	
8.5.3	Some Aspects of Protein Hydration, 277	
	References, 279	
	<b>Author Index</b>	<b>284</b>
	<b>Subject Index</b>	<b>294</b>

---

# 1

---

## INTRODUCTION

### 1.1 THE SIGNIFICANCE AND PHENOMENOLOGY OF IONS IN SOLUTION

Chemistry is for a large part conducted in solutions involving ions and such solutions are ubiquitous in nature. Oceans are vast aqueous solutions of salts, consisting mainly of sodium chloride, but other salts and minor components are also present in ocean water. Lakes, rivers, and brackish water are dilute solutions of ions and are essential to survival, since they provide drinking water and water for irrigation. Rain and other precipitates may remove ionic species from the atmosphere that arrived there as spray from oceans and seas or from human activities, for example, acid rain. Physiological fluids consist mostly of water in which colloidal substances, but also ions essential to their function, are dissolved.

It appears from the above that water is the only medium in which ions play a role, but this picture is too narrow because human endeavors utilize many other liquid media in which ions are present and have an active role. The manufacture of organic substances, as raw materials or intermediates in many industries, such as textiles, drugs, and food additives, generally involves reactions carried out in mixed aqueous-organic or completely nonaqueous liquid media in which ions participate. In chemical analysis, such media have long been of invaluable use, for instance in electroanalytical measurements or chromatographic separations. Industrial uses of nonaqueous media involving ions include solvent extraction in hydrometallurgy or in nuclear fuel reprocessing and nuclear waste disposal (Chapter 8).

The ions involved in these as well as other systems and applications interact with each other and with nonionized solutes that may be present. These interactions are of prime interest to the chemist, but the extent, intensity, and rate of proceeding of these interactions depend heavily on the solvent or solvent mixture present, a fact that is not always clearly recognized by the operator. The ion-solvent interactions should be understood in order to make the best use of the solutions of the ions, since it is the solvated ions that take part in the interactions of interest. If a free choice of the solvent or the solvent mixture to be used is possible, the most suitable one for the purpose should be selected on the basis of the knowledge available on the interactions that take place, bearing in mind also costs, ecology, and hazards. If the solvent is prescribed, this knowledge is still needed in order to select the proper reaction conditions or the additives that could be useful. So-called "bare" or nonsolvated ions occur in gas-phase reactions (Chapter 2) but not in condensed media, that is, in solutions. A seeming exception to this generalization is the use of room temperature ionic liquids (RTILs) as reaction media, where all the ions are surrounded by ions of the opposite charge sign rather than by a nonionic liquid medium. Whether the ions of RTILs are called "bare" or "solvated" is a semantic question. In common situations, which are the subject of this book, there is always an excess of a nonionic liquid medium in which the ions find themselves, the molecules of which surround the ions more or less completely, unless some other species, be it another ion (of opposite sign) or a solute molecule (a ligand) replaces some of the solvent molecules in the ionic solvation shell.

Ions cannot be added individually to any major extent to a solvent or a solution; it is always an electrolyte consisting of ions of both signs in a combination that makes the electrolyte electrically neutral, which is added to form a solution involving ions. Many commonly used and studied electrolytes are crystalline solids, such as NaCl or  $(C_4H_9)_4NClO_4$ . The electrostatic energy that holds the ions constituting such crystals together, the lattice energy that must be invested in order to separate the ions in the solution, is compensated by the solvation energy that is gained in the process of dissolution, with some effect also of the entropic changes encountered in the process. Some potential electrolytes are gaseous, for example, HCl, but they produce ions only on reaction with the solvent in which the covalent H-Cl bonds are broken and replaced with others to compensate for the energy involved. On the other hand, ions may leave the solution, if not individually then as a small combination of ions, in electrospray experiments, in which they are then monitored in the gas phase by mass spectrometry. The results of such experiments have some bearing on the state of the liquid ionic solutions, but this subject is outside the scope of this book.

It should be kept in mind that, connected with such ion solvation reactions with crystalline or gaseous electrolytes, a further reaction takes place, which is not always recognized, namely the breaking of some solvent-solvent molecular interactions, required to produce the space to accommodate the ions in the solution. It is the balance of all the (Gibbs) energies that have to be invested and those that are gained that determine the extent to which an electrolyte will dissolve in a given solvent (Chapter 4).

Some electrolytes are completely dissociated into “free,” that is, solvated, positively charged cations, and negatively charged anions. Other electrolytes are only partly so dissociated, depending on the concentration and on the nature of the solvent. Some substances are ionogenic, in the sense that some dissociation into ions occurs only under specific conditions, and these include also so-called “weak electrolytes” such as many acids and essentially basic substances in aqueous solutions.

A solution of a single, individual ion in a very large amount of solvent could presumably be the basis of a study of ion solvation which is not encumbered by other interactions. This situation cannot be achieved in the laboratory but can be dealt with as a thought process and now for many years also in computer simulations. The results of the latter have by now consolidated into a large body of knowledge that is constantly not only extended but also improved by the level of sophistication that can nowadays be achieved in such simulations. Such results are incorporated into the discussions in the present book, where they are compared with laboratory experimental data obtained on electrolytes, extrapolated to infinite dilution (Chapters 4 and 5). At such high dilutions, each ion is surrounded by solvent molecules only and does not interact with the very remote ions of the opposite charge sign that must be present somewhere in the solution. Still, the allocation of the extrapolated values of the electrolyte properties to its constituent individual ions is a problem that must be solved.

For the purpose of only reducing the number of items in the properties list from the many electrolytes (combination of cations and anions), that have been derived at infinite dilution to the much smaller number of individual ions that constitute them, it is sufficient to employ the so-called “conventional” values. These are based on assigning to one ion, say the solvated hydrogen ion, an arbitrary value (generally zero) and rely on the additivity of individual infinite dilution ionic values to derive values of all other ions. The sum of the conventional ionic values, weighted according to the stoichiometric coefficients (the numbers of ions of each kind constituting the electrolyte), expresses correctly the infinite dilution property of the electrolyte. Within a given charge sign series of ions, say cations only, comparisons between conventional values of diverse ions can throw some light on the effects of the individual ionic properties, such as size and valency, but the cation and the anion series cannot be compared with each other.

The problem of assignment of the so-called “absolute” individual ionic values to these infinite dilution electrolyte data is solved mainly on the basis of chemical intuition (Chapter 4) that can be assisted by the results from computer simulations. Once individual ionic values of their properties in a given solvent (or solvent mixture) at a given thermodynamic state [temperature and pressure, usually specified as 298.15 K (25°C) and 0.1 MPa (less commonly now 1 atm=0.101325 MPa)] have been established, they may be compared with other properties of the ions (e.g., their sizes) or with theoretical expectations (models). The latter are the main incentives to obtaining the absolute values. Such comparisons and correlations provide insights into the ion–solvent interactions that take place and form the basis for understanding interactions of ions with other solutes, be they ionic themselves or nonionic.

There are some experimental measurements that can be made on solutions of ions that pertain directly to individual ions. These include transport properties, such as the

ionic conductivities that are obtained from specific conductivities of electrolytes in conjunction with transport number measurements. Diffusivities of individual ionic species can also be measured by the use of isotopically labeled ions and should be compatible with the mobilities deduced from the ionic conductivities. Spectroscopic data can also in certain cases be due to individual ionic species, such as NMR chemical shifts and relaxation rates of the signals from appropriate nuclei (e.g.,  $^7\text{Li}$  or  $^{27}\text{Al}$ ). Such information may be used as a guide for the "chemical intuition" mentioned earlier needed for obtaining absolute individual ionic values from measurements on electrolytes extrapolated to infinite dilution.

As already mentioned at the beginning, nonaqueous and mixed aqueous-organic solvents play important roles in chemistry, but water is still the most studied solvent for ions, not only at ambient conditions but also under diverse conditions of temperature, pressure, and the existence of external fields, such as electrical ones. Their solvation by water and their properties in aqueous solutions are therefore useful as a reference basis for evaluation of the effects occurring on the exchange of the water, in part or completely, by another solvent. The situation is complicated by the fact that water is a unique solvent in many respects, and effects encountered in aqueous solutions of ions may be absent in other media. For instance, the effects that ions have on the hydrogen-bonded structure of water, breaking it or enhancing it, have no counterpart in most other solvents, but there are exceptions. The so called "hydrophobic interactions" applicable to ionic species with extended alkyl chains or aromatic rings are practically unique for aqueous solutions. These interactions do not allude only to "model" ions such as  $(\text{C}_4\text{H}_9)_4\text{N}^+$  but more importantly to biomolecules in general and specifically also to side chains of proteins that are ionized due to the  $-\text{CO}_2^-$  or  $-\text{NH}_3^+$  groups that they carry.

The ion solvation efficacy of nonaqueous solvents, alone or mixed with water, has beneficial results for some uses, such as enhancement of the solubility of ionic substances, but may be detrimental in other aspects, for example, in the availability of ions as reaction partners. Poorly solvated ions are more reactive than strongly solvated ones and this is manifested in the rates of organic reactions. Aprotic solvents (such that do not provide hydrogen bonding) have been found as optimal media for reactions involving anions that are only poorly solvated by them. Nonpolar solvents, on the other hand, have a low solvating power for the ions and therefore play a very minor role in solutions of ions, because of the low solubility of electrolytes in such solvents. Polar solvents, whether protic or aprotic, interact with ions of both charge signs by means of their dipole moments. Protic solvents carry a hydrogen atom capable of hydrogen-bonding to anions whereas both protic and aprotic ones provide a pair of nonbonded electrons to form coordinate bonds with mainly the cations (Chapter 3). The electron pair donor-acceptor properties of the solvents are generally of more importance than the dipole moments of their molecules. The permittivity of the solvent, important for the ionic dissociation of the electrolytes, plays a minor role in the solvation of the ions or properties of the solvated ions, once formed in the solution.

Mixture of solvents poses additional problems to the study of ion solvation and the properties of the solvated ions: the interactions between the molecules of the different solvents with each other besides those of molecules of each solvent among

themselves are significant. Also, the selective solvation of the ions by the molecules of the component solvents may affect the solvated ions profoundly: the ions may “see” around them nearly only molecules of the favored component (selective solvation). The preferential solvation of ions in solvent mixture needs to be described quantitatively, and once this is done should be explained in terms of both the solvating properties of the solvents and the properties of the ions (Chapter 6).

Much emphasize has been provided in this introduction to solution of electrolytes at infinite dilution, where individual ionic properties are manifested and may be obtained and discussed. However, “real life” encounters solutions of ions that involve finite, sometimes quite large, concentrations of ions, where interactions of the ions among themselves are important. The number of solvent molecules per formula unit of the electrolyte diminishes with increasing concentrations, and hence the integrity of the ionic solvation shells is eventually broken. The ion–ion interactions then compete with the ion–solvent interactions. Ion pairs, consisting of couples of ions of opposite charge sign, may have transient existence, but an equilibrium concentration of them may result and needs to be taken into account. Similarly, the presence of non-ionic solutes (already referred to earlier text, where solvent mixtures are considered), which may be solid, liquid, or gaseous solutes, poses new forms of interactions that have to be dealt with (Chapter 7).

Several aspects of ion solvation and the properties of solvated ions are dealt with in this introduction. On the whole, one goal of the discussion of ion solvation is the provision of small sets of properties of the ions and of the solvents, from which the solvation can be predicted for any ion/solvent combination. This has been attempted in this book, with a view to be useful for the many applications of ions in solution, examples of which are shown in Chapter 8, with no attempt to exhaust this subject.

## 1.2 LIST OF SYMBOLS AND ABBREVIATIONS

Chemical species and units of physical quantities are denoted by Roman type characters, whereas physical quantities that can be expressed by numerical values are denoted by Greek or *italic* characters. Mathematical symbols have their usual meaning and are not listed here. The same symbol is used for an extensive property of a system and for the molar quantity of a constituent of the system. The SI system of physical units is used throughout, but some extra SI units commonly used in the physicochemical literature are also included where they simplify the notation. These include the symbols °C for centigrade temperatures ( $T/K - 273.15$ ), M for mol·dm<sup>-3</sup>, and m for mol (kg·solvent)<sup>-1</sup>.

## PRINCIPAL LATIN CHARACTERS

- A<sup>z-</sup> a generalized anion
- A coefficient in the Debye–Hückel expression for activity coefficients
- AN acceptor number of a solvent
- a* distance of closest approach of ions in solution (nm)



$a_x$	thermodynamic activity of species X
(aq)	an ion in aqueous solution, generally at infinite dilution
$B$	coefficient in the Debye–Hückel expression for activity coefficients
$B_\eta$	$B$ -coefficient of the Jones–Dole viscosity expression ( $M^{-1}$ )
$b$	coefficient of the expression of the electric field dependence of the permittivity
$b$	parameter in the Bjerrum expression for ion pairing
$C^{z+}$	a generalized cation
$C_p$	molar heat capacity at constant pressure ( $J \cdot K^{-1} \cdot mol^{-1}$ )
$c_x$	molar concentration of species X (M)
(cr)	crystalline phase
$D$	Debye unit of dipole moments ( $3.33564 \times 10^{-30} C \cdot m$ )
$D$	diffusion coefficient ( $m^2 \cdot s^{-1} \cdot mol^{-1}$ )
$DN$	donor number of a solvent
$d$	interatomic distance (nm)
$E$	generalized electrolyte
$E$	energy, molar energy ( $J \cdot mol^{-1}$ )
$E$	electric field strength ( $V \cdot m^{-1}$ )
$E$	electromotive force of an electrochemical cell (V)
$E_j$	liquid junction potential
$E_T$	polarity index of a solvent ( $kcal \cdot mol^{-1}$ , $1 cal = 4.184 J$ )
$e$	elementary charge ( $1.6022 \times 10^{-19} C$ )
$F$	Faraday constant ( $9.6485 \times 10^4 C \cdot mol^{-1}$ )
$f_x$	fraction of species X
$G$	Gibbs energy, molar Gibbs energy ( $J \cdot mol^{-1}$ )
$g$	Kirkwood dipole orientation parameter
(g)	gas phase
$g(r)$	pair correlation function
$H$	enthalpy, molar enthalpy ( $J \cdot mol^{-1}$ )
$h$	solvation (hydration) number
$I^{\pm}$	generalized ion
$I$	ionic strength (M or m)
$K$	equilibrium constant
$K_a, K_b$	acid, base dissociation constant in aqueous solutions
$K_{ass}$	ion pair association constant (M)
$K_w$	ion product of water ( $M^2$ )
$k_B$	Boltzmann constant ( $1.3807 \times 10^{-23} J \cdot K^{-1}$ )
$k$	rate constant of specified reaction ( $s^{-1}$ for unimolecular reactions, $M^{-1} \cdot s^{-1}$ for bimolecular reactions)
(l)	liquid phase
$M^{z+}$	metal ion of charge $z+$
$M_x$	molar mass of species X (in $kg \cdot mol^{-1}$ )
$m_x$	molal concentration of species X (m)
$N$	generalized nonelectrolyte solute
$N_A$	Avogadro's number ( $6.0221 \times 10^{23} mol^{-1}$ )