

Kung-Ching Chang

Springer  
Monographs  
in  
Mathematics

# Methods in Nonlinear Analysis

非线性分析方法



Springer

世界图书出版公司  
[www.wpcbj.com.cn](http://www.wpcbj.com.cn)

Kung-Ching Chang

# Methods in Nonlinear Analysis



**Kung-Ching Chang**

School of Mathematical Sciences  
Peking University  
100871 Beijing  
People's Republic of China  
E-mail: kcchang@math.pku.edu.cn

Library of Congress Control Number: 2005931137

Mathematics Subject Classification (2000): 47H00, 47J05, 47J07, 47J25, 47J30, 58-01, 58C15, 58E05, 49-01, 49J15, 49J35, 49J45, 49J53, 35-01

ISSN 1439-7382

ISBN-10 3-540-24133-7 Springer Berlin Heidelberg New York

ISBN-13 978-3-540-24133-1 Springer Berlin Heidelberg New York

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer. Violations are liable for prosecution under the German Copyright Law.

Springer is a part of Springer Science+Business Media  
springeronline.com

© Springer-Verlag Berlin Heidelberg 2005

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Reprint from English language edition:

Methods in Nonlinear Analysis

by Kung-Ching Chang

Copyright © 2005, Springer-Verlag Berlin Heidelberg

Springer-Verlag Berlin Heidelberg is a part of Springer Science+Business Media

All Rights Reserved

This reprint has been authorized by Springer Science & Business Media for distribution in China Mainland only and not for export therefrom.

***Springer Monographs in Mathematics***



---

## Preface

Nonlinear analysis is a new area that was born and has matured from abundant research developed in studying nonlinear problems. In the past thirty years, nonlinear analysis has undergone rapid growth; it has become part of the mainstream research fields in contemporary mathematical analysis.

Many nonlinear analysis problems have their roots in geometry, astronomy, fluid and elastic mechanics, physics, chemistry, biology, control theory, image processing and economics. The theories and methods in nonlinear analysis stem from many areas of mathematics: Ordinary differential equations, partial differential equations, the calculus of variations, dynamical systems, differential geometry, Lie groups, algebraic topology, linear and nonlinear functional analysis, measure theory, harmonic analysis, convex analysis, game theory, optimization theory, etc. Amidst solving these problems, many branches are intertwined, thereby advancing each other.

The author has been offering a course on nonlinear analysis to graduate students at Peking University and other universities every two or three years over the past two decades. Facing an enormous amount of material, vast numbers of references, diversities of disciplines, and tremendously different backgrounds of students in the audience, the author is always concerned with how much an individual can truly learn, internalize and benefit from a mere semester course in this subject.

The author's approach is to emphasize and to demonstrate the most fundamental principles and methods through important and interesting examples from various problems in different branches of mathematics. However, there are technical difficulties: Not only do most interesting problems require background knowledge in other branches of mathematics, but also, in order to solve these problems, many details in argument and in computation should be included. In this case, we have to get around the real problem, and deal with a simpler one, such that the application of the method is understandable. The author does not always pursue each theory in its broadest generality; instead, he stresses the motivation, the success in applications and its limitations.

## VI Preface

The book is the result of many years of revision of the author's lecture notes. Some of the more involved sections were originally used in seminars as introductory parts of some new subjects. However, due to their importance, the materials have been reorganized and supplemented, so that they may be more valuable to the readers.

In addition, there are notes, remarks, and comments at the end of this book, where important references, recent progress and further reading are presented.

The author is indebted to Prof. Wang Zhiqiang at Utah State University, Prof. Zhang Kewei at Sussex University and Prof. Zhou Shulin at Peking University for their careful reading and valuable comments on Chaps. 3, 4 and 5.

Peking University  
September, 2003

*Kung Ching Chang*

---

# Contents

<b>1</b>	<b>Linearization</b>	<b>1</b>
1.1	Differential Calculus in Banach Spaces	1
1.1.1	Frechet Derivatives and Gateaux Derivatives	2
1.1.2	Nemytski Operator	7
1.1.3	High-Order Derivatives	9
1.2	Implicit Function Theorem and Continuity Method	12
1.2.1	Inverse Function Theorem	12
1.2.2	Applications	17
1.2.3	Continuity Method	23
1.3	Lyapunov-Schmidt Reduction and Bifurcation	30
1.3.1	Bifurcation	30
1.3.2	Lyapunov-Schmidt Reduction	33
1.3.3	A Perturbation Problem	43
1.3.4	Gluings	47
1.3.5	Transversality	49
1.4	Hard Implicit Function Theorem	54
1.4.1	The Small Divisor Problem	55
1.4.2	Nash-Moser Iteration	62
<b>2</b>	<b>Fixed-Point Theorems</b>	<b>71</b>
2.1	Order Method	72
2.2	Convex Function and Its Subdifferentials	80
2.2.1	Convex Functions	80
2.2.2	Subdifferentials	84
2.3	Convexity and Compactness	87
2.4	Nonexpansive Maps	104
2.5	Monotone Mappings	109
2.6	Maximal Monotone Mapping	120



<b>3</b>	<b>Degree Theory and Applications</b>	127
3.1	The Notion of Topological Degree	128
3.2	Fundamental Properties and Calculations of Brouwer Degrees	137
3.3	Applications of Brouwer Degree	148
3.3.1	Brouwer Fixed-Point Theorem	148
3.3.2	The Borsuk-Ulam Theorem and Its Consequences	148
3.3.3	Degrees for $S^1$ Equivariant Mappings	151
3.3.4	Intersection	153
3.4	Leray-Schauder Degrees	155
3.5	The Global Bifurcation	164
3.6	Applications	175
3.6.1	Degree Theory on Closed Convex Sets	175
3.6.2	Positive Solutions and the Scaling Method	180
3.6.3	Krein-Rutman Theory for Positive Linear Operators	185
3.6.4	Multiple Solutions	189
3.6.5	A Free Boundary Problem	192
3.6.6	Bridging	193
3.7	Extensions	195
3.7.1	Set-Valued Mappings	195
3.7.2	Strict Set Contraction Mappings and Condensing Mappings	198
3.7.3	Fredholm Mappings	200
<b>4</b>	<b>Minimization Methods</b>	205
4.1	Variational Principles	206
4.1.1	Constraint Problems	206
4.1.2	Euler-Lagrange Equation	209
4.1.3	Dual Variational Principle	212
4.2	Direct Method	216
4.2.1	Fundamental Principle	216
4.2.2	Examples	217
4.2.3	The Prescribing Gaussian Curvature Problem and the Schwarz Symmetric Rearrangement	223
4.3	Quasi-Convexity	231
4.3.1	Weak Continuity and Quasi-Convexity	232
4.3.2	Morrey Theorem	237
4.3.3	Nonlinear Elasticity	242
4.4	Relaxation and Young Measure	244
4.4.1	Relaxations	245
4.4.2	Young Measure	251
4.5	Other Function Spaces	260
4.5.1	BV Space	260
4.5.2	Hardy Space and BMO Space	266
4.5.3	Compensation Compactness	271
4.5.4	Applications to the Calculus of Variations	274

4.6	Free Discontinuous Problems .....	279
4.6.1	$\Gamma$ -convergence .....	279
4.6.2	A Phase Transition Problem .....	280
4.6.3	Segmentation and Mumford–Shah Problem .....	284
4.7	Concentration Compactness .....	289
4.7.1	Concentration Function .....	289
4.7.2	The Critical Sobolev Exponent and the Best Constants .....	295
4.8	Minimax Methods .....	301
4.8.1	Ekeland Variational Principle .....	301
4.8.2	Minimax Principle .....	303
4.8.3	Applications .....	306
<b>5</b>	<b>Topological and Variational Methods .....</b>	<b>315</b>
5.1	Morse Theory .....	317
5.1.1	Introduction .....	317
5.1.2	Deformation Theorem .....	319
5.1.3	Critical Groups .....	327
5.1.4	Global Theory .....	334
5.1.5	Applications .....	343
5.2	Minimax Principles (Revisited) .....	347
5.2.1	A Minimax Principle .....	347
5.2.2	Category and Ljusternik–Schnirelmann Multiplicity Theorem .....	349
5.2.3	Cap Product .....	354
5.2.4	Index Theorem .....	358
5.2.5	Applications .....	363
5.3	Periodic Orbits for Hamiltonian System and Weinstein Conjecture .....	371
5.3.1	Hamiltonian Operator .....	373
5.3.2	Periodic Solutions .....	374
5.3.3	Weinstein Conjecture .....	376
5.4	Prescribing Gaussian Curvature Problem on $S^2$ .....	380
5.4.1	The Conformal Group and the Best Constant .....	380
5.4.2	The Palais–Smale Sequence .....	387
5.4.3	Morse Theory for the Prescribing Gaussian Curvature Equation on $S^2$ .....	389
5.5	Conley Index Theory .....	392
5.5.1	Isolated Invariant Set .....	393
5.5.2	Index Pair and Conley Index .....	397
5.5.3	Morse Decomposition on Compact Invariant Sets and Its Extension .....	408
	<b>Notes .....</b>	<b>419</b>
	<b>References .....</b>	<b>425</b>



## Linearization

The first and the easiest step in studying a nonlinear problem is to linearize it. That is, to approximate the initial nonlinear problem by a linear one. Nonlinear differential equations and nonlinear integral equations can be seen as nonlinear equations on certain function spaces. In dealing with their linearizations, we turn to the differential calculus in infinite-dimensional spaces. The implicit function theorem for finite-dimensional space has been proved very useful in all differential theories: Ordinary differential equations, differential geometry, differential topology, Lie groups etc. In this chapter we shall see that its infinite-dimensional version will also be useful in partial differential equations and other fields; in particular, in the local existence, in the stability, in the bifurcation, in the perturbation problem, and in the gluing technique etc. This is the contents of Sects. 1.2 and 1.3. Based on Newton iterations and the smoothing operators, the Nash–Moser iteration, which is motivated by the isometric embeddings of Riemannian manifolds into Euclidean spaces and the KAM theory, is now a very important tool in analysis. Limited in space and time, we restrict ourselves to introducing only the spirit of the method in Sect. 1.4.

### 1.1 Differential Calculus in Banach Spaces

There are two kinds of derivatives in the differential calculus of several variables, the gradients and the directional derivatives. We shall extend these two to infinite-dimensional spaces.

Let  $X$ ,  $Y$  and  $Z$  be Banach spaces, with norms  $\|\cdot\|_X$ ,  $\|\cdot\|_Y$ ,  $\|\cdot\|_Z$ , respectively. If there is no ambiguity, we omit the subscripts. Let  $U \subset X$  be an open set, and let  $f : U \rightarrow Y$  be a map.

### 1.1.1 Fréchet Derivatives and Gateaux Derivatives

**Definition 1.1.1** (*Fréchet derivative*) Let  $x_0 \in U$ ; we say that  $f$  is Fréchet differentiable (or F-differentiable) at  $x_0$ , if  $\exists A \in L(X, Y)$  such that

$$\|f(x) - f(x_0) - A(x - x_0)\|_Y = o(\|x - x_0\|_X).$$

Let  $f'(x_0) = A$ , and call it the Fréchet (or F-) derivative of  $f$  at  $x_0$ .

If  $f$  is F-differentiable at every point in  $U$ , and if  $x \mapsto f'(x)$ , as a mapping from  $U$  to  $L(X, Y)$ , is continuous at  $x_0$ , then we say that  $f$  is continuously differentiable at  $x_0$ . If  $f$  is continuously differentiable at each point in  $U$ , then we say that  $f$  is continuously differentiable on  $U$ , and denote it by  $f \in C^1(U, Y)$ .

Parallel to the differential calculus of several variables, by definition, we may prove the following:

1. If  $f$  is F-differentiable at  $x_0$ , then  $f'(x_0)$  is uniquely determined.
2. If  $f$  is F-differentiable at  $x_0$ , then  $f$  must be continuous at  $x_0$ .
3. (Chain rule) Assume that  $U \subset X, V \subset Y$  are open sets, and that  $f$  is F-differentiable at  $x_0$ , and  $g$  is F-differentiable at  $f(x_0)$ , where

$$U \xrightarrow{f} V \xrightarrow{g} Z$$

Then

$$(g \circ f)'(x_0) = g' \circ f(x_0) \cdot f'(x_0).$$

**Definition 1.1.2** (*Gateaux derivative*) Let  $x_0 \in U$ ; we say that  $f$  is Gateaux differentiable (or G-differentiable) at  $x_0$ , if  $\forall h \in X, \exists df(x_0, h) \in Y$ , such that

$$\|f(x_0 + th) - f(x_0) - tdf(x_0, h)\|_Y = o(t) \text{ as } t \rightarrow 0$$

for all  $x_0 + th \in U$ . We call  $df(x_0, h)$  the Gateaux derivative (or G-derivative) of  $f$  at  $x_0$ .

We have

$$\frac{d}{dt}f(x_0 + th) \big|_{t=0} = df(x_0, h),$$

if  $f$  is G-differentiable at  $x_0$ .

By definition, we have the following properties:

1. If  $f$  is G-differentiable at  $x_0$ , then  $df(x_0, h)$  is uniquely determined.
2.  $df(x_0, th) = tdf(x_0, h) \quad \forall t \in \mathbb{R}^1$ .
3. If  $f$  is G-differentiable at  $x_0$ , then  $\forall h \in X, \forall y^* \in Y^*$ , the function  $\varphi(t) = \langle y^*, f(x_0 + th) \rangle$  is differentiable at  $t = 0$ , and  $\varphi'(t) = \langle y^*, df(x_0, h) \rangle$ .
4. Assume that  $f : U \rightarrow Y$  is G-differentiable at each point in  $U$ , and that the segment  $\{x_0 + th \mid t \in [0, 1]\} \subset U$ , then

$$\|f(x_0 + h) - f(x_0)\|_Y \leq \sup_{0 < t < 1} \|df(x_0 + th, h)\|_Y$$

*Proof.* Let

$$\varphi_{y^*}(t) = \langle y^*, f(x_0 + th) \rangle \quad t \in [0, 1], \forall y^* \in Y^*$$

$$\begin{aligned} |\langle y^*, f(x_0 + h) - f(x_0) \rangle| &= |\varphi_{y^*}(1) - \varphi_{y^*}(0)| \\ &= |\varphi'_{y^*}(t^*)| \\ &= |\langle y^*, df(x_0 + t^*h, h) \rangle| \end{aligned}$$

for some  $t^* \in (0, 1)$  depending on  $y^*$ . The conclusion follows from the Hahn-Banach theorem.  $\square$

5. If  $f$  is F-differentiable at  $x_0$ , then  $f$  is G-differentiable at  $x_0$ , with  $df(x_0, h) = f'(x_0)h \quad \forall h \in X$ .

Conversely it is not true, but we have:

**Theorem 1.1.3** Suppose that  $f : U \rightarrow Y$  is G-differentiable, and that  $\forall x \in U, \exists A(x) \in L(X, Y)$  satisfying

$$df(x, h) = A(x)h \quad \forall h \in X.$$

If the mapping  $x \mapsto A(x)$  is continuous at  $x_0$ , then  $f$  is F-differentiable at  $x_0$ , with  $f'(x_0) = A(x_0)$ .

*Proof.* With no loss of generality, we assume that the segment  $\{x_0 + th \mid t \in [0, 1]\}$  is in  $U$ . According to the Hahn-Banach theorem,  $\exists y^* \in Y^*$ , with  $\|y^*\| = 1$ , such that

$$\|f(x_0 + h) - f(x_0) - A(x_0)h\|_Y = \langle y^*, f(x_0 + h) - f(x_0) - A(x_0)h \rangle.$$

Let

$$\varphi(t) = \langle y^*, f(x_0 + th) \rangle.$$

From the mean value theorem,  $\exists \xi \in (0, 1)$  such that

$$\begin{aligned} |\varphi(1) - \varphi(0) - \langle y^*, A(x_0)h \rangle| &= |\varphi'(\xi) - \langle y^*, A(x_0)h \rangle| \\ &= |\langle y^*, df(x_0 + \xi h, h) - A(x_0)h \rangle| \\ &= |\langle y^*, [A(x_0 + \xi h) - A(x_0)]h \rangle| \\ &= o(\|h\|), \end{aligned}$$

i.e.,  $f'(x_0) = A(x_0)$ .  $\square$

The importance of Theorem 1.1.3 lies in the fact that it is not easy to write down the F-derivative for a given map directly, but the computation of G-derivative is reduced to the differential calculus of single variables. The same situation occurs in the differential calculus of several variables: Gradients are reduced to partial derivatives, and partial derivatives are reduced to derivatives of single variables.

*Example 1.* Let  $A \in L(X, Y)$ ,  $f(x) = Ax$ . Then  $f'(x) = A \quad \forall x$ .

*Example 2.* Let  $X = \mathbb{R}^n$ ,  $Y = \mathbb{R}^m$ , and let  $\varphi_1, \varphi_2, \dots, \varphi_m \in C^1(\mathbb{R}^n, \mathbb{R}^1)$ . Set

$$f(x) = \begin{pmatrix} \varphi_1(x) \\ \vdots \\ \varphi_m(x) \end{pmatrix}, \text{ i.e., } f : X \rightarrow Y.$$

Then

$$f'(x_0) = \left( \frac{\partial \varphi_i(x_0)}{\partial x_j} \right)_{m \times n}.$$

*Example 3.* Let  $\Omega \subset \mathbb{R}^n$  be an open bounded domain. Denote by  $C(\overline{\Omega})$  the continuous function space on  $\overline{\Omega}$ . Let

$$\varphi : \overline{\Omega} \times \mathbb{R}^1 \longrightarrow \mathbb{R}^1,$$

be a  $C^1$  function. Define a mapping  $f : C(\overline{\Omega}) \rightarrow C(\overline{\Omega})$  by

$$u(x) \mapsto \varphi(x, u(x)).$$

Then  $f$  is F-differentiable, and  $\forall u_0 \in C(\overline{\Omega})$ ,

$$(f'(u_0) \cdot v)(x) = \varphi_u(x, u_0(x)) \cdot v(x) \quad \forall v \in C(\overline{\Omega}).$$

*Proof.*  $\forall h \in C(\overline{\Omega})$

$$t^{-1}[f(u_0 + th) - f(u_0)](x) = \varphi_u(x, u_0(x) + t\theta(x)h(x))h(x),$$

where  $\theta(x) \in (0, 1)$ .  $\forall \varepsilon > 0$ ,  $\forall M > 0$ ,  $\exists \delta = \delta(M, \varepsilon) > 0$  such that

$$|\varphi_u(x, \xi) - \varphi_u(x, \xi')| < \varepsilon, \quad \forall x \in \overline{\Omega},$$

as  $|\xi|, |\xi'| \leq M$  and  $|\xi - \xi'| \leq \delta$ . We choose  $M = \|u_0\| + \|h\|$ , then for  $|t| < \delta < 1$ ,

$$|\varphi_u(x, u_0(x) + t\theta(x)h(x)) - \varphi_u(x, u_0(x))| < \varepsilon.$$

It follows that  $df(u_0, h)(x) = \varphi_u(x, u_0(x))h(x)$ .

Noticing that the multiplication operator  $h \mapsto A(u)h = \varphi_u(x, u(x)) \cdot h(x)$  is linear and continuous, and the mapping  $u \mapsto A(u)$  from  $C(\overline{\Omega})$  into  $L(C(\overline{\Omega}), C(\overline{\Omega}))$  is continuous, from Theorem 1.1.3,  $f$  is F-differentiable, and

$$(f'(u_0) \cdot v)(x) = \varphi_u(x, u_0(x)) \cdot v(x) \quad \forall v \in C(\overline{\Omega}).$$

□

We investigate nonlinear differential operators on more general spaces. Let  $\Omega \subset \mathbb{R}^n$  be a bounded open set, and let  $m$  be a nonnegative integer,  $\gamma \in (0, 1)$ .  $C^m(\bar{\Omega})$  (and the Hölder space  $C^{m,\gamma}(\bar{\Omega})$ ) is defined to be the function space consisting of  $C^m$  functions (with  $\gamma$ -Hölder continuous  $m$ -order partial derivatives).

The norms are defined as follows:

$$\|u\|_{C^m} = \max_{x \in \bar{\Omega}} \sum_{|\alpha| \leq m} |\partial^\alpha u(x)|,$$

and

$$\|u\|_{C^{m,\gamma}} = \|u\|_{C^m} + \max_{x,y \in \bar{\Omega}} \sum_{|\alpha|=m} \frac{|\partial^\alpha u(x) - \partial^\alpha u(y)|}{|x-y|^\gamma},$$

where  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$  is a multi-index,  $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$ ,  $\partial^\alpha = \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \dots \partial_{x_n}^{\alpha_n}$ .

We always denote by  $m^*$  the number of the index set  $\{\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \mid |\alpha| \leq m\}$ , and  $D^m u$  the set  $\{\partial^\alpha u \mid |\alpha| \leq m\}$ .

Suppose that  $r$  is a nonnegative integer, and that  $\varphi \in C^\infty(\bar{\Omega} \times \mathbb{R}^{r^*})$ . Define a differentiable operator of order  $r$ :

$$f(u)(x) = \varphi(x, D^r u(x)).$$

Suppose  $m \geq r$ , then  $f : C^m(\bar{\Omega}) \rightarrow C^{m-r}(\bar{\Omega})$  (and also  $C^{m,\gamma}(\bar{\Omega}) \rightarrow C^{m-r,\gamma}(\bar{\Omega})$ ) is F-differentiable. Furthermore

$$(f'(u_0)h)(x) = \sum_{|\alpha| \leq r} \varphi_\alpha(x, D^r u_0(x)) \cdot \partial^\alpha h(x), \quad \forall h \in C^m(\bar{\Omega}),$$

where  $\varphi_\alpha$  is the partial derivative of  $\varphi$  with respect to the variable index  $\alpha$ .

The proof is similar to Example 3.

*Example 4.* Suppose  $\varphi \in C^\infty(\bar{\Omega} \times \mathbb{R}^{r^*})$ . Define

$$f(u) = \int_{\bar{\Omega}} \varphi(x, D^r u(x)) dx \quad \forall u \in C^r(\bar{\Omega}).$$

Then  $f : C^r(\bar{\Omega}) \rightarrow \mathbb{R}^1$  is F-differentiable. Furthermore

$$\langle f'(u_0), h \rangle = \int_{\bar{\Omega}} \sum_{|\alpha| \leq r} \varphi_\alpha(x, D^r u_0(x)) \partial^\alpha h(x) dx \quad \forall h \in C^r(\bar{\Omega}).$$

*Proof.* Use the chain rule:

$$C^r(\bar{\Omega}) \xrightarrow{\varphi(\cdot, D^r u(\cdot))} C(\bar{\Omega}) \xrightarrow{\int_{\bar{\Omega}}} \mathbb{R}^1,$$

and combine the results of Examples 1 and 3. □



In particular, the following functional occurs frequently in the calculus of variations ( $r = 1, r^* = n + 1$ ). Assume that  $\varphi(x, u, p)$  is a function of the form:

$$\varphi(x, u, p) = \frac{1}{2}|p|^2 + \sum_{i=1}^n a_i(x)p_i + a_0(x)u,$$

where  $p = (p_1, p_2, \dots, p_n)$ , and  $a_i(x)$ ,  $i = 0, 1, \dots, n$ , are in  $C(\overline{\Omega})$ .

Set

$$f(u) = \int_{\Omega} \left[ \frac{1}{2} |\nabla u(x)|^2 + \sum_{i=1}^n a_i(x) \partial_{x_i} u + a_0(x) u(x) \right] dx,$$

we have

$$\langle f'(u), h \rangle = \int_{\Omega} \left[ \nabla u(x) \cdot \nabla h(x) + \sum_{i=1}^n a_i(x) \partial_{x_i} h(x) + a_0(x) h(x) \right] dx$$

$$\forall h \in C^1(\overline{\Omega}).$$

*Example 5.* Let  $X$  be a Hilbert space, with inner product  $(\cdot, \cdot)$ . Find the F-derivative of the norm  $f(x) = \|x\|$ , as  $x \neq \theta$ .

Let  $F(x) = \|x\|^2$ . Since

$$t^{-1}(\|x + th\|^2 - \|x\|^2) = 2(x, h) + t\|h\|^2,$$

we have  $dF(x, h) = 2(x, h)$ . It is continuous for all  $x$ , therefore  $F$  is F-differentiable, and

$$F'(x)h = 2(x, h).$$

Since  $f = F^{\frac{1}{2}}$ , by the chain rule

$$F'(x) = 2\|x\| \cdot f'(x).$$

As  $x \neq \theta$ ,

$$f'(x)h = \left( \frac{x}{\|x\|}, h \right).$$

In the applications to PDE as well as to the calculus of variations, Sobolev spaces are frequently used. We should extend the above studies to nonlinear operators defined on Sobolev spaces.

$\forall p \geq 1$ ,  $\forall$  nonnegative integer  $m$ , let

$$W^{m,p}(\Omega) = \{u \in L^p(\Omega) \mid \partial^\alpha u \in L^p(\Omega) \mid |\alpha| \leq m\},$$

where  $\partial^\alpha u$  stands for the  $\alpha$ -order generalized derivative of  $u$ , i.e., the derivative in the distribution sense. Define the norm

$$\|u\|_{W^{m,p}} = \left( \sum_{|\alpha| \leq m} \|\partial^\alpha u\|_{L^p(\Omega)}^p \right)^{\frac{1}{p}}.$$

The Banach space is called the Sobolev space of index  $\{m, p\}$ .

$W^{m,2}(\Omega)$  is denoted by  $H^m(\Omega)$ , and the closure of  $C_0^\infty(\Omega)$  under this norm is denoted by  $H_0^m(\Omega)$ .