


METHODOLOGICAL SURVEYS IN BIOCHEMISTRY AND ANALYSIS
Series Editor: Eric Reid • Volume 12

Drug Metabolite Isolation and Determination



Edited by

Eric Reid and J. P. Leppard

Drug Metabolite Isolation and Determination

METHODOLOGICAL SURVEYS IN BIOCHEMISTRY AND ANALYSIS

Series Editor: Eric Reid

Guildford Academic Associates
72 The Chase
Guildford GU2 5UL, United Kingdom

The series is divided into Subseries A: Analysis, and B: Biochemistry
Enquiries concerning Volumes 1-11 should be sent to the above address.

Volumes 1-10 edited by Eric Reid

Volume 1 (B): Separations with Zonal Rotors

Volume 2 (B): Preparative Techniques

Volume 3 (B): Advances with Zonal Rotors

Volume 4 (B): Subcellular Studies

Volume 5 (A): Assay of Drugs and Other Trace Compounds in Biological Fluids

Volume 6 (B): Membranous Elements and Movement of Molecules

Volume 7 (A): Blood Drugs and Other Analytical Challenges

Volume 8 (B): Cell Populations

Volume 9 (B): Plant Organelles

Volume 10 (A): Trace-Organic Sample Handling

Volume 11 (B): Cancer-Cell Organelles
edited by Eric Reid, G. M. W. Cook, and D. J. Morré

Volume 12 (A): Drug Metabolite Isolation and Determination
edited by Eric Reid and J. P. Leppard
(includes a cumulative compound-type index)

A Continuation Order Plan is available for this series. A continuation order will bring delivery of each new volume immediately upon publication. Volumes are billed only upon actual shipment. For further information please contact the publisher.

Drug Metabolite Isolation and Determination

Edited by

Eric Reid

and

J. P. Leppard

Formerly of the University of Surrey

PLENUM PRESS • NEW YORK AND LONDON

Library of Congress Cataloging in Publication Data

Main entry under title:

Drug metabolite isolation and determination.

(Methodological surveys in biochemistry and analysis. Subseries A, Analysis, v. 12)

Based on the Fourth International Bioanalytical Forum, held Sept. 7-10, 1981, at the University of Surrey, Guildford, United Kingdom.

Includes bibliographical references and indexes.

1. Drugs—Analysis—Congresses. 2. Drugs—Metabolism—Congresses. I. Reid, Eric, 1922— II. Leppard, J. P. III. International Bioanalytical Forum (4th: 1981: University of Surrey) IV. Series.

RS189.D784 1983

615.7'028

83-2214

ISBN 0-306-41265-9

Based on the Fourth International Bioanalytical Forum held September 7-10, 1981, at the University of Surrey, Guildford, United Kingdom

©1983 Plenum Press, New York

A Division of Plenum Publishing Corporation

233 Spring Street, New York, N.Y. 10013

All rights reserved

No part of this book may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, microfilming, recording, or otherwise, without written permission from the Publisher

Printed in the United States of America

Senior Editor's Preface

In drug metabolism, as in kindred fields, sound conclusions hinge on good methodology with special skills which, however, have had less than their share of recognition and 'kudos'. This series of books is earning regard for its description and encouragement of good methods, not least because of emphasis on difficulties and on rationale rather than recipes such as prevail in the literature.

Each book is based on a Forum - in this case the 4th Bioanalytical Forum held at the University of Surrey in September 1981 - and mirrors some of the cut-and-thrust of the debates, but is not to be regarded as a symposium record. There has in fact been strong editing, partly for the sake of clarity and of relevance to analytical practice (thus, pharmacokinetic data have been excluded). The aim has been to produce a balanced desk book, well cross-referenced and complementing earlier 'Analysis' volumes, whose material on particular analytes is cumulatively re-indexed here (Index of Compound Types); the concluding article and the General Index aid retrieval of lore on problems and approaches.

The evolution of the *Methodological Surveys* twin series (outline facing title page) has undergone yet another discontinuity, in respect of publisher and format. For Vols. 10 and 11, conventional typesetting by the publisher was obligatory; it led to a generally good appearance, although there were errors that the page-proofing should have eliminated, but it contributed to the slowness of publication. Moreover, it entailed an intolerable editorial burden, now reduced by reversion to 'camera-ready' text produced at the Guildford end. The initiative of switching to Plenum Press, with marketing and other strengths and with pricing that appeals to American and other customers, has entailed sad parting from the U.K. publisher whose involvement could have started with Vol. 5 (illness precluded this) and did start with Vol. 6. If the now widening readership can get past volumes back-ordered, this will help not only bibliographic searching but also future Forum financing. Here too there is a discontinuity: from 1983 each Forum will be under the auspices of a new Trust (an 'educational charity'), although the normal venue will still be the nearby campus of the well-liked University of Surrey. Hopefully there will still be company support, which for the 1981 Forum came from Ciba-Geigy (U.K.), Glaxo and ICI Pharmaceuticals. It has been a boon that many contributors made little claim on Forum funds.

Scope of the book, and acknowledgements.— The articles are focused on 'real problems' in body-fluid analysis, typically with a final chromatographic separation of μg or often ng amounts if the aim is quantitation. The pitfalls may not be realized by a typical chemist (cf. remarks in #A-3), but he may excel in metabolite identification — which this book covers to a fair extent. Where identity is known, and the metabolite could interfere in therapeutic drug monitoring or in diagnosis or itself have clinical relevance, useful guidance will come from articles that follow. Authors have gone to much trouble, and are not to be blamed by any reader who would have liked an introduction to chromatography or to metabolic pathways (cf. list of conjugation reactions at end of concluding article). Appreciation is also expressed for permission to reproduce published material; the acknowledged sources include *J. Chromatog.* (Elsevier; e.g. in #A-1), *Anal. Chem.* (American Chemical Society; #A-2) and Wiley.

HPLC nomenclature.— With regard to the terms 'NP' and 'RP' (normal-, reverse-phase) the Editors have respected authors' preferences (cf. comment following Table 2 in #A-2), but share misgivings expressed by J.H. Knox and (in *LC in Practice*) by P.A. Bristow, who writes: "Reverse phase is a term which could well be retired gracefully". Thus, an 'uncapped' packing with a low content of bonded alkyl groups may behave adsorptively. Some authors exclude unbonded ('straight') silica from the 'NP' category that includes polar liquid stationary phases and polar bonded phases such as cyano. The designations 'hydrophobic' and 'hydrophilic' chromatography (Knox) have merit, and the concept of 'surface zone' rather than 'phase'. Some comment on so-called ion-pairing is made in #E (cf. p. 91).

Various abbreviations.— Nowadays the terms TLC (HPTLC = 'high performance' variant) and HPLC need no definition. In GC (gas-liquid in the present context), detector types include FID = flame ionization, AFID = alkali flame ionization ('nitrogen detector') which can detect phosphorus (hence NPD, N-PD), and ECD = electron capture but unfortunately connoting electrochemical detection in HPLC work. Mass spectrometry is denoted MS; EI = electron-impact, and CI = chemical ionization. In HPLC, t_R (preferred; otherwise R_t) = retention time.

Non-chromatographic abbreviations include i.s. = internal standard (usually 'processed'; see #A-6 in Vol. 7), RIA = radioimmunoassay, UV = ultraviolet [absorptiometry], IR = infrared, and NMR = nuclear magnetic resonance.

An example of metabolite jargon is a dialkylated amine drug, 'D', that has lost one alkyl group: it might variously be termed de(s)alkyld, monodesalkyld, desmonoalkyld, norD..... A conjugate is 'Phase II'.

Guildford Academic Associates
72 The Chase, Guildford GU2 5UL
Surrey, U.K.

ERIC REID
1 April 1982

List of Authors

'Notes' (headed 'NC.....' in the text) are not distinguished from main contributions.

- | | |
|---|--|
| C.D. Bevan - pp. 111-118
as for J. Chamberlain | W. Dieterle - pp. 13-22
Ciba-Geigy, Switzerland |
| A. Bhatti - pp. 181-187
as for J. Chamberlain | B.F.H. Drenth - pp. 75-80
as for R.A. de Zeeuw |
| G.R. Bourne - pp. 97-102
ICI Pharmaceuticals, U.K. | C.V. Eadsforth - pp. 119-133
Shell Res., U.K. |
| M.A. Brooks - pp. 201-206, 225-230
as for J.A.F. de Silva | J.W. Faigle - pp. 13-22
as for W. Dieterle |
| T.A. Bryce - pp. 181-187
as for J. Chamberlain | Y.Y.Z. Farid - pp. 141-142
as for M.J. Stewart |
| J. Caldwell - pp. 161-179
St. Mary's Hosp. Med. School,
London | |
| J. Chamberlain - pp. 111-118,
181-187, 249
Hoechst Pharmaceutical Res. Labs.,
U.K. | R.T. Ghijsen - pp. 75-80
as for R.A. de Zeeuw |
| Anne Chambers - pp. 103-110
as for M.J. Stewart | B.L. Goodwin - pp. 47-53, 71-73
Queen Charlotte's Hosp., London |
| F.G. Coppin - p. 249
as for J. Chamberlain | M.R. Hackman - pp. 201-206
as for J.A.F. de Silva |
| | N.G.L. Harding - pp. 141-142
as for M.J. Stewart |
| H. de Bree - p. 69
Duphar Res. Labs., The Netherlands | M.P. Harrison - pp. 191-196
ICI Pharmaceuticals, U.K. |
| D. Dell - p. 249
Hofmann-La Roche, Switzerland | A.J. Hutt - pp. 161-179
as for J. Caldwell |
| J.A.F. de Silva - pp. 201-206,
225-230.
Hofmann-La Roche, U.S.A. | J.B. Houston - pp. 207-214
Univ. of Manchester, U.K. |
| R.A. de Zeeuw - pp. 75-80
State Univ., Groningen,
The Netherlands | H.P.A. Illing - pp. 181-187
as for J. Chamberlain |
| | T. Jagersma - pp. 75-80
as for R.A. de Zeeuw |
| | D.R. Jarvie - pp. 103-110
Royal Infirmary, Edinburgh, U.K. |

- C.R. Jones - pp. 33-40
Wellcome Res. Labs., U.K.
- L.J. King - pp. 247-248
Univ. of Surrey, U.K.
- Mary V. Marsh - pp. 161-179
as for J. Caldwell
- S.R. Moss - pp. 97-102
as for G.R. Bourne
- H. Munzer - pp. 215-223
as for J. Rosenthaler
- R.G. Muusze - pp. 241-242
Algemeen Psychiatrisch Inst.,
Delft, The Netherlands
- E.A.M. Neill - pp. 243-246
Wellcome Res. Labs., U.K.
- F. Overzet - pp. 75-80
as for R.A. de Zeeuw
- J. Pao - pp. 225-230
as for J.A.F. de Silva
- E. Reid - pp. 255-270
Univ. of Surrey, U.K.
- J.C. Rhodes - pp. 207-214
as for J.B. Houston
- W. Ritter - pp. 55-62, 231-238
Bayer, Wuppertal, W. Germany
- D.R. Roberts - pp. 81-88
ICI Pharmaceuticals, U.K.
- J.D. Robinson - pp. 111-118
as for J. Chamberlain
- J. Rosenthaler - pp. 215-223
Sandoz, Basle, Switzerland
- K.A. Sinclair - pp. 161-179
as for J. Caldwell
- G.G. Skellern - pp. 3-11
Univ. of Strathclyde, U.K.
- M.J. Stewart
Royal Infirmary, Glasgow, U.K.
- D.A. Stopher - pp. 65-67
Pfizer Central Res., U.K.
- Jelka Tomašić - pp. 137-139, 149-160
'Ruder Bošković' Inst., Yugoslavia
- R. Voges - pp. 215-223
as for J. Rosenthaler
- I.D. Watson - pp. 103-110, 141-142
Royal Infirmary, Glasgow, U.K.
- R.E. Weinfeld - pp. 201-206
as for J.A.F. de Silva
- I.D. Wilson - pp. 111-118, 181-187
as for J. Chamberlain

Contents

The NOTES & COMMENTS ('NC' items) at the end of each section comprise some comments made at the Forum on which the book is based, together with some supplementary points besides those in the concluding survey.

	Senior Editor's Preface (with comments on HPLC nomenclature, and list of abbreviations)	v
	List of Authors	xi
#A	TECHNIQUES APPLICABLE TO METABOLITE INVESTIGATION	1
#A-1	Strategies centered on HPLC - G.G. SKELLERN	3
#A-2	Preparative liquid-chromatographic methods in drug metabolism studies - W. DIETERLE & J.W. FAIGLE	13
#A-3	Problems arising from assaying low concentrations by HPLC - C.R. JONES	33
#A-4	Extraction of drugs and metabolites from plasma and urine - M.J. STEWART	41
#A-5	The use of long capillary GC columns for urinary phenolic acids from catecholamines and analogues - B.L. GOODWIN	47
#A-6	Quantitative HPTLC in assaying body fluids for drugs and metabolites - W. RITTER	55
#NC(A)	NOTES and COMMENTS related to the foregoing topics	63†
	<i>including Notes on</i>	
#NC(A)-1	TLC densitometry for drug and metabolite determination - D.A. STOPHER	65
#NC(A)-2	Taking aliquots from small volumes of upper phases - H. DE BREE	69
#NC(A)-3	Rapid removal of fine particles from alumina for column chromatography - B.L. GOODWIN	71

†Sub-listed here: comments on particular articles in the section; likewise on pp. 135, 189 & 231.

#NC(A)-4	Assessment of radioactivity measurements on HPLC effluents - B.F.H. DRENTH, T. JAGERSMA, F. OVERZET, R.T. GHIJSEN & R.A. DE ZEEUW	75
#NC(A)-5	Quantitation of ^{14}C radioactivity in HPLC eluates - D.W. ROBERTS	81
#B	INVESTIGATION OF METABOLITES, ESPECIALLY PHASE I	95
#B-1	*Examples of approaches to isolating and identifying Phase I metabolites - G.R. BOURNE & S.R. MOSS	97
#B-2	The identification of amitriptyline and benzodiazepine metabolites - IAN D. WATSON, DAVID R. JARVIE, ANNE CHAMBERS & MICHAEL J. STEWART	103
#B-3	A novel use of RIA and HPLC in the identification of plasma metabolites of HR 158 - J.D. ROBINSON, I.D. WILSON, C.D. BEVAN & J. CHAMBERLAIN	111
#B-4	Investigation of pesticide metabolites in the human - C.V. EADSFORTH	119
#NC(B)	NOTES and COMMENTS relating to the foregoing topics <i>including Notes on</i>	135
#NC(B)-1	Metabolic fate of the immunoadjuvant peptidoglycan monomer - JELKA TOMASIĆ	137
#NC(B)-2	HPLC as a means of following an <i>in vitro</i> synthesis of 7-hydroxymethotrexate - YAHYA Y.Z. FARID, IAN D. WATSON, NIGEL G.L. HARDING & MICHAEL J. STEWART	141
#C	INVESTIGATION OF CONJUGATES	147
#C-1	Isolation of conjugates and identification of the attached groups - JELKA TOMASIĆ	149
#C-2	Isolation and characterization of amino acid and sugar conjugates of xenobiotic carboxylic acids - JOHN CALDWELL, ANDREW J. HUTT, MARY VARWELL MARSH & KEITH A. SINCLAIR	161
#C-3	Chromatography, hydrolysis and rearrangement of the glucuronide conjugates of isoxepac, as affecting its assay in urine - I.D. WILSON, A. BHATTI, H.P.A. ILLING, T.A. BRYCE & J. CHAMBERLAIN	181

* Analytical case histories

#NC(C)	NOTE and COMMENTS relating to the foregoing topics	189
	<i>including a Note on</i>	
#NC(C)-1	*Isolation and identification of a drug conjugate - M.P. HARRISON	191
#D	DETERMINATION OF PARTICULAR DRUGS AND METABOLITES	199
#D-1	*HPLC analysis of the major urinary metabolite of flurazepam - J.A.F. DE SILVA, M.A. BROOKS, M.R. HACKMAN & R.E. WEINFELD	201
#D-2	*Quantification of metabolites of aminopyrine and antipyrine in plasma and urine - J.B. HOUSTON & J.C. RHODES	207
#D-3	Immunoassay of bromcriptine and specificity of antibody: criteria for choice of antiserum and marker compound - J. ROSENTHALER, H. MUNZER & R. VOGES	215
#D-4	*Determination of 3-phenoxy-N-methylmorphinan and metabolites in plasma - J.A.F. DE SILVA, J. PAO & M.A. BROOKS	225
#D-5	*Assays where high sensitivity was achieved, especially by HPTLC with automatic spotting and colour reactions - W. RITTER	231
#NC(D)	NOTES and COMMENTS on drug and metabolite determinations	239
	<i>including Notes on</i>	
#NC(D)-1	Selective column-extraction of psycho-active drugs and metabolites - R.G. MUUSZE	241
#NC(D)-2	*Determination of atracurium and its metabolites - E.A.M. NEILL	243
#NC(D)-3	Discrimination between carbon monoxide as a metabolite of carbon tetrachloride and as a degradation product of the biological material - L.J. KING	247
#NC(D)-4	*HPLC determination of cefotaxime and its metabolites - J. CHAMBERLAIN, D. DELL & F.G. COPPIN	249
#E	METABOLITE METHODOLOGY: THE STATE-OF-PLAY - E. Reid (p. 267 : Tabulation of conjugate types, with indexing)	255
	CUMULATIVE INDEX OF COMPOUND TYPES, covering Vols. 5, 7 & 10 also ..	271
	GENERAL INDEX	281

Section #A

TECHNIQUES APPLICABLE TO METABOLITE INVESTIGATION

#A-1

STRATEGIES CENTERED ON HPLC

- G.G. Skellern

Drug Metabolism Research Unit
Department of Pharmaceutical Chemistry
University of Strathclyde
Glasgow G1 1XW, U.K.

The choice of an HPLC system will depend on its application and on the nature of the metabolites to be measured. Normal-phase (NP) chromatography is limited to the measurement of drugs whose physico-chemical properties are similar to the parent drug, whereas reverse-phase (RP) chromatography in its various forms has made possible the simultaneous measurement of metabolites varying widely in pK_a , lipophilicity and polarity. Either alone or in combination with other chromatographic techniques, HPLC has aided the isolation of some polar metabolites which may be difficult to isolate from biological material. In contrast with other types of chromatography, the *in vitro* formation of metabolites can be monitored directly by HPLC, with minimal sample preparation. Thus HPLC will aid the study of enzyme kinetics and reaction mechanisms.

The biotransformation of a compound may result in a diversity of metabolites with widely differing physico-chemical properties. It is therefore desirable to know the properties of the parent compound and its possible metabolites in order that a suitable HPLC system may be chosen. Its choice also depends on whether it is being used primarily as an analytical method for the identification of a metabolite and the determination of its concentration, or preparatively, when substantial quantities of one or more of the metabolites are to be isolated for further structural characterization. HPLC has been used extensively for studying the metabolism of drugs, particularly drugs of high relative molecular mass and polar metabolites [1].

Recently the 'State of the Art' of HPLC has been reviewed [2,

3], and the theoretical and applied aspects discussed, in addition to newer developments. Consequently I will focus upon particular ways in which HPLC has been exploited in metabolic studies.

The subject of sample preparation and clean-up is discussed by other contributors (e.g. #A-2, #E). It suffices to say that RP materials are useful for sample clean-up and enrichment. Thus, a liquid-solid extraction with ODS material in cartridges removed the sparingly water-soluble mebendazole and its metabolites from plasma [4], yielding a chromatographically cleaner extract in a single operation without substantial loss of compound. The authors state that they could re-use the cartridges without loss of performance.

The first decision to be made is the type of chromatography to be adopted. NP (adsorption and bonded-phases) and RP systems are applicable to the study of Phase I metabolites and methylated and acetylated metabolites, whereas RP systems with either ion-pair or ion-suppression modes are suitable for separating the polar, water-soluble conjugates (glucuronides, sulphaconjugates and amino-acid conjugates). This article will concentrate more on the latter type.


SELECTION OF HPLC SYSTEM

Paracetamol is an example of a drug which is eliminated from the body predominantly as its highly polar and water-soluble glucuronide and sulphate conjugates. The cysteine and *N*-acetylcysteine conjugates are two other polar metabolites which are excreted in significant amounts when a substantial overdose of drug (20 g) is ingested. In this case filtered aliquots of urine can be directly

Fig. 1. Comparison of ODS silica (upper) with ODS-TMS silica packing (lower; TMS = trimethylsilyl) for separating paracetamol and its metabolites ([5], by permission).

Eluant: water-methanol-formic acid (86:14:0.1 v/v/v).

S = sulphate conjugate; G = glucuronide; P = paracetamol; C = cysteine conjugate; M = mercapturate.






Fig. 2. Separation of paracetamol metabolites on ODS-TMS silica in the absence (A) and presence (B) of an ion-pairing agent ([6], by permission).

Eluant: A as for Fig. 1

B as for Fig. 1 but with dioctylamine (DOA; 0.7 mg/l) and KNO_3 (3 g/l).

injected onto the HPLC column, since there is a sufficient concentration of metabolites. With methanol-water-formic acid, Knox & Jurand [5] compared conventional ODS-silica with ODS-silica capped with TMS, and found the latter superior for the determination of paracetamol and its polar metabolites (Fig. 1). The effect of varying the concentration of methanol, formic acid and added salts on the k' values of the 5 compounds was thoroughly examined. The presence of acid in the eluting solvent markedly increased k' for these compounds, partly by suppressing the degree of ionization of the acidic metabolites. Acid in the eluant also decreased the possibility of hydrolysis of the conjugates. Slight pH changes of the eluant around the pK_a value of the metabolites can markedly alter k' . With straight methanol, water and formic acid mixtures as eluant, the elution order was sulphoconjugate, glucuronide, paracetamol, cysteine and *N*-acetylcysteine conjugates. However, upon the addition of salt (KH_2PO_4) to the eluant the capacity factor of the sulphoconjugate increased from zero to 0.8, and the elution order was glucuronide, sulphoconjugate, cysteine conjugate, paracetamol and mercapturate.

The addition of either dioctylamine (DOA) or tetrabutylammonium hydroxide (TBAOH) to the eluant to form hydrophobic ion-pairs with the highly ionized sulphate conjugate dramatically increased its k' value [6], the elution order now being glucuronide, cysteine conjugate, paracetamol, mercapturate and sulphoconjugate. To obtain reasonable values of k' for the mercapturate and sulphoconjugate it was necessary (Fig. 2) to add a salt (KNO_3) to the eluant when DOA