

**International  
Review of Science**

---

**Carbohydrates**

---

**Organic Chemistry  
Series Two  
Volume 7**

---

**Consultant Editor  
D H Hey FRS  
Volume Editor  
G O Aspinall**


**Butterworths**

0.2  
V.7  
V.2/7  
International Review of Science

# Organic Chemistry Series Two

## Volume 7 Carbohydrates

Edited by **G. O. Aspinall**  
York University, Ontario



E7952468

**BUTTERWORTHS**  
LONDON - BOSTON  
Sydney - Wellington - Durban - Toronto

- |                |                                                                                                                             |
|----------------|-----------------------------------------------------------------------------------------------------------------------------|
| United Kingdom | Butterworth & Co (Publishers) Ltd<br>London: 88 Kingsway, WC2B 6AB                                                          |
| Australia      | Butterworths Pty Ltd<br>Sydney: 586 Pacific Highway, Chatswood, NSW 2067<br>Also at Melbourne, Brisbane, Adelaide and Perth |
| Canada         | Butterworth & Co (Canada) Ltd<br>Toronto: 2265 Midland Avenue, Scarborough,<br>Ontario, M1P 4S1                             |
| New Zealand    | Butterworths of New Zealand Ltd<br>Wellington: T & W Young Building, 77-85 Customhouse<br>Quay, 1, CPO Box 472              |
| South Africa   | Butterworth & Co (South Africa) (Pty) Ltd<br>Durban: 152-154 Gale Street                                                    |
| USA            | Butterworth (Publishers) Inc<br>Boston: 19 Cummings Park, Woburn, Mass. 01801                                               |

All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means, including photocopying and recording, without the written permission of the copyright holder, application for which should be addressed to the Publishers. Such written permission must also be obtained before any part of this publication is stored in a retrieval system of any nature.

This book is sold subject to the Standard Conditions of Sale of Net Books and may not be re-sold in the UK below the net price given by the Publishers in their current price list.

First published 1976  
Reprinted 1978

ISBN 0 408 70619 8

©Butterworth & Co (Publishers) Ltd 1976

**Library of Congress Cataloging in Publication Data**

Aspinall, Gerald O.  
Carbohydrates.  
(Organic chemistry, series two; v. 7) (International  
review of science)  
Includes index.  
I. Carbohydrates. II. Title. III. Series:  
International review of science.  
QD245.073 vol. 7 [QD321] 547'.008s [547'.78]  
ISBN 0 408 70619 8 75-19372

Typeset, printed and bound in Great Britain by  
REDWOOD BURN LIMITED  
Trowbridge & Esher

International Review of Science

**Organic Chemistry**  
**Series Two**

Consultant Editor  
**D. H. Hey, F.R.S.**

## Publisher's Note

The International Review of Science is an important venture in scientific publishing presented by Butterworths. The basic concept of the Review is to provide regular authoritative reviews of entire disciplines. Chemistry was taken first as the problems of literature survey are probably more acute in this subject than in any other. Biochemistry and Physiology followed naturally. As a matter of policy, the authorship of the Review of Science is international and distinguished, the subject coverage is extensive, systematic and critical.

The Review has been conceived within a carefully organised editorial framework. The overall plan was drawn up and the volume editors appointed by seven consultant editors. In turn, each volume editor planned the coverage of his field and appointed authors to write on subjects which were within the area of their own research experience. No geographical restriction was imposed. Hence the 500 or so contributions to the Review of Science come from many countries of the world and provide an authoritative account of progress.

The publication of Organic Chemistry Series One was completed in 1973 with ten text volumes and one index volume; in accordance with the stated policy of issuing regular reviews to keep the series up to date, volumes of Series Two will be published between the middle of 1975 and early 1976; Series Two of Physical Chemistry will be published at the same time, while Inorganic Chemistry Series Two was published during the first half of 1975. Volume titles are the same as in Series One but the articles themselves either cover recent advances in the same subject or deal with a different aspect of the main theme of the volume. In Series Two an index is incorporated in each volume and there is no separate index volume.

Butterworth & Co. (Publishers) Ltd.

## ORGANIC CHEMISTRY SERIES TWO

Consultant Editor  
D. H. Hey, F.R.S., *formerly of  
the Department of Chemistry,  
King's College, University  
of London*

*Volume titles and Editors*

- 1 STRUCTURE  
DETERMINATION IN  
ORGANIC CHEMISTRY**  
Professor L. M. Jackman,  
*Pennsylvania State  
University*
- 2 ALIPHATIC COMPOUNDS**  
Professor N. B. Chapman,  
*University of Hull*

- 3 AROMATIC COMPOUNDS**  
Professor H. Zollinger,  
*Eidgenössische Technische  
Hochschule, Zurich*
- 4 HETEROCYCLIC  
COMPOUNDS**  
Professor K. Schofield,  
*University of Exeter*
- 5 ALICYCLIC COMPOUNDS**  
Professor D. Ginsburg,  
*Technion-Israel Institute  
of Technology, Haifa*
- 6 AMINO ACIDS,  
PEPTIDES AND  
RELATED COMPOUNDS**  
Professor H. N. Rydon,  
*University of Exeter*

- 7 CARBOHYDRATES**  
Professor G. O. Aspinall,  
*York University, Ontario*
- 8 STEROIDS**  
Dr. W. F. Johns, G. D.  
*Searle & Co., Chicago*
- 9 ALKALOIDS**  
Professor K. Wiesner, F.R.S.,  
*University of New Brunswick*
- 10 FREE RADICAL  
REACTIONS**  
Professor W. A. Waters, F.R.S.,  
*University of Oxford*

## INORGANIC CHEMISTRY SERIES TWO

Consultant Editor  
H. J. Emeléus, C.B.E., F.R.S.  
*Department of Chemistry  
University of Cambridge*

*Volume titles and Editors*

- 1 MAIN GROUP  
ELEMENTS—HYDROGEN  
AND GROUPS I-III**  
Professor M. F. Laopert,  
*University of Sussex*
- 2 MAIN GROUP  
ELEMENTS—GROUPS  
IV AND V**  
Dr. D. B. Sowerby,  
*University of Nottingham*

- 3 MAIN GROUP  
ELEMENTS—GROUPS  
VI AND VII**  
Professor V. Gutmann,  
*Technical University of  
Vienna*
- 4 ORGANOMETALLIC  
DERIVATIVES OF THE  
MAIN GROUP ELEMENTS**  
Professor B. J. Aylett,  
*Westfield College,  
University of London*
- 5 TRANSITION METALS—  
PART 1**  
Professor D. W. A. Sharp,  
*University of Glasgow*
- 6 TRANSITION METALS—  
PART 2**  
Dr. M. J. Mays, *University  
of Cambridge*

- 7 LANTHANIDES AND  
ACTINIDES**  
Professor K. W. Bagnall,  
*University of Manchester*
- 8 RADIOCHEMISTRY**  
Dr. A. G. Maddock,  
*University of Cambridge*
- 9 REACTION  
MECHANISMS  
IN INORGANIC  
CHEMISTRY**  
Professor M. L. Tobe,  
*University College,  
University of London*
- 10 SOLID STATE  
CHEMISTRY**  
Dr. L. E. J. Roberts, *Atomic  
Energy Research Establish-  
ment, Harwell*

## PHYSICAL CHEMISTRY SERIES TWO

Consultant Editor  
A. D. Buckingham, F.R.S.,  
*Department of Chemistry  
University of Cambridge*

*Volume titles and Editors*

- 1 THEORETICAL  
CHEMISTRY**  
Professor A. D. Buckingham,  
*F.R.S., University of Cambridge*  
and Professor C. A. Coulson,  
*F.R.S., University of Oxford*
- 2 MOLECULAR  
STRUCTURE AND  
PROPERTIES**  
Professor A. D. Buckingham,  
*F.R.S., University of Cambridge*
- 3 SPECTROSCOPY**  
Dr. D. A. Ramsay, F.R.S.C.,  
*National Research Council  
of Canada*
- 4 MAGNETIC RESONANCE**  
Professor C. A. McDowell,  
*F.R.S.C., University of  
British Columbia*

- 5 MASS SPECTROMETRY**  
Professor A. Maccoll,  
*University College,  
University of London*
- 6 ELECTROCHEMISTRY**  
Professor J. O'M. Bockris,  
*The Flinders University of  
S. Australia*
- 7 SURFACE CHEMISTRY  
AND COLLOIDS**  
Professor M. Kerker,  
*Clarkson College of  
Technology, New York*
- 8 MACROMOLECULAR  
SCIENCE**  
Professor C. E. R. Bawn,  
*C.B.E., F.R.S., formerly of the  
University of Liverpool*
- 9 CHEMICAL KINETICS**  
Professor D. R. Herschbach  
*Harvard University*
- 10 THERMOCHEMISTRY  
AND THERMO-  
DYNAMICS**  
Dr. H. A. Skinner, *University  
of Manchester*
- 11 CHEMICAL  
CRYSTALLOGRAPHY**  
Professor J. M. Robertson,  
*C.B.E., F.R.S., formerly of the  
University of Glasgow*
- 12 ANALYTICAL CHEMISTRY  
—PART 1**  
Professor T. S. West,  
*Macaulay Institute for  
Soil Research, Aberdeen*
- 13 ANALYTICAL CHEMISTRY  
—PART 2**  
Professor T. S. West,  
*Macaulay Institute for  
Soil Research, Aberdeen*

## BIOCHEMISTRY SERIES ONE

Consultant Editors

H. L. Kornberg, F.R.S.

*Department of Biochemistry*

*University of Leicester and*

*D. C. Phillips, F.R.S., Department of  
Zoology, University of Oxford*

*Volume titles and Editors*

### 1 CHEMISTRY OF MACRO- MOLECULES

Professor H. Gutfreund, *University of  
Bristol*

### 2 BIOCHEMISTRY OF CELL WALLS AND MEMBRANES

Dr. C. F. Fox, *University of California,  
Los Angeles*

### 3 ENERGY TRANSDUCING MECHANISMS

Professor E. Racker, *Cornell University  
New York*

### 4 BIOCHEMISTRY OF LIPIDS

Professor T. W. Goodwin, F.R.S.,  
*University of Liverpool*

### 5 BIOCHEMISTRY OF CARBO- HYDRATES

Professor W. J. Whelan, *University  
of Miami*

### 6 BIOCHEMISTRY OF NUCLEIC ACIDS

Professor K. Burton, F.R.S., *University of  
Newcastle upon Tyne*

### 7 SYNTHESIS OF AMINO ACIDS AND PROTEINS

Professor H. R. V. Arnstein, *King's  
College, University of London*

### 8 BIOCHEMISTRY OF HORMONES

Professor H. V. Rickenberg, *National  
Jewish Hospital & Research Center,  
Colorado*

### 9 BIOCHEMISTRY OF CELL DIFFER- ENTIATION

Professor J. Paul, *The Beatson Institute  
for Cancer Research, Glasgow*

### 10 DEFENCE AND RECOGNITION

Professor R. R. Porter, F.R.S., *University  
of Oxford*

### 11 PLANT BIOCHEMISTRY

Professor D. H. Northcote, F.R.S.,  
*University of Cambridge*

### 12 PHYSIOLOGICAL AND PHARMACO- LOGICAL BIOCHEMISTRY

Dr. H. K. F. Blaschko, F.R.S., *University  
of Oxford*

---

## PHYSIOLOGY SERIES ONE

Consultant Editors

A. C. Guyton,

*Department of Physiology and*

*Biophysics, University of Mississippi*

*Medical Center and*

*D. F. Horrobin,*

*Department of Physiology, University  
of Newcastle upon Tyne*

*Volume titles and Editors*

### 1 CARDIOVASCULAR PHYSIOLOGY

Professor A. C. Guyton and Dr. C. E. Jones,  
*University of Mississippi Medical Center*

### 2 RESPIRATORY PHYSIOLOGY

Professor J. G. Widdicombe, *St. George's  
Hospital, London*

### 3 NEUROPHYSIOLOGY

Professor C. C. Hunt, *Washington  
University School of Medicine, St. Louis*

### 4 GASTROINTESTINAL PHYSIOLOGY

Professor E. D. Jacobson and Dr. L. L.  
Shanbour, *University of Texas Medical  
School*

### 5 ENDOCRINE PHYSIOLOGY

Professor S. M. McCann, *University of  
Texas*

### 6 KIDNEY AND URINARY TRACT PHYSIOLOGY

Professor K. Thurau, *University of Munich*

### 7 ENVIRONMENTAL PHYSIOLOGY

Professor D. Robertshaw, *University  
of Nairobi*

### 8 REPRODUCTIVE PHYSIOLOGY

Professor R. O. Greep, *Harvard Medical  
School*

## Consultant Editor's Note

The ten volumes in Organic Chemistry in the Second Series of the biennial reviews in the International Review of Science follow logically from those of the First Series. No major omissions have come to light in the overall coverage of the First Series. The titles of the ten volumes therefore remain unchanged but there are three new Volume Editors. The volume on Structure Determination in Organic Chemistry has been taken over by Professor Lloyd M. Jackman of Pennsylvania State University, that on Alicyclic Compounds by Professor D. Ginsburg of Technion-Israel Institute of Technology, and that on Amino Acids, Peptides and Related Compounds by Professor H. N. Rydon of the University of Exeter. The international character of the Series is thus strengthened with four Volume Editors from the United Kingdom, two each from Canada and the U.S.A., and one each from Israel and Switzerland. An even wider pattern is shown for the authors, who now come from some sixteen countries. The reviews in the Second Series are mainly intended to cover work published in the years 1972 and 1973, although relevant results published in 1974 and 1975 are included in some cases, and earlier work is also covered where applicable.

It is my pleasure once again to thank all the Volume Editors for their helpful cooperation in this venture.

London

D. H. Hey



## Preface

The second volume devoted to carbohydrates in the Organic Chemistry Section of the International Review of Science continues with the objective of publishing critical and well-documented articles with carefully selected rather than necessarily exhaustive lists of reference to the recent literature. Again a balance is maintained between those chapters dealing with the chemistry of monosaccharides and those dealing with carbohydrate polymers. Certain chapters extend the subject matter coverage of the first volume over the intervening two-year period, whilst other chapters review topics not previously covered in this series. The general chemistry of carbohydrates is again discussed in articles dealing with, general carbohydrate synthesis, displacement, elimination and rearrangement reactions, and amino and nitro sugars. In addition, an article on sugar-containing antibiotics deals with a topic that both is of considerable biological importance and continues to provide much of the driving force for the development of new carbohydrate syntheses. The review on  $^{13}\text{C}$  n.m.r. spectroscopy is of importance for applications both in monosaccharide chemistry and increasingly for structure determination of polysaccharides which have fairly regular 'repeating units'.

Topics of particular interest to biochemists are discussed in articles on membrane glycolipids (S. Hakomori), iduronic acid-containing glycosaminoglycans (U. Lindahl) and blood group substances and related glycoproteins (K. O. Lloyd). These articles are of importance in stressing the need for properly evaluated chemical data as a basis for understanding biological phenomena at a molecular level. Attention may be drawn to the forthcoming appearance of a volume on 'Biochemistry of Carbohydrates' in the Biochemistry Section of the International Review of Science. This should be regarded as a companion volume in which the editors (Dr. W. J. Whelan and I) have chosen topics for review which are mutually complementary rather than overlapping in content.

Ontario

G. O. Aspinall



## Contents

<b>Carbon-13 n.m.r. spectroscopy of carbohydrates</b> A. S. Perlin, <i>McGill University, Montreal</i>	1
<b>Displacement, elimination and rearrangement reactions</b> R. J. Ferrier, <i>Victoria University of Wellington</i>	35
<b>General carbohydrate synthesis</b> W. A. Szarek and D. M. Vyas, <i>Queen's University, Kingston, Ontario</i>	89
<b>The nitro sugars</b> A. C. Richardson, <i>Queen Elizabeth College, University of London</i>	131
<b>Sugar-containing antibiotics</b> S. Umezawa, <i>Keio University, Yokohama</i>	149
<b>Polysaccharide methodology</b> G. O. Aspinall, <i>York University, Toronto, Ontario</i>	201
<b>Glycolipids of animal cell membranes</b> S. Hakomori, <i>University of Washington</i>	223
<b>Glycoproteins with blood group activity</b> K. O. Lloyd, <i>Columbia University, New York and Texas Tech. University, Lubbock</i>	251
<b>Structure and biosynthesis of iduronic acid-containing glycosaminoglycans</b> U. Lindahl, <i>Royal Veterinary College of Sweden</i>	283
<b>Index</b>	313



# 1 Carbon-13 N.M.R. Spectroscopy of Carbohydrates

A. S. PERLIN

McGill University, Montreal

1.1	INTRODUCTION	2
1.2	CHEMICAL SHIFTS	2
1.2.1	<i>General characteristics</i>	2
1.2.2	<i>Substituent effects</i>	4
1.2.2.1	<i>Chemical shifts of substituents</i>	5
1.2.3	<i>Configurational and conformational effects on pyranoses</i>	5
1.2.4	<i><math>^{13}\text{C}</math> Chemical shifts of furanoses</i>	13
1.2.5	<i>Factors influencing sterically induced <math>^{13}\text{C}</math> chemical shifts: charge density, bond polarisation, diamagnetic contribution</i>	14
1.3	THE ANALYSIS OF MIXTURES—EQUILIBRIA OF KETOSES	15
1.4	$^{13}\text{C}$ N.M.R. SPECTRA OF HIGHER SACCHARIDES	16
1.4.1	<i>Disaccharides and oligosaccharides</i>	16
1.4.2	<i>Polysaccharides: polymer sequence; mucopolysaccharides; derivatives; spin relaxation</i>	18
1.5	$^{13}\text{C}$ — $^1\text{H}$ SPIN-SPIN COUPLING	21
1.5.1	<i>General characteristics</i>	21
1.5.2	<i>Direct, one-bond, coupling (<math>^1J</math>): orientational dependence</i>	23
1.5.3	<i>Two-bond coupling (<math>^2J</math>): orientational dependence; sign of coupling</i>	23
1.5.4	<i>Vicinal coupling (<math>^3J</math>): orientational dependence: factors affecting magnitude of splitting</i>	24
1.6	STEREOCHEMICAL APPLICATIONS OF $^{13}\text{C}$ — $^1\text{H}$ COUPLING	26
1.6.1	<i>Substituent orientation: hydroxyl, O-alkyl, nucleosides</i>	26
1.6.1.1	<i>Hydroxyl groups</i>	26

1.6.1.2	O-Alkyl groups	27
1.6.1.3	Nucleoside bases	27
1.6.2	Orientation of glycosidic bonds in disaccharides	27
1.6.3	The configuration of branched sugars	29
1.6.4	Other applications: orientations of rotamers; conformation of lactones	30

ACKNOWLEDGEMENTS	32
------------------	----

---

## 1.1 INTRODUCTION

The proliferation of studies on the  $^{13}\text{C}$  n.m.r. spectra of carbohydrates during the past four or five years attests to the current strong interest of chemists in exploring applications of this area of spectroscopy. It is now clear that  $^{13}\text{C}$  n.m.r. is acquiring a status not only as a useful adjunct to  $^1\text{H}$  n.m.r. spectroscopy, but one characterised by its own unique contributions.

Two comprehensive texts on  $^{13}\text{C}$  n.m.r.<sup>1, 2</sup> deal at some length with spectra of the carbohydrates, and recent reviews<sup>3-5</sup> cover the more current literature and include<sup>5b</sup> a detailed compilation of data. The intention of this article is to present a relatively broad description of the characteristics of these  $^{13}\text{C}$  spectra, and to emphasise applications — demonstrated or potential — which yield information (or may do so) not readily attainable by other techniques.

## 1.2 CHEMICAL SHIFTS


### 1.2.1 General characteristics

Chemical shifts of the  $^{13}\text{C}$  nuclei of carbohydrates and derivatives, aside from those of substituent groups that may be introduced, encompass most regions of the 200 p.p.m. range covered by organic compounds. It is common practice to obtain chemical shifts from proton-decoupled spectra\*, in which the signal produced by a  $^{13}\text{C}$  nucleus appears as a narrow singlet†. This simplified form of display, together with a resolution of 0.1 p.p.m. afforded by present instruments, usually ensures an overall excellent separation of resonance signals even with highly complex molecules or mixtures. Herein lies much of the power of  $^{13}\text{C}$  n.m.r. spectroscopy.

\* Normal irradiation of the sample at the  $^{13}\text{C}$  resonance frequency is accompanied by an intense broadband irradiation at the appropriate  $^1\text{H}$  frequency, thereby eliminating coupling between the  $^{13}\text{C}$  and  $^1\text{H}$  nuclei. In this process also, the signal intensity of a carbon bearing a proton is augmented by a factor of *ca.* 3 owing to nuclear Overhauser enhancement<sup>1, 2</sup> and also by coalescence of the components of the split signal. This is an important gain, because of the low natural abundance (1.1%) and low sensitivity of  $^{13}\text{C}$  nuclei.

†  $^{13}\text{C}$ - $^{13}\text{C}$  coupling causes no significant interference, because at the natural abundance level two  $^{13}\text{C}$  nuclei rarely occur in the same molecule.

A good example of the signal separation that can be achieved with a complicated mixture of closely related compounds is illustrated in Figure 1.1 by the pulsed Fourier transform (FT)<sup>1,2,6,7</sup> spectrum of 3-*O*-methyl-*D*-psicose<sup>8</sup>: two pyranose and two furanose forms are in equilibrium (see Section 1.3), corresponding to 28 individual <sup>13</sup>C signals, each of which is



**Figure 1.1** FT <sup>13</sup>C n.m.r. proton-decoupled spectrum (at 25.15 MHz) of 3-*O*-methyl *D*-psicose in D<sub>2</sub>O at 30°C. Signals of the anomeric carbons (C-2) are displayed to low field of 95 p.p.m. Inset formulae represent the anomers of the pyranose and furanose forms (From Herve du Penhoat and Perlin<sup>8</sup>, by courtesy of Elsevier)

detectable. In common with other FT <sup>13</sup>C spectra<sup>1,2,7</sup>, quantitative aspects of this spectrum are not as generally satisfactory as those of protons. The peak areas are highly sensitive to the relaxation properties of the various <sup>13</sup>C nuclei and hence to the experimental pulsing procedure adopted, and vary also with the extent of Overhauser enhancement. Since spin-lattice relaxation is more efficient for a carbon that has an attached proton than for a non-protonated carbon, the small areas of the anomeric (C-2) signals in Figure 1.1 relative to those of the other carbons are a reflection of the fact that C-2 bears no hydrogen. Despite these limitations, however, the error in estimating the composition of a mixture will probably be  $< \pm 5\%$  if comparisons of the integrated intensities are based on signals representing the same class of carbon (such as the four anomeric carbons in Figure 1.1).

Resonances of simple aldoses and ketoses<sup>8-16</sup> are distinguishable for the

most part in terms of anomeric, secondary and primary carbons: relative to tetramethylsilane (which is being widely adopted as a standard for  $^{13}\text{C}$  as well as  $^1\text{H}$ <sup>1,2,5b</sup>), these classes of nuclei usually resonate at  $98 \pm 6$ ,  $75 \pm 5$  and  $65 \pm 5$  p.p.m., respectively. Variations within each class are associated with changes in configuration or ring size according to generally recognisable patterns, as described below (Sections 1.2.3 and 1.2.4).

### 1.2.2 Substituent effects

The effects on chemical shift owing to substituent groups or to other modifications of carbohydrates have as yet received limited systematic study. However, general experience with various classes of organic compounds shows a high degree of consistency in applying substituent effects to predict chemical shifts<sup>1,2</sup>, and the same is therefore to be expected for carbohydrates. An outline of the data available on the main types of transformations within the sugar series\* is given below; the impact on the carbon in question is usually accompanied by a much smaller one on neighbouring carbons, and it is noteworthy also that some of the effects (especially those of *O*-alkylation and *O*-acylation) are quite different from those of protons.

Most marked is the change observed when a carbinol group is converted into a carbonyl group (aldehyde, ketone or carboxyl), in which case the particular  $^{13}\text{C}$  resonance is moved downfield into the region of 175–200 p.p.m.<sup>1,2</sup>. Uronic acid carboxyls, for example, produce signals at 176–180 p.p.m.<sup>17, 18</sup>.

Deoxy groups are characterised by an increase in shielding relative to the carbinol carbon of 25 p.p.m. (C-1 of 1,5-anhydro-D-glucitol) or 35–40 p.p.m. (C-2 and C-6 of 2-deoxy-D-glucose)<sup>9, 12</sup>.

A distinctive upfield shift is observed for aminodeoxy derivatives: C-2 of 2-acetamido-2-deoxyaldoses resonates 15–20 p.p.m. upfield of secondary carbinol signals<sup>19</sup>; the 2-deoxy-2-sulphamino group is associated with increased shielding of *ca.* 13 p.p.m.<sup>17</sup> and  $^{13}\text{C}$ —NHMe by 10 p.p.m.<sup>20</sup>.

For alkenes the chemical shift of an enolic carbon should be close to 145 p.p.m. (C-1 of 3,4,6-tri-*O*-acetyl-D-glucal resonates at 144.6 p.p.m.), and of other  $\text{sp}^2$  carbons in the region of 100 p.p.m. [ $\delta(\text{C-2})$  of 3,4,6-tri-*O*-acetyl-D-glucal is 98.2 p.p.m.]<sup>21</sup>.

A deuterium atom markedly reduces the effective intensity of the signal of the  $^{13}\text{C}$  nucleus to which it is attached (owing to splitting into a triplet, modified relaxation characteristics and loss of Overhauser enhancement), which essentially removes that signal from the spectrum<sup>13, 22</sup>. In some instances an upfield shift of about 0.05 p.p.m. of signals produced by the  $\beta$   $^{13}\text{C}$  nuclei is observed<sup>23</sup>. These effects aid in the analysis of spectra.

*O*-Methylation to yield ethers usually induces deshielding of the  $\alpha$ -C by 8–10 p.p.m.; adjacent carbons may show an increase in shielding of 0.5–1.0 p.p.m. or occasionally more<sup>11, 12</sup>. However, it has been found<sup>8</sup> that C-2 of D-psicosides is deshielded by only 2–4 p.p.m. on methyl glycoside formation.

\* Obviously, some of the functional groups listed are components of naturally-occurring carbohydrates; for convenience they are treated here as if a simple sugar is being chemically modified.

An anomeric phenyl or nitrophenyl group promotes deshielding of C-1 by 5–7 p.p.m.<sup>24</sup>. C-1 of 1,6-anhydrohexopyranoses is deshielded relative to the free sugar by 5–6 p.p.m.<sup>18</sup>. *O*-Tritylation has been found to produce an upfield shift of the C-6 signal of 1,2,3,4-tetra-*O*-acetyl- $\beta$ -D-glucose or -D-mannose<sup>25</sup> by 1 p.p.m.

*O*-Acetylation to give esters has a relatively small, and diverse, effect<sup>11</sup> varying from deshielding by a few p.p.m. to increased shielding by a similar amount. An *O*-phosphate group promotes deshielding by 2–3 p.p.m.<sup>26, 27</sup> and an *O*-sulphate deshielding by 6–7 p.p.m. (primary) or 8–10 p.p.m. (secondary)<sup>17, 28</sup>.

Strong deshielding of  $\alpha$  carbons usually accompanies the formation of cyclic acetals. An *O*-isopropylidene group promotes a downfield displacement of 5–10 p.p.m. In *trans*-fused 4,6-*O*-benzylidene derivatives, C-4 is deshielded by *ca.* 10 p.p.m. and C-6 by *ca.* 7 p.p.m.; by contrast, C-5 shows an increased shielding of 11 p.p.m. and C-3 of 4 p.p.m.<sup>29</sup>.

The formation of a glycosidic bond between two sugar residues causes deshielding both of the anomeric carbon and the other <sup>13</sup>C nucleus of the bond by 5–10 p.p.m.<sup>17, 30</sup>.

Lanthanide shift reagents have a relatively small impact on <sup>13</sup>C chemical shifts, especially compared with that on protons, but nevertheless are sufficiently large to offer substantial promise for use in spectral analysis<sup>31, 31a</sup>.


### 1.2.2.1 Chemical shifts of substituents

Chemical shifts of the carbons of a number of common substituents show small variations within each group, and as yet no consistent structural or stereochemical correlations with these differences have been detected. The approximate ranges of chemical shifts reported in the literature for some of these substituents are as follows: *O*-CH<sub>3</sub>, 55–60 p.p.m.<sup>5b, 11, 12, 32</sup>; *O*-COCH<sub>3</sub>, 19–20 p.p.m. (CH<sub>3</sub>) and 165–170 p.p.m. (C=O)<sup>5b, 31</sup>; *N*-COCH<sub>3</sub>, 22–24 p.p.m. (CH<sub>3</sub>) and 175–177 p.p.m. (C=O)<sup>19</sup>; *O*-isopropylidene 26–30 p.p.m. (CH<sub>3</sub>) and 128–135 p.p.m. (O—C—O)<sup>5b</sup>; *O*-benzylidene, 99–102 p.p.m. (O—C—O)<sup>29</sup>.

### 1.2.3 Configurational and conformational effects on pyranoses

The proton-decoupled spectrum of  $\alpha$ - and  $\beta$ -D-glucose (1) and (2) (Figure 1.2) illustrates a number of features of natural-abundance <sup>13</sup>C spectra of pyranoses<sup>9–16</sup>. Although a few of the signals overlap, the twelve resonance signals required for a mixture of two anomers are readily accounted for, and their relative intensities correspond to the expected  $\alpha$ : $\beta$  ratio of close to 2:3.

Carbons 1, 2, 3 and 5 of  $\alpha$ -D-glucose all resonate 2.5–4.5 p.p.m. upfield of the corresponding nuclei of the  $\beta$ -anomer. That is, the carbon (C-1 $\alpha$ ) bearing the axial OH, carbons adjacent to it (C-2) and those involved in a *syn*-diaxial C—H, C—O interaction (C-3 and C-5) all show an increase in shielding; C-4 and C-6, which are more remote from the axial hydroxyl



**Figure 1.2** FT  $^{13}\text{C}$  n.m.r. proton-decoupled spectrum (at 22.6 MHz) of  $\alpha$ - and  $\beta$ -D-glucose in  $\text{D}_2\text{O}$  at  $30^\circ\text{C}$

group, have the same chemical shifts as those of the  $\beta$ -anomer. It is convenient to describe the differences observed as  $\alpha$ ,  $\beta$ ,  $\gamma$  or  $\delta$  shielding effects<sup>11, 33</sup>: an axial hydroxyl group is associated with a notable increase in shielding of the carbon to which it is bonded, and of the  $\beta$  and  $\gamma$  carbons, but usually not of the  $\delta$  and more distant carbons. As a further example, the fact that the axial C-2—O bond of  $\beta$ -D-mannose introduces an additional orientational factor of this kind appears to be reflected in increased shielding (by an average of *ca.* 2.5 p.p.m.) of carbons 1–4 compared with  $\beta$ -D-glucose. Such shielding effects are described for  $\alpha$ - and  $\beta$ -D-lyxose in Table 1.1.

An analogous relationship between chemical shift and orientation is observed for inositols<sup>34</sup>, 2-hexuloses<sup>8, 16</sup>, 2-acetamido-2-deoxyhexoses<sup>19</sup>, deoxynitro derivatives<sup>35</sup> and 4,6-*O*-benzylidene glycosides<sup>29</sup>. In all these instances, chemical shifts for the various members of the series can be interrelated with reasonable accuracy by taking account of the orientational differences expected, and adding or subtracting the appropriate shielding values (Table 1.1). Such data furnish a general reference base from which structural and stereochemical information may be derived when applied to other kinds of derivatives or higher molecular weight carbohydrates.



Although it seems reasonable to associate such shielding effects with steric interactions (see Section 1.2.4), the evidence for this is mainly circumstantial, and there are a number of inconsistencies. Thus, less shielding is observed for a *syn*-axial interaction between two OH groups than between an H and OH group, i.e. a shielding value of only 0–2 p.p.m.\*, rather than 2.5–3 p.p.m., accounts for the chemical shifts observed<sup>12, 16, 34</sup>, which is the reverse of the order of steric compressions. Among other examples<sup>12, 16</sup> is the unexpectedly small impact on the shielding of C-4 and C-5 by axial C-4 of *L-arabino* isomers.

In comparing the chemical shifts of configurational isomers as above, one begins with the assumption that the basic architecture (bond angles, etc.)

\* A value of 0 Hz is used (Table 1.1) for inositols (entry IV), and of 2.0 Hz for hexuloses (entries V and VI). More striking examples of this kind have been observed in the alicyclic series, in which some *syn*-axial OH, CH<sub>3</sub> orientations are associated with *deshielding*<sup>36</sup>.



**Table 1.1**  $^{13}\text{C}$  Shielding differences between isomers

	Inversion*	C	$\delta\text{A}$ (expt.)	Relationship†	$\Delta\delta$	$\delta\text{B}$ (calc.)	$\delta\text{B}$ (expt.)
I		1	95.0	+ax. O, -O-1/O-2	+2.5, -2.5	95.0	95.0
		2	70.8	+adj. O-1, -O-2/O-1	+2.5, -2.5	70.8	71.5
		3	73.7	+H//O-1	+2.5	71.2	71.1
		4	67.6	—	0	67.6	68.5
		5	64.9	+H//O-1	+2.5	62.4	63.9
II		1	96.2	+adj. N-2	+2.5	93.7	94.3
		2	58.0	+ax. N	+2.5	55.5	55.3
		3	75.2	+adj. N-2	+2.5	72.7	73.2
		4	71.2	+H//N-2	+2.5	68.7	67.8
		5	77.2	—	0	77.2	77.5
		6	62.0	—	0	62.0	61.7