


EMS Textbooks in Mathematics



Fabrice Baudoin

Diffusion Processes and Stochastic Calculus



European Mathematical Society

Fabrice Baudoin

**Diffusion
Processes and
Stochastic
Calculus**



European Mathematical Society

Author:

Fabrice Baudoin
Department of Mathematics
Purdue University
150 N. University Street
West Lafayette, IN 47907
USA

E-mail: fbaudoin@purdue.edu

2010 Mathematics Subject Classification: 60-01, 60G07, 60J60, 60J65, 60H05, 60H07

Key words: Brownian motion, diffusion processes, Malliavin calculus, rough paths theory, semigroup theory, stochastic calculus, stochastic processes

ISBN 978-3-03719-133-0

The Swiss National Library lists this publication in The Swiss Book, the Swiss national bibliography, and the detailed bibliographic data are available on the Internet at <http://www.helvetica.ch>.

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, re-use of illustrations, recitation, broadcasting, reproduction on microfilms or in other ways, and storage in data banks. For any kind of use permission of the copyright owner must be obtained.

© European Mathematical Society 2014

Contact address:

European Mathematical Society Publishing House
Seminar for Applied Mathematics
ETH-Zentrum SEW A27
CH-8092 Zürich
Switzerland

Phone: +41 (0)44 632 34 36

Email: info@ems-ph.org

Homepage: www.ems-ph.org

Typeset using the author's T_EX files: I. Zimmermann, Freiburg

Printing and binding: Beltz Bad Langensalza GmbH, Bad Langensalza, Germany

∞ Printed on acid free paper

9 8 7 6 5 4 3 2 1



EMS Textbooks in Mathematics

EMS Textbooks in Mathematics is a series of books aimed at students or professional mathematicians seeking an introduction into a particular field. The individual volumes are intended not only to provide relevant techniques, results, and applications, but also to afford insight into the motivations and ideas behind the theory. Suitably designed exercises help to master the subject and prepare the reader for the study of more advanced and specialized literature.

Jørn Justesen and Tom Høholdt, *A Course In Error-Correcting Codes*

Markus Stroppel, *Locally Compact Groups*

Peter Kunkel and Volker Mehrmann, *Differential-Algebraic Equations*

Dorothee D. Haroske and Hans Triebel, *Distributions, Sobolev Spaces, Elliptic Equations*

Thomas Timmermann, *An Invitation to Quantum Groups and Duality*

Oleg Bogopolski, *Introduction to Group Theory*

Marek Jarnicki and Peter Pflug, *First Steps in Several Complex Variables: Reinhardt Domains*

Tammo tom Dieck, *Algebraic Topology*

Mauro C. Beltrametti et al., *Lectures on Curves, Surfaces and Projective Varieties*

Wolfgang Woess, *Denumerable Markov Chains*

Eduard Zehnder, *Lectures on Dynamical Systems. Hamiltonian Vector Fields and Symplectic Capacities*

Andrzej Skowroński and Kunio Yamagata, *Frobenius Algebras I. Basic Representation Theory*

Piotr W. Nowak and Guoliang Yu, *Large Scale Geometry*

Joaquim Bruna and Julià Cufí, *Complex Analysis*

Eduardo Casas-Alvero, *Analytic Projective Geometry*

Preface

This book originates from a set of lecture notes for graduate classes I delivered since 2005, first at the Paul Sabatier University in Toulouse, and then at Purdue University in West Lafayette. The lecture notes benefited much from the input and criticism from several students and have been modified and expanded numerous times before reaching the final form of this book.

My motivation is to present at the graduate level and in a concise but complete way the most important tools and ideas of the classical theory of continuous time processes and at the same time introduce the readers to more advanced theories: the theory of Dirichlet forms, the Malliavin calculus and the Lyons rough paths theory.

Several exercises of various levels are distributed throughout the text in order to test the understanding of the reader. Results proved in these exercises are sometimes used in later parts of the text so I really encourage the reader to have a dynamic approach in his reading and to try to solve the exercises.

I included at the end of each chapter a short section listing references for the reader wishing to complement his reading or looking for more advanced theories and topics.

Chapters 1, 2, 5 and 6 are essentially independent from Chapters 3 and 4. I often used the materials in Chapters 1, 2, 5 and 6 as a graduate course on stochastic calculus and Chapters 3 and 4 on their own as a course on Markov processes and Markov semigroups assuming some of the basic results of Chapter 1. Chapter 7 is almost entirely independent from the other chapters. It is an introduction to Lyons rough paths theory which offers a deterministic approach to understand differential equations driven by very irregular processes including as a special case Brownian motion.

To conclude, I would like to express my gratitude to the students who pointed out typos and inaccuracies in various versions of the lecture notes leading to this book and to my colleague Cheng Ouyang for a detailed reading of an early draft of the manuscript. Of course, all the remaining typos and mistakes are my own responsibility and a list of corrections will be kept online on my personal blog. Finally, I thank Igor Kortchmeski for letting me use his nice picture of a random stable tree on the cover of the book.

West Lafayette, May 2014

Fabrice Baudoin

Conventions and frequently used notations

Unless specified otherwise, the Borel measures we consider on \mathbb{R}^n are assumed to be Radon measures, that is, are finite on compact sets.

$\mathbb{R}_{\geq 0}$	$[0, +\infty)$
$\mathbb{R}^{n \times p}$	set of $n \times p$ matrices
cA or A^c	complement of the set A
$[x]$	integer part of x
$x \wedge y$	minimum between x and y
$\Gamma(x)$	$\Gamma(x) = \int_0^{+\infty} t^{x-1} e^{-t} dt$
$\mathcal{A}(A, B)$	set of functions $A \rightarrow B$
$\mathcal{C}(A, B)$	set of continuous functions $A \rightarrow B$
$\mathcal{C}^k(A, B)$	functions $A \rightarrow B$, k -times continuously differentiable
$\mathcal{C}_c(A, B)$	functions $A \rightarrow B$, smooth and compactly supported inside A
$\mathcal{C}_0(A, B)$	continuous functions $A \rightarrow B$ whose limit at ∞ is 0
$\mathcal{C}^{k,l}(A \times B, C)$	functions $A \times B \rightarrow C$ which are k -times continuously differentiable in the first variable and l times in the second
$\mathcal{T}(A, B)$	σ -field on $\mathcal{A}(A, B)$ generated by cylinders
$\mathcal{B}(A, B)$	σ -field on $\mathcal{C}(A, B)$ generated by cylinders
$\mathcal{B}(A)$	Borel σ -field on A
$L^p_\mu(A, B)$	L^p space of functions $A \rightarrow B$ for the measure μ
$L^p(\mathcal{F}, \mathbb{P})$	real L^p space of \mathcal{F} measurable random variables
$\Delta_n[0, t]$	$\{0 = t_0^n \leq t_1^n \leq \dots \leq t_n^n = t\}$
$\int H_s dM_s$	Itô integral
$\int H_s \circ dM_s$	Stratonovitch integral
$\mathcal{H}_s(\mathbb{R}^n)$	Sobolev space of order s
$\mathcal{H}_s^0(\Omega)$	closure of $\mathcal{C}_c(\Omega, \mathbb{C})$ in $\mathcal{H}_s(\mathbb{R}^n)$
\mathbf{D}	Malliavin derivative
δ	divergence operator
$\mathbb{D}^{k,p}$	domain of \mathbf{D}^k in $L^p(\mathcal{F}, \mathbb{P})$
$\ \cdot\ _{p\text{-var}, [s, t]}$	p -variation norm on $[s, t]$
$\ \cdot\ _{\infty, [s, t]}$	supremum norm on $[s, t]$
$C^{p\text{-var}}([s, t], \mathbb{R}^d)$	continuous paths $[s, t] \rightarrow \mathbb{R}^d$ with bounded p -variation
$\int_{\Delta^k[s, t]} dx^I$	$\int_{s \leq t_1 \leq t_2 \leq \dots \leq t_k \leq t} dx^{i_1}(t_1) \dots dx^{i_k}(t_k)$
$\Omega^p([0, T], \mathbb{R}^d)$	space of p -rough paths

Contents

Preface	v
Conventions and frequently used notations	xi
Introduction	1
1 Stochastic processes	6
1.1 Measure theory in function spaces	6
1.2 Stochastic processes	8
1.3 Filtrations	9
1.4 The Daniell–Kolmogorov extension theorem	11
1.5 The Kolmogorov continuity theorem	16
1.6 Stopping times	18
1.7 Martingales	20
1.8 Martingale inequalities	32
Notes and comments	34
2 Brownian motion	35
2.1 Definition and basic properties	35
2.2 Basic properties	39
2.3 The law of iterated logarithm	42
2.4 Symmetric random walks	45
2.5 Donsker theorem	55
Notes and comments	62
3 Markov processes	63
3.1 Markov processes	63
3.2 Strong Markov processes	72
3.3 Feller–Dynkin diffusions	75
3.4 Lévy processes	87
Notes and comments	96
4 Symmetric diffusion semigroups	97
4.1 Essential self-adjointness, spectral theorem	97
4.2 Existence and regularity of the heat kernel	106
4.3 The sub-Markov property	109
4.4 L^p -theory: The interpolation method	112

4.5	L^p -theory: The Hille–Yosida method	114
4.6	Diffusion semigroups as solutions of a parabolic Cauchy problem	125
4.7	The Dirichlet semigroup	127
4.8	The Neumann semigroup	131
4.9	Symmetric diffusion processes	132
	Notes and comments	135
5	Itô calculus	138
5.1	Variation of the Brownian paths	138
5.2	Itô integral	140
5.3	Square integrable martingales and quadratic variations	146
5.4	Local martingales, semimartingales and integrators	154
5.5	Döblin–Itô formula	161
5.6	Recurrence and transience of the Brownian motion in higher dimensions	166
5.7	Itô representation theorem	168
5.8	Time changed martingales and planar Brownian motion	171
5.9	Burkholder–Davis–Gundy inequalities	175
5.10	Girsanov theorem	178
	Notes and comments	183
6	Stochastic differential equations and Malliavin calculus	185
6.1	Existence and uniqueness of solutions	185
6.2	Continuity and differentiability of stochastic flows	190
6.3	The Feynman–Kac formula	193
6.4	The strong Markov property of solutions	199
6.5	Stratonovitch stochastic differential equations and the language of vector fields	201
6.6	Malliavin calculus	205
6.7	Existence of a smooth density	216
	Notes and comments	219
7	An introduction to Lyons’ rough paths theory	221
7.1	Continuous paths with finite p -variation	221
7.2	The signature of a bounded variation path	226
7.3	Estimating iterated integrals	228
7.4	Rough differential equations	239
7.5	The Brownian motion as a rough path	245
	Notes and comments	254
	Appendix A Unbounded operators	257

Appendix B Regularity theory	262
References	269
Index	275

Introduction

The first stochastic process that has been extensively studied is the Brownian motion, named in honor of the botanist Robert Brown, who observed and described in 1828 the random movement of particles suspended in a liquid or gas. One of the first mathematical studies of this process goes back to the mathematician Louis Bachelier, in 1900, who presented in his thesis [2] a stochastic modelling of the stock and option markets. But, mainly due to the lack of rigorous foundations of probability theory at that time, the seminal work of Bachelier has been ignored for a long time by mathematicians. However, in his 1905 paper, Albert Einstein brought this stochastic process to the attention of physicists by presenting it as a way to indirectly confirm the existence of atoms and molecules. The rigorous mathematical study of stochastic processes really began with the mathematician Andrei Kolmogorov. His monograph [46] published in Russian in 1933 built up probability theory in a rigorous way from fundamental axioms in a way comparable to Euclid's treatment of geometry. From this axiomatic, Kolmogorov gives a precise definition of stochastic processes. His point of view stresses the fact that a stochastic process is nothing else but a random variable valued in a space of functions (or a space of curves). For instance, if an economist reads a financial newspaper because he is interested in the prices of barrel of oil for last year, then he will focus on the curve of these prices. According to Kolmogorov's point of view, saying that these prices form a stochastic process is then equivalent to saying that the curve that is seen is the realization of a random variable defined on a suitable probability space. This point of view is mathematically quite deep and provides existence results for stochastic processes as well as pathwise regularity results.

Joseph Doob writes in the introduction to his famous book "Stochastic processes" [19]:

A stochastic process is any process running along in time and controlled by probability laws...more precisely any family of random variables where a random variable ... is simply a measurable function ...

Doob's point of view, which is consistent with Kolmogorov's and built on the work by Paul Lévy, is nowadays commonly given as a definition of a stochastic process. Relying on this point of view that emphasizes the role of time, Doob's work, developed during the 1940s and the 1950s has quickly become one of the most powerful tools available to study stochastic processes.

Let us now describe the seminal considerations of Bachelier. Let X_t denote the price at time t of a given asset on a financial market (Bachelier considered a given quantity of wheat). We will assume that $X_0 = 0$ (otherwise, we work with $X_t - X_0$). The first observation is that the price X_t can not be predicted with absolute certainty.

It seems therefore reasonable to assume that X_t is a random variable defined on some probability space. One of the initial problems of Bachelier was to understand the distribution of prices at given times, that is, the distribution of the random variable $(X_{t_1}, \dots, X_{t_n})$, where t_1, \dots, t_n are fixed.

The two following fundamental observations of Bachelier were based on empirical observations:

- If τ is very small then, in absolute value, the price variation $X_{t+\tau} - X_t$ is of order $\sigma\sqrt{\tau}$, where σ is a positive parameter (nowadays called the volatility of the asset).
- The expectation of a speculator is always zero¹ (nowadays, a generalization of this principle is called the absence of arbitrage).

Next, Bachelier assumes that for every $t > 0$, X_t has a density with respect to the Lebesgue measure, let us say $p(t, x)$. It means that if $[x - \varepsilon, x + \varepsilon]$ is a small interval around x , then

$$\mathbb{P}(X_t \in [x - \varepsilon, x + \varepsilon]) \simeq 2\varepsilon p(t, x).$$

The two above observations imply that for τ small,

$$p(t + \tau, x) \simeq \frac{1}{2}p(t, x - \sigma\sqrt{\tau}) + \frac{1}{2}p(t, x + \sigma\sqrt{\tau}).$$

Indeed, due to the first observation, if the price is x at time $t + \tau$, it means that at time t the price was equal to $x - \sigma\sqrt{\tau}$ or to $x + \sigma\sqrt{\tau}$. According to the second observation, each of these cases occurs with probability $\frac{1}{2}$.

Now Bachelier assumes that $p(t, x)$ is regular enough and uses the following approximations coming from a Taylor expansion:

$$\begin{aligned} p(t + \tau, x) &\simeq p(t, x) + \tau \frac{\partial p}{\partial t}(t, x), \\ p(t, x - \sigma\sqrt{\tau}) &\simeq p(t, x) - \sigma\sqrt{\tau} \frac{\partial p}{\partial x}(t, x) + \frac{1}{2}\sigma^2\tau \frac{\partial^2 p}{\partial x^2}(t, x), \\ p(t, x + \sigma\sqrt{\tau}) &\simeq p(t, x) + \sigma\sqrt{\tau} \frac{\partial p}{\partial x}(t, x) + \frac{1}{2}\sigma^2\tau \frac{\partial^2 p}{\partial x^2}(t, x). \end{aligned}$$

This yields the equation

$$\frac{\partial p}{\partial t} = \frac{1}{2}\sigma^2 \frac{\partial^2 p}{\partial x^2}(t, x).$$

This is the so-called heat equation, which is the primary example of a diffusion equation. Explicit solutions to this equation are obtained using the Fourier transform, and by using the fact that at time 0, p is the Dirac distribution at 0, it is

¹Quoted and translated from the French: *It seems that the market, the aggregate of speculators, can believe in neither a market rise nor a market fall, since, for each quoted price, there are as many buyers as sellers.*

computed that

$$p(t, x) = \frac{e^{-\frac{x^2}{2\sigma^2 t}}}{\sigma \sqrt{2\pi t}}.$$

It means that X_t has a Gaussian distribution with mean 0 and variance σ^2 . Let now $0 < t_1 < \dots < t_n$ be fixed times and I_1, \dots, I_n be fixed intervals. In order to compute $\mathbb{P}(X_{t_1} \in I_1, \dots, X_{t_n} \in I_n)$ the next step is to assume that the above analysis did not depend on the origin of time, or, more precisely, that the best information available at time t is given by the price X_t . This leads to the following computation:

$$\begin{aligned} \mathbb{P}(X_{t_1} \in I_1, X_{t_2} \in I_2) &= \int_{I_1} \mathbb{P}(X_{t_2} \in I_2 | X_{t_1} = x_1) p(t_1, x_1) dx_1 \\ &= \int_{I_1} \mathbb{P}(X_{t_2-t_1} + x_1 \in I_2 | X_{t_1} = x_1) p(t_1, x_1) dx_1 \\ &= \int_{I_1 \times I_2} p(t_2 - t_1, x_2 - x_1) p(t_1, x_1) dx_1 dx_2, \end{aligned}$$

which is easily generalized to

$$\begin{aligned} &\mathbb{P}(X_{t_1} \in I_1, \dots, X_{t_n} \in I_n) \\ &= \int_{I_1 \times \dots \times I_n} p(t_n - t_{n-1}, x_n - x_{n-1}) \dots p(t_2 - t_1, x_2 - x_1) p(t_1, x_1) dx_1 dx_2 \dots dx_n. \end{aligned} \tag{0.1}$$

In many regards, the previous computations were not rigorous but heuristic. One of our first motivations is to provide a rigorous construction of this object X_t on which Bachelier worked and which is called a Brownian motion.

From a rigorous point of view the question is: Does there exist a sequence of random variables $\{X_t, t \geq 0\}$ such that $t \rightarrow X_t$ is continuous and such that the property (0.1) is satisfied? Chapter 1 will give a positive answer to this question. We will see how to define and construct processes. In particular we will prove the existence of Brownian motion and then study several of its properties.

Chapter 1 sets the foundations. It deals with the basic definitions and results that are required to rigorously deal with stochastic processes. We introduce the relevant σ -fields and prove the fundamental Daniell–Kolmogorov theorem which may be seen as an infinite-dimensional version of the Carathéodory extension of measure theorem. It is the basic theorem to prove the existence of a stochastic process. However, despite its importance and usefulness, the Daniell–Kolmogorov result relies on the axiom of choice and as such is non-constructive and does not give any information or insight about the stochastic process that has been proved to exist. The

Kolmogorov continuity theorem fills one of these gaps and gives a useful criterion to ensure that we can work with a process whose sample paths are continuous. Chapter 1 also includes a thorough study of continuous martingales. We focus on Doob's theorems: The stopping theorem, the regularization result and the maximal inequalities. Martingale techniques are essential to study stochastic processes. They give the tools to handle stopping times which are naturally associated to processes and provide the inequalities which are the cornerstones of the theory of stochastic integration which is presented in Chapter 5.

Chapter 2 is devoted to the study of the most important stochastic process: The Brownian motion. As a consequence of the Daniell–Kolmogorov and Kolmogorov continuity theorems, we prove the existence of this process and then study some of its most fundamental properties. From many point of views, Brownian motion can be seen as the continuous random walk in continuous time. This is made precise at the end of the chapter where we give an alternative proof of the existence of the Brownian motion as a limit of suitably rescaled random walks.

Chapter 3 is devoted to the study of Markov processes. Our goal is to emphasize the role of the theory of semigroups when studying Markov processes. More precisely, we wish to understand how one can construct a Markov process from a semigroup and then see what are the properties inherited from the semigroup to the sample path properties of the process. We will particularly focus on the class of Feller–Dynkin Markov processes which are a class of Markov processes enjoying several nice properties, like the existence of regular versions and the strong Markov property. We finish the chapter with a study of the Lévy processes which form an important subset of the class of Feller–Dynkin Markov processes.

Chapter 4 can be thought an introduction to the theory of symmetric Dirichlet forms. As we will see, this theory and the set of tools attached to it belong much more to functional analysis than to probability theory. The basic problem is the construction of a Markov semigroup and of a Markov process only from the generator. More precisely, the question is: Given a diffusion operator L , does L generate a Markov semigroup P_t and if yes, is this semigroup the transition semigroup of a continuous Markov process? We will answer positively this question in quite a general framework under the basic assumption that L is elliptic and essentially self-adjoint with respect to some Borel measure.

Chapter 5 is about stochastic calculus and its applications to the study of Brownian motion. Stochastic calculus is an integral and differential calculus with respect to Brownian motion or more generally with respect to martingales. It allows one to give a meaningful sense to integrals with respect to Brownian motion paths and to define differential equations driven by such paths. It is not a straightforward extension of the usual calculus because Brownian motion paths are not regular enough, they are only γ -Hölder continuous with $\gamma < 1/2$. The range of applications of the stochastic calculus is huge and still growing today. We mention in particular that the applications to mathematical finance have drawn a lot of attention on this