

Josephine Barnes
**Lecture Notes on
Gynaecology**

Third edition

LECTURE N GYNAECOLOGY

JOSEPHINE BARNES

D.B.E., M.A., D.M., F.R.C.P., F.R.C.S., F.R.C.O.G.

*Consultant Obstetrician and Gynaecologist
Charing Cross Hospital and
Elizabeth Garrett Anderson Hospital*

*Examiner in Obstetrics and Gynaecology
Universities of Oxford and London and
the Royal College of Obstetricians and Gynaecologists
Formerly Examiner to the Queen's University of
Belfast and the Conjoint Board*

THIRD EDITION

CKWELL SCIENTIFIC PUBLICATIONS
OXFORD LONDON EDINBURGH MELBOURNE

TO PENNY, AMANDA
AND ANTONY

© 1971, 1975 BY BLACKWELL SCIENTIFIC PUBLICATIONS
Osney Mead, Oxford
85 Marylebone High Street, London W1M 3DE
9 Forest Road, Edinburgh
P.O. Box 9, North Balwyn, Victoria, Australia

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without the prior permission of the copyright owner.

ISBN 0 632 00197 6

FIRST PUBLISHED 1966
REVISED REPRINT 1969
SECOND EDITION 1971
THIRD EDITION 1975

Distributed in the United States of America by
J. B. Lippincott Company, Philadelphia
and in Canada by
J. B. Lippincott Company of Canada Ltd, Toronto

Printed in Great Britain by
Western Printing Services Ltd, Bristol

PREFACE TO THE THIRD EDITION

This book has appeared to fulfil its role as a comprehensive survey of the field of gynaecology intended especially for medical students preparing for their final examinations.

It has been pleasant to discover that the book has achieved a wider scope and has proved of use to post-graduates and to student nurses.

In preparing the third edition the aim has been to bring the book as closely up to date as is possible and this has meant that some new material has had to be included. Nevertheless the book has not increased greatly in size and the aim has always been to keep it within a reasonable scope and yet to make it comprehensive and up to date.

PREFACE TO THE FIRST EDITION

This book has grown out of lectures and tutorials given over the years, mainly to medical students but also to postgraduates and student nurses. In giving tutorials in gynaecology to final year students I have always felt aware of the large syllabus they have to cover in medicine, surgery and obstetrics as well. Tutorials are thus designed to give a comprehensive survey of the subject without omitting any condition of importance to the student or postgraduate, but with a minimum of detailed study of rare conditions.

Great changes have occurred in the field of gynaecology in recent years. Some of these have resulted from general scientific progress in all fields of medical science and from improvements in operating techniques. The discovery of hormones has also changed the treatment of many conditions and a certain amount of space has therefore been devoted to endocrinology including its application in oral contraception. But in spite of these advances the basic needs of the doctor and patient remain unchanged; the student must have a basic knowledge of anatomy, physiology and pathology, and must learn how to make a clinical examination, to interpret his findings and to understand the significance of symptoms and signs.

In compiling these lecture notes I have constantly felt indebted to my own teachers and especially to those from whom I learnt the importance of thorough examination and accurate diagnosis. In particular I must mention Professor F. J. Browne who taught me above all the importance of teaching in simple and easily understood terms. I must also mention another great teacher, Clifford White, whose tutorials to final year

students at University College Hospital helped so many to reach the standard required for the final examinations in obstetrics and gynaecology.

This book is to a great extent based on what I learnt from these two great men and from many others who devote their professional lives to the training of good doctors.

ACKNOWLEDGMENTS

My thanks for help in the production of this book are due to my two secretaries, Miss Doris Le Bas and Miss Susan Horsey, who typed the manuscript. I am indebted to Dr Ella Prieskel for the X-ray reproduced in Figure 11 and to Mrs E. M. Lucas at Charing Cross Hospital Medical School, who compiled the Index.

I must again thank the staff of Blackwell Scientific Publications for their continued encouragement in the production of this book.

I should also like to thank Mr James Smibert, F.R.C.O.G., Melbourne, Australia, who read the second edition and made many helpful comments and corrections. In the preparation of the third edition I was greatly helped by three medical students from Charing Cross Hospital Medical School, C. M. Backhouse, A. J. Green and V. J. Green. For all the help I have received I am deeply grateful.

I have to thank the Conjoint Board, the University of London and the University of Oxford for permission to reproduce specimen examination questions from recent Final Medical examinations.

J. B.

CONTENTS

Preface to the Third Edition	v
Preface to the First Edition	vii
Acknowledgments	viii
1 Essential Gynaecological Anatomy	1
2 Clinical Examinations in Gynaecology	28
3 Menstruation and its Disorders	38
4 Infections—Vaginitis and Cervicitis	58
5 Infections—Salpingitis and Endometritis	69
6 Female Genital Tuberculosis	77
7 Benign Tumours of the Uterus	82
8 Malignant Tumours of the Uterus	94
9 Pelvic Endometriosis	108
10 Tumours of the Ovary	116
11 Displacements of the Uterus	132
12 Prolapse	139

13	Diseases of the Vulva	151
14	Problems of Fertility and Marriage (Infertility, Contraception, Sterilization, Dyspareunia)	163
15	Complications of Early Pregnancy	187
16	Essential Gynaecological Endocrinology	205
17	Gynaecological Operations	220
	Specimen Examination Questions	234
	Index	242

CHAPTER 1. ESSENTIAL GYNAECOLOGICAL ANATOMY

THE OVARY

The *ovaries* are situated in the pelvis one on each side of the uterus and in close relation to the Fallopian tube. Each ovary is attached to the back of the broad ligament by a peritoneal fold called the *mesovarium*; the blood supply, venous and lymphatic drainage of the ovary and its nerve supply pass through the mesovarium. A fibromuscular cord, the *round ligament of the ovary* attaches it to the back of the uterus. It represents the remains of the *gubernaculum*.

The ovary is shaped like an almond and its surface is often wrinkled. The ovaries vary much in size in different women and even in the same woman; the approximate dimensions are 3.5 to 5 cm. long, 2.5 cm. wide and 1.5 cm. thick. Each ovary weighs 5 to 10 grammes.

STRUCTURE OF THE OVARY

The *ovary* is described as having two main parts, the *cortex* and the *medulla* (Fig. 1). In fact it consists of large numbers of ova supported by a connective tissue stroma.

The outer surface of the ovary is covered by a single layer of cubical epithelium, the *germinal epithelium*; this layer is often missing from ovaries removed from adult women. Beneath this is the fibrous capsule of the ovary, the *tunica albuginea*, a protective layer derived from the fibrous connective tissue.

The cortex of the ovary contains thousands of *primordial follicles*. Each consists of an *oögonium* or egg with a single layer of granulosa cells, specially differentiated stroma cells which




FIG. 1. Histology of the normal ovary

secrete hormones. The primordial follicle is 0.25 to 1 mm. in diameter.

During each menstrual cycle one primordial follicle undergoes full ripening to become a Graafian follicle (Fig. 2). The granulosa cells multiply and secrete clear fluid, the *liquor folliculi*. This pushes the oögonium, now 0.1 mm. in diameter and called the oöcyte, to one side. The oöcyte is surrounded by a clear area, the *zona pellucida*, and invested with granulosa cells, the *corona radiata*. The oöcyte with its granulosa layer projects into the follicle as the discus proligerus (cumulus oöphori). The follicle is lined with more granulosa cells, the *membrana granulosa*. The stroma cells outside the membrana granulosa become differentiated to form hormone secreting cells, the *theca interna*. The stroma cells outside this form a false capsule, the *theca externa*, which has no hormone secreting function.

The ripe follicle is 7 to 8 mm. in diameter. At ovulation it ruptures, releasing the oöcyte with its corona radiata. This is called *ovulation*.

Shortly before ovulation the process of *meiosis* is completed in the primary oöcyte. The oöcyte casts off the *first polar body* and as a result the number of chromosomes in the nucleus is reduced from 46 to 23, the primary oöcyte and the first polar body both containing the haploid number (23) of chromosomes. At fertilization the primary oöcyte unites with a spermatozoön,




FIG. 2. Maturing Graafian follicle

which also has the haploid number of chromosomes (23); this secondary oöcyte, now properly called an ovum, has 46 chromosomes. It casts off the *second polar body* and becomes a *zygote*. The first polar body also divides, making three in all. If the primary oöcyte is not fertilized it quickly shrivels and degenerates.

The Graafian follicle now collapses and the cells undergo luteinization; the granulosa cells become *luteal cells* and the theca interna *theca lutein cells*. A *corpus luteum* develops and projects from the surface of the ovary. It is recognized by its crenated outline and the yellow appearance of the cells which secrete *oestrogen* and *progesterone*. If the ovum is not fertilized

the corpus luteum degenerates in about 10 days. A small amount of bleeding occurs into the cavity, the cells undergo hyaline degeneration and a *corpus albicans* is formed. If pregnancy occurs, the corpus luteum grows and may reach 3 cm. in diameter. It persists for 80 to 120 days and then gradually degenerates.

Some follicles mature but do not rupture and no corpus luteum is formed. If this is the case the liquor is gradually absorbed and the follicle regresses forming a *corpus fibrosum*.

As the ovary ages the number of follicles diminishes and the number of corpora albicantia increases, while the stroma gradually becomes more fibrous.

RELATIONS OF THE OVARY

The *ovary* lies free in the *peritoneal cavity* and attached to the back of the *broad ligament* which forms its anterior relation




FIG. 3. Anatomy of the Fallopian tube

(Fig. 3). The *Fallopian tube* curls over the ovary and is *medial*, *superior* and *lateral* to it. The ovary itself lies in the *ovarian fossa* (fossa ovarica); its posterior relations are the *peritoneum* lining the posterior wall of the true pelvis. Behind this is the *common iliac artery and vein*. The common iliac artery divides at this point into external and internal iliac (hypogastric) arteries and the ovary is more often related to the *internal iliac vessels*. The *ureter* enters the pelvis in front of the bifurcation

of the common iliac artery and is thus a close posterior relation of the ovary. Lateral to the ovary is the lateral wall of the pelvis with the *obturator internus muscle* and the *obturator vessels and nerve*. Above the Fallopian tube and ovary are loops of bowel. On the left side the ovary is closely related to the *pelvic colon* and its *mesentery*. On the right side it may be related to the *caecum* and to a *pelvic appendix*.

The *infundibulo-pelvic ligament* is a double fold of peritoneum, continuous with the outer end of the broad ligament. It contains the ovarian vessels, lymphatics and nerves.

THE FALLOPIAN TUBE

The *Fallopian tube* or *uterine tube* is embryologically part of the uterus. It is the oviduct conveying ova to the uterine cavity. Fertilization usually takes place in the tube. It consists of four parts: the *interstitial* or *intramural* part traverses the wall of the uterus; it is 1 to 2 cm. long and about 1 mm. in diameter; the *isthmus* is narrow and lies near the uterus; it is 2 to 3 cm. long and its diameter is 1 to 2 mm.; the *ampulla* is wide, thin-walled and curved, measuring 5 cm. in length; the *infundibulum* is the outer end of the tube and is trumpet shaped with the *fimbriae*, frond-like processes that surround the outer opening of the tube or *abdominal ostium*. One fimbria is longer than the others and is attached to the ovary, the *fimbria ovarica*. In life the fimbriae have a sucking action which wafts the ovum into the tube.

STRUCTURE OF THE FALLOPIAN TUBE

The tube has three coats, the outer *serous* layer consisting of peritoneum; this covers the tube except in its intramural part and over a small area over its attachment to the broad ligament. The *muscle* layer consists of *outer longitudinal* and *inner circular* layers of *smooth muscle*. A delicate layer of connective tissue separates the muscle layer from the mucosa, the *endosalpinx* which lines the tube and is thrown into numerous

longitudinal folds or *rugae*. The rugae consist of a core of connective tissue covered with a tall columnar epithelium. In the inner part of the tube the epithelium is thrown into only three or four folds and the epithelial cells are smaller.

HISTOLOGY

Three main types of cell are found: 1. Ciliated cells. 2. Non-ciliated or 'secretory' cells which may be pear-shaped or dumb-bell shaped. 3. 'Peg' or intercalary cells. These are cells with long narrow nuclei, squeezed between the other cells. There are rhythmic changes in the epithelium during the menstrual cycle. In the follicular or proliferative phase the cells increase in height and activity.

RELATIONS OF THE FALLOPIAN TUBE

Anterior to the Fallopian tube is the top of the bladder and the utero-vesical peritoneal pouch. *Superior* to it are coils of intestine with the caecum on the right and the pelvic colon on the left. *Posterior* to it are the ovary and the pouch of Douglas (utero-rectal pouch) and its contents. *Lateral* to it are the structures in the lateral pelvic wall, the peritoneum covering the obturator internus muscle with the obturator vessels and nerve.

Inferior to the Fallopian tube lie the structures contained within the layers of the *broad ligament*; in addition to the tube and the round ligament of the uterus these include the following: the anastomosis between the ovarian and uterine arteries, a plexus of veins, draining into the uterine and ovarian veins: lymphatic vessels and a lymph node lying close to the cervix: a plexus of nerves and the *cervical nerve ganglion*: certain vestigial structures, namely the *epoöphoron*, the *paroöphoron* and *Gärtner's duct* (mesonephric duct).

FUNCTIONS OF THE FALLOPIAN TUBE

The functions of the Fallopian tube are complex and incompletely understood. It actively sucks the oöcyte into its ampulla

where fertilization takes place. Tubal secretion, which contains among other constituents lactate, pyruvate and bicarbonate, is essential for 'capacitation' of the spermatozoön and the oöcyte. After fertilization the ovum must remain in the tube for 3 days to undergo the early stages of division. It is then transported to the uterus, probably by the movement of the cilia in the tubal epithelium, and becomes implanted about $5\frac{1}{2}$ days after fertilization.

THE UTERUS

The *uterus* (Fig. 4) is a hollow muscular organ which lies at about the centre of the true pelvis and is covered over most of its surface with peritoneum. It consists of the following parts:

1. The *fundus* is the part above the opening of the Fallopian tubes.
2. The *cornu* is the part into which the Fallopian tube opens.
3. The *body* is the main part of the cavity of the uterus.
4. The *isthmus* is the narrow lowest part of the body and is bounded above by the *anatomical* internal os and below by the *histological* internal os of the uterus.
5. The *cervix* (neck) runs from the histological internal os to the external os which opens into the vagina.

DIMENSIONS

The nulliparous uterus is about 9 cm. long, 6 cm. wide and 4 cm. thick and weighs about 50 grammes.

The uterus changes in size and shape during a woman's life. In the child the cervix exceeds the body in length but at puberty there is growth of the muscle until the uterus attains adult size. After childbirth the uterus is always larger. At the menopause and after, the body of the uterus atrophies to a varying degree.




FIG. 4. Uterus with cervix at different ages

a—28 weeks fetus

b—at birth

c—1 month

d—2 years

e—10 years

f—2 years, before menarche

g—14 years, after menarche

h—22 years nulliparous

STRUCTURE OF THE UTERUS

The *body*, with the *fundus* of the uterus, is covered with peritoneum except at its sides where the broad ligaments are attached to it. There are two layers of smooth muscle, a thin outer longitudinal layer continuous with that of the Fallopian tube and the vagina, and an inner layer arranged in oblique spirals. There is fibrous tissue among the muscle fibres. The

epithelium is the *endometrium* and rests directly on the muscle layer with no intervening submucous layer.

The *cervix* is a fusiform canal, one-third of which lies above the attachment of the vagina while two-thirds of it projects into the vagina. It is made up mainly of fibrous tissue and elastic tissue; the upper part contains smooth muscle which is condensed in a circular pattern to form the *internal os* or sphincter of the uterus.

The *cervical canal* is lined by columnar epithelium; this is thrown up into several longitudinal folds and numerous lateral folds the *arbor vitae*. The epithelium is continued into the compound racemose glands (distinct from the simple tubular glands of the endometrium) which dip deep into the stroma of the cervix; these glands produce the mucous secretion of the cervix.

The vaginal part of the cervix (*portio vaginalis*) is covered with squamous epithelium continuous with that of the vagina. The junction between the columnar and squamous epithelium is at the *external os*.

RELATIONS OF THE UTERUS

The *body and fundus* of the uterus are covered with peritoneum; in front this is reflected on to the upper surface of the bladder at the level of the internal os. Here the peritoneum is loosely attached, but over the rest of the uterus the attachment is so dense that it cannot be stripped off the uterine muscle. The body of the uterus is related *anteriorly* to the *utero-vesical pouch* and *bladder*; as the bladder fills the uterus becomes more vertical in position. *Laterally* the *broad ligaments* with their contents are attached to the side of the uterus. The *round ligaments* are fibro-muscular cords which contain blood vessels and lymphatics; the round ligament ends by passing through the inguinal canal.

Posteriorly the peritoneum passes down into the pouch of Douglas; this varies in depth in different women; it contains