


SMM 8


Surveys of Modern Mathematics



Lie-Bäcklund-Darboux Transformations

李-巴克兰-达布变换

Y. Charles Li · Artyom Yurov



Lie-Bäcklund-Darboux Transformations

李-巴克兰-达布变换

LI BAKELAN DABU BIANHUAN

Y. Charles Li · Artyom Yurov



Authors

Y. Charles Li
Department of Mathematics
University of Missouri
Columbia, MO 65211, USA

Artyom Yurov
Immanuel Kant Baltic Federal University
236041, Al. Nevsky Street, 14,
Kaliningrad, Russia

Copyright © 2014 by
Higher Education Press
4 Dewai Dajie, Beijing 100120, P. R. China, and
International Press
387 Somerville Ave, Somerville, MA, U. S. A.

All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage and retrieval system, without permission.

图书在版编目 (C I P) 数据

李 - 巴克兰 - 达布变换 = Lie-Bäcklund-Darboux
transformations: 英文 / 李沿光, (俄罗斯) 优洛夫
(Yurov, A.) 著. -- 北京: 高等教育出版社, 2014. 1
ISBN 978-7-04-039056-8

I. ①李… II. ①李… ②优… III. ①微分方程 - 英文 IV. ①O175

中国版本图书馆 CIP 数据核字 (2013) 第 301587 号

策划编辑 王丽萍
责任校对 窦丽娜

责任编辑 王丽萍
责任印制 韩 刚

封面设计 张申申

版式设计 余 杨

出版发行 高等教育出版社
社 址 北京市西城区德外大街4号
邮政编码 100120
印 刷 北京汇林印务有限公司
开 本 787 mm×1092 mm 1/16
印 张 10.75
字 数 200 千字
购书热线 010-58581118

咨询电话 400-810-0598
网 址 <http://www.hep.edu.cn>
<http://www.hep.com.cn>
网上订购 <http://www.landaco.com>
<http://www.landaco.com.cn>
版 次 2014 年 1 月第 1 版
印 次 2014 年 1 月第 1 次印刷
定 价 49.00 元

本书如有缺页、倒页、脱页等质量问题, 请到所购图书销售部门联系调换
版权所有 侵权必究
物 料 号 39056-00

Surveys of Modern Mathematics

Surveys of Modern Mathematics

Mathematics has developed to a very high level and is still developing rapidly. An important feature of the modern mathematics is strong interaction between different areas of mathematics. It is both fruitful and beautiful. For further development in mathematics, it is crucial to educate students and younger generations of mathematicians about important theories and recent developments in mathematics. For this purpose, accessible books that instruct and inform the reader are crucial. This new book series "Surveys of Modern Mathematics" (SMM) is especially created with this purpose in mind. Books in SMM will consist of either lecture notes of introductory courses, collections of survey papers, expository monographs on well-known or developing topics.

With joint publication by Higher Education Press (HEP) inside China and International Press (IP) in the West with affordable prices, it is expected that books in this series will broadly reach out to the reader, in particular students, around the world, and hence contribute to mathematics and the world mathematics community.

Series Editors

Shing-Tung Yau

Department of Mathematics
Harvard University
Cambridge, MA 02138, USA

Lizhen Ji

Department of Mathematics
University of Michigan
530 Church Street
Ann Arbor, MI, USA

Yat-Sun Poon

Department of Mathematics
Surge Building, 202 Surge
University of California at Riverside
Riverside, CA 92521, USA

Jie Xiao

Department of Mathematics
Tsinghua University
Beijing 100084, China

Jean-Pierre Demailly

Institut Fourier
100 rue des Maths
38402 Saint-Martin d'Hères, France

Eduard J.N. Looijenga

Mathematics Department
Universiteit Utrecht
Postbus 80.010 3508 TA
Utrecht Nederland

Neil Trudinger

Centre for Mathematics
and Its Applications
Mathematical Sciences Institute
Australian National University
Canberra, ACT 0200, Australia

Preface

One of the mathematical miracles of the 20th century was the discovery of a group of nonlinear wave equations being integrable. These integrable systems are the infinite dimensional counterpart of the finite dimensional integrable Hamiltonian systems of classical mechanics. Icons of integrable systems are the KdV equation, sine-Gordon equation, nonlinear Schrödinger equation etc. The beauty of the integrable theory is reflected by the explicit formulas of nontrivial solutions to the integrable systems. These explicit solutions bear the iconic names of soliton, multi-soliton, breather, quasi-periodic orbit, homoclinic orbit (the focus of this book) etc. There are several ways now available for obtaining these explicit solutions: Bäcklund transformation, Darboux transformation, and inverse scattering transform. The clear connection among these transforms is still an open question although they are certainly closely related. These transformations can be regarded as the counterpart of the canonical transformation of the finite dimensional integrable Hamiltonian system. Bäcklund transformation originated from a quest for Lie's second type invariant transformation rather than his tangent transformation. That brings the title of this book: Lie-Bäcklund-Darboux Transformations which refer to both Bäcklund transformations and Darboux transformations.

The most famous mathematical miracle of the 20th century was probably the discovery of chaos. When the finite dimensional integrable Hamiltonian systems are under perturbations, their regular solutions can turn into chaotic solutions. For such near integrable systems, existence of chaos can sometimes be proved mathematically rigorously. Following the same spirit, one may attempt to prove the existence of chaos for near integrable nonlinear wave equations viewed as near integrable Hamiltonian partial differential equations. This has been accomplished as summarized in the book [69]. The key ingredients in this theory of chaos in partial differential equations are the explicit formulas for the homoclinic orbit and Melnikov integral. The first author's taste is to use Darboux transformation to obtain the homoclinic orbit and Melnikov integral. This will be the focus of the first part of this book.

The second author's taste is to use Darboux transformation in a diversity of applications especially in higher spatial dimensions. The range of applications crosses many different fields of physics. This will be the focus of the second part of

this book. This book is a result of the second author's several visits at University of Missouri as a Miller scholar.

The first author would like to thank his wife Sherry and his son Brandon, and the second author would like to thank his wife Alla and his son Valerian, for their loving support during this work.

Contents

Chapter 1	Introduction	1
Chapter 2	A Brief Account on Bäcklund Transformations	3
2.1	A Warm-Up Approach	3
2.2	Chen’s Method	4
2.3	Clairin’s Method	5
2.4	Hirota’s Bilinear Operator Method	6
2.5	Wahlquist-Estabrook Procedure	7
Chapter 3	Nonlinear Schrödinger Equation	9
3.1	Physical Background	9
3.2	Lax Pair and Floquet Theory	10
3.3	Darboux Transformations and Homoclinic Orbit	12
3.4	Linear Instability	15
3.5	Quadratic Products of Eigenfunctions	16
3.6	Melnikov Vectors	16
3.7	Melnikov Integrals	20
Chapter 4	Sine-Gordon Equation	23
4.1	Background	23
4.2	Lax Pair	23
4.3	Darboux Transformations	24
4.4	Melnikov Vectors	25
4.5	Heteroclinic Cycle	28
4.6	Melnikov Vectors Along the Heteroclinic Cycle	31
Chapter 5	Heisenberg Ferromagnet Equation	35
5.1	Background	35
5.2	Lax Pair	36
5.3	Darboux Transformations	36
5.4	Figure Eight Structures Connecting to the Domain Wall	37
5.5	Floquet Theory	40
5.6	Melnikov Vectors	42

5.7	Melnikov Vectors Along the Figure Eight Structures	44
5.8	A Melnikov Function for Landau-Lifshitz-Gilbert Equation	46
Chapter 6	Vector Nonlinear Schrödinger Equations	47
6.1	Physical Background	47
6.2	Lax Pair	47
6.3	Linearized Equations	48
6.4	Homoclinic Orbits and Figure Eight Structures	49
6.5	A Melnikov Vector	51
Chapter 7	Derivative Nonlinear Schrödinger Equations	53
7.1	Physical Background	53
7.2	Lax Pair	53
7.3	Darboux Transformations	54
7.4	Floquet Theory	56
7.5	Strange Tori	56
7.6	Whisker of the Strange $\mathbb{T}^2$	59
7.7	Whisker of the Circle	60
7.8	Diffusion	61
7.9	Diffusion Along the Strange $\mathbb{T}^2$	63
7.10	Diffusion Along the Whisker of the Circle	64
Chapter 8	Discrete Nonlinear Schrödinger Equation	67
8.1	Background	67
8.2	Hamiltonian Structure	67
8.3	Lax Pair and Floquet Theory	68
8.4	Examples of Floquet Spectra	70
8.5	Melnikov Vectors	71
8.6	Darboux Transformations	73
8.7	Homoclinic Orbits and Melnikov Vectors	75
Chapter 9	Davey-Stewartson II Equation	79
9.1	Background	79
9.2	Linear Stability	80
9.3	Lax Pair and Darboux Transformations	80
9.4	Homoclinic Orbits	82
9.5	Melnikov Vectors	86
9.5.1	Melnikov Integrals	87
9.5.2	An Example	88
9.6	Extra Comments	91
Chapter 10	Acoustic Spectral Problem	93
10.1	Physical Background	93
10.2	Connection with Linear Schrödinger Operator	94
10.3	Discrete Symmetries of the Acoustic Problem	95

10.4	Crum Formulae and Dressing Chains for the Acoustic Problem · · · ·	95
10.5	Harry-Dym Equation · · · · ·	99
10.6	Modified Harry-Dym Equation · · · · ·	101
10.7	Moutard Transformations · · · · ·	103
Chapter 11	SUSY and Spectrum Reconstructions · · · · ·	107
11.1	SUSY in Two Dimensions · · · · ·	108
11.2	The Level Addition · · · · ·	109
11.3	Potentials with Cylindrical Symmetry · · · · ·	111
11.4	Extended Supersymmetry · · · · ·	112
Chapter 12	Darboux Transformations for Dirac Equation · · · ·	115
12.1	Dirac Equation · · · · ·	115
12.2	Crum Law · · · · ·	117
Chapter 13	Moutard Transformations for the 2D and 3D Schrödinger Equations · · · · ·	123
13.1	A 2D Moutard Transformation · · · · ·	123
13.2	A 3D Moutard Transformation · · · · ·	124
Chapter 14	BLP Equation · · · · ·	127
14.1	The Darboux Transformations for the BLP Equation · · · · ·	127
14.2	Crum Law · · · · ·	128
14.3	Exact Solutions · · · · ·	132
14.4	Dressing From Burgers Equation · · · · ·	134
Chapter 15	Goursat Equation · · · · ·	137
15.1	The Reduction Restriction · · · · ·	139
15.2	Binary Darboux Transformations · · · · ·	141
15.3	Moutard Transformations for 2D-MKdV Equation · · · · ·	143
Chapter 16	Links Among Integrable Systems · · · · ·	147
16.1	Borisov-Zykov's Method · · · · ·	147
16.2	Higher Dimensional Systems · · · · ·	149
16.3	Modified Nonlinear Schrödinger Equations · · · · ·	151
16.4	NLS and Toda Lattice · · · · ·	152
Bibliography	· · · · ·	155
Index	· · · · ·	161

Chapter 1

Introduction

The so-called Bäcklund transformation originated from the studies by S. Lie [80–82] and A. V. Bäcklund [11–15] on the Lie’s second question on the existence of invariant multi-valued surface transformations [5]. Lie’s first question was on the well-known Lie’s tangent transformations. The first example of a Bäcklund transformation was studied on the Bianchi’s geometrical construction of surfaces of constant negative curvatures – pseudospheres [18]. The Gauss equation of a pseudosphere can be rewritten as the sine-Gordon equation. The Bäcklund transformation for the sine-Gordon equation is an invariant transformation with a so-called Bäcklund parameter first introduced by Bäcklund. The Bäcklund parameter is particularly important in Bianchi’s diagram of iterating the Bäcklund transformation to generate a so-called nonlinear superposition law [18, 19]. Immediate further studies on Bäcklund transformations were conducted by J. Clairin [26] and E. Goursat [40].

Darboux transformation was first introduced by Gaston Darboux [29] for the nowadays well-known one-dimensional linear Schrödinger equation – a special form of the Sturm-Liouville equations [84]. Darboux found a covariant transformation for the eigenfunction and the potential. The covariant transformation was built upon a particular eigenfunction at a particular value of the spectral parameter.

At the beginning, it seemed that Bäcklund transformation and Darboux transformation are irrelevant. The first link of the two came about in 1967 when Gardner, Greene, Kruskal, and Miura related KdV equation to its Lax pair of which the spatial part is the one-dimensional linear Schrödinger equation [38]. Soon afterwards, the Bäcklund transformation for the KdV equation was found. This was the beginning of a renaissance of Bäcklund transformations and Darboux transformations. It turned out that the existence of a Lax pair for a nonlinear wave equation, the solvability of the Cauchy problem for the nonlinear wave equation by the inverse scattering transform [38], the existence of a Bäcklund transformation for the nonlinear wave equation, and the existence of a Darboux transformation for the nonlinear wave equation and its Lax pair are closely related (although clear

relation is still not known). Up to now, Bäcklund transformations and Darboux transformations for most of the nonlinear wave equations solvable by the inverse scattering transform, have been found [84, 96]. The potential lies at utilizing these transformations. All the earlier books [3, 5, 27, 41, 84, 91, 95, 96] focus on using Bäcklund or Darboux or inverse scattering transformation to construct multi-soliton solutions. Such multi-soliton solutions are defined on the whole spatial space with decaying boundary conditions. When the integrable system is posed with periodic boundary conditions, the solutions are temporally quasi-periodic or periodic or homoclinic. The first part of this book will focus on homoclinic orbit. Chapters 3–9 contain many valuable formulas for homoclinic orbits and Melnikov integrals. Here the Darboux transformations are not only used to generate explicit formulas for the homoclinic orbits but also interlaced with the isospectral theory of the corresponding Lax pairs to generate Melnikov vectors crucial for building the Melnikov integrals. The integrable systems studied in Chapters 3–9 are the so-called canonical systems each of which models a variety of different phenomena. The formulas can be directly used by the readers to study their own near integrable systems. In Chapter 2, we briefly summarize various methods for deriving Bäcklund transformations. These are all “experimental” or “trial-correction” methods. For a more detailed account on these methods, we refer the readers to the book [96]. There are not many methods for deriving Darboux transformations. The commonly used one is the dressing method [84]. Sometimes Chen’s method in Chapter 2 can be effective too. Again these are all “trial-correction” methods. Chapters 10–16 contain applications of Darboux transformations in more specific physics problems, and various connections among different systems. Here no specific boundary condition is posed.

The future of Lie-Bäcklund-Darboux transformations is very bright. Besides the potential of their important applications and new transformations, it is possible to broaden their notion and still end up with useful transformations. This broadening process had begun long ago, e.g. the group notion in [5], the jet bundle in [96], and the Moutard transformation in this book. The broadened transformations even reached the Euler equations of fluids [63, 79].

Chapter 2

A Brief Account on Bäcklund Transformations

Bäcklund transformation is an invariant transformation that transforms one solution into another of a differential equation. It usually involves partial derivatives, and is easily solvable once the initial solution is given. Since J. Clairin [26], researchers have been trying to find an effective systematic way of obtaining a Bäcklund transformation. It was not successful. The commonly employed approaches are Clairin's method [5, 26, 96], Chen's method [25, 91], Hirota's bilinear operator approach [96], and Wahlquist-Estabrook procedure [96, 103, 104]. These are all "trial and correction" approaches. The most common use of Bäcklund transformations is to obtain multi-soliton solutions to integrable systems. For a detailed account, we refer the readers to [96]. Here we shall briefly go over the above mentioned derivation methods.

2.1 A Warm-Up Approach

Take the sine-Gordon equation for example

$$u_{xy} = \sin u, \quad (2.1)$$

where u is a real-valued function of two variables x and y . Let us assume that a Bäcklund transformation transforms u into v where v also satisfies the same sine-Gordon equation

$$v_{xy} = \sin v, \quad (2.2)$$

and our goal is to find the Bäcklund transformation. Let $w^+ = \frac{1}{2}(u + v)$ and $w^- = \frac{1}{2}(u - v)$ [59], then w^+ and w^- satisfy

$$w_{xy}^+ = \sin w^+ \cos w^-, \quad w_{xy}^- = \cos w^+ \sin w^-. \quad (2.3)$$

We assume that the Bäcklund transformation has the trial form

$$w_x^- = f(w^+), \quad (2.4)$$

then by the second equation of (2.3), one has

$$f'w_y^+ = \cos w^+ \sin w^-. \quad (2.5)$$

By substituting (2.5) into the first equation of (2.3), one has

$$-\frac{w_x^+ \sin w^-}{f'} \left(\frac{f''}{f'} \cos w^+ + \sin w^+ \right) + \cos w^- \left(\frac{f}{f'} \cos w^+ - \sin w^+ \right) = 0,$$

which can be satisfied if one demands

$$\frac{f''}{f'} \cos w^+ + \sin w^+ = 0, \quad \text{and} \quad \frac{f}{f'} \cos w^+ - \sin w^+ = 0.$$

A solution of this over-determined system can be found

$$f = a \sin w^+,$$

where a is an arbitrary constant called a Bäcklund parameter. (2.4) and (2.5) now take the form

$$w_x^- = a \sin w^+, \quad w_y^+ = \frac{1}{a} \sin w^-. \quad (2.6)$$

(2.6) implies (2.3) which is equivalent to (2.1) and (2.2). (2.6) is the Bäcklund transformation for sine-Gordon equation (2.1). The above approach of derivation was developed in [59]. After experimenting with different forms of assumptions like (2.4), one hopes to find a Bäcklund transformation for a given equation.

2.2 Chen's Method

Chen's method [25] is a quite efficient method. Take the KdV equation for example,

$$u_t + 6uu_x + u_{xxx} = 0,$$

its Lax Pair is given by

$$L\psi = \lambda\psi, \quad \partial_t \psi = A\psi,$$

where

$$L = \partial_x^2 + u, \quad A = -4\partial_x^3 - 3(u\partial_x + \partial_x u).$$

Let $v = \psi_x/\psi$, one gets

$$v_x + v^2 + u = \lambda, \quad (2.7)$$

$$\begin{aligned} -v_t &= 4v_{xxx} + 12vv_{xx} + 12v^2v_x + 12v_x^2 \\ &\quad + 6u_xv + 6uv_x + 3u_{xx}. \end{aligned} \quad (2.8)$$

Eliminating u , one finds that v satisfies the modified KdV equation

$$v_t - 6v^2v_x + 6\lambda v_x + v_{xxx} = 0.$$

Relation (2.7) is the Miura transformation between KdV and modified KdV. If v solves the modified KdV, so does $-v$, then one can find $\hat{u}$ by solving KdV such that

$$-v_x + v^2 + \hat{u} = \lambda, \quad (2.9)$$

$$\begin{aligned} v_t = & -4v_{xxx} + 12vv_{xx} - 12v^2v_x + 12v_x^2 \\ & - 6\hat{u}_xv - 6\hat{u}v_x + 3\hat{u}_{xx}. \end{aligned} \quad (2.10)$$

Subtracting (2.9) from (2.7), one gets $v_x = \frac{1}{2}(\hat{u} - u)$. Let $w_x = \frac{1}{2}u$ and $\hat{w}_x = \frac{1}{2}\hat{u}$, then from (2.7) and (2.8), one gets the Bäcklund transformation for KdV equation

$$\begin{aligned} (\hat{w} + w)_x &= \lambda - (\hat{w} - w)^2, \\ (w - \hat{w})_t &= 4(\hat{w} - w)_{xxx} + 12(\hat{w} - w)(\hat{w} - w)_{xx} \\ &\quad + 12(\hat{w} - w)^2(\hat{w} - w)_x + 12(\hat{w} - w)_x^2 \\ &\quad + 12(\hat{w} - w)w_{xx} + 12w_x(\hat{w} - w)_x + 6w_{xxx}, \end{aligned}$$

where λ is the Bäcklund parameter.

2.3 Clairin's Method

Clairin's method is a direct trial method which involves tedious calculations. In general, let $u = u(t, x)$ satisfy some equations, and let $v = v(t, x)$ be another variable linked to u through a transformation of the form

$$u_x = f(u, v, v_x, v_t), \quad u_t = g(u, v, v_x, v_t).$$

The compatibility condition

$$f_t = g_x,$$

hopefully can lead to an equation for v , of course, by virtue of the equation satisfied by u . Consider the focusing cubic nonlinear Schrödinger equation

$$iq_t + q_{xx} + |q|^2q = 0,$$

Lamb started with a transformation of the form [55]

$$\begin{aligned} q_x &= f(q, \bar{q}, p, \bar{p}, p_x, \bar{p}_x), \\ q_t &= g(q, \bar{q}, p, \bar{p}, p_x, \bar{p}_x, p_t, \bar{p}_t). \end{aligned}$$

After some lengthy calculation, Lamb obtained the Bäcklund transformation [55],

$$\begin{aligned} q_x &= p_x - \frac{1}{2}iw\xi + ikv, \\ q_t &= p_t + \frac{1}{2}\xi w_x - k\zeta + \frac{1}{4}iv(|w|^2 + |v|^2), \end{aligned}$$

where

$$\begin{aligned} \xi &= \pm i(b - 2|v|^2)^{1/2}, \quad w = q + p, \\ v &= q - p, \quad \zeta = -\frac{1}{2}iw\xi + ikv, \end{aligned}$$

b and k are arbitrary real constants called Bäcklund parameters, and p and q satisfy the same equation.

2.4 Hirota's Bilinear Operator Method

Hirota [45, 46] introduced certain bilinear operators to convert the nonlinear wave equations into bilinear equations for which Bäcklund transformations can be constructed. The Hirota bilinear operators D_t and D_x act according to

$$D_t^m D_x^n f \circ g = (\partial_t - \partial_{\hat{t}})^m (\partial_x - \partial_{\hat{x}})^n f(t, x) g(\hat{t}, \hat{x})|_{\hat{t}=t, \hat{x}=x}.$$

Consider the KdV equation

$$u_t + 6uu_x + u_{xxx} = 0,$$

by setting $u = 2(\ln f)_{xx}$, one obtains

$$u_t + 6uu_x + u_{xxx} = \partial_x \left[\frac{1}{f^2} D_x (D_t + D_x^3) f \circ f \right].$$

Thus, if f solves the associated bilinear equation

$$D_x (D_t + D_x^3) f \circ f = 0,$$

then u solves the KdV equation. A Bäcklund transformation for the bilinear equation can be obtained as [46]

$$(D_x^2 - \beta) f \circ \tilde{f} = 0, \quad (D_t + 3\beta D_x + D^3) f \circ \tilde{f} = 0,$$

where β is the Bäcklund parameter.