

Cambridge Texts in Chemistry and Biochemistry

Some modern methods of organic synthesis

W. CARRUTHERS

Thrid edition

有机合成中的现代方法 第3版

CAMBRIDGE

世界图书出版公司

SOME MODERN METHODS OF ORGANIC SYNTHESIS

W. CARRUTHERS

*Chemistry Department
University of Exeter*

THIRD EDITION

CAMBRIDGE
UNIVERSITY PRESS

世界图书出版公司

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge CB2 1RP, United Kingdom

CAMBRIDGE UNIVERSITY PRESS

The Edinburgh Building, Cambridge CB2 2RU, UK <http://www.cup.cam.ac.uk>

40 West 20th Street, New York, NY 10011-4211, USA <http://www.cup.org>

10 Stamford Road, Oakleigh, Melbourne 3166, Australia

© Cambridge University Press 1971, 1978, 1987

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 1971

Second edition 1978

Reprinted 1985

Third edition 1986

Reprinted 1987 (with corrections), 1988, 1989, 1990, 1992, 1993, 1996, 1998

A catalogue record for this book is available from the British Library

Library of Congress Cataloguing in Publication data

Carruthers, W.

Some modern methods of organic synthesis.

(Cambridge texts in chemistry and biochemistry)

Includes index.

I. Chemistry, Organic—Synthesis. I. Title.

II. Series

QD262.C33 1986 547'.2 85-21270

ISBN 0 521 32234 0 hardback

ISBN 0 521 31117 9 paperback

This reprint edition is published with the permission of the Syndicate of the Press of the
University of Cambridge, Cambridge, England.

THIS EDITION IS LICENSED FOR DISTRIBUTION AND SALE IN THE PEOPLE'S
REPUBLIC OF CHINA ONLY, EXCLUDING TAIWAN, HONG KONG AND MACAO
AND MAY NOT BE DISTRIBUTED AND SOLD ELSEWHERE.

书 名: Some Modern Methods of Organic Synthesis Third Edition
作 者: W. Carruthers
中译名: 有机合成中的现代方法 第3版
出 版 者: 世界图书出版公司北京公司
印 刷 者: 北京世图印刷厂
发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)
联系电话: 010-64015659, 64038347
电子信箱: kjsk@vip.sina.com
开 本: 24 印 张: 23
出版年代: 2004 年 11 月
书 号: 7-5062-7250-4/O · 519
版权登记: 图字: 01-2004-5387
定 价: 98.00 元

世界图书出版公司北京公司已获得 Cambridge University Press 授权在中国大陆
独家重印发行。

Some modern methods of organic synthesis

Cambridge Texts in Chemistry and Biochemistry

GENERAL EDITORS

S. J. Benkovic
Professor of Chemistry
Pennsylvania State University

D. T. Elmore
Professor of Biochemistry
The Queen's University of Belfast

J. Lewis
Professor of Inorganic Chemistry
University of Cambridge

K. Schofield
Professor of Organic Chemistry
University of Exeter

J. M. Thomas
Professor of Physical Chemistry
University of Cambridge

B. A. Thrush
Professor of Physical Chemistry
University of Cambridge

Preface to the first edition

This book is addressed principally to advanced undergraduates and to graduates at the beginning of their research careers, and aims to bring to their notice some of the reactions used in modern organic syntheses. Clearly, the whole field of synthesis could not be covered in a book of this size, even in a cursory manner, and a selection has had to be made. This has been governed largely by consideration of the usefulness of the reactions, their versatility and, in some cases, their selectivity.

A large part of the book is concerned with reactions which lead to the formation of carbon-carbon single and double bonds. Some of the reactions discussed, such as the alkylation of ketones and the Diels-Alder reaction, are well established reactions whose scope and usefulness has increased with advancing knowledge. Others, such as those involving phosphorus ylids, organoboranes and new organometallic reagents derived from copper, nickel, and aluminium, have only recently been introduced and add powerfully to the resources available to the synthetic chemist. Other reactions discussed provide methods for the functionalisation of unactivated methyl and methylene groups through intramolecular attack by free radicals at unactivated carbon-hydrogen bonds. The final chapters of the book are concerned with the modification of functional groups by oxidation and reduction, and emphasise the scope and limitations of modern methods, particularly with regard to their selectivity.

Discussion of the various topics is not exhaustive. My object has been to bring out the salient features of each reaction rather than to provide a comprehensive account. In general, reaction mechanisms are not discussed except in so far as is necessary for an understanding of the course or stereochemistry of a reaction. In line with the general policy in the series references have been kept to a minimum. Relevant reviews are noted but, for the most part, references to the original literature are given only for points of outstanding interest and for very recent work. Particular reference is made here to the excellent book by H. O. House, *Modern Synthetic Reactions* which has been my guide at several points and on which I have tried to build, I feel all too inadequately.

I am indebted to my friend and colleague, Dr K. Schofield, for much helpful comment and careful advice which has greatly assisted me in writing the book.

W. CARRUTHERS
26 October 1970

Preface to the third edition

The general plan of this third edition follows that of the earlier editions, but the opportunity has been taken to bring the book up to date as far as possible and to take account of advances in knowledge and of new synthetic methods which have come into use since publication of the second edition. Perhaps the most striking trend in synthesis since then has been the development of highly stereoselective reactions and their application in complex syntheses. These reactions include the stereoselective alkylation of carbonyl compounds, stereoselective aldol condensations and stereoselective oxidations, epoxidations and reductions, and these are among the topics discussed in this edition. New methods for the stereoselective formation of carbon-carbon double bonds, and modern applications of the Diels-Alder reaction, particularly its use in the control of stereochemistry in the synthesis of natural products, and the related class of 1,3-dipolar cyclo-addition reactions are also considered. Other sections of the book are concerned with the increasingly important application in synthesis of organo-metallic reagents, including organoboranes and organosilanes and reagents derived from copper, nickel and palladium, and with the continuing interest in selective reactions at unactivated carbon-hydrogen bonds.

The book is addressed principally to advanced undergraduates and to graduates at the beginning of their research careers, and my aim has been to bring out the salient features of the reactions and reagents rather than to provide a comprehensive account. Reaction mechanisms are not discussed except in so far as is necessary for an understanding of the course or stereochemistry of a reaction. To prevent the book from becoming too big some material of less immediate interest which appeared in earlier editions has been excised from this one. Discussion of new reactions is supported by references.

W. CARRUTHERS
May 1985

Contents

	Preface to the first edition	page xi
	Preface to the third edition	xiii
1	Formation of carbon-carbon single bonds	1
	1.1. Alkylation: importance of enolate anions	1
	1.2. Alkylation of relatively acidic methylene groups	4
	1.3. γ -Alkylation of 1,3-dicarbonyl compounds; dianions in synthesis	8
	1.4. Alkylation of ketones	12
	1.5. The enamine and related reactions	26
	1.6. Alkylation of α -thio and α -seleno carbanions	38
	1.7. Umpolung (dipole inversion)	41
	1.8. The aldol reaction	48
	1.9. Allylic alkylation of alkenes	63
	1.10. The dihydro-1,3-oxazine synthesis of aldehydes and ketones	70
	1.11. Coupling of organonickel and organocopper com- plexes	73
	1.12. Reactions of lithium organocuprates: copper-cata- lysed reactions of Grignard reagents	78
	1.13. Synthetic applications of carbenes and carbenoids	87
	1.14. Formation of carbon-carbon bonds by addition of free radicals to alkenes	97
	1.15. Some photocyclisation reactions	104
2	Formation of carbon-carbon double bonds	111
	2.1. β -Elimination reactions	111
	2.2. Pyrolytic <i>syn</i> eliminations	115
	2.3. Sulphoxide-sulphenate rearrangement; synthesis of allyl alcohols	124
	2.4. The Wittig and related reactions	125
	2.5. Alkenes from sulphones	144
	2.6. Decarboxylation of β -lactones	146

2.7.	Stereoselective synthesis of tri- and tetra- substituted alkenes	148
2.8.	Fragmentation reactions	156
2.9.	Oxidative decarboxylation of carboxylic acids	159
2.10.	Alkenes from arylsulphonylhydrazones	162
2.11.	Stereospecific synthesis from 1,2-diols	165
2.12.	Claisen rearrangement of allyl vinyl ethers	167
2.13.	Reductive dimerisation of carbonyl compounds	180
3	The Diels–Alder and related reactions	184
3.1.	General	184
3.2.	The dienophile - heterodienophiles, oxygen as a dienophile	184
3.3.	The diene - acyclic dienes, heterodienes, 1,2-dimethylenecycloalkanes, vinylcycloalkenes and vinylarenes, aromatic hydrocarbons, cyclic dienes, cyclopentadienones and <i>ortho</i> -quinones, furans	198
3.4.	Intramolecular Diels–Alder reactions	215
3.5.	The retro Diels–Alder reaction	219
3.6.	Catalysis by Lewis acids	221
3.7.	Regiochemistry of the Diels–Alder reaction	225
3.8.	Stereochemistry of the Diels–Alder reaction	229
3.9.	Asymmetric Diels–Alder reactions	237
3.10.	Mechanism of the Diels–Alder reaction	240
3.11.	Photosensitised Diels–Alder reactions	243
3.12.	The ene reaction	244
3.13.	Cyclo-addition reactions with allyl cations and allyl anions	253
3.14.	1,3-Dipolar cyclo-addition reactions	256
4	Reactions at unactivated C—H bonds	263
4.1.	The Hoffmann–Loeffler–Freitag reaction	264
4.2.	Cyclisation reactions of nitrenes	268
4.3.	The Barton reaction and related processes - photolysis of organic nitrites and hypohalites, photolysis of <i>N</i> -nitrosoamides	269
4.4.	Reaction of monohydric alcohols with lead tetra-acetate	279
4.5.	Miscellaneous reactions - unsaturated alcohols from hydroperoxides, cyclobutanols by photolysis of ketones, long-range functionalisation of unactivated carbons on the steroid nucleus	280

5	Synthetic applications of organoboranes and organosilanes	288
5.1.	Hydroboration	288
5.2.	Reactions of organoboranes - protonolysis, oxidation, enantioselective synthesis of secondary alcohols from alkenes, isomerisation and cyclisation of alkylboranes	294
5.3.	Formation of carbon-carbon bonds - carbonylation of organoboranes, reaction with α -bromoketones and α -bromoesters, reaction with diazo compounds	302
5.4.	Reactions of alkenylboranes and trialkylalkynylborates	312
5.5.	Free-radical reactions of organoboranes	315
5.6.	Applications of organosilicon compounds in synthesis - protection of functional groups, trimethylsilyl ethers, silyl enol ethers	317
5.7.	Alkenylsilanes and allylsilanes	324
5.8.	Control of rearrangement of carbonium ions by silicon - the β -effect	333
5.9.	α -Silyl carbanions	335
5.10.	β -Silylcarbonyl compounds	339
5.11.	Trimethylsilyl cyanide	341
5.12.	Trimethylsilyl iodide and trimethylsilyl triflate	342
6	Oxidation	344
6.1.	Oxidation of hydrocarbons	346
6.2.	Oxidation of alcohols - chromic acid, chromiumVI oxide-pyridine complexes, manganeseIV oxide, silver carbonate, oxidation via alkoxysulphonium salts, other methods	351
6.3.	Oxidation of carbon-carbon double bonds - diols, epoxides, enantioselective epoxidation of allylic alcohols (Sharpless epoxidation), diastereoselective epoxidation of homoallylic alcohols, synthetic reactions of epoxides, ozonolysis	364
6.4.	Photosensitised oxidation of alkenes	392
6.5.	Palladium-catalysed oxidation of alkenes	397
6.6.	Oxidation of ketones - conversion into $\alpha\beta$ -unsaturated ketones, Baeyer-Villiger oxidation	398
6.7.	Oxidations with ruthenium tetroxide	407
6.8.	Oxidations with thallium(III) nitrate	408

7	Reduction	411
7.1.	Catalytic hydrogenation - the catalyst, selectivity of reduction, reduction of functional groups, stereochemistry and mechanism, homogeneous hydrogenation	411
7.2.	Reduction by dissolving metals - reduction with metal and acid, reduction of carbonyl compounds, reductions with metal in liquid ammonia (Birch reduction), reductive fission of alcohols and halides	431
7.3.	Reduction by hydride-transfer reagents - aluminium alkoxides, lithium aluminium hydride and sodium borohydride, lithium hydridoalkoxyaluminates, lithium aluminium hydride-aluminium chloride reagents, di-isobutylaluminium hydride, sodium cyanoborohydride, trialkylborohydrides	457
7.4.	Reductions with borane and dialkylboranes	478
7.5.	Other methods - Wolff-Kishner reduction, desulphurisation of thio-acetals, di-imide, low-valent titanium species, trialkyltin hydrides, trialkylsilanes	482
	References	492
	Index	515

1 Formation of carbon-carbon single bonds

In spite of the fundamental importance in organic synthesis of the formation of carbon-carbon single bonds there are comparatively few general methods available for effecting this process, and fewer still which proceed in good yield under mild conditions. Many of the most useful procedures involve carbanions, themselves derived from organometallic compounds, or from compounds containing 'activated' methyl or methylene groups. They include reactions which proceed by attack of the carbanion on a carbonyl or conjugated carbonyl group, as in the Grignard reaction, the aldol and Claisen ester condensations and the Michael reaction, and reactions which involve nucleophilic displacement at a saturated carbon atom, as in the alkylation of ketones and the coupling reactions of some organometallic compounds. Other reactions employed in the formation of carbon-carbon bonds involve carbonium ions and pericyclic processes and recently free-radical reactions have been finding useful application. Examples of all of these procedures will be discussed in this chapter.

1.1. Alkylation: importance of enolate anions

It is well known that certain unsaturated groups attached to a saturated carbon atom render hydrogen atoms attached to that carbon relatively acidic, so that the compound can be converted into an anion on treatment with an appropriate base. Table 1.1, taken from House (1965), shows the pK_a values for some compounds of this type and for some common solvents and reagents.

The acidity of the C—H bonds in these compounds is due to a combination of the inductive electron-withdrawing effect of the unsaturated groups and resonance stabilisation of the anion formed by removal of a proton (1.1). Not all groups are equally effective in 'activating' a neighbouring CH_2 or CH_3 ; nitro is the most powerful of the common groups and thereafter the series follows the approximate order $NO_2 > COR > SO_2R > CO_2R > CN > C_6H_5$. Two activating groups reinforce


Table 1.1. Approximate acidities of active methylene compounds and other common reagents

Compound	pK_a	Compound	pK_a
$\text{CH}_3\text{CO}_2\text{H}$	5	$\text{C}_6\text{H}_5\text{COCH}_3$	19
$\text{CH}_2(\text{CN})\text{CO}_2\text{C}_2\text{H}_5$	9	CH_3COCH_3	20
$\text{CH}_2(\text{CO}_2\text{CH}_3)_2$	9	$\text{CH}_3\text{SO}_2\text{CH}_3$	~23
CH_3NO_2	10	$\text{CH}_3\text{CO}_2\text{C}_2\text{H}_5$	~24
$\text{CH}_3\text{COCH}_2\text{CO}_2\text{C}_2\text{H}_5$	11	$\text{CH}_3\text{CO}_2\text{H}$	~24
$\text{CH}_2(\text{CO}_2\text{C}_2\text{H}_5)_2$	13	CH_3CN	~25
CH_3OH	16	$\text{C}_6\text{H}_5\text{NH}_2$	~30
$\text{C}_2\text{H}_5\text{OH}$	18	$(\text{C}_6\text{H}_5)_3\text{CH}$	~40
$(\text{CH}_3)_3\text{COH}$	19	CH_3SOCH_3	~40

(Acidic hydrogen atoms are underlined)


H. O. House, *Modern synthetic reactions*, copyright 1972, W. A. Benjamin, Inc. Menlo Park, California.

each other, as can be seen by comparing diethyl malonate ($pK_a \approx 13$) with ethyl acetate ($pK_a \approx 24$). Acidity is also increased slightly by electron-withdrawing substituents (e.g. sulphide), and decreased by alkyl groups, so that diethyl methylmalonate, for example, has a slightly less acidic C—H group than diethyl malonate itself.




By far the most important activating groups in synthesis are the carbonyl and carboxylic ester groups. Removal of a proton from the α -carbon atom of a carbonyl compound with base gives the corresponding enolate anion, and it is these anions which are involved in base-catalysed condensation reactions of carbonyl compounds, such as the aldol condensation, and in bimolecular nucleophilic displacements (alkylations) (1.2).


The enolate anions should be distinguished from the enols themselves, which are always present in equilibrium with the carbonyl compound. Most monoketones and esters contain only small amounts of enol (< 1



per cent) at equilibrium, but with 1,2- and 1,3-dicarbonyl compounds much higher amounts of enol (> 50 per cent) may be present. In the presence of acid catalysts monoketones may be converted largely into the enol form, and enols are concerned in many acid-catalysed condensations of carbonyl compounds (1.3).



The formation of the enolate anion results from an equilibrium reaction between the carbonyl compound and the base. A competing equilibrium involves the enolate anion and the solvent. Thus, with diethyl malonate in solvent SolH in presence of base B⁻, we have



and to ensure an adequate concentration of the enolate anion at equilibrium clearly both the solvent and the conjugate acid of the base must be much weaker acids than the active methylene compound. The correct choice of base and solvent is thus of great importance if the subsequent alkylation, or other, reaction is to be successful. Reactions must normally be effected under anhydrous conditions since water is a much stronger acid than the usual activated methylene compounds and, if present, would instantly protonate any carbanion produced. Another point of importance is that the solvent must not be a much stronger acid than the conjugate acid of the base, otherwise the equilibrium



will lie too far to the right and lower the concentration of B⁻. For example,

sodamide can be used as base in liquid ammonia or in benzene, but, obviously, not in ethanol. Base-solvent combinations commonly used to convert active methylene compounds into the corresponding anions include sodium methoxide, sodium ethoxide and sodium or potassium t-butoxide in solution in the corresponding alcohol, or as suspensions in ether, benzene or dimethoxyethane. Potassium t-butoxide is a particularly useful reagent, since it is a poor nucleophile and its solutions in different solvents have widely different basic strengths; it is most active in solution in dry dimethyl sulphoxide (Pearson and Buehler, 1974). Metallic sodium or potassium, or sodium hydride, in suspension in benzene, ether or dimethoxyethane, sodamide in suspension in an inert solvent or in solution in liquid ammonia, and solutions of sodium or potassium triphenylmethyl in ether or benzene have also been used with the less 'active' compounds.

For many purposes, however, these traditional bases have now been superseded by the lithium salts of certain sterically hindered secondary amines, particularly lithium diisopropylamide and lithium 2,2,6,6-tetramethylpiperidide (Olofson and Dougherty, 1973) or the alkali metal salts of bis(trimethylsilyl)amine, $\text{HN}(\text{SiMe}_3)_2$ (Colvin, 1978; Smith and Richmond, 1983). These strong amide bases are only weakly nucleophilic, so that they do not themselves attack susceptible functional groups, and they have the added advantage that they are soluble in non-polar, even hydrocarbon, solvents. The insolubility of the traditional bases in most common organic solvents seriously limits their usefulness.


1.2. Alkylation of relatively acidic methylene groups

In order to effect a reasonably rapid reaction it is, of course, necessary to have a high concentration of the appropriate carbanion. Because of their relatively high acidity (see Table 1.1) compounds in which a C—H bond is activated by a nitro group or by two or more carbonyl, ester or cyano groups can be converted largely into their anions with a comparatively weak base such as a solution of sodium ethoxide in ethanol. An alternative procedure is to prepare the enolate in benzene or ether, using finely divided sodium or potassium metal or sodium hydride, which react irreversibly with compounds containing active methylene groups with formation of the metal salt and evolution of hydrogen. β -Diketones can often be converted into their enolates with alkali metal hydroxides or carbonates in aqueous alcohol or acetone.


Much faster alkylation of enolate anions can often be achieved in dimethylformamide, dimethyl sulphoxide, 1,2-dimethoxyethane or hexamethylphosphoramide than in the usual protic solvents. This appears to be due to the fact that the former solvents do not solvate the enolate

anion and thus do not diminish its reactivity as a nucleophile. At the same time they are able to solvate the cation, separating it from the cation-enolate ion pair and leaving a relatively free enolate ion which would be expected to be a more reactive nucleophile than the ion pair (Parker, 1962). Reactions effected with aqueous alkali as base are often improved in the presence of a phase-transfer catalyst such as a tetra-alkylammonium salt (cf. Makosza and Jończyk, 1976).

Alkylation of enolate anions is readily effected with alkyl halides or other alkylating agents. Both primary and secondary alkyl, allyl or benzyl halides may be used successfully, but with tertiary halides poor yields of alkylated product often result because of competing dehydrohalogenation of the halide. It is often advantageous to proceed by way of the toluene-*p*-sulphonate or methanesulphonate rather than a halide. The sulphonates are excellent alkylating agents, and can usually be obtained from the alcohol in a pure condition more readily than the corresponding halides. Epoxides have also been used as alkylating agents, generally reacting at the less substituted carbon atom. Attack of the enolate anion on the alkylating agent takes place by an S_N2 pathway and thus results in inversion of configuration at the carbon atom of the alkylating agent.



With secondary and tertiary allylic halides or sulphonates, reaction of an enolate anion may give mixtures of products formed by competing attack at the α - and γ -positions (1.7).



A difficulty sometimes encountered in the alkylation of active methylene compounds is the formation of unwanted dialkylated products. During the alkylation of diethyl sodiomalonate, the monoalkyl derivative formed initially is in equilibrium with its anion as indicated in the first equation of (1.8). In ethanol solution, dialkylation does not take place to any appreciable extent because ethanol is sufficiently acidic to reduce