

V. P. WIGGLESWORTH

The Physiology of Insect Metamorphosis

THE PHYSIOLOGY OF INSECT METAMORPHOSIS

BY

V. B. WIGGLESWORTH

C.B.E., M.D., F.R.S.

Fellow of Gonville and Caius College

Quick Professor of Biology in the University of Cambridge

*Director, Agricultural Research Council Unit of
Insect Physiology*



CAMBRIDGE
AT THE UNIVERSITY PRESS
1954

CAMBRIDGE
UNIVERSITY PRESS

University Printing House, Cambridge CB2 8BS, United Kingdom

Cambridge University Press is part of the University of Cambridge.

It furthers the University's mission by disseminating knowledge in the pursuit of education, learning and research at the highest international levels of excellence.

www.cambridge.org

Information on this title: www.cambridge.org/9781107502376

© Cambridge University Press 1954

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 1954

First paperback edition 2015

A catalogue record for this publication is available from the British Library

ISBN 978-1-107-50237-6 Paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

CAMBRIDGE MONOGRAPHS IN
EXPERIMENTAL BIOLOGY

No. 1

EDITORS:

M. ABERCROMBIE, P. M. MEDAWAR
GEORGE SALT (*General Editor*)
M. M. SWANN, V. B. WIGGLESWORTH

THE
PHYSIOLOGY OF INSECT
METAMORPHOSIS

LIST OF PLATES

PLATE I

facing page 44

- (a) *Rhodnius* 4th-stage larva, unfed.
- (b) *Rhodnius* 5th-stage larva, unfed.
- (c) *Rhodnius* adult.
- (d) *Rhodnius* 6th-stage larva produced by implantation of corpus allatum of a 4th-stage larva into a 5th-stage.

PLATE II

facing page 45

- (a) *Rhodnius* larva in 4th stage decapitated one week after feeding, and connected by a capillary tube to a 4th-stage larva decapitated one day after feeding.
- (b) Young larva of *Triatoma* (above) decapitated at one day after feeding, in parabiosis with a 5th-stage larva of *Rhodnius*, decapitated ten days after feeding.
- (c) *Rhodnius* with characters intermediate between larva and adult produced by implanting corpus allatum of a young larva into the abdomen of a 5th-stage larva.
- (d) Giant adult of *Rhodnius* resulting from the metamorphosis of a 6th-stage larva.

PLATE III

facing page 74

- (a) Front view of normal mature larva of silkworm (5th stage). (After Fukuda.)
- (b) Front view of 6th-stage larva of silkworm produced by the implantation of prothoracic glands into a 5th-stage larva early in the instar. It has developed pupal antennae. (After Fukuda.)
- (c) Cocoons of the silkworm: from larva in which the corpora allata were extirpated in the 3rd stage; from larva similarly treated in the 4th stage; from normal larva pupating after the 5th stage. (After Fukuda.)
- (d) Pupae from the cocoons shown in (c). (After Fukuda.)
- (e) Silkmoths derived from the pupae shown in (d). (After Fukuda.)

Illustrations

PLATE IV

facing page 75

- (a) Small patch of larval cuticle on the abdomen of an adult *Rhodnius*, overlying an implanted corpus allatum.
- (b) *Rhodnius* 5th-stage larva showing 'adultoid' characters, following the implantation of the corpus allatum of a newly moulted adult into a 4th-stage larva.
- (c) Normal 4th-stage larva of *Rhodnius*.
- (d) *Rhodnius* 5th-stage larva produced from a 4th-stage to which a 3rd-stage had been joined (fig. 24B). It has developed 4th-instar characters again.
- (e) *Rhodnius* 5th-stage larva produced from a 4th-stage to which another 4th-stage larva seven days after feeding had been joined (fig. 24C). It has developed characters intermediate between a 4th- and a 5th-stage larva.
- (f) Normal 5th-stage larva of *Rhodnius*.

CONTENTS

LIST OF PLATES

page vii

I INTRODUCTION

I

Polymorphism. Embryology and metamorphosis. Origin and evolution of metamorphosis. Origin of the pupa. Imaginal discs. Metamorphosis in *Rhodnius*.

2 HISTOLOGICAL CHANGES DURING MOULTING AND METAMORPHOSIS

II

Growth and moulting in the integument. Chromatolysis. New cuticle formation. Metamorphosis in the integument. Imaginal discs and cell size. Metamorphosis in internal organs: phagocytosis. Growth ratios.

3 PHYSIOLOGY OF GROWTH AND MOULTING

2I

Humoral control of moulting. The brain and moulting. The brain and natural arrest of growth. Thoracic growth centre. Activation of thoracic gland. Thoracic glands in other insects. Growth and moulting in higher Diptera. Growth of imaginal discs in Diptera. Neurosecretion and endocrine organs. Mode of action of thoracic gland hormone. Nature of thoracic gland hormone. Control of hormone production. Moulting of adult insects. Wound healing and moulting.

4 PHYSIOLOGY OF METAMORPHOSIS

5I

The latent imaginal pattern. Humoral control of metamorphosis. Corpus allatum as source of juvenile hormone. Disappearance of thoracic gland. Control of stages towards metamorphosis. Control of pupal characters. Prothetely and metathetely. Hormone balance in normal development. Competence of tissues to undergo metamorphosis. Endocrine glands and reproduction. Endocrine glands and metabolism. Reversal of metamorphosis. Mode of action of corpus allatum hormone. Quantitative action of hormones concerned in growth. Control of function of corpus allatum.

Contents

5	DIFFERENTIATION AND POLYMORPHISM	<i>page</i> 96
	Integration of growth. Relative growth of parts of the body. Nature of the 'substrate'. Determination in epidermis of <i>Rhodnius</i> . Differentiation in epidermis of <i>Rhodnius</i> . Homœosis. Regeneration. Polymorphism. Conclusion.	
	REFERENCES	130
	INDEX	151

CHAPTER I

INTRODUCTION

FROM the earliest times the insect larva has been regarded as an embryo enjoying an independent life. For Aristotle taught that the embryonic life of insects continues until the formation of the perfect insect or imago: 'the larva while it is yet in growth', he writes, 'is a soft egg.' The insect egg, according to William Harvey (1651), contains so little yolk that the embryo is forced to leave it before development is complete: it requires a larval stage in which to store up food material, until it reverts once more to the egg form or pupa. Ramdohr (1811) still retains the same idea and calls the caterpillar a 'moving, growing and feeding egg'; and so, indeed, does Viallanes (1882), who writes of 'those sorts of eggs that are called nymphs or pupae'.

Lubbock (1883) likewise traces the origin of metamorphosis to the immaturity in the condition of the insect at the time of hatching. Such an insect as the cockroach, as it develops in the egg, goes through a protopod phase, a polypod phase and an oligopod phase. According to the conception outlined by Lubbock the different groups of insects hatch from the egg at one or other of these developmental stages.

This idea, which was characterized by Pérez (1902) as 'bizarre', was later elaborated by Berlese (1913) and has been given wide currency in the writings of Imms (1925, 1937). It is often referred to as the Berlese theory. According to this 'theory' the newly hatched young of such insects as Adephaga, Staphylinidae, Meloidae, Strepsiptera or *Mantispa*, which bear a general resemblance to the primitive *Campodea*, do not leave the egg until they have reached the oligopod phase. The caterpillars of Lepidoptera, Tenthredinidae or Mecoptera, with abdominal appendages, are regarded as having hatched in the polypod phase. The larvae of other Hymenoptera and Coleoptera, with no abdominal appendages, are pictured as being between the

Physiology of Insect Metamorphosis

polypod and oligopod phases, and the larvae of Diptera as being oligopod but highly modified. All these forms, which are supposed to leave the egg at an early stage of morphological development, are commonly termed 'larvae'; whereas such forms as Orthoptera, Hemiptera, Odonata, etc., which are supposed to leave the egg at a relatively advanced stage, are commonly termed 'nymphs'.

POLYMORPHISM. This approach to the problem of metamorphosis, in which the larva is seen as some kind of embryo, may have a certain value for descriptive purposes; but it is very different from the conception of insect growth that comes from a study of developmental physiology. Growth in the insect is not so very different from that in other animals. There are changes in form as growth proceeds; and since the form of the insect body is defined by an external cuticle, which remains unchanged throughout each moulting stage, it follows that any alteration in form must occur suddenly as the cuticle is shed, or must at least take place in a series of steps each more or less abrupt. This has the effect of making metamorphosis appear more catastrophic than it really is (Lubbock, 1883). But even allowing for this the changes in form are spectacular enough.

Now difference in form or polymorphism is almost universal among animals. There are differences between young and mature, between male and female, between the forms appearing in the cold or wet season and those appearing in the hot season or the dry. There are differences in caste between workers, soldiers and reproductives in ants and termites, or differences in successive or alternate generations. Metamorphosis is merely one type of polymorphism.

The polymorphic organism generally contains within it the potentialities for all its diverse forms. Sometimes the form to be developed is determined by the genetic constitution; but not uncommonly the genetic constitution is overridden by the environmental conditions experienced during growth. The 'penetrance' of genes may be influenced by temperature, or by nutrition. The genetically determined sex of Hymenoptera or Homoptera may be reversed in the course of development as the result of parasitism by Strepsiptera or dryinids. In ants and

Introduction

termites the potentialities to form all the female castes exist in every female egg; which castes appear is determined by the needs of their societies, acting by way of stimuli that are very incompletely understood.

Lubbock drew a distinction between animals with different terminal or mature forms (polymorphism) and animals which pass through a succession of different forms in the course of their development (polyeidism). It is doubtful whether there is anything fundamental about this distinction. The essential feature of polymorphism is multiple potentiality: one form alone is realized; the others remain latent or suppressed. The characteristic of metamorphosis is that this suppression is only temporary; so that one form replaces another during the normal life of the individual.

EMBRYOLOGY AND METAMORPHOSIS. It is obvious, as Swammerdam (1758) rightly maintained, that the latent form must exist, in some state which we do not understand, within the undeveloped organism. This applies equally to the embryonic germ in the egg; and the egg of insects has proved particularly well suited for studying the progressive determination and differentiation of this latent organism. When the egg is laid the single nucleus or germinal vesicle lies at the centre of the yolk and the egg plasma radiates from it, enclosing in its meshes the reserve substances of the yolk, to condense around the surface of the egg as the cortical plasma. One limited region of the cortical plasma, sometimes (as in *Tenebrio* (Ewest, 1937)) distinguishable visually from the remainder, and lying in the postero-ventral region of the egg, is already destined to become the embryo.

At this stage the non-cellular germ of the organism is highly plastic; it is still capable of 'regulation'. If one part is removed it is made good and a complete organism develops from the remainder. If it is divided, two complete organisms are differentiated. Then the nucleus undergoes cleavage; the daughter nuclei as they divide repeatedly move outwards and eventually reach the cortical plasma. Those nuclei which fall within the germ band zone are destined to contribute to the embryo; those which arrive elsewhere form only the extra-embryonic blastoderm.

Physiology of Insect Metamorphosis

At some stage in this process the germ band loses its capacity for complete regulation. Each part becomes committed to form some part of the embryo. The egg becomes an invisible mosaic of determined zones. This change to the mosaic state is gradual, and in many eggs some capacity for regulation persists until a late stage; but the main change can be recognized as occurring in different eggs at quite different stages of cleavage or blastoderm formation. In the dragonfly *Platycnemis*, regulation is still possible in the late blastoderm stage (Seidel, 1936); in *Sialis* determination occurs between the fourth and fifth cleavages, at the moment when the blastema is formed by peripheral migration of the cytoplasm (Du Bois, 1938); in *Bruchus* the posterior cytoplasmic regions of the egg are determined before the entrance of the cleavage nuclei, the more anterior parts shortly after the cleavage cells arrive (Brauer and Taylor, 1936); whereas in *Musca* determination of the cortical plasma is complete at the time of laying before cleavage has even begun (Reith, 1925).

In *Drosophila*, as in *Musca*, the egg at the time of laying is a mosaic egg. Local injuries effected by burning with a pencil of ultra-violet light immediately after laying cause local defects in the resulting larva. But the adult characters are unaffected; in respect to imaginal characters the egg is still capable of 'regulation'. If the egg is similarly treated 7 hours after laying or later, localized defects appear in the adult—often without any visible effect during larval development (fig. 1) (Geigy, 1931). Similar results can be obtained by puncturing the egg at different levels with a needle (Howland and Child, 1935; Howland and Sonnenblick, 1936). Likewise in *Tineola*, by burning different points in the egg with ultra-violet irradiation at different times it is possible to obtain purely larval or purely imaginal defects (Lüscher, 1944). It is clear that even at this early stage of embryonic development certain parts of the germ have already become determined to form specialized parts of the adult insect.

ORIGIN AND EVOLUTION OF METAMORPHOSIS. These observations serve to demonstrate the independent existence within the embryo of two latent organisms, larval and adult. As development proceeds the larval organism becomes differen-

Introduction

tiated, hatches and grows, while the adult organism remains in its latent, mosaic, invisible condition. This state of affairs persists until the larva is full grown; it then disappears and the adult organism becomes visibly differentiated. It is this change that is called metamorphosis. It is often regarded as a renewal of embryonic development, in that the process of transition




Fig. 1. *Drosophila* adult showing complete suppression of left hind-leg with deformation of first and second leg on the same side, following unilateral irradiation of the egg with ultra-violet light. Larval stages showed no visible defects. (Geigy, 1931.)

from a state of invisible determination to a state of visible differentiation is the same in both.

The important point at this stage of the argument is the independence of the alternative forms in which polymorphic organisms can exist. It is by virtue of this independence that these forms are capable of separate evolution. The same applies to the successive forms that exist in insects showing a metamorphosis. The larva is subject to the influence of its environment, and by the action of natural selection (or whatever other agencies control organic change) it may undergo an evolution of

Physiology of Insect Metamorphosis

its characters which leaves the adult insect untouched. Larvae of the silkworm or of the gipsy moth *Lymantria* (Klatt, 1919) exist in many very different genetic racial forms which cannot be differentiated in the adult state. Indeed, certain species of Lepidoptera (such as *Acronycta psi* and *A. tridens*) distinguishable with great difficulty in the adult have very different larvae. On the other hand, highly polymorphic butterflies such as *Papilio polytes* show no corresponding differences in their caterpillars. A similar independence is seen in the pupa: the female of *P. dardanus* has no tails on the hind-wing, but pockets are provided for them in the pupal wing cases (Lamborn, 1914), and in the wingless mutant of the silkworm the wing cases of the pupa are formed as usual (Goldschmidt, 1927).

It follows, therefore, that as soon as there are differences in the mode of life of an adult insect and its larva the characters of the two forms will tend to diverge, and such divergence may reach the extreme that separates the maggot from the fly. Lubbock recognized that the form of the larva is affected by phylogenetic influences on the one hand and by adaptive influences on the other. Writing in 1883, at a time when the theory of recapitulation exerted a greater influence than it does to-day, he was inclined to give greater weight to the importance of phylogeny. At the present day we are inclined to reverse this emphasis: to regard the larva as a highly specialized form, closely adapted to the conditions of its life, whose resemblance to an embryo is superficial and adventitious (cf. Müller, 1869; Schindler, 1902; Snodgrass, 1953).

ORIGIN OF THE PUPA. The origin of metamorphosis is to be sought, therefore, in the divergent evolution of a polymorphic organism. The same considerations apply to the pupa in holometabolous insects. Perhaps the most obvious function of the pupa is that it bridges the morphological gap between the larva and the fully developed adult. But here again the best guide is to regard the pupa as the product of independent evolution in one form in a polymorphic organism to suit the ecological or anatomical needs of the insect in question.

Attempts have been made to find a single overriding final cause which has necessitated the development of the pupa. One of the most attractive is the view put forward by Poyarkoff (1914)

Introduction

and developed by Hinton (1948) that the skeletal muscles require a 'mould' in which the cuticular parts have approximately the same spatial relations as in the adult, and that a pupal mould becomes necessary when the anatomical differences are so great that the muscles required by the adult cannot be formed in the larva. But as Weber (1952) points out, the aleurodids are able to develop directly at a single moult from an extremely flattened scale-like larva to a fully winged and long-legged adult.

IMAGINAL DISCS. The classic discussions of insect metamorphosis have always centred around the more extreme examples, such as the Lepidoptera, Hymenoptera or Diptera. The anatomical study of these forms led to the discovery of the 'imaginal discs' (Lyonet, 1762), the significance of which was first appreciated by Weismann (1864), in his work on *Calliphora*, as nests of embryonic cells set aside for the formation of the adult. They represent the latent imaginal organism in visible form; and physiological discussion turned upon what factors inhibit the development of the imaginal discs in the young stages, or cause the larval tissues to disintegrate and activate the imaginal buds at metamorphosis.

It has been suggested, for example, (i) that the imaginal discs are held quiescent by the excretory products of the active larval tissues, and when these begin to degenerate the adult tissues can begin to grow (Anglas, 1901); (ii) that the larval tissues give out substances which keep the phagocytic blood cells at bay; when they age they become a prey to phagocytes and the imaginal discs develop in their place (Kowalevsky, 1887; v. Rees, 1888); (iii) that asphyxia in the larval tissues causes their degeneration (Bataillon, 1893); (iv) that the oxidase associated with melanin formation, which is at a maximum at the time of pupation, is in fact the *cause* of metamorphosis (Dewitz, 1916; cf. Agrell, 1951); (v) that the increasing size of the larval tissue cells set a physical limit to their ability to gain nourishment by diffusion (Murray and Tiegs, 1935; Tiegs, 1922); (vi) that secretions from the growing sex glands activate the imaginal discs (Pérez, 1902, 1910); (vii) that in the larva, the development of the imaginal discs is checked by secretions from the brain (Kopeč, 1924; but cf. Kopeč, 1927).

Physiology of Insect Metamorphosis

It will be observed that all these hypotheses relate to the inhibition or activation of imaginal discs. But many parts of the adult body, even in such insects as Lepidoptera or Hymenoptera which suffer a spectacular metamorphosis, are not formed from imaginal discs but are laid down by the same cells as have formed the larva. The general integument of the abdomen in adult Lepidoptera is very different from that of the caterpillar, but the same cells are concerned in its formation. In some insects the imaginal discs appear very early: in Diptera some are already evident in the late embryo (Enzmann and Haskins, 1938); in *Vanessa* the wing discs are present in the first larval stage, although those for the limbs do not appear until the end of the fourth stage (Bodenstein, 1935). But in every insect, if they are traced back far enough in development, the imaginal discs will be found to merge into the general epidermis of the larva or the embryo. There must, therefore, be a time, even in Lepidoptera or Diptera, when the same cells which are forming the visible larval structures carry latent within them the capacity to form imaginal structures.

We have already seen that this is so in the developing egg of *Drosophila* and *Tineola*. The purpose in elaborating the argument here is to emphasize that imaginal discs are not a necessary feature of metamorphosis, and that the essential problems of metamorphosis—the latency and subsequent realization of an imaginal organism—can be studied equally well in an insect in which imaginal discs do not occur.

METAMORPHOSIS IN RHODNIUS. It was on these grounds that the Hemipteron *Rhodnius prolixus* was chosen for the study of metamorphosis (Plate Ia-c). *Rhodnius* has five larval stages throughout which the structural characters of the cuticle and the pigment pattern show little change. There is progressive growth and differentiation in the wing lobes, and in the later larval stages the rudiments of the genitalia begin to differentiate. But these changes are very slight in comparison with the metamorphosis which occurs when the 5th-stage larva moults to become adult.

Elaborate genital appendages are then developed: the thorax becomes elaborately shaped, with fully formed flight muscles; the wing lobes are transformed into large functional wings;