

SOBOLEV SPACES

索伯列夫空间
第2版

SECOND
EDITION

ROBERT A. ADAMS
JOHN J.F. FOURNIER



Elsevier (Singapore) Pte Ltd.

世界图书出版公司
www.wpcbj.com.cn

SOBOLEV SPACES

Second Edition

Robert A. Adams and John J. F. Fournier

Department of Mathematics
The University of British Columbia
Vancouver, Canada

 **ACADEMIC PRESS**
An imprint of Elsevier Science

Amsterdam Boston Heidelberg London New York Oxford
Paris San Diego San Francisco Singapore Sydney Tokyo

图书在版编目 (CIP) 数据

索伯列夫空间: 第2版=Sobolev Spaces: 英文/(加)亚当斯著.—北京: 世界图书出版公司北京公司, 2009.8

ISBN 978-7-5100-0537-4

I. 索… II. 亚… III. 索伯列夫空间—英文 IV. 0177.3

中国版本图书馆 CIP 数据核字 (2009) 第 127400 号

书 名: Sobolev Spaces 2nd ed.

作 者: Rotert A. Adams, John J. F. Fournier

中译名: 索伯列夫空间 第2版

责任编辑: 高蓉

出 版 者: 世界图书出版公司北京公司

印 刷 者: 北京集惠印刷有限责任公司

发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010-64021602, 010-64015659

电子信箱: kjb@wpcbj.com.cn

开 本: 24 开

印 张: 13.5

版 次: 2009 年 08 月

版权登记: 图字:01-2009-4134

书 号: 978-7-5100-0537-4/0•753

定 价: 69.00 元

Sobolev Spaces 2nd Edition

Robert A. Adams, John J. F. Fournier

ISBN: 978-0-12-044143-3

Copyright © 2003, Elsevier Ltd. All rights reserved.

Authorized English language reprint edition published by the Proprietor.

Reprint ISBN: 978-981-272383-3

Copyright © 2009 by Elsevier (Singapore) Pte Ltd. All rights reserved.

Elsevier (Singapore) Pte Ltd.

3 Killiney Road

#08-01 Winsland House I

Singapore 239519

Tel: (65) 6349-0200

Fax: (65) 6733-1817

First Published 2009

2009 年初版

Printed in China by Elsevier (Singapore) Pte Ltd. under special arrangement with Beijing World Publishing Corporation. This edition is authorized for sale in China only, excluding Hong Kong SAR and Taiwan. Unauthorized export of this edition is a violation of the Copyright Act. Violation of this Law is subject to Civil and Criminal Penalties.

本书英文影印版由 Elsevier (Singapore) Pte Ltd. 授权世界图书出版公司北京公司在中国大陆境内独家发行。本版仅限在中国境内（不包括香港特别行政区及台湾）出版及标价销售。未经许可出口，视为违反著作权法，将受法律制裁。

SOBOLEV SPACES

Second Edition

To Anne and Frances

who had to put up with it all

PREFACE

This monograph presents an introductory study of the properties of certain Banach spaces of weakly differentiable functions of several real variables that arise in connection with numerous problems in the theory of partial differential equations, approximation theory, and many other areas of pure and applied mathematics. These spaces have become associated with the name of the late Russian mathematician S. L. Sobolev, although their origins predate his major contributions to their development in the late 1930s.

Even by 1975 when the first edition of this monograph was published, there was a great deal of material on these spaces and their close relatives, though most of it was available only in research papers published in a wide variety of journals. The monograph was written to fill a perceived need for a single source where graduate students and researchers in a wide variety of disciplines could learn the essential features of Sobolev spaces that they needed for their particular applications. No attempt was made even at that time for complete coverage. To quote from the Preface of the first edition:

The existing mathematical literature on Sobolev spaces and their generalizations is vast, and it would be neither easy nor particularly desirable to include everything that was known about such spaces between the covers of one book. An attempt has been made in this monograph to present all the core material in sufficient generality to cover most applications, to give the reader an overview of the subject that is difficult to obtain by reading research papers, and finally . . . to provide a ready reference for someone requiring a result about Sobolev spaces for use in some application.

This remains as the purpose and focus of this second edition. During the intervening twenty-seven years the research literature has grown exponentially, and there

are now several other books in English that deal in whole or in part with Sobolev spaces. (For example, see [Ad2], [Bu1], [Mz1], [Tr1], [Tr3], and [Tr4].) However, there is still a need for students in other disciplines than mathematics, and in other areas of mathematics than just analysis to have available a book that describes these spaces and their core properties based only a background in mathematical analysis at the senior undergraduate level. We have tried to make this such a book.

The organization of this book is similar but not identical to that of the first edition: Chapter 1 remains a potpourri of standard topics from real and functional analysis, included, mainly without proofs, because they provide a necessary background for what follows.

Chapter 2 on the Lebesgue Spaces $L^p(\Omega)$ is much expanded and reworked from the previous edition. It provides, in addition to standard results about these spaces, a brief treatment of mixed-norm L^p spaces, weak- L^p spaces, and the Marcinkiewicz interpolation theorem, all of which will be used in a new treatment of the Sobolev Imbedding Theorem in Chapter 4. For the most part, complete proofs are given, as they are for much of the rest of the book.

Chapter 3 provides the basic definitions and properties of the Sobolev spaces $W^{m,p}(\Omega)$ and $W_0^{m,p}(\Omega)$. There are minor changes from the first edition.

Chapter 4 is now completely concerned with the imbedding properties of Sobolev Spaces. The first half gives a more streamlined presentation and proof of the various imbeddings of Sobolev spaces into L^p spaces, including traces on subspaces of lower dimension, and spaces of continuous and uniformly continuous functions. Because the approach to the Sobolev Imbedding Theorem has changed, the roles of Chapters 4 and 5 have switched from the first edition. The latter part of Chapter 4 deals with situations where the regularity conditions on the domain Ω that are necessary for the full Sobolev Imbedding Theorem do not apply, but some weaker imbedding results are still possible.

Chapter 5 now deals with interpolation, extension, and approximation results for Sobolev spaces. Part of it is expanded from material in Chapter 4 of the first edition with newer results and methods of proof.

Chapter 6 deals with establishing compactness of Sobolev imbeddings. It is only slightly changed from the first edition.

Chapter 7 is concerned with defining and developing properties of scales of spaces with fractional orders of smoothness, rather than the integer orders of the Sobolev spaces themselves. It is completely rewritten and bears little resemblance to the corresponding chapter in the first edition. Much emphasis is placed on real interpolation methods. The J-method and K-method are fully presented and used to develop the theory of Lorentz spaces and Besov spaces and their imbeddings, but both families of spaces are also provided with intrinsic characterizations. A key theorem identifies lower dimensional traces of functions in Sobolev spaces

as constituting certain Besov spaces. Complex interpolation is used to introduce Sobolev spaces of fractional order (also called spaces of Bessel potentials) and Fourier transform methods are used to characterize and generalize these spaces to yield the Triebel Lizorkin spaces and illuminate their relationship with the Besov spaces.

Chapter 8 is very similar to its first edition counterpart. It deals with Orlicz and Orlicz-Sobolev spaces which generalize L^p and $W^{m,p}$ spaces by allowing the role of the function t^p to be assumed by a more general convex function $A(t)$. An important result identifies a certain Orlicz space as a target for an imbedding of $W^{m,p}(\Omega)$ in a limiting case where there is an imbedding into $L^p(\Omega)$ for $1 \leq p < \infty$ but not into $L^\infty(\Omega)$.

This monograph was typeset by the authors using T_EX on a PC running Linux-Mandrake 8.2. The figures were generated using the mathematical graphics software package *MG* developed by R. B. Israel and R. A. Adams.

RAA & JJFF

Vancouver, August 2002

List of Spaces and Norms

Space	Norm	Paragraph
$B^{s;p,q}(\Omega)$	$\ \cdot; B^{s;p,q}(\Omega)\ $	7.32
$B^{s;p,q}(\mathbb{R}^n)$	$\ \cdot; B^{s;p,q}(\mathbb{R}^n)\ $	7.67
$\dot{B}^{s;p,q}(\mathbb{R}^n)$		7.68
$C^m(\Omega), C^\infty(\Omega)$		1.26
$C_0(\Omega), C_0^\infty(\Omega)$		1.26
$C^m(\overline{\Omega})$	$\ \cdot; C^m(\overline{\Omega})\ $	1.28
$C^{m,\lambda}(\overline{\Omega})$	$\ \cdot; C^{m,\lambda}(\overline{\Omega})\ $	1.29
$C_B^m(\Omega)$	$\ \cdot; C_B^m(\Omega)\ $	1.27, 4.2
$C^j(\overline{\Omega})$	$\ \cdot; C^j(\overline{\Omega})\ $	4.2
$C^{j,\lambda}(\overline{\Omega})$	$\ \cdot; C^{j,\lambda}(\overline{\Omega})\ $	4.2
$C^{j,\lambda,q}(\overline{\Omega})$	$\ \cdot; C^{j,\lambda,q}(\overline{\Omega})\ $	7.35
$\mathcal{D}(\Omega)$		1.56
$\mathcal{D}'(\Omega)$		1.57
$E_A(\Omega)$	$\ \cdot\ _A = \ \cdot\ _{A,\Omega}$	8.14

$F^{s;p,q}(\Omega)$	$\ \cdot; F^{s;p,q}(\Omega)\ $	7.69
$F^{s;p,q}(\mathbb{R}^n)$	$\ \cdot; F^{s;p,q}(\mathbb{R}^n)\ $	7.65
$\dot{F}^{s;p,q}(\mathbb{R}^n)$		7.66
$H^{m,p}(\Omega)$	$\ \cdot\ _{m,p} = \ \cdot\ _{m,p,\Omega}$	3.2
$L_A(\Omega)$	$\ \cdot\ _A = \ \cdot\ _{A,\Omega}$	8.9
$L^p(\Omega)$	$\ \cdot\ _p = \ \cdot\ _{p,\Omega}$	2.1, 2.3
$L^p(\mathbb{R}^n)$	$\ \cdot\ _p$	2.48
$L^\infty(\Omega)$	$\ \cdot\ _\infty = \ \cdot\ _{\infty,\Omega}$	2.10
$L^q(a,b;d\mu,X)$	$\ \cdot; L^q(a,b;d\mu,X)\ $	7.4
L_*^q	$\ \cdot; L_*^q\ $	7.5
$L_{\text{loc}}^1(\Omega)$		1.58
$L^{p,q}(\Omega)$	$\ \cdot; L^{p,q}(\Omega)\ $	7.25
ℓ^p	$\ \cdot; \ell^p\ $	2.27
$\mathcal{S} = \mathcal{S}(\mathbb{R}^n)$		7.59
weak- $L^p(\Omega)$	$[\cdot]_p = [\cdot]_{p,\Omega}$	2.55
$W^{m,p}(\Omega)$	$\ \cdot\ _{m,p} = \ \cdot\ _{m,p,\Omega}$	3.2
$W_0^{m,p}(\Omega)$	$\ \cdot\ _{m,p} = \ \cdot\ _{m,p,\Omega}$	3.2
$W^{-m,p'}(\Omega)$	$\ \cdot\ _{-m,p'}$	3.12, 3.13
$W^m E_A(\Omega)$	$\ \cdot\ _{m,A} = \ \cdot\ _{m,A,\Omega}$	8.30
$W^m L_A(\Omega)$	$\ \cdot\ _{m,A} = \ \cdot\ _{m,A,\Omega}$	8.30
$W^{s,p}(\Omega)$	$\ \cdot; W^{s,p}(\Omega)\ $	7.57
$W^{s,p}(\mathbb{R}^n)$	$\ \cdot; W^{s,p}(\mathbb{R}^n)\ $	7.64
X	$\ \cdot; X\ $	1.7
$X_0 \cap X_1$	$\ \cdot\ _{X_0 \cap X_1}$	7.7
$X_0 + X_1$	$\ \cdot\ _{X_0 + X_1}$	7.7
$(X_0, X_1)_{\theta,q;J}$	$\ \cdot\ _{\theta,q;J}$	7.13
$(X_0, X_1)_{\theta,q;K}$	$\ \cdot\ _{\theta,q;K}$	7.10
$[X_0, X_1]_\theta$	$\ u\ _{[X_0, X_1]_\theta}$	7.51
$X_0^{1-\theta} X_1^\theta$	$\ \cdot; X_0^{1-\theta} X_1^\theta\ $	7.54

CONTENTS

Preface	ix
List of Spaces and Norms	xii
1. PRELIMINARIES	1
Notation	1
Topological Vector Spaces	3
Normed Spaces	4
Spaces of Continuous Functions	10
The Lebesgue Measure in \mathbb{R}^n	13
The Lebesgue Integral	16
Distributions and Weak Derivatives	19
2. THE LEBESGUE SPACES $L^p(\Omega)$	23
Definition and Basic Properties	23
Completeness of $L^p(\Omega)$	29
Approximation by Continuous Functions	31
Convolutions and Young's Theorem	32
Mollifiers and Approximation by Smooth Functions	36
Precompact Sets in $L^p(\Omega)$	38
Uniform Convexity	41
The Normed Dual of $L^p(\Omega)$	45
Mixed-Norm L^p Spaces	49
The Marcinkiewicz Interpolation Theorem	52

3. THE SOBOLEV SPACES $W^{m,p}(\Omega)$	59
Definitions and Basic Properties	59
Duality and the Spaces $W^{-m,p'}(\Omega)$	62
Approximation by Smooth Functions on Ω	65
Approximation by Smooth Functions on \mathbb{R}^n	67
Approximation by Functions in $C_0^\infty(\Omega)$	70
Coordinate Transformations	77
4. THE SOBOLEV IMBEDDING THEOREM	79
Geometric Properties of Domains	81
Imbeddings by Potential Arguments	87
Imbeddings by Averaging	93
Imbeddings into Lipschitz Spaces	99
Sobolev's Inequality	101
Variations of Sobolev's Inequality	104
$W^{m,p}(\Omega)$ as a Banach Algebra	106
Optimality of the Imbedding Theorem	108
Nonimbedding Theorems for Irregular Domains	111
Imbedding Theorems for Domains with Cusps	115
Imbedding Inequalities Involving Weighted Norms	119
Proofs of Theorems 4.51–4.53	131
5. INTERPOLATION, EXTENSION, AND APPROXIMATION THEOREMS	135
Interpolation on Order of Smoothness	135
Interpolation on Degree of Sumability	139
Interpolation Involving Compact Subdomains	143
Extension Theorems	146
An Approximation Theorem	159
Boundary Traces	163
6. COMPACT IMBEDDINGS OF SOBOLEV SPACES	167
The Rellich-Kondrachov Theorem	167
Two Counterexamples	173
Unbounded Domains — Compact Imbeddings of $W_0^{m,p}(\Omega)$	175
An Equivalent Norm for $W_0^{m,p}(\Omega)$	183
Unbounded Domains — Decay at Infinity	186
Unbounded Domains — Compact Imbeddings of $W^{m,p}(\Omega)$	195
Hilbert-Schmidt Imbeddings	200

7. FRACTIONAL ORDER SPACES	205
Introduction	205
The Bochner Integral	206
Intermediate Spaces and Interpolation — The Real Method	208
The Lorentz Spaces	221
Besov Spaces	228
Generalized Spaces of Hölder Continuous Functions	232
Characterization of Traces	234
Direct Characterizations of Besov Spaces	241
Other Scales of Intermediate Spaces	247
Wavelet Characterizations	256
 8. ORLICZ SPACES AND ORLICZ-SOBOLEV SPACES	 261
Introduction	261
N-Functions	262
Orlicz Spaces	266
Duality in Orlicz Spaces	272
Separability and Compactness Theorems	274
A Limiting Case of the Sobolev Imbedding Theorem	277
Orlicz-Sobolev Spaces	281
Imbedding Theorems for Orlicz-Sobolev Spaces	282
 References	 295
 Index	 301

1

PRELIMINARIES

1.1 (Introduction) Sobolev spaces are vector spaces whose elements are functions defined on domains in n -dimensional Euclidean space \mathbb{R}^n and whose partial derivatives satisfy certain integrability conditions. In order to develop and elucidate the properties of these spaces and mappings between them we require some of the machinery of general topology and real and functional analysis. We assume that readers are familiar with the concept of a vector space over the real or complex scalar field, and with the related notions of dimension, subspace, linear transformation, and convex set. We also expect the reader will have some familiarity with the concept of topology on a set, at least to the extent of understanding the concepts of an open set and continuity of a function.

In this chapter we outline, mainly without any proofs, those aspects of the theories of topological vector spaces, continuity, the Lebesgue measure and integral, and Schwartz distributions that will be needed in the rest of the book. For a reader familiar with the basics of these subjects, a superficial reading to settle notations and review the main results will likely suffice.

Notation

1.2 Throughout this monograph the term *domain* and the symbol Ω will be reserved for a nonempty open set in n -dimensional real Euclidean space \mathbb{R}^n . We shall be concerned with the differentiability and integrability of functions defined on Ω ; these functions are allowed to be complex-valued unless the contrary is

explicitly stated. The complex field is denoted by \mathbb{C} . For $c \in \mathbb{C}$ and two functions u and v , the scalar multiple cu , the sum $u + v$, and the product uv are always defined pointwise:

$$\begin{aligned}(cu)(x) &= cu(x), \\ (u + v)(x) &= u(x) + v(x), \\ (uv)(x) &= u(x)v(x)\end{aligned}$$

at all points x where the right sides make sense.

A typical point in \mathbb{R}^n is denoted by $x = (x_1, \dots, x_n)$; its norm is given by $|x| = (\sum_{j=1}^n x_j^2)^{1/2}$. The inner product of two points x and y in \mathbb{R}^n is $x \cdot y = \sum_{j=1}^n x_j y_j$.

If $\alpha = (\alpha_1, \dots, \alpha_n)$ is an n -tuple of nonnegative integers α_j , we call α a *multi-index* and denote by x^α the monomial $x_1^{\alpha_1} \cdots x_n^{\alpha_n}$, which has degree $|\alpha| = \sum_{j=1}^n \alpha_j$. Similarly, if $D_j = \partial/\partial x_j$, then

$$D^\alpha = D_1^{\alpha_1} \cdots D_n^{\alpha_n}$$

denotes a differential operator of order $|\alpha|$. Note that $D^{(0, \dots, 0)} u = u$.

If α and β are two multi-indices, we say that $\beta \leq \alpha$ provided $\beta_j \leq \alpha_j$ for $1 \leq j \leq n$. In this case $\alpha - \beta$ is also a multi-index, and $|\alpha - \beta| + |\beta| = |\alpha|$. We also denote

$$\alpha! = \alpha_1! \cdots \alpha_n!$$

and if $\beta \leq \alpha$,

$$\binom{\alpha}{\beta} = \frac{\alpha!}{\beta!(\alpha - \beta)!} = \binom{\alpha_1}{\beta_1} \cdots \binom{\alpha_n}{\beta_n}.$$

The reader may wish to verify the Leibniz formula

$$D^\alpha(uv)(x) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} D^\beta u(x) D^{\alpha - \beta} v(x)$$

valid for functions u and v that are $|\alpha|$ times continuously differentiable near x .

1.3 If $G \subset \mathbb{R}^n$ is nonempty, we denote by \overline{G} the closure of G in \mathbb{R}^n . We shall write $G \Subset \Omega$ if $\overline{G} \subset \Omega$ and \overline{G} is a compact (that is, closed and bounded) subset of \mathbb{R}^n . If u is a function defined on G , we define the *support* of u to be the set

$$\text{supp}(u) = \overline{\{x \in G : u(x) \neq 0\}}.$$

We say that u has *compact support* in Ω if $\text{supp}(u) \Subset \Omega$. We denote by “bdry G ” the boundary of G in \mathbb{R}^n , that is, the set $\overline{G} \cap \overline{G}^c$, where G^c is the complement of G in \mathbb{R}^n ; $G^c = \mathbb{R}^n - G = \{x \in \mathbb{R}^n : x \notin G\}$.

If $x \in \mathbb{R}^n$ and $G \subset \mathbb{R}^n$, we denote by “ $\text{dist}(x, G)$ ” the distance from x to G , that is, the number $\inf_{y \in G} |x - y|$. Similarly, if $F, G \subset \mathbb{R}^n$ are both nonempty,

$$\text{dist}(F, G) = \inf_{y \in F} \text{dist}(y, G) = \inf_{\substack{x \in G \\ y \in F}} |y - x|.$$

Topological Vector Spaces

1.4 (Topological Spaces) If X is any set, a *topology* on X is a collection \mathcal{O} of subsets of X which contains

- (i) the whole set X and the empty set \emptyset ,
- (ii) the union of any collection of its elements, and
- (iii) the intersection of any finite collection of its elements.

The pair (X, \mathcal{O}) is called a *topological space* and the elements of \mathcal{O} are the *open sets* of that space. An open set containing a point x in X is called a *neighbourhood* of x . The complement $\bar{X} - U = \{x \in X : x \notin U\}$ of any open set U is called a *closed set*. The closure \bar{S} of any subset $S \subset X$ is the smallest closed subset of X that contains S .

Let \mathcal{O}_1 and \mathcal{O}_2 be two topologies on the same set X . If $\mathcal{O}_1 \subset \mathcal{O}_2$, we say that \mathcal{O}_2 is *stronger* than \mathcal{O}_1 , or that \mathcal{O}_1 is *weaker* than \mathcal{O}_2 .

A topological space (X, \mathcal{O}) is called a *Hausdorff space* if every pair of distinct points x and y in X have disjoint neighbourhoods.

The *topological product* of two topological spaces (X, \mathcal{O}_X) and (Y, \mathcal{O}_Y) is the topological space $(X \times Y, \mathcal{O})$, where $X \times Y = \{(x, y) : x \in X, y \in Y\}$ is the Cartesian product of the sets X and Y , and \mathcal{O} consists of arbitrary unions of sets of the form $\{O_X \times O_Y : O_X \in \mathcal{O}_X, O_Y \in \mathcal{O}_Y\}$.

Let (X, \mathcal{O}_X) and (Y, \mathcal{O}_Y) be two topological spaces. A function f from X into Y is said to be *continuous* if the preimage $f^{-1}(O) = \{x \in X : f(x) \in O\}$ belongs to \mathcal{O}_X for every $O \in \mathcal{O}_Y$. Evidently the stronger the topology on X or the weaker the topology on Y , the more such continuous functions f there will be.

1.5 (Topological Vector Spaces) We assume throughout this monograph that all vector spaces referred to are taken over the complex field unless the contrary is explicitly stated.

A *topological vector space*, hereafter abbreviated TVS, is a Hausdorff topological space that is also a vector space for which the vector space operations of addition and scalar multiplication are continuous. That is, if X is a TVS, then the mappings

$$(x, y) \rightarrow x + y \quad \text{and} \quad (c, x) \rightarrow cx$$