

国家出版基金项目  
NATIONAL PUBLICATION FOUNDATION



*Series in Information and Computational Science*

**55**

# Effective Condition Number for Numerical Partial Differential Equations

Zi-Cai Li    Hung-Tsai Huang  
Yimin Wei    Alexander H.-D. Cheng

(偏微分方程数值解的有效条件数)



SCIENCE PRESS  
Beijing



国家出版基金项目  
NATIONAL PUBLICATION FOUNDATION


Series in Information and Computational Science 55

Zi-Cai Li   Hung-Tsai Huang   Yimin Wei

Alexander H.-D. Cheng

# Effective Condition Number for Numerical Partial Differential Equations

(偏微分方程数值解的有效条件数)



 Science Press  
Beijing

Responsible Editor: Li Xin

Copyright© 2013 by Science Press  
Published by Science Press  
16 Donghuangchenggen North Street  
Beijing 100717, China

Printed in Beijing

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the copyright owner.

ISBN 978-7-03-036753-2 (Beijing)

*This book is dedicated to our families.*

## Editorial Board

**Editor-in-Chief:** Shi Zhongci

**Vice Editor-in-Chief:** Wang Xinghua Yu Dehao

<b>Members:</b>	Bai Fengshan	Bai Zhongzhi	Chen Falai
	Chen Zhiming	Chen Zhongying	Cheng Jin
	E Weinan	Guo Benyu	He Bingsheng
	Hou Yizhao	Shu C.-W.	Song Yongzhong
	Tang Tao	Wu Wei	Xu Jinchao
	Xu Zongben	Yang Danping	Zhang Pingwen

## **Preface to the Series in Information and Computational Science**

Since the 1970s, Science Press has published more than thirty volumes in its series Monographs in Computational Methods. This series was established and led by the late academician, Feng Kang, the founding director of the Computing Center of the Chinese Academy of Sciences. The monograph series has provided timely information of the frontier directions and latest research results in computational mathematics. It has had great impact on young scientists and the entire research community, and has played a very important role in the development of computational mathematics in China.

To cope with these new scientific developments, the Ministry of Education of the People's Republic of China in 1998 combined several subjects, such as computational mathematics, numerical algorithms, information science, and operations research and optimal control, into a new discipline called Information and Computational Science. As a result, Science Press also reorganized the editorial board of the monograph series and changed its name to Series in Information and Computational Science. The first editorial board meeting was held in Beijing in September 2004, and it discussed the new objectives, and the directions and contents of the new monograph series.

The aim of the new series is to present the state of the art in Information and Computational Science to senior undergraduate and graduate students, as well as to scientists working in these fields. Hence, the series will provide concrete and systematic expositions of the advances in information and computational science, encompassing also related interdisciplinary developments.

I would like to thank the previous editorial board members and assistants, and all the mathematicians who have contributed significantly to the monograph series on Computational Methods. As a result of their contributions the monograph series achieved an outstanding reputation in the community. I sincerely wish that we will extend this support to the new Series in Information and Computational Science, so that the new series can equally enhance the scientific development in information and computational science in this century.

Shi Zhongci  
2005.7



# Preface

For numerical methods, the stability is a crucial issue in the sense that the unstable numerical methods are useless in practical applications. The Lax's principle<sup>[74]</sup> for initial problems states that under the consistent condition, the convergence and the stability are equivalent to each other. When the truncation errors are derived, which are not very difficult, the errors of the numerical solutions can be obtained. However, the final numerical solutions also include rounding errors, which are related to stability. Since for the given algorithms of partial differential equations (PDE), the stability proof is often difficult and challenging, error analysis provides an easier pathway to answer the stability question.

Let us consider the finite element method (FEM) for elliptic boundary problems. The uniformly  $V_h$  elliptic inequality is important for a priori error estimates<sup>[40]</sup>, and it also implies stability, because the solutions of elliptic problems are not very sensitive to the perturbation of the data involved. The linear algebraic equations obtained from the FEM can be solved by the direct methods such as Gaussian elimination, or iterative methods such as the conjugate gradient methods, or the multigrid methods. Since all computations are completed in computer, the rounding errors are inevitable. Since the double precision has only 16 significant decimal digits, the final numerical solutions must have the extra errors from rounding errors. Even when certain software, such as the computer algebra software *Mathematica*, is used with more working digits, it is also finite. The more working digits are used, the more CPU time and the more computer storage are needed. Hence, the perturbation errors, such as rounding errors, are important to numerical methods for PDE.

Consider the overdetermined linear algebraic equations resulting from numerical PDE,

$$F\mathbf{x} = \mathbf{b}, \quad (0.0.1)$$

where  $F \in \mathbf{R}^{m \times n}$ ,  $m \geq n$ , and  $\mathbf{x} \in \mathbf{R}^n$  and  $\mathbf{b} \in \mathbf{R}^m$  are the unknown and the known vectors, respectively. The traditional condition number in the 2-norm is defined by

$$\text{Cond} = \frac{\sigma_{\max}}{\sigma_{\min}}, \quad (0.0.2)$$

where  $\sigma_{\max}$  and  $\sigma_{\min}$  are the maximal and the minimal singular values, respectively.

The new effective condition number in this book is defined by

$$\text{Cond\_eff} = \frac{\|\mathbf{b}\|}{\sigma_{\min}\|\mathbf{x}\|}, \quad (0.0.3)$$

where  $\|\mathbf{x}\|$  is the 2-norm. When there exist the perturbation of  $\mathbf{b}$  and  $\mathbf{F}$ , the practical computation for (0.0.1) is carried out by

$$\mathbf{F}(\mathbf{x} + \Delta\mathbf{x}) = \mathbf{b} + \Delta\mathbf{b}, \quad (0.0.4)$$

$$(\mathbf{F} + \Delta\mathbf{F})(\mathbf{x} + \Delta\mathbf{x}) = \mathbf{b} + \Delta\mathbf{b}, \quad (0.0.5)$$

where  $\Delta\mathbf{F} \in \mathbf{R}^{m \times n}$  ( $m \geq n$ ),  $\Delta\mathbf{x} \in \mathbf{R}^n$  and  $\Delta\mathbf{b} \in \mathbf{R}^m$ . Suppose that  $\Delta\mathbf{F}$  is small so that  $\text{rank}(\mathbf{F}) = \text{rank}(\mathbf{F} + \Delta\mathbf{F}) = n$ . For (0.0.4) (i.e.,  $\Delta\mathbf{F} = \mathbf{0}$ ), there exist the bounds of relative errors,

$$\frac{\|\Delta\mathbf{x}\|}{\|\mathbf{x}\|} \leq \text{Cond} \times \frac{\|\Delta\mathbf{b}\|}{\|\mathbf{b}\|}, \quad \frac{\|\Delta\mathbf{x}\|}{\|\mathbf{x}\|} \leq \text{Cond\_eff} \times \frac{\|\Delta\mathbf{b}\|}{\|\mathbf{b}\|}. \quad (0.0.6)$$

Equations (0.0.6) indicate the errors from the perturbation, e.g., from the rounding errors. More specifically, the relative errors of the solution  $\mathbf{x}$  may be enlarged from the rounding errors by a factor of Cond, and Cond has often been used to provide a stability analysis of numerical methods (see Wilkinson [226]). In fact, since the upper bound Cond in (0.0.6) is the worst case, it rarely happens in most PDE problems. The error bound of  $\mathbf{x}$  can be improved by Cond\_eff and shown in (0.0.6). Cond\_eff in (0.0.3) is smaller, or even much smaller than Cond in (0.0.2). Such a conclusion has been proved by the analysis and computation in this entire book. Since the algorithms of (0.0.3) are so simple, easy and straightforward in computation, Cond\_eff is strongly recommended, to replace Cond. This is one objective of this book.

The idea of effective condition number was first studied in Rice [186], and Chan and Foulser [27], and the formula (0.0.3) of Cond\_eff was first used in Christiansen and Saranen [38]. Only a few papers [27, 37, 38, 51] follow this trend for stability analysis. Recently, we have carried out a systematic study on effective condition number of various numerical methods for PDE and the boundary integral equation (BIE). Interestingly, the Cond\_eff is significantly smaller than Cond for numerical methods of PDE, but only fairly smaller than Cond for numerical methods of BIE<sup>[98]</sup>. Comparing (0.0.3) with (0.0.2), the minimal singular value  $\sigma_{\min}$  is crucial for both Cond\_eff and Cond, but the maximal singular value  $\sigma_{\max}$  is necessary only to Cond. Hence when  $\sigma_{\max}$  is large, Cond is large, but Cond\_eff may remain small. This happens for the finite element method (FEM), the finite difference method (FDM), the Trefftz method (TM) and the spectral method (SM) for elliptic boundary value problems. In particular, when the maximal boundary length  $h$  of grids and elements

is small in FDM and FEM, the traditional condition number  $\text{Cond}$  is large (or even huge for local refinements of partitions). However, the effective condition number is small, to display a good stability of numerical methods. This is particularly important to the local refinements used in FDM and FEM for singularity problems in Li [116], explored in Chapters 8 and 9.

The previous study [27, 37, 186] for effective condition number was active until Banoczi et al. [14] in 1998, where a number of numerical examples of linear algebraic equations display insignificance of effective condition number. In fact,  $\text{Cond\_eff}$  is significant for numerical PDE, not for linear algebraic equations<sup>[134]</sup>. For the perturbations in (0.0.5), from Section 1.8 there exists the bound,

$$\frac{\|\Delta \mathbf{x}\|}{\|\mathbf{x}\|} \leq \frac{\text{Cond\_eff}}{1 - \delta} \left[ \frac{1 + \sqrt{5}}{2} \text{Cond} \times \frac{\|\Delta \mathbf{F}\|}{\|\mathbf{F}\|} + \frac{\|\Delta \mathbf{b}\|}{\|\mathbf{b}\|} \right], \quad (0.0.7)$$

where  $\delta = \|\mathbf{F}^\dagger\| \|\Delta \mathbf{F}\| < 1$ , and  $\mathbf{F}^\dagger$  is the pseudo-inverse of  $\mathbf{F}$ . In (0.0.7), the condition number is defined by  $\text{Cond} = \|\mathbf{F}^\dagger\| \|\mathbf{F}\|$ , and the effective condition number by  $\text{Cond\_eff} = \|\mathbf{F}^\dagger\| \frac{\|\mathbf{b}\|}{\|\mathbf{x}\|}$ . For linear algebraic equations, the rounding errors  $\frac{\|\Delta \mathbf{b}\|}{\|\mathbf{b}\|}$  are smaller than the errors  $\frac{\|\Delta \mathbf{F}\|}{\|\mathbf{F}\|}$  of solution methods, such as Gaussian elimination method, so that the condition number plays a dominant role in (0.0.7). However, for numerical PDE, the discretization and the truncation errors  $\frac{\|\Delta \mathbf{b}\|}{\|\mathbf{b}\|}$  are larger than  $\frac{\|\Delta \mathbf{F}\|}{\|\mathbf{F}\|}$ , so that the effective condition number plays a dominant role in (0.0.7).

Here, let us mention the most important references of condition number. The definition of the traditional condition number was given in Wilkinson [227], and then used in many books and papers, see Atkinson [3], Atkinson and Han [5], Christiansen [36], Cucker et al. [42], Geurts [64], Golub and van Loan [66], Laub [108], Parlett [180], Quarteroni and Valli [184], and Schwarz [191]. The condition number for eigenvalues was reported in Parlett [180], and Frayssé and Toumazou [61], and more discussions on condition number were given in Gulliksson and Wedin [71], Elsner et al. [55], Rice [185] and Rohn [187].

This book is a summary of our recent study of effective condition numbers, and the most significant results are selected from more than twenty papers, published in international journals in mathematics and engineering. There are a number of characteristics of this book. The effective condition number is a new criterion for numerical stability of numerical PDE, and this book covers the newest discoveries on this subject. The first characteristic is its novelty. Since the analysis of effective condition number involves two disciplines: linear algebra and partial differ-

mulas of  $\text{Cond}$  and  $\text{Cond}_{\text{eff}}$  are explored, and error bounds are derived. When the minimal singular value  $\sigma_{\min}$  is infinitesimal, there exists a severe subtraction cancellation. This is the other stability. The  $\text{Cond}$  can be regarded as the global stability:  $\text{Cond}_{\text{eff}}$  plus the subtraction. This chapter provides a new stability analysis on TSVD and TR for numerical PDE.

Various problems by different numerical methods for different applications demonstrate the outstanding advantages of the effective condition number over the traditional condition number.

Zi-Cai Li  
Hung-Tsai Huang  
Yimin Wei  
Alexander H.-D. Cheng  
October, 2012

# Acknowledgments

We sincerely appreciated Professors Q. Lin, E. Jiang and Z.H. Cao for their encouragements and supports during our study of effective condition number. We are indebted to Professors T.T. Lu, J. Huang, J.T. Chen, M.G. Lee, C.S. Chien, S.C. Huang, Q. Fang, L.P. Zhang and L. Ling for their collaboration on the publications related to this book. We also express our thanks to Professors C.S. Chen, D.L. Young, O. Axelsson, T.F. Chan, S. Christiansen, P.C. Hansen, L.M. Yeh, Ching Y. Suen and N. Yan, for their valuable suggestions on the research works related to this book.

We are grateful to X. Li and the editorial staff at Academic Science Press, Beijing for their support and assistance in publishing this book. We also wish to thank Professors H.K. Xu and N.C. Wong at the Research Center of Nonlinear Analysis and Discrete Mathematics, National Sun Yat-sen University, Taiwan, for their support during the writing process.

This book is the summary of the international collaborated research among National Sun Yat-sen University, I-Shou University, Taiwan, Fudan University, China, and University of Mississippi, USA. This book and the related works were supported in part by the National Science Council of Taiwan, and by National Natural Science Foundation of China under grant 11271084 and Shanghai Key Laboratory of Contemporary Applied Mathematics.



# Contents

## Preface

## Acknowledgments

<b>Chapter 1</b>	<b>Effective Condition Number</b>	1
1.1	Introduction	1
1.2	Preliminary	3
1.3	Symmetric Matrices	5
1.3.1	Definitions of effective condition numbers	6
1.3.2	A posteriori computation	8
1.4	Overdetermined Systems	9
1.4.1	Basic algorithms	9
1.4.2	Refinements of (1.4.10)	13
1.4.3	Criteria	16
1.4.4	Advanced refinements	17
1.4.5	Effective condition number in $p$ -norms	19
1.5	Linear Algebraic Equations by GE or QR	21
1.6	Application to Numerical PDE	23
1.7	Application to Boundary Integral Equations	33
1.8	Weighted Linear Least Squares Problems	40
1.8.1	Effective condition number	41
1.8.2	Perturbation bounds	44
1.8.3	Applications and comparisons	45
<b>Chapter 2</b>	<b>Collocation Trefftz Methods</b>	47
2.1	Introduction	47
2.2	CTM for Motz's Problem	48
2.3	Bounds of Effective Condition Number	51
2.4	Stability for CTM of $R_p = 1$	55
2.5	Numerical Experiments	57
2.5.1	Choice of $R_p$	57
2.5.2	Extreme accuracy of $D_0$	59
2.6	GCTM Using Piecewise Particular Solutions	60
2.7	Stability Analysis of GCTM	63
2.7.1	Trefftz methods	63
2.7.2	Collocation Trefftz methods	66

2.8	Method of Fundamental Solutions	67
2.9	Collocation Methods Using RBF	70
2.10	Comparisons Between $\text{Cond}_{\text{eff}}$ and $\text{Cond}$	72
2.10.1	CTM using particular solutions for Motz's problem	72
2.10.2	MFS and CM-RBF	73
2.11	A Few Remarks	73
<b>Chapter 3</b>	<b>Simplified Hybrid Trefftz Methods</b>	<b>75</b>
3.1	The Simplified Hybrid TM	75
3.1.1	Algorithms	75
3.1.2	Error analysis	79
3.1.3	Integration approximation	79
3.2	Stability Analysis for Simplified Hybrid TM	80
<b>Chapter 4</b>	<b>Penalty Trefftz Method Coupled with FEM</b>	<b>87</b>
4.1	Introduction	87
4.2	Combinations of TM and Adini's Elements	89
4.2.1	Algorithms	89
4.2.2	Basic theorem	92
4.2.3	Global superconvergence	93
4.3	Bounds of $\text{Cond}_{\text{eff}}$ for Motz's Problem	98
4.4	Effective Condition Number of One and Infinity Norms	104
4.5	Concluding Remarks	105
<b>Chapter 5</b>	<b>Trefftz Methods for Biharmonic Equations with Crack Singularities</b>	<b>107</b>
5.1	Introduction	107
5.2	Collocation Trefftz Methods	107
5.2.1	Three crack models	107
5.2.2	Description of the method	110
5.2.3	Error bounds	111
5.3	Stability Analysis	112
5.3.1	Upper bound for $\sigma_{\max}(\mathbf{F})$	112
5.3.2	Lower bound for $\sigma_{\min}(\mathbf{F})$	113
5.3.3	Upper bound for $\text{Cond}_{\text{eff}}$ and $\text{Cond}$	116
5.4	Proofs of Important Results Used in Section 5.3	117
5.4.1	Basic theorem	117
5.4.2	Proof of Lemma 5.4.3	121
5.4.3	Proof of Lemma 5.4.4	123
5.5	Numerical Experiments	128

5.6	Concluding Remarks	131
<b>Chapter 6</b>	<b>Finite Difference Method</b>	<b>133</b>
6.1	Introduction	133
6.2	Shortley-Weller Difference Approximation	133
6.2.1	A Lemma	135
6.2.2	Bounds for Cond <sub>EE</sub>	137
6.2.3	Bounds for Cond <sub>eff</sub>	141
<b>Chapter 7</b>	<b>Boundary Penalty Techniques of FDM</b>	<b>145</b>
7.1	Introduction	145
7.2	Finite Difference Method	146
7.2.1	Shortley-Weller difference approximation	147
7.2.2	Superconvergence of solution derivatives	147
7.2.3	Bounds for Cond <sub>eff</sub>	148
7.3	Penalty-Integral Techniques	149
7.4	Penalty-Collocation Techniques	154
7.5	Relations Between Penalty-Integral and Penalty-Collocation Techniques	160
7.6	Concluding Remarks	161
<b>Chapter 8</b>	<b>Boundary Singularly Problems by FDM</b>	<b>162</b>
8.1	Introduction	162
8.2	Finite Difference Method	163
8.3	Local Refinements of Difference Grids	164
8.3.1	Basic results	165
8.3.2	Nonhomogeneous Dirichlet and Neumann boundary conditions	171
8.3.3	A remark	173
8.3.4	A view on assumptions A1-A4	176
8.3.5	Discussions and comparisons	177
8.4	Numerical Experiments	177
8.5	Concluding Remarks	183
<b>Chapter 9</b>	<b>Finite Element Method Using Local Mesh Refinements</b>	<b>185</b>
9.1	Introduction	185
9.2	Optimal Convergence Rates	186
9.3	Homogeneous Boundary Conditions	191
9.4	Nonhomogeneous Boundary Conditions	196
9.5	Intrinsic View of Assumption A2 and Improvements of Theorem	
9.4.1		200
9.5.1	Intrinsic view of assumption A2	200

9.5.2	Improvements of Theorem 9.4.1	201
9.6	Numerical Experiments	203
<b>Chapter 10</b>	<b>Hermite FEM for Biharmonic Equations</b>	<b>208</b>
10.1	Introduction	208
10.2	Description of Numerical Methods	209
10.3	Stability Analysis	211
10.3.1	Bounds of Cond	211
10.3.2	Bounds of Cond <sub>eff</sub>	211
10.4	Numerical Experiments	215
<b>Chapter 11</b>	<b>Truncated SVD and Tikhonov Regularization</b>	<b>221</b>
11.1	Introduction	221
11.2	Algorithms of Regularization	224
11.3	New Estimates of Cond and Cond <sub>eff</sub>	225
11.4	Brief Error Analysis	231
<b>Appendix</b>	<b>Definitions and Formulas</b>	<b>236</b>
A.1	Square Systems	236
A.1.1	Symmetric and positive definite matrices	237
A.1.2	Symmetric and nonsingular matrices	239
A.1.3	Nonsingular matrices	239
A.2	Overdetermined Systems	240
A.3	Underdetermined Systems	241
A.4	Method of Fundamental Solutions	242
A.5	Regularization	243
A.5.1	Truncated singular value decomposition	244
A.5.2	Tikhonov regularization	244
A.6	$p$ -Norms	245
A.7	Conclusions	246
<b>Epilogue</b>		<b>247</b>
<b>Bibliography</b>		<b>249</b>
<b>Index</b>		<b>263</b>