

NONLINEAR
PHYSICAL
SCIENCE

L.V. Ovsyannikov

Lectures on the Theory of Group Properties of Differential Equations 微分方程群性质理论讲义

Edited by N. H. Ibragimov



高等教育出版社
HIGHER EDUCATION PRESS

L.V. Ovsyannikov

Lectures on the Theory of Group Properties of Differential Equations

微分方程群性质理论讲义

Weifen Fangcheng Qunxingzhi Lilun Jiangyi

Edited by N.H. Ibragimov
Translated by E.D. Avdonina, N.H. Ibragimov



Author

L.V. Ovsyannikov
Councilor of Russian Academy of Sciences
Lavrentyev Institute of Hydrodynamics
Lavrentyev Prosp. 15
Novosibirsk, 630090, Russia
E-mail: ovs@hydro.nsc.ru

Editor

Nail H. Ibragimov
Department of Mathematics and Science
Blekinge Institute of Technology
371 79 Karlskrona, Sweden
E-mail: nib@bth.se

© 2013 Higher Education Press Limited Company, 4 Dewai Dajie, 100120, Beijing, P. R. China

图书在版编目(CIP)数据

微分方程群性质理论讲义 = Lectures on the theory of
group properties of differential equations : 英文 / (俄罗斯)
奥夫斯亚尼科夫著. — 北京: 高等教育出版社, 2013.4
(非线性物理科学 / 罗朝俊, (瑞典) 伊布拉基莫夫
主编)

ISBN 978-7-04-036944-1

I. ①微… II. ①奥… III. ①微分方程 - 研究 - 英文
②李群 - 研究 - 英文 IV. ①O175 ②O152.5

中国版本图书馆 CIP 数据核字 (2013) 第 029566 号

策划编辑 王丽萍
责任校对 陈旭颖

责任编辑 李华英
责任印制 韩 刚

封面设计 杨立新

版式设计 王艳红

出版发行 高等教育出版社
社 址 北京市西城区德外大街 4 号
邮政编码 100120
印 刷 涿州市星河印刷有限公司
开 本 787mm × 1092mm 1/16
印 张 9.75
字 数 210 千字
购书热线 010-58581118

咨询电话 400-810-0598
网 址 <http://www.hep.edu.cn>
<http://www.hep.com.cn>
网上订购 <http://www.landaco.com>
<http://www.landaco.com.cn>
版 次 2013 年 4 月第 1 版
印 次 2013 年 4 月第 1 次印刷
定 价 49.00 元

本书如有缺页、倒页、脱页等质量问题, 请到所购图书销售部门联系调换

版权所有 侵权必究

物 料 号 36944-00

Sales only inside the mainland of China

{仅限中国大陆地区销售}

本书海外版由 WSP 负责在中国大陆地区之外区域销售, ISBN 为 978-981-4460-81-1

NONLINEAR PHYSICAL SCIENCE

非线性物理科学

NONLINEAR PHYSICAL SCIENCE

Nonlinear Physical Science focuses on recent advances of fundamental theories and principles, analytical and symbolic approaches, as well as computational techniques in nonlinear physical science and nonlinear mathematics with engineering applications.

Topics of interest in *Nonlinear Physical Science* include but are not limited to:

- New findings and discoveries in nonlinear physics and mathematics
- Nonlinearity, complexity and mathematical structures in nonlinear physics
- Nonlinear phenomena and observations in nature and engineering
- Computational methods and theories in complex systems
- Lie group analysis, new theories and principles in mathematical modeling
- Stability, bifurcation, chaos and fractals in physical science and engineering
- Nonlinear chemical and biological physics
- Discontinuity, synchronization and natural complexity in the physical sciences

SERIES EDITORS

Albert C.J. Luo

Department of Mechanical and Industrial Engineering
Southern Illinois University Edwardsville
Edwardsville, IL 62026-1805, USA
Email: aluo@siue.edu

Nail H. Ibragimov

Department of Mathematics and Science
Blekinge Institute of Technology
S-371 79 Karlskrona, Sweden
Email: nib@bth.se

INTERNATIONAL ADVISORY BOARD

Ping Ao, University of Washington, USA; Email: aoping@u.washington.edu

Jan Awrejcewicz, The Technical University of Lodz, Poland; Email: awrejcew@p.lodz.pl

Eugene Benilov, University of Limerick, Ireland; Email: Eugene.Benilov@ul.ie

Eshel Ben-Jacob, Tel Aviv University, Israel; Email: eshel@tamar.tau.ac.il

Maurice Courbage, Université Paris 7, France; Email: maurice.courbage@univ-paris-diderot.fr

Marian Gidea, Northeastern Illinois University, USA; Email: mgidea@neiu.edu

James A. Glazier, Indiana University, USA; Email: glazier@indiana.edu

Shijun Liao, Shanghai Jiaotong University, China; Email: sjliao@sjtu.edu.cn

Jose Antonio Tenreiro Machado, ISEP-Institute of Engineering of Porto, Portugal; Email: jtm@dee.isep.ipp.pt

Nikolai A. Magnitskii, Russian Academy of Sciences, Russia; Email: nmag@isa.ru

Josep J. Masdemont, Universitat Politècnica de Catalunya (UPC), Spain; Email: josep@barquins.upc.edu

Dmitry E. Pelinovsky, McMaster University, Canada; Email: dmpeli@math.mcmaster.ca

Sergey Prants, V.I.II'ichev Pacific Oceanological Institute of the Russian Academy of Sciences, Russia;
Email: prants@poi.dvo.ru

Victor I. Shrira, Keele University, UK; Email: v.i.shrira@keele.ac.uk

Jian Qiao Sun, University of California, USA; Email: jqsun@ucmerced.edu

Abdul-Majid Wazwaz, Saint Xavier University, USA; Email: wazwaz@sxu.edu

Pei Yu, The University of Western Ontario, Canada; Email: pyu@uwo.ca

Editor's preface

When I studied at Novosibirsk State University (Russian) I was lucky to have such brilliant teachers in mathematics as M.A. Lavrentyev, S.L. Sobolev, A.I. Mal'tsev, Yu.G. Reshetnyak and others. But it were L.V. Ovsyannikov's lectures in Ordinary differential equations, Partial differential equations, Gas dynamics and Group properties of differential equations that were of the most benefit for me. I attended his course "Group properties of differential equations" when I was a third-year student. His lectures provided a clear introduction to Lie group methods for determining symmetries of differential equations and a variety of their applications in gas dynamics and other nonlinear models as well as Ovsyannikov's remarkable contribution to this classical subject. His lectures were spectacular not only due to the brilliant presentation of the material but also due to absolutely new discoveries for us. I remember one of our most emotional student's repeated exclamations like "Wonderful, . . . Incredible!" every time when Ovsyannikov revealed most unusual properties of symmetries or unexpected methods.

His lecture notes of this course were published in 1966 with the print of 300 copies only. Since then the Notes were neither reprinted nor translated into English, though they contain the material that is useful for students and teachers but cannot be found in modern texts. For example, theory of partially invariant solutions developed by Ovsyannikov and presented in §3.5, §3.6 is useful for investigating mathematical models described by systems of nonlinear differential equations. It is important to make this classical text available to everyone interested in modern group analysis.

In order to adapt the text for modern students I made several minor changes in the English translation. In particular, sections have been divided into subsections and few misprints have been corrected. Part of the problems formulated in §3.7 have been completely or partially solved since 1966. But we did not make any comments on this matter in the present translation.

Preface

The theory of differential equations has two aspects of investigation, namely local and global, no matter whether the equations arise from applied problems of physics and mechanics or from abstract speculations (which is rather frequent in modern mathematics). The local aspect is characterized by dealing with the inner structure of a family of solutions and its investigation in a neighborhood of a certain point. The global approach deals with solutions defined in some domain and having a given behavior on its boundary.

It would certainly be erroneous to oppose these directions to each other. However, it is no good to ignore the differences in approaches either. While the global approach necessitates the functional analytic apparatus, the local viewpoint allows one to get along with algebraic means only. A brilliant example of a profound local consideration is the famous Cauchy-Kovalevskaya theorem which is, in fact, an algebraic statement. Moreover, it is an easy matter to notice that the theory of boundary value problems also makes an essential application of various algebraic properties of the whole family of solutions. Therefore, the local aspect of the algebraic theory of differential equations is quite vital.

The theory of group properties of differential equations described in the present lecture notes is a typical example of a local theory. It is especially valuable in investigating nonlinear differential equations, for its algorithms act here as reliably as for linear cases.

In spite of the fact that the fundamentals of the theory of group properties were elaborated in works of the Norwegian mathematician Sophus Lie more than a hundred years ago, its development is desirable nowadays as well.

Methodological peculiarity of the present text is that its first chapter uses only the simplest algebraic apparatus of one-parameter groups, which is especially advisable for researchers engaged in applied fields. This allows one to solve the problem of finding a group admitted by a given system of differential equations completely. The second chapter is tailored to provide a deeper insight into the subject resulting from solving determining equations. The group structure of the family of solutions itself is discussed in the third chapter, which also suggests some new elements for the theory. The latter are related to the notions of a partially invariant manifold of the

group, its defect of invariance, and the problem of reduction of partially invariant solutions. The final section §3.7 suggests a qualitative formulation of several problems demonstrating possibilities for further development of the theory with no claim to be complete.

The present lecture notes are written hot on the traces of a special course given by the author in Novosibirsk State University during the 1965/1966 academic year. Such a prompt decision was made to have the lecture notes published by the spring examinations. Therefore, the lectures may appear to be “raw” to many extent, and the author is ready to be completely responsible for that.

The quick release of the lecture notes would be impossible without the support of administration of the university. A major technical work in preparing the manuscript was done by the students V.G. Firsov, E.Z. Borovskaya, T.E. Kuzmina, N.I. Naumenko, M.L. Kochubievskaya and others. The author is sincerely grateful to all these people.

Novosibirsk, Russia, May 1966

L. V. Ovsyannikov

Contents

Editor's preface	v
Preface	vii
1 One-parameter continuous transformation groups admitted by differential equations	1
1.1 One-parameter continuous transformation group	1
1.1.1 Definition	1
1.1.2 Canonical parameter	3
1.1.3 Examples	4
1.1.4 Auxiliary functions of groups	5
1.2 Infinitesimal operator of the group	7
1.2.1 Definition and examples	7
1.2.2 Transformation of functions	9
1.2.3 Change of coordinates	10
1.3 Invariants and invariant manifolds	12
1.3.1 Invariants	12
1.3.2 Invariant manifolds	14
1.3.3 Invariance of regularly defined manifolds	15
1.4 Theory of prolongation	17
1.4.1 Prolongation of the space	17
1.4.2 Prolonged group	18
1.4.3 First prolongation of the group operator	19
1.4.4 Second prolongation of the group operator	22
1.4.5 Properties of prolongations of operators	24
1.5 Groups admitted by differential equations	28
1.5.1 Determining equations	28
1.5.2 First-order ordinary differential equations	30
1.5.3 Second-order ordinary differential equations	32
1.5.4 Heat equation	34
1.5.5 Gasdynamic equations	42

1.6	Lie algebra of operators	47
1.6.1	Commutator. Definition of a Lie algebra	47
1.6.2	Properties of commutator	49
1.6.3	Lie algebra of admitted operators	51
2	Lie algebras and local Lie groups	53
2.1	Lie algebra	53
2.1.1	Definition and examples	53
2.1.2	Subalgebra and ideal	54
2.1.3	Structure of finite-dimensional Lie algebras	55
2.2	Adjoint algebra	57
2.2.1	Inner derivation	57
2.2.2	Adjoint algebra	58
2.2.3	Inner automorphisms of a Lie algebra	60
2.3	Local Lie group	61
2.3.1	Coordinates in a group	61
2.3.2	Subgroups	62
2.3.3	Canonical coordinates of the first kind	64
2.3.4	First fundamental theorem of Lie	65
2.3.5	Second fundamental theorem of Lie	69
2.3.6	Properties of canonical coordinate systems of the first kind	70
2.3.7	Third fundamental theorem of Lie	71
2.3.8	Lie algebra of a local Lie group	73
2.4	Subgroup, normal subgroup and factor group	75
2.4.1	Lemma on commutator	75
2.4.2	Subgroup	77
2.4.3	Normal subgroup	79
2.4.4	Factor group	80
2.5	Inner automorphisms of a group and of its Lie algebra	82
2.5.1	Inner automorphism	82
2.5.2	Lie algebra of G^A and adjoint algebra of L_r	83
2.6	Local Lie group of transformations	85
2.6.1	Introduction	85
2.6.2	Lie's first theorem	85
2.6.3	Lie's second theorem	87
2.6.4	Canonical coordinates of the second kind	88
3	Group invariant solutions of differential equations	91
3.1	Invariants of the group G_r^N	91
3.1.1	Invariance criterion	91
3.1.2	Functional independence	92
3.1.3	Linearly unconnected operators	93
3.1.4	Integration of Jacobian systems	95
3.1.5	Computation of invariance	96

3.2	Invariant manifolds	98
3.2.1	Invariant manifolds criterion	98
3.2.2	Induced group and its Lie algebra	98
3.2.3	Theorem on representation of nonsingular invariant manifolds	100
3.2.4	Differential invariant manifolds	102
3.3	Invariant solutions of differential equations	103
3.3.1	Definition of invariant solutions	103
3.3.2	The system (S/H)	106
3.3.3	Examples from one-dimensional gas dynamics	109
3.3.4	Self-similar solutions	111
3.4	Classification of invariant solutions	116
3.4.1	Invariant solutions for similar subalgebras	116
3.4.2	Classes of similar subalgebras	118
3.5	Partially invariant solutions	121
3.5.1	Partially invariant manifolds	121
3.5.2	Defect of invariance	123
3.5.3	Construction of partially invariant solutions	125
3.6	Reduction of partially invariant solutions	129
3.6.1	Statement of the reduction problem	129
3.6.2	Two auxiliary lemmas	132
3.6.3	Theorem on reduction	136
3.7	Some problems	139

Chapter 1

One-parameter continuous transformation groups admitted by differential equations

1.1 One-parameter continuous transformation group

1.1.1 Definition

Let us consider transformations T of an N -dimensional Euclidian space E^N into itself, so that one has

$$x' = Tx \in E^N$$

for $x \in E^N$. If the point x has coordinates x^1, \dots, x^N then this transformation can be given by the system of equalities

$$x'^i = f^i(x) = f^i(x^1, \dots, x^N) \quad (i = 1, \dots, N). \quad (1.1.1)$$

The functions f are assumed to be thrice continuously differentiable and locally invertible, which means that there exists an inverse transformation T^{-1} for the transformation T in some neighborhood of the point $x' = Tx$, so that $x = T^{-1}x'$.

The product of transformations $T_1 T_2$ is the transformation T consisting in successive transformations T_2 and then T_1 . The identity transformation plays the role of the unit transformation with respect to the product.

The above product is written in terms of functions f by the following equations:

$$f^i(x) = f_1^i(f_2^1(x), \dots, f_2^N(x)) \quad (i = 1, \dots, N).$$

This operation of multiplication is associative, i.e.

$$T_1(T_2 T_3) = (T_1 T_2) T_3.$$

Note that the inversion of the product of transformations is given by

$$(T_1 T_2)^{-1} = T_2^{-1} T_1^{-1}.$$

We will consider a family of transformations $\{T_a\}$ with the above properties depending on a real parameter a that varies within an interval Δ .

The family $\{T_a\}$ is said to be *locally closed with respect to the product* if there exists a subinterval $\Delta' \subset \Delta$ such that

$$T_b T_a \in \{T_a\}$$

for any $a, b \in \Delta'$. This leads to a function $c = \varphi(a, b)$ which determines the *multiplication law* for transformations of $\{T_a\}$ according to the formula

$$T_b T_a = T_c.$$

The transformation T_a is written in coordinates in the following form similar to Eqs. (1.1.1):

$$T_a : x^i = f^i(x, a) \quad (i = 1, \dots, N). \quad (1.1.2)$$

The product of transformations is written in terms of the functions (1.1.2) and the multiplication law $\varphi(a, b)$ as follows:

$$T_b T_a = T_{\varphi(a, b)} : f^i(f(x, a), b) = f^i(x, \varphi(a, b)) \quad (i = 1, \dots, N). \quad (1.1.3)$$

Definition 1.1. The family $\{T_a\}$ is called a *local one-parameter continuous transformation group* if it is locally closed with respect to the product and if the interval Δ' can be chosen so that the following conditions hold.

1° There exists the unique value $a_0 \in \Delta'$ such that T_{a_0} is an identity transformation.

2° The function $\varphi(a, b)$ is thrice continuously differentiable and the equation $\varphi(a, b) = a_0$ has the unique solution $b = a^{-1}$ for any $a \in \Delta'$.

Condition 2° means that the operation of inversion of transformations $(T_a)^{-1} = T_{a^{-1}}$ is possible in $\{T_a\}$.

Hereafter the symbol a^{-1} indicates only a definite value of the parameter and not the inverse value of the number a , so that $a^{-1} \neq \frac{1}{a}$.

The choice of the interval Δ' is not unique, generally speaking. If such an interval is selected one can take any smaller interval instead of Δ' . It means that we are interested only in some sufficiently small neighborhood of a_0 . The operations of multiplication and inversion of transformations T_a are feasible only for values of the parameter a from the above neighborhood. Therefore, the object introduced by Definition 1.1 is termed as a *local group*. In what follows the sufficient closeness of all considered values of the parameters a, b, \dots to the value a_0 is provided.

Further on, the term “group G_1 ” will be used to indicate a local one-parameter continuous transformation group.

1.1.2 Canonical parameter

Generally introduction of the new parameter $\bar{a} = \bar{a}(a)$, where $\bar{a}(a)$ is a thrice continuously differentiable monotonous function, changes φ, Δ and Δ' .

In what follows, we assume that $a_0 = 0$ without loss of generality. Note that in this case the definition leads to the following properties of the function $\varphi(a, b)$:

$$\begin{aligned}\varphi(0, 0) &= 0, & \varphi(a, 0) &= a, & \varphi(0, b) &= b, \\ \varphi(a, a^{-1}) &= \varphi(a^{-1}, a) = 0.\end{aligned}\tag{1.1.4}$$

The parameter a is said to be canonical if the multiplication law is given by

$$\varphi(a, b) = a + b.$$

Then $a^{-1} = -a$ and equations (1.1.3) take the form

$$f^i(f(x, a), b) = f^i(x, a + b) \quad (i = 1, \dots, N).\tag{1.1.5}$$

Theorem 1.1. A canonical parameter can be introduced in any one-parameter group.

Proof. Let $T_c = T_b T_a$, so that $c = \varphi(a, b)$. Let us give a small increment Δb to the parameter b , then c receives a small increment Δc , so that

$$\varphi(a, b + \Delta b) = c + \Delta c.$$

For transformations it is written by the formula

$$T_{b+\Delta b} T_a = T_{c+\Delta c}.$$

Multiplying the right-hand side by

$$T_c^{-1} = T_a^{-1} T_b^{-1},$$

one obtains

$$T_{c+\Delta c} T_c^{-1} = T_{b+\Delta b} T_b^{-1}$$

due to the associative multiplication law. The equality has the form

$$\varphi(c^{-1}, c + \Delta c) = \varphi(b^{-1}, b + \Delta b)\tag{1.1.6}$$

in terms of the function φ .

Let

$$V(b) = \left. \frac{\partial \varphi(a, b)}{\partial b} \right|_{a=b^{-1}}.$$

Taylor's formula and the equation $\varphi(b^{-1}, b) = 0$ yield

$$\varphi(b^{-1}, b + \Delta b) = V(b)\Delta b + O(|\Delta b|^2).$$

Applying the above equation to Eq. (1.1.6) and invoking that $|\Delta c| = O|\Delta b|$ one obtains

$$V(c)\Delta c = V(b)\Delta b + O(|\Delta b|^2). \quad (1.1.7)$$

Dividing both sides of Eq. (1.1.7) by Δb and taking the limit $\Delta b \rightarrow 0$, one arrives at the differential equation

$$V(c) \frac{dc}{db} = V(b) \quad (1.1.8)$$

with the initial condition

$$c|_{b=0} = a.$$

Furthermore, equations (1.1.4) show that $V(0) = 1$.

Let us introduce the function

$$\bar{a}(a) = \int_0^a V(s)ds.$$

Then the function $c = \varphi(a, b)$, determined by the relations

$$\bar{a}(c) = \bar{a}(a) + \bar{a}(b), \quad (1.1.9)$$

is a solution to Eq. (1.1.8).

The function $\bar{a}(a)$ is obviously monotonous and thrice continuously differentiable with respect to a . Taking it as a new parameter, one obtains that the new parameter is canonical due to Eq. (1.1.9).

Corollary 1.1. Any one-parameter transformation group is Abelian. Indeed, if a is a canonical parameter then, according to the definition, one has

$$T_b T_a = T_{a+b} = T_{b+a} = T_a T_b.$$

1.1.3 Examples

Example 1.1. *Translations on a straight line:*

$$x' = x + a.$$

Here

$$\varphi(a, b) = a + b.$$

Translations in an N -dimensional space in the direction of the vector $\lambda = (\lambda^1, \dots, \lambda^N)$ are given by

$$x'^i = x^i + \lambda^i a \quad (i = 1, \dots, N).$$