

美国医师执照考试

High-Yield™ *Histopathology*

组织病理学

(第2版)


RONALD W. DUDEK

High-Yield™ Histopathology
is designed to:

- Provide an uncomplicated review of histology and pathology
- Help equip you for the histology and pathology questions on the USMLE
- Clarify difficult concepts through simple illustrations and key images



北京大学医学出版社



美国医师执照考试

High-Yield™ 组织病理学

Histopathology

(第2版)

Ronald W. Dudek, PhD


Professor

Department of Anatomy and Cell Biology

Brody School of Medicine

East Carolina University

Greenville, North Carolina



北京大学医学出版社

图书在版编目 (CIP) 数据

High-Yield™ 组织病理学 = High-Yield™ Histopathology: 第2版: 英文 / (美) 杜德克 (Dudek, R. W.) 编写. — 北京: 北京大学医学出版社, 2013. 1

(美国医师执照考试丛书)

ISBN 978-7-5659-0480-6

I. ①H… II. ①杜… III. ①组织病理学—医师—资格考核—美国—自学参考资料—英文 IV. ①R361
中国版本图书馆CIP数据核字(2012)第255507号

北京市版权局著作权合同登记号: 01-2012-5817

Ronald W. Dudek: Histopathology 2nd ed

ISBN: 978-1-60913-015-2

Copyright©2011 by Lippincott Williams & Wilkins, a Wolters Kluwer business.

All Rights reserved.

This is a reprint under copublishing agreement with Lippincott Williams & Wilkins/Wolters Kluwer Health, Inc., USA

本书限在中华人民共和国境内 (不包括香港、澳门特别行政区及台湾) 销售。

本书封底贴有 Wolters Kluwer Health 激光防伪标签, 无标签者不得销售。

本书提供了药物的准确的适应证、副作用和疗程剂量, 但有可能发生改变。读者须阅读药商提供的外包装上的用药信息。作者、编辑、出版者或发行者对因使用本书信息所造成的错误、疏忽或任何后果不承担责任, 对出版物的内容不做明示的或隐含的保证。作者、编辑、出版者或发行者对由本书引起的任何人身伤害或财产损失不承担任何法律责任。

美国医师执照考试: High-Yield™ 组织病理学 (第2版)

编 写: Ronald W. Dudek

出版发行: 北京大学医学出版社(电话: 010-82802230)

地 址: (100191)北京市海淀区学院路38号 北京大学医学部院内

网 址: <http://www.pumpress.com.cn>

E-mail: booksale@bjmu.edu.cn

印 刷: 北京京华虎彩印刷有限公司

经 销: 新华书店

责任编辑: 冯智勇 责任印制: 张京生

开 本: 787×1092mm 1/16 印张: 21.5 字数: 546千字

版 次: 2013年1月第2版 2013年1月第1次印刷

书 号: ISBN 978-7-5659-0480-6

定 价: 89.00元

版权所有, 违者必究

(凡属质量问题请与本社发行部联系退换)

出版说明

High-Yield™ 系列丛书是针对美国医师执照考试 (United States Medical Licensing Examination, USMLE) 的知名品牌图书, 受到世界各地读者的欢迎。该系列丛书具有以下特色:

1. 内容高度概括, 重点突出, 有利于读者快速掌握学科的核心知识。
2. 编排新颖, 既有基础知识要点的介绍, 又有以疾病为核心的综合归纳, 并体现了相关学科的横向联系。
3. 语言规范、地道, 既有利于读者快速掌握专业词汇, 又有利于医学英语思维的培养。

本系列丛书是参加美国医师执照考试的必备辅导用书, 也可作为我国医学院校从事双语教学的教材和参考用书, 对教师进行英语授课, 学生学习、参加考试具有重要的参考价值。

北京大学医学出版社

I would like to dedicate this book to my mother, Lottie Dudek, who was born on November 11, 1918. Through the years my mother raised her children, maintained a loving marriage, and worked 40 hours per week. In the year 2004, society would describe such a person as a "liberated woman" or "supermom." I would like to acknowledge that my mother was a "supermom" 20 years before the word was fashionable. A son cannot repay a mother. My hope is that "I love you and thank you" will suffice.

Preface

High-Yield Histopathology does more than just review histology. The questions on the USMLE Step 1 cross traditional course boundaries, making it difficult to identify a question that is “strictly histology.” Many USMLE Step 1 questions fall into the categories such as histopathology, histophysiology, histomicrobiology, and histopharmacology. To write a review book on basic, traditional histology would not be helpful to the student preparing for the USMLE Step 1 since there are no basic traditional histology questions on the exam. In this regard, *High-Yield Histopathology* reviews important histology concepts as a gateway to the pathology, physiology, microbiology, and pharmacology of clinically relevant topics.

In addition, many students have commented that cell biology topics have been well represented on the USMLE Step 1. To this end, I have included Chapter 1 (Nucleus), Chapter 2 (Cytoplasm and Organelles), and Chapter 3 (Cell Membrane) with up-to-date and clinically relevant information.

I would appreciate any comments or suggestions concerning *High-Yield Histopathology*, especially after you have taken the USMLE Step 1 exam, that you think might improve the book. You may contact me at dudekr@ecu.edu.

Contents

Preface	vii
---------------	-----

1 Nucleus	1
I. Nuclear Envelope	1
II. Apoptosis	1
III. Nucleolus	1
IV. Chromatin	2
V. Chromosomes	2
VI. Types of DNA Damage and DNA Repair	3
VII. Clinical Importance of DNA Repair	3
VIII. Cell Cycle	4
IX. Proto-Oncogenes and Oncogenes	8
X. Tumor Suppressor Genes	10
XI. Oncofetal Antigens and Tumor Markers	11
XII. Transcription in Protein Synthesis	11
XIII. Processing the RNA Transcript into mRNA	11
XIV. Selected Photomicrographs	13

2 Cytoplasm and Organelles	18
I. Cytoplasm	18
II. Ribosomes	18
III. Rough Endoplasmic Reticulum (rER)	19
IV. Translation	20
V. Golgi Complex	22
VI. Smooth Endoplasmic Reticulum (sER)	22
VII. Mitochondria	22
VIII. Lysosomes	25
IX. Peroxisomes	25
X. Cytoskeleton	27
XI. Lipofuscin	28
XII. Hemosiderin	28
XIII. Glycogen	29
XIV. Selected Photomicrographs	31

3 The Cell Membrane: Eicosanoids and Receptors/Signal Transduction	36
I. The Lipid Component of the Cell Membrane	36
II. The Protein Component of the Cell Membrane	38
III. Membrane Transport Proteins	38

IV. G Protein-linked Receptors	42
V. Types of G Protein-linked Receptors	44
VI. Enzyme-linked Receptors	46
VII. Low-density Lipoprotein (LDL) Receptor	48

4 Epithelium 49

I. Introduction	49
II. Classification of Epithelium	49
III. Apical Region	50
IV. Lateral Region	51
V. Basal Region	52
VI. Clinical Considerations	52
VII. Selected Photomicrographs	54

5 Connective Tissue 55

I. Introduction	55
II. Ground Substance	55
III. Fibers	55
IV. Cells	56
V. Clinical Considerations	58
VI. Selected Photomicrographs	60

6 Cartilage 62

I. Introduction	62
II. Ground Substance	62
III. Fibers	62
IV. Cells	62
V. Blood Vessels and Nerves	62
VI. Chondrogenesis	63
VII. Hormonal Influence	63
VIII. Repair	63

7 Bone 64

I. Introduction	64
II. Ground Substance	64
III. Fibers	64
IV. Cells	64
V. Blood Vessels and Nerves	65
VI. Osteogenesis	65
VII. Bone Repair	66
VIII. Hormonal Influence	66
IX. Cartilage and Bone Comparison	67
X. Clinical Considerations of Bone	67
XI. Clinical Considerations of Joints	68
XII. Selected Photomicrographs	69

8 Muscle 73

I. Skeletal Muscle	73
II. Cardiac Muscle	77

III. Smooth Muscle	77
IV. Comparisons and Contrasts of Skeletal, Cardiac, and Smooth Muscle	79
V. Selected Photomicrographs	80

9 Nervous Tissue83

I. The Neuron	83
II. Neurotransmitters	85
III. Parasympathetic Pharmacology	86
IV. Sympathetic Pharmacology	86
V. Neuroglial Cells	87
VI. The Blood-Brain Barrier (BB)	88
VII. Nerve Degeneration and Regeneration	88
VIII. Clinical Considerations	88
IX. Selected Photomicrographs	92

10 Heart and Blood Vessels97

I. Heart Layers	97
II. Cardiac Myocytes	97
III. Purkinje Myocytes	99
IV. Myocardial Endocrine Cells	99
V. Conduction System	100
VI. Parasympathetic Regulation of Heart Rate	102
VII. Sympathetic Regulation of Heart Rate	102
VIII. Clinical Consideration: Myocardial Infarction (MI)	103
IX. Tunics of Blood Vessels	105
X. Types of Blood Vessels	107
XI. Functions of Endothelium	107
XII. Clinical Considerations	108

11 Blood111

I. Plasma	111
II. Red Blood Cells	111
III. Hemoglobin (Hb)	113
IV. Clinical Considerations	114
V. White Blood Cells	117
VI. Platelets (Thrombocytes)	118
VII. Hemostasis (Blood Clotting)	119
VIII. Clinical Considerations	121
IX. Red Bone Marrow (Myeloid Tissue)	122
X. Selected Photomicrographs	123

12 Thymus130

I. General Features	130
II. Thymic Cortex	130
III. Thymic Medulla	130
IV. Types of Mature T Cells	131
V. Blood-Thymus Barrier	132
VI. T Cell Lymphopoiesis (T Cell Formation)	132
VII. Clinical Considerations	134
VIII. Selected Photomicrographs	134

13	Lymph Node	135
	I. General Features	135
	II. Outer Cortex	136
	III. Inner Cortex	136
	IV. Medulla	136
	V. Flow of Lymph	136
	VI. Flow of Blood	136
	VII. B-Cell Lymphopoiesis (B-cell Formation)	136
	VIII. Cytokines	138
	IX. Clinical Consideration	141
14	Spleen	143
	I. General Features	143
	II. White Pulp	143
	III. Marginal Zone	143
	IV. Red Pulp	143
	V. Blood Flow	145
	VI. Clinical Considerations	145
	VII. Hypersensitivity Reactions	145
15	Esophagus and Stomach	147
	Esophagus	147
	I. General Features	147
	II. Mucosa	147
	III. Submucosa	147
	IV. Muscularis Externa	147
	V. Gastroesophageal (GE) Junction	147
	VI. Clinical Considerations	148
	Stomach	148
	I. General Features	148
	II. Gastric Mucosa	148
	III. Gastric Glands	148
	IV. Clinical Considerations	150
16	Small Intestine	152
	I. General Features	152
	II. Intestinal Mucosa	152
	III. Intestinal Glands (Crypts of Lieberkühn)	154
	IV. Gut-Associated Lymphatic Tissue (GALT; Peyer Patches)	155
	V. Clinical Considerations	156
17	Large Intestine (Colon)	160
	I. General Features	160
	II. Large Intestinal Mucosa	160
	III. Intestinal Glands	160
	IV. GUT-Associated Lymphatic Tissue (GALT)	160
	V. Anal Canal	160
	VI. Clinical Considerations	162

18	Liver and Gallbladder	165
	I. Hepatocytes	165
	II. Kupffer Cells	168
	III. Hepatic Stellate Cells (Fat-Storing Cells; ITO Cells)	168
	IV. Classic Liver Lobule	168
	V. Liver Acinus	169
	VI. Repair (Regeneration)	170
	VII. Clinical Considerations	170
	VIII. Selected Photomicrographs	172
19	Exocrine Pancreas and Islets of Langerhans	175
	I. Exocrine Pancreas	175
	II. Endocrine Pancreas	175
	III. Clinical Considerations	177
	IV. Selected Photomicrographs	179
20	Respiratory System	183
	I. General Features	183
	II. Trachea	183
	III. Bronchi	183
	IV. Bronchioles	184
	V. Terminal Bronchioles	184
	VI. Respiratory Bronchioles	184
	VII. Alveolar Ducts	184
	VIII. Alveoli	184
	IX. Surfactant	186
	X. Blood-Air Barrier	186
	XI. Air Flow	186
	XII. Clinical Considerations	187
	XIII. Selected Photomicrographs	190
21	Urinary System	196
	I. General Features	196
	II. Internal Structure of the Kidney	196
	III. Nephrons	197
	IV. Collecting Duct (CD)	199
	V. Renal Vasculature	202
	VI. Hormonal Control of the Kidney	204
	VII. Glomerular Filtration Barrier (GFB)	206
	VIII. Juxtaglomerular (JG) Complex	208
	IX. Pharmacology of Diuretics	208
	X. Clinical Considerations	211
22	Hypophysis	223
	I. The Adenohypophysis	223
	II. Hormonal Secretion	224
	III. The Neurohypophysis	224

23	Thyroid	226
	I. Thyroid Follicles	226
	II. Follicular Cells	226
	III. Functions of T ₃ and T ₄	228
	IV. Parafollicular Cells	228
	V. Clinical Considerations	228
	VI. Pharmacology of the Thyroid	230
	VII. Selected Photomicrographs	230
24	Parathyroid	234
	I. Chief Cells	234
	II. Oxyphil Cells	234
	III. Calcium Homeostasis	234
	IV. Clinical Considerations	236
	V. Pharmacology of Calcium Homeostasis	237
	VI. Selected Photomicrographs	237
25	Adrenal	238
	I. Cortex	238
	II. The Medulla	242
	III. Selected Photomicrographs	244
26	Female Reproductive System	251
	I. Ovary	251
	II. Corpus Luteum	252
	III. Uterine Tubes (fallopian tubes; oviducts)	254
	IV. Uterus	255
	V. The Menstrual Cycle	256
	VI. Cervix	259
	VII. Ectocervix	259
	VIII. Vagina	260
	IX. Histopathology of the Vagina	261
	X. Breast	263
	XI. Selected Photomicrographs	265
27	Male Reproductive System	269
	I. Testes	269
	II. Duct System	277
	III. Accessory Glands	278
28	Skin	284
	I. General Features	284
	II. Epidermis	284
	III. Dermis	286
	IV. Glands	286
	V. Nerves	286
	VI. Clinical Considerations	287

29	Eye	290
	I. General Features	290
	II. Cornea	290
	III. Sclera	291
	IV. Limbus	291
	V. Iris	291
	VI. Ciliary Body	292
	VII. Lens	292
	VIII. Retina	294
	IX. Clinical Considerations	295
	X. Selected Photomicrographs	295
30	Ear	298
	I. General Features	298
	II. External Ear	298
	III. Middle Ear	298
	IV. Internal Ear	299
	V. Clinical Considerations	301
	Credits	302
	Index	307

Nucleus

I Nuclear Envelope. The nuclear envelope is a two-membrane structure. The **inner membrane** is associated with a network of **intermediate filaments (lamins A, B, C)** called the **nuclear lamina**, which plays a role in the disassembly of the nuclear envelope during prometaphase of mitosis by phosphorylation of the lamins by **lamin kinase** and in the reassembly of the nuclear envelope during telophase. The **outer membrane** is studded with ribosomes and is continuous with the rough endoplasmic reticulum (rER). The inner and outer membranes are separated by a **perinuclear cisterna**. The nuclear envelope contains many pores that allow passage of molecules between the nucleus and cytoplasm (e.g., ions, messenger RNA (mRNA), transfer RNA (tRNA), ribosomal RNA (rRNA), gene regulatory proteins, DNA polymerases, RNA polymerases). The pores are associated with a **nuclear pore complex** that consists of many different proteins arranged in octagonal symmetry with a central channel.

II Apoptosis. Apoptosis is a noninflammatory programmed cell death ("cell suicide") that is characterized by DNA fragmentation, a decrease in cell volume, loss of mitochondrial function, cell membrane blebbing, and formation of apoptotic bodies, which are rapidly phagocytosed without an inflammatory response.

- A.** The chromatin is eventually cleaved by a specific endonuclease into DNA fragments that generate a distinctive **180-bp ladder** that is pathognomonic of apoptotic cell death.
- B.** Apoptosis is related to a family of proteases called **caspases**, which are found in all cells.
- C.** Caspases are activated by either extracellular death signals (e.g., killer lymphocytes produce **Fas ligand**, which binds to the **Fas death receptor** on the target cell; **tumor necrosis factor [TNF]** binds to the **TNF death receptor** on the target cell) or intracellular death signals (e.g., mitochondria release **cytochrome c** into the cytoplasm where it activates **Apaf-1 adaptor protein**, which in turn activates caspases).
- D.** The **Bcl-2** and the **IAP (inhibitor of apoptosis) family of proteins** are the main regulators of apoptosis.
- E.** Apoptosis occurs in hormone-dependent involution of cells during the menstrual cycle, embryogenesis, toxin-induced injury (e.g., diphtheria), viral cell death (e.g., Councilman bodies in yellow fever), and cell death via cytotoxic T cells or other immune cells.

III Nucleolus

- A.** The nucleolus consists of portions of five pairs of chromosomes (i.e., 13, 14, 15, 21, and 22) that contain about **200 copies of rRNA genes** per haploid genome that code for rRNA.

- B.** In humans, RNA polymerase I catalyzes the formation of 45S rRNA and RNA polymerase III catalyzes the formation of 5S RNA.

IV

Chromatin. Chromatin is double-helical DNA associated with histones and nonhistone proteins.

- A. HETEROCHROMATIN** is condensed chromatin and is **transcriptionally inactive**. In electron micrographs, heterochromatin is electron dense (i.e., very black). An example of heterochromatin is the **Barr body**, which is found in female cells and represents the inactive X chromosome. Heterochromatin makes up ~10% of the total chromatin.
- 1. Constitutive heterochromatin** is always condensed (i.e., transcriptionally inactive) and consists of repetitive DNA found near the centromere and other regions.
 - 2. Facultative heterochromatin** can be either condensed (i.e., transcriptionally inactive) or dispersed (i.e., transcriptionally active). An example of facultative heterochromatin is the **XY body**, which forms when both the X and Y chromosome are inactivated for ~15 days during male meiosis.
- B. EUCHROMATIN** is dispersed chromatin and makes up ~90% of the total chromatin. Of this 90%, 10% is transcriptionally active and 80% is transcriptionally inactive. When chromatin is transcriptionally active, there is weak binding to the H1 histone protein and acetylation of the H2A, H2B, H3, and H4 histone proteins.
- C. NUCLEOSOME**
- The most fundamental unit of packaging of DNA is the nucleosome.
 - A nucleosome consists of a histone protein octamer (two each of H2A, H2B, H3, and H4 histone proteins) around which 146 bp of DNA is coiled in 1.75 turns. The nucleosomes are connected by spacer DNA, which results in 10-nm-diameter fiber that resembles a “beads on a string” appearance by electron microscopy.
 - Histones are small proteins containing a high proportion of lysine and arginine that impart a positive charge to the proteins that enhances their binding to negatively charged DNA. Histones bind to DNA in A-T-rich regions of DNA.
 - Histone proteins have exposed N-terminal amino acid tails that are subject to modification and are crucial in regulating nucleosome structure.
 - Histone acetylation** reduces the affinity between histones and DNA. An increased acetylation of histone proteins will make a DNA segment more likely to be transcribed into RNA and hence any genes in that DNA segment will be expressed (i.e., $\uparrow$ acetylation of histones = expressed genes).
- D. 30-nm CHROMATIN FIBER.** The 10-nm nucleosome fiber is joined by H1 histone protein to form a 30-nm chromatin fiber. When the general term “chromatin” is used, it refers specifically to the 30-nm chromatin fiber.

V

Chromosomes. The human genome refers to the total DNA content in the cell, which is divided into two genomes: the very complex **nuclear genome** and the relatively simple **mitochondrial genome**. The human nuclear genome consists of 24 different chromosomes (22 autosomes; X and Y sex chromosomes). The human nuclear genome codes for ~30,000 genes (precise number is uncertain), which make up ~2% of the human nuclear genome. There are ~27,000 protein-coding genes and ~3000 RNA-coding genes. The fact that the ~30,000 genes make up only ~2% of the human nuclear genome means that ~2% of the human nuclear genome consists of coding DNA and ~98% of the human nuclear genome consists of noncoding DNA.

A. CENTROMERE

- A centromere is a specialized nucleotide DNA sequence that binds to the mitotic spindle during cell division.

2. Chromosomes have a single centromere that is observed microscopically as a **primary constriction**, which is the region where sister chromatids are joined.
3. During prometaphase, a pair of protein complexes called **kinetochores** forms at the centromere where one kinetochore is attached to each sister chromatid.
4. Microtubules produced by the **centrosome** of the cell attach to the kinetochore (called **kinetochore microtubules**) and pull the two sister chromatids toward opposite poles of the mitotic cell.

B. THE TELOMERE

1. The human telomere is a 3- to 20-kb repeating nucleotide sequence (TTAGGG) located at the end of a chromosome.
2. The telomere allows replication of linear DNA to its full length. Since DNA polymerases *cannot* synthesize in the 3' → 5' direction or start synthesis *de novo*, removal of the RNA primers will always leave the 5' end of the newly synthesized lagging strand shorter than the lagging strand template. If the 5' end of the newly synthesized lagging strand is not lengthened, a chromosome would get progressively shorter as the cell goes through a number of cell divisions. This would lead to cell death, which some investigators believe may be related to the aging process in humans.
3. This problem is solved by a special RNA-directed DNA polymerase or reverse transcriptase called **telomerase**, which adds many repeats of TTAGGG to the newly synthesized lagging strand.
4. Telomerase is present in **human germline cells** (i.e., spermatogonia, oogonia) and **stem cells** (e.g., in skin, bone marrow, and gut), but is absent from most other somatic cells.

VI

Types of DNA Damage and DNA Repair. Chromosomal breakage refers to breaks in chromosomes due to sunlight (or ultraviolet [UV]) irradiation, ionizing irradiation, DNA cross-linking agents, or DNA-damaging agents. These insults may cause **depurination** of DNA, **deamination** of cytosine to uracil, or **pyrimidine dimerization**, which must be repaired by **DNA repair enzymes**. The system that detects and signals DNA damage is a multiprotein complex called **BASC** (**BRCA1-associated genome surveillance complex**).

- A. DEPURINATION.** About 5000 purines (As or Gs) per day are lost from DNA of each human cell when the *N*-glycosyl bond between the purine and deoxyribose sugar phosphate is broken. This is the most frequent type of lesion and leaves the deoxyribose sugar phosphate with a missing purine base.
- B. DEAMINATION OF CYTOSINE TO URACIL.** About 100 cytosines (C) per day are spontaneously deaminated to uracil (U). If the U is not corrected back to a C, then upon replication, instead of the occurrence of a correct C-G base pairing, a U-A base pairing will occur instead.
- C. PYRIMIDINE DIMERIZATION.** Sunlight (UV radiation) can cause covalent linkage of adjacent pyrimidines forming, for example, **thymine dimers**.

VII

Clinical Importance of DNA Repair (Table 1-1). The clinical importance of DNA repair enzymes is illustrated by some rare inherited diseases that involve genetic defects in DNA repair enzymes such as xeroderma pigmentosa (XP), ataxia-telangiectasia (AT), Fanconi anemia (FA), and Bloom syndrome (BS), as indicated in Table 1-1.

TABLE 1-1

DNA REPAIR ENZYME PATHOLOGY

Genetic Disorder	Gene Gene Product Chromosome	Clinical Features
Xeroderma pigmentosum (XP) is an autosomal recessive genetic disorder caused by mutations in nucleotide excision repair enzymes that results in the inability to remove pyrimidine dimers and individuals who are hypersensitive to sunlight (ultraviolet [UV] radiation)	<i>XPA</i> gene DNA repair enzyme 9q22.3 <i>XPC</i> gene DNA repair enzyme 3p25	Sunlight (UV radiation) hypersensitivity with sunburnlike reaction, severe skin lesions around the eyes and eyelids, and malignant skin cancers (basal and squamous cell carcinomas and melanomas) whereby most individuals die by 30 years of age
Ataxia-telangiectasia (AT) is an autosomal recessive genetic disorder caused by mutations in DNA recombination repair enzymes that results in individuals who are hypersensitive to ionizing radiation	<i>ATM</i> gene PI-3 kinase and a DNA repair enzyme/cell cycle checkpoint protein 11q22-q23	Ionizing radiation hypersensitivity; cerebellar ataxia with depletion of Purkinje cells; progressive nystagmus; slurred speech; oculocutaneous telangiectasia initially in the bulbar conjunctiva followed by ear, eyelid, cheeks, and neck; immunodeficiency; and death in the second decade of life. A high frequency of structural rearrangements of chromosomes 7 and 14 is the cytogenetic observation with this disease.
Fanconi anemia (FA) is an autosomal recessive genetic disorder caused by mutations in DNA recombination repair that results in individuals who are hypersensitive to DNA cross-linking agents	<i>FA-A</i> gene A protein that normalizes cell growth, corrects sensitivity to chromosomal breakage in the presence of mitomycin C, and generally promotes genomic stability 16q24	DNA cross-linking agent hypersensitivity, short stature, hypopigmented spots, café-au-lait spots, hypogonadism, microcephaly, hypoplastic or aplastic thumbs, renal malformation including unilateral aplasia or horseshoe kidney, acute leukemia, progressive aplastic anemia, head and neck tumors, and medulloblastoma; is the most common form of congenital aplastic anemia
Bloom syndrome (BS) is an autosomal recessive genetic disorder caused by mutations in DNA repair enzymes that results in individuals who are hypersensitive to DNA-damaging agents	<i>BLM</i> gene RecQ helicase 15q26	Hypersensitivity to DNA-damaging agents; long, narrow face; erythema with telangiectasias in butterfly distribution over the nose and cheeks; high-pitched voice; small stature; small mandible; protuberant ears; absence of upper lateral incisors; well-demarcated patches of hypopigmentation and hyperpigmentation; immunodeficiency with decreased immunoglobulin A (IgA), IgM, and IgG levels; and predisposition to several types of cancers

VIII Cell Cycle

A. PHASES OF THE CELL CYCLE (TABLE 1-2)

1. **G₀ (Gap) Phase.** The G₀ phase is the resting phase of the cell where the cell cycle is suspended.

试读结束，需要全本PDF请购买 www.ertongbook.com