

Cerebral Angiomas

Advances in Diagnosis and Therapy

Edited by

H.W. Pia · J.R.W. Gleave · E. Grote · J. Zierski



Springer-Verlag Berlin Heidelberg New York

Cerebral Angiomas

Advances in Diagnosis and Therapy

Edited by

H. W. Pia J. R. W. Gleave E. Grote J. Zierski

With 161 Figures

Springer-Verlag

Berlin · Heidelberg · New York 1975

Symposium held in Gießen January 10–12, 1974 on the occasion of 20 years in Gießen

ISBN 3-540-07073-7 Springer-Verlag Berlin Heidelberg New York
ISBN 0-387-07073-7 Springer-Verlag New York Heidelberg Berlin

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically those of translations, reprinting, re-use of illustrations, broadcasting, reproduction by photocopying machine or similar means, and storage in data banks.

Under § 54 of the German Copyright Law where copies are made for other than private use, a fee is payable to the publisher, the amount of the fee to be determined by agreement with the publisher.

© by Springer-Verlag Berlin · Heidelberg 1975. Printed in Germany.

Library of Congress Cataloging in Publication Data. Main entry under title: Cerebral angiomas.

Papers presented at a meeting held Jan. 11–12, 1974 at Giessen. Bibliography: p. Includes index.

1. Brain--Tumors--Congresses. 2. Angioma--Congresses. I. Pia, Hans Werner, ed. [DNLM:

[DNLM: 1. Brain neoplasms--Congresses. 2. Hemangioma--Congresses. WL358' I58' 1974]
RC280.B7I56 616.9'92'81 74-28227

The use of general descriptive names, trade marks, etc. in this publication, even if the former are not especially identified, is not to be taken as a sign that such names, as understood by the Trade Marks and Merchandise Marks Act, may accordingly be used freely by anyone.

Printing and bookbinding: Offsetdruckerei Julius Beltz, KG, Hemsbach



Preface

The basic principles of the management of cerebral arteriovenous malformations were established during the first phase of the neurological attack on these problems between 1930 and 1960. The leaders were CUSHING, BAILEY and DANDY, but principally OLIVECRONA, and in Germany TÖNNIS. The experience gained showed that complete excision of the arteriovenous angioma was the only certain cure, and therefore was the procedure of choice.

In the present second phase important advances should be made and indeed are occurring. New diagnostic techniques such as total angiography, selective and superselective angiography, intraoperative and fluorescein angiography, and the EMI-scanner have been developed. The pathophysiological aspects have been further investigated by indirect and direct measurement of local and general cerebral blood flow.

Parallel with these developments operative technique itself has been improved and modified by new methods. A more aggressive attitude has been stimulated towards those angiomas, which had to be regarded as inoperable only a few years ago. Among these many improvements and technical advances include microsurgical techniques, combined stereotactic and microsurgical procedures, artificial embolization of different kinds and the cryosurgical management.

Multiple variables such as the age of the patient, the type, localization, and size of the angioma, its clinical picture and the possible complications, such as hemorrhage have been analysed and are understood better. These factors influence the indication for, and choice of, the appropriate procedure to a great extent.

The 20th anniversary of my work in Giessen, after 20 years of neurosurgery at Giessen University, has given the occasion for the discussion of the present position with regard to assessment and treatment of inoperable angiomas in a workshop together with experts in this field. The workshop, which was held Jan. 11.–12. 1974, was possible thanks to the generosity of Mr. and Mrs. H. KRÖNER of Fresenius Company, Bad Homburg. This volume contains the papers and discussions of this meeting. The interest, which it aroused, led to rapid publication. The editors take great pleasure in expressing their thanks to the contributors for their participation and cooperation, and to Springer-Verlag for personal and technical aid in preparing and publishing the proceedings.

ipation and cooperation, and to Springer-Verlag for personal and technical aid in preparing and publishing the proceedings.

We hope that they may contribute to giving a more precise answer to the problem of how to deal with patients who suffer from the so-called inoperable cerebral angiomas.

Giessen, November 1974

HANS WERNER PIA

List of Editors and Contributors

AGNOLI, A. L.	Gießen, Zentrum für Neurochirurgie der Universität Gießen
BAUER, B. L.	Gießen, Zentrum für Neurochirurgie der Universität Gießen
BECK, B.	Bad Nauheim, W. G. Kerckhoff-Herzforschungsinstitut der Max-Planck-Gesellschaft
BEKS, J. W. F.	Groningen/Niederlande, Kliniek voor Neurochirurg., Academisch Ziekenhuis
BERGLEITER, R.	Stuttgart, Neuroradiologische Abteilung des Katharinenhospitals der Stadt Stuttgart
BRAUN, W.	Wuppertal-Elberfeld, Neurochirurgische Abteilung des Krankenhauses Bethesda
BURNEY, R. E.	APO New York 09220, USAF Hospital Box 582
BUSHE, K.-A.	Göttingen, Neurochirurgische Universitätsklinik
DJINDJIAN, R.	Paris/France, Département de Neuroradiologie Hôpital Lariboisière
ERBSLÖH, F.	Gießen, Zentrum für Neurologie der Universität Gießen
FEINDEL, W.	Montreal/Canada, Department of Neurology and Neurosurgery, McGill University and Montreal Neurological Hospital
GLEAVE, J. R. W.	Cambridge/Great Britain, Neurosurgical Department, Addenbrooke's Hospital
GROTE, E.	Gießen, Zentrum für Neurochirurgie am Klinikum der Justus-Liebig-Universität Gießen
HEKSTER, R. E. M.	Leiden/Niederlande, Neuroradiologische Abteilung, Academisch Ziekenhuis
HEMMER, R.	Freiburg i. Brsg., Neurochirurgische Universitätsklinik
JELLINGER, K.	Wien/Austria, Neurologisches Institut der Universität Wien
JOHNSON, R.	Manchester/Great Britain, Department of Neurological Surgery, Royal Infirmary

KOHLMEYER, K.	Gießen, Zentrum für Neurologie der Universität Gießen
KUHLEND AHL, H. KUNC, Z.	Düsseldorf, Neurochirurgische Universitätsklinik Prague/Czechoslovakia, Department of Neurosurgery, Charles University
LAPRAS, C. LOEW, F. LUESSENHOP, A. J.	Lyon/France, Hôpital Neurologique Homburg, Neurochirurgische Universitätsklinik Washington/U.S.A., Division of Neurological Surgery, Georgetown University Hospital
NADJMI, M.	Würzburg, Neuroradiologische Abteilung der Neurolog. u. Neurochirurg. Universitätskliniken
PAMPUS, F.	Stuttgart, Neurochirurgische Klinik des Katharinenhospitals der Stadt Stuttgart
PEETERS, F.	Nijmegen/Niederlande, Neuroradiologie – Neurochirurgische Abteilung des St. Radbond-Ziekenhuis und St. Canisius-Ziekenhuis der Universität Nijmegen
PENNING, L.	Groningen/Niederlande, Department of Neuroradiology, University Hospital
PENZHOLZ, H. PERRET, G.	Heidelberg, Neurochirurgische Universitätsklinik Iowa City/U.S.A., Division of Neurosurgery, University of Iowa Hospitals and Clinics
PIA, H. W.	Gießen, Zentrum für Neurochirurgie der Universität Gießen
RIECHERT, T.	Freiburg i. Brsg., Neurochirurgische Universitätsklinik
SANO, K.	Tokyo/Japan, Department of Neurosurgery, Univ. of Tokyo
SCHEPELMANN, F.	Gießen, Zentrum für Neurochirurgie der Universität Gießen
SCHÜRMANN, K. SEEGER, W.	Mainz, Neurochirurgische Universitätsklinik Gießen, Zentrum für Neurochirurgie der Universität Gießen
TZONOS, T.	Stuttgart. Neurochirurgische Klinik des Katharinenhospitals der Stadt Stuttgart
VOGELSANG, H.	Hannover, Abteilung für Neuroradiologie, Institut für klinische Radiologie der Medizinischen Hochschule Hannover
WALDER, H. A. D.	Nijmegen/Niederlande, Neurochirurgische Abteilung des St. Radbond-Ziekenhuis und St. Canisius-Ziekenhuis der Universität Nijmegen
WALTER, W. WÜLLENWEBER, R.	Erlangen, Neurochirurgische Universitätsklinik Bonn, Neurochirurgische Universitätsklinik
ZIERSKI, J.	Gießen, Zentrum für Neurochirurgie der Universität Gießen

Contents

The Indications and Contraindications for Treatment or Assessment	1
H. W. PIA: Introduction	1
 Chapter I. Morphological Aspects	9
K. JELLINGER: The Morphology of Centrally-situated Angiomas	9
Discussion on Chapter I	19
 Chapter II. Clinical Aspects	21
G. PERRET: The Epidemiology and Clinical Course of Arteriovenous Malformations	21
W. WALTER: The Influence of the Type and Localization of the An- gioma on the Clinical Syndrome	27
Discussion on Chapter II	34
 Chapter III. Angiographical Aspects	38
H. VOGELSANG: The Influence of Modern Angiographic Techniques .	38
R. DJINDJIAN: Superselective Arteriography of the Branches of the External Carotid Artery	47
F. PEETERS: Use of Intraoperative Angiography	58
A. L. AGNOLI: Extracranial and Extra-Intracranial Arteriovenous Angiomas	66
Discussion on Chapter III	86
 Chapter IV. Pathophysiological Aspects	87
W. FEINDEL: The Influence of Cerebral Steal: Demonstration by Fluorescein Angiography and Focal Cerebral Blood Flow Meas- urement	87
B. L. BAUER, B. BECK, and U. MÖSSELER: Investigation on Volume and Pressure Overloading of the Heart in Cerebral Angiomas	101
Discussion on Chapter IV	111
	VII

Chapter V. Operative Macro- and Microsurgical Treatment	114
H. W. PIA: Introduction	114
Z. KUNC: Extensive Cerebellar Arteriovenous Malformations	116
K.-A. BUSHE, J. BOCKHORN and E. R. SCHÄFER: Macro- and Micro- surgery of Central Angiomas	123
T. RIECHERT: Stereotactic Treatment of Central Angiomas	129
C. LAPRAS: Angiomas of Cerebellum and Brain Stem	136
F. SCHEPELMANN: Treatment of an Angioma of the Left Cerebral Hemisphere	142
J. ZIERSKI: Treatment of Cranio-Cervical Angioma. Case Report	148
H. W. PIA: The Acute Treatment of Cerebral Arteriovenous Angio- mas Associated with Hematomas	155
Discussion on Chapter V	178
Chapter VI. Cryosurgical Treatment	183
H. A. D. WALDER: Freezing Arteriovenous Anomalies in the Brain	183
Discussion on Chapter VI	194
Chapter VII. Artificial Embolization	198
A. J. LUESSENHOP: Artificial Embolization of Inoperable Arteriove- nous Malformations	198
T. TZONOS, R. BERGLEITER, F. PAMPUS: Experiences in the Use of Artificial Embolization as a Method of Treating Cerebral Angio- mas	206
W. SEEGER: The Artificial Embolization of Inoperable Angiomas	213
K. SANO, M. JIMBO, I. SAITO, and N. BASUGI: Artificial Embolization of Inoperable Angioma with Polymerizing Substance	222
R. DJINDJIAN: Extra-Cerebral Embolization	230
R. E. M. HEKSTER: Treatment of Inoperable Glomus Jugulare Tumors	244
Discussion on Chapter VII	249
Chapter VIII. Radiotherapy	256
R. T. JOHNSON: Radiotherapy of Cerebral Angiomas. With a note on some problems in Diagnosis	256
Discussion on Chapter VIII	267
Chapter IX. The Natural History	268
G. E. PERRET: Conservative Management of Inoperable Arteriove- nous Malformations	268
W. WALTER: The Conservative Treatment of Cerebral Arterio- venous Angiomas	271
Summary and Conclusions	279
Subject Index	283

The Indications and Contraindications for Treatment or Assessment

Introduction

H. W. PIA

It is believed that Fedor KRAUSE was one of the first neurosurgeons to operate upon cerebral angiomas when in 1908 in Berlin he tried to ligate the feeding vessels. However, the era of operative treatment did not start until the thirties during which period CUSHING and BAI-LEY reported 9 cases, DANDY 8 cases, and OLIVECRONA, TÖNNIS, and BERG-STRAND published their classical monograph in 1936. They established the principles of diagnosis and treatment: diagnosis based on the angiographic evaluation of the structure of the malformation, the number, localization, and function of the feeding arterial vessels; and treatment based on the complete extirpation of the malformation where possible. Out of 20 intracranial arterio-venous angiomas 3 were totally excised, the first by OLIVECRONA in 1932. In 2 cases the attempt had to be abandoned, 4 cases were only explored, 3 patients had the intracranial arterial feeding vessels ligated; in 4 patients ligation of the internal carotid artery was performed, and 3 patients were treated by X-ray therapy.

Total excision was limited to small angiomas, in whom the occlusion of the feeding vessels was possible before the draining veins were approached. Angiomas situated in the motor area, the Sylvian fissure, and the speech areas were considered inoperable. "The risk of rendering a patient with seizures and perhaps only slight disturbance of function hemiplegic and aphasic is not abolished by the technique of vessel ligation which is of doubtful value". The intracranial ligation of vessels was successful in only one case with a single feeding artery.

This pioneering work of OLIVECRONA and TÖNNIS is particularly impressive even on review of their results 20 years later. OLIVECRONA and LADENHEIM reported in 1957 a series of 125 angiomas, out of which 81 were totally excised. 7 patients died, 50 were cured, 15 improved, and 7 became worse. TÖNNIS reported a series of 134 angiomas, out of which 56 were totally excised. 5 patients died. 61% of the survivors had full working capacity, 22% were partially employed, and 11% were invalids. Other large series from KRAYENBÜHL (15a) with 72 cases, MCKISSOCK and HANKINSON (21) with 106 cases, NORLEN (23) with 57 cases, KRENCHER (16) with 98 cases, and FRENCH, CHOU and STORY (12) with 82 cases confirmed such results.

OLIVECRONA summarized his experiences in 1957 with the following words: "As matters stand today, if a patient with a lesion, which meets the standards of operability undergoes extirpative surgery under hypotensive anesthesia, the likelihood of surgical mortality should be negligible. About 85% of these patients will be clinically improved with

complete excision of the angiomatous tissue and freedom from future hemorrhage. Few of the remaining 15% will suffer any detriment.

1. The lesion may be totally removed, but the preexisting symptoms will be somewhat aggravated. This is a small premium to pay for insurance against further vascular accidents.
2. A small residue of angiomatous tissue may remain, but the symptoms will be improved.
3. Residual tissue may persist with aggravation of the preoperative symptoms. This last group may require further surgical intervention at a later date".

In spite of the overwhelming evidence of the effectiveness of extirpation of operable angiomas even in such cases many questions remain unsolved. This concerns particularly the prognosis of such important symptoms as epilepsy and hemorrhage which of course has a bearing on the question of the indications for surgery. Epileptic seizures were abolished in approximately 50% of patients (MCKISSOCK, TÖNNIS, NORLEN). In a few cases extirpative surgery led to epilepsy. Elimination of the danger of hemorrhage by total extirpation must be considered in relation to its prognosis which in spite of a recurrent hemorrhage rate between 40 and 60% is often quite good.

Palliative surgical procedures proved to be unsatisfactory. Out of 44 cases treated by OLIVECRONA 14 died, 2 following surgery, and 12 probably from later intracranial hemorrhage. 30 patients survived, 14 in good condition, but 9 were partial and 7 were complete invalids.

Afferent ligation, as performed by DANDY, should be considered only in cases where one single artery feeds the malformation, but even here the temptation should be resisted as arteriography is not infallible, and contributing vessels may not be demonstrated.

The electrocoagulation of surface vessels (TRUPP and SACHS) gave unsatisfactory results.

Carotid ligation should be given up. Not only was there no effect on the angioma, but the procedure was dangerous.

After ligation the blood flow through the angioma was derived from collateral sources, thus increasing, as we would say now, the steal effect. Radiation, as employed by CUSHING, was without the slightest discernible benefit. OLIVECRONA took a strong stand against the revival of this ineffectual form of therapy for arterio-venous malformations.

The indications for operation followed the rules of tumor surgery. Angiomas located in the functionally less important areas were considered to be surgically approachable. Small angiomas in the motor cortex were also removed; but in larger lesions the postoperative deficit constituted a greater handicap than the preoperative occasional seizure. Angiomas in the Sylvian fissure were often removed.

Intracerebral hemorrhage per se did not contraindicate prompt surgery when the condition of the patient might otherwise be endangered by delay, for the removal of the clot often exposes a partially mobilized lesion.

In angiomas in the region of the basal ganglia and internal capsule, in large malformations of the dominant hemisphere, and in most angio-

mas of the cerebellum and cerebello-pontine angle, surgery was contraindicated. The same was true in cases of epilepsy of long standing, as there was little to gain from operation.

"In the absence of an absolute rule to guide his decision, the surgeon must carefully plot each borderline case against the abscissa of danger and the ordinate of benefit".

The last decade has brought important advances. The clinical symptomatology and course, and last but not least the natural history of unoperated cases has been studied and understood better, especially in the cooperative study by PERRET and NISHIOKA (26) and by WALTER (1970). Further progress was made by the introduction of preoperative total angiography and peroperative angiography, superselective angiography (DJINDJIAN), and magnification angiography. These investigations improved the diagnosis of small malformations and our knowledge of the pathomorphology of angiomas.

The pathophysiological aspects were advanced among others by SCHIEFER and TÖNNIS (39) and FEINDEL (9, 10) who elaborated upon the general and local cerebral blood flow disturbances. Current investigations by BECK and BAUER on the influence of the cerebral shunt on the heart and circulation may contribute to the better understanding of local and general compensation and decompensation mechanisms.

Parallel to these developments operative technique itself has been improved and this has stimulated a more aggressive attitude towards angiomas in functionally important areas. KUNC (18) reported on 97 angiomas out of which 58 were operated on; 7 were located in the speech area, 19 in the sensori-motor cortex, 5 in the Sylvian fissure, and 27 close to the above regions. Radical removal was possible in 48 cases with 8 deaths. On the other hand SANO, AIBA and JIMBO (36) stressed the high morbidity. In their series of 108 patients, 81 were operated upon. There was only one postoperative death, but 81% of patients had definite though variable residual neurological deficits.

The indications for radical removal of medial and paramedial angiomas are far from well defined and considerable reluctance is apparent. An important contribution was made by RIECHERT and MUNDINGER (35) who introduced a combined stereotactic and conventional procedure. The advantage of their method is the precise localization of the angioma with the possibility of an approach which causes the least functional disability. RIECHERT reported in 1972 16 cases of deep lying malformations on which successful operations have been carried out.

The use of the operating microscope was an undubitable and important step forward in the treatment of angiomas as was shown by RIECHERT and many others at the microsurgical symposia in Zürich (1968), Vienna (1972), and Kyoto (1973).

BUSHE (3) in a series of 42 cases treated 6 angiomas located in the midline, 7 in the ventricular and paraventricular region, and 2 in the basal ganglia. Radical removal was performed in 10 cases, vessel ligation in 2, and no operation in 3 cases. 2 patients died. DRAKE reported recently on 6 angiomas in the midbrain and pons, all successfully removed. The review of WALTER and BISCHOF (45) comprising 72 cases with angiomas located in those regions revealed that there were only 16 patients in whom the angioma was completely extirpated and 3 of them died. 16 further patients had partial ligation and out of this number 6 died.

Angiomas of the cerebellum were reported less frequently in earlier series. Latterly VERBIEST (42), PIA (29), LAPRAS (1972) and YASARGIL (46) and others have shown that even huge angiomas of this area have become accessible due to improved technique especially the use of the microscope.

Huge angiomas of the external and internal carotid system are regarded as inoperable in most cases.

My own contributions to the emergency treatment of angiomas associated with intracerebral and intraventricular hematomas (PIA 1968, 1972) have shown that this can be a life-saving procedure with considerable benefit also on morbidity. The operation risk is minimal, a contraindication exceptional.

It is to LUESSENHOP that the credit is due for introducing artificial embolization in 1960 for the treatment of inoperable angiomas. He also showed the effectiveness of such treatment. Incomplete embolization of the angioma may facilitate or permit a total extirpation at a second stage. SANO in 1965, and in Germany TZONOS, BERGLEITER and PAMPUS, as well as SEEGER, have applied and modified the method of LUESSENHOP. A further step forward was made by DJINDJIAN in 1970 who combined super-selective angiography with embolization in a one stage procedure and reported excellent results in arteriovenous malformations of the skull and soft tissues of the face as well as in spinal angiomas. A glomus tumor was successfully treated by embolization by HEKSTER, LUYENDIJK and MATRICALI (13).

The freezing technique was developed by WALDER (1970) in order to thrombose the pathological vessels of inoperable and deep seated angiomas. Cryosurgery may also be used as a complementary procedure after incomplete extirpation of a deep lying angioma.

X-ray therapy of arteriovenous angiomas is almost generally regarded as ineffective. However, in 1955 POTTER reported on 10 patients treated by X-ray therapy out of which one died and 9 survived, 4 of them for more than 20 years.

The similar experience of JOHNSON (14) suggests that X-ray therapy may exert a favourable effect at least upon some forms of arteriovenous malformations, so that we may have to reconsider the indications for X-ray therapy in selected inoperable cases.

The basic principles of management of cerebral arteriovenous malformations were laid down during the first phase of therapy between 1932 and the mid-sixties. Experience showed that complete excision of the angioma was the only method of cure, and therefore was the procedure of choice.

In the present second phase not only have new techniques been developed but also multiple variables such as the age of the patient, the type, location, and size of the angioma; its influence on the cerebral and general circulation, the clinical picture and possible complications, such as hemorrhage and deficits produced by surgery, were analysed and understood better. These factors influence the indication for and choice of the appropriate procedure to a great extent.

The subject of this workshop is the treatment of inoperable angiomas. I think that starting from this borderline position, we should try to define the principles of management of patients with such lesions. I hope also that we will try to define objective criteria for the choice

of treatment of different types of angiomas in general. The participation of so many distinguished experts from different specialties makes me confident that we will come closer to our aim of answering more precisely the question of how to deal with those patients who have so-called "inoperable angiomas".

R e f e r e n c e s

1. BERGSTRAND, H., OLIVECRONA, H., TÖNNIS, W.: Gefäßmißbildungen und Gefäßgeschwülste des Gehirns. pp. 181. Leipzig: Georg Thieme 1936.
2. BROOKS, B.: The treatment of traumatic arterio-venous fistula. Sth. med. J. (Birmingham) 23, 100-106 (1930).
3. BUSHE, K. A., PETERSON, E., SCHÄFER, E. R.: Surgical Indication for Arteriovenous Angiomas in Functionally Important Regions and in Case of Spreading within the Area of the Ventricular System and of the Basal Ganglia. In Present Limits of Neurosurgery, Avicenum, Czechoslov. Med. Press, 299-302. Prague 1972.
4. CUSHING, H., BAILEY, P.: Tumors arising from the blood vessels of the brain. Baltimore: C. C. Thomas 1928.
5. DANDY, W. E.: Arteriovenous aneurysms of the brain. Arch. Surg. 17, 190-243 (1928).
6. DJINDJIAN, R., COPHIGNON, J., THERON, J., MERLAND, J. J., HOUDART, R.: L'embolisation en neuroradiologie vasculaire. Technique et indications à propos de 30 cas. Nouv. Presse méd. 33, 2153-2158 (1972).
7. DJINDJIAN, R., HOUDART, R., COPHIGNON, J.: Premiers essais d'embolisation par voie fémorale de fragments de muscle dans un cas d'angiome médullaire et dans un cas d'angiome alimenté par la carotide externe. Rev. Neurol. 125, 119-130 (1971).
8. DJINDJIAN, R., MERLAND, J. J., REY, A., THUREL, J., HOUDART, R.: Artériographie super-sélective de la carotide externe. Neuro-Chirurgie Paris 19, 165-171 (1973).
9. FEINDEL, W., GARRETSON, H., YAMAMOTO, Y. L., PEROT, P., RUMIN, N.: Blood flow patterns in the cerebral vessels and cortex in man studied by intracarotid injection of radioisotopes and Coomassie Blue dye. J. Neurosurg. 23, 12-22 (1965).
10. FEINDEL, W., HODGE, Ch. P., YAMAMOTO, Y. L.: Epicerebral angiography by Fluorescein during craniotomy. In Progr. Brain Research 30, 471-477. Amsterdam-London-New York: Elsevier 1968.
11. FRENCH, L. A., CHOU, S. N.: Conventional Methods of Treating Intracranial Arteriovenous Malformations. Progr. Neurosurg. 3, 275-361. Basel-New York: S. Karger 1969.
12. FRENCH, L. A., CHOU, S. N., STORY, J. L.: Cerebrovascular malformations. Clinical Neurosurg., Vol. XI, Chapter 14. Baltimore: Williams and Wilkins Co. 1964.
13. HEKSTER, R. E. M., LEUYENDIJK, W., MATRICALI, B.: Transfemoral Catheter Embolization: A Method of Treatment of Glomus Jugulare Tumors. Neuroradiology 5, 208-214 (1973).

14. JOHNSON, R. T.: Surgery of Cerebral Hemorrhage. In Recent Advances in Neurology, 124-126. (Eds. Lord Brain and Maria Wilkinson). London: Churchill 1969.
15. KRAUSE, F.: Krankenvorstellung aus der Hirnchirurgie (Fall 4. Ang. venosum racemosum). Zbl. Chir. 35, 61-63 (1908).
- 15a. KRAYENBÜHL, H.: Discussion des rapports sur les angiomas supratentoriels. 1. Congr. Internat. de Neurochirurgie, 263-267. Bruxelles 1957.
16. KRENCHEL, N. J.: Intracranial racemose angiomas, a clinical study. Universitetsforlaget I. Aarhus (1961).
17. KUNC, Z.: Operability of Arteriovenous Malformations in Functionally Important Cortical Regions. In Present Limits of Neurosurgery, Avicenum, Czechoslov. Med. Press, 293-298. Prague 1972.
18. KUNC, Z.: Possibility of surgery in arteriovenous malformations in the anatomically important and dangerous regions of the brain. J. Neurol. Neurosurg. Psychiat. 28, 183 (1965).
- 18a. LAPRAS, C., TUSINI, G., THIERRY, A.: A propos de 8 observations cliniques d'angiomes de la fosse cérébrale postérieure. Congrès Soc. Neurochirurgie langue Franc. Lyon 1963.
19. LUESSENHOP, A.: Artificial Embolization for Cerebral Arteriovenous Malformations. Progr. neurol. Surg. 3, 320-362. Basel-New York: Karger 1969.
20. LUESSENHOP, A. J., SPENCE, W. T.: Artificial embolization of cerebral arteries. Report of use in a case of arteriovenous malformation. J. amer. med. Ass. 172, 1153-1155 (1960).
21. MCKISSOCK, W., HANKINSON, J.: The surgical treatment of the supratentorial angiomas. Rapports et discussions. 1er Congrès International de Neurochirurgie, Bruxelles 1957.
22. NORLÉN, G.: Arteriovenous aneurysms of the brain: Report of ten cases of total removal of the lesion. J. Neurosurg. 6, 475-494 (1949).
23. NORLÉN, G.: Die chirurgische Behandlung intracranieller Gefäßmißbildungen. A. Angiome. Handbuch der Neurochirurgie. Hrsg. H. Olivecrona und W. Tönnis, Bd. 4/II, S. 147-206. Berlin-Heidelberg-New York: Springer 1966.
24. OLIVECRONA, H., LADENHEIM, J.: Congenital arteriovenous aneurysms of the carotid and vertebral arterial systems. Berlin: Springer 1957.
25. OLIVECRONA, H., RIIVES, J.: Arteriovenous aneurysms of the brain: their diagnosis and treatment. Arch. Neurol. Psychiat. 59, 567-602 (1948).
26. PERRET, G., NISHIOKA, H.: Arteriovenous malformations. An analysis of 545 cases of cranio-cerebral arteriovenous malformations and fistulae reported to the cooperative study. J. Neurosurg. 25, 467-490 (1966).
27. PIA, H. W.: The Diagnosis and Treatment of Intraventricular haemorrhages. Progr. in Brain Research 30, 463-470 (1968).

28. PIA, H. W.: Diagnose und Therapie der Hirnblutungen im Kindesalter. *Tijdschr. Geneesk.* 24, 765-770 (1968) und *Beiträge Neurochir.* 15, 245-251 (1969).
29. PIA, H. W.: Vasculäre Mißbildungen und Erkrankungen der hinteren Schädelgrube und ihre Behandlung. *HNO* 17, 165-171 (1969).
30. PIA, H. W.: The Operative Treatment of Intracerebral and Intraventricular Hematomas. *Acta Neurochir.* 27, 149-164 (1972).
31. PIA, H. W.: The Surgical Treatment of Intracerebral Hematomas. In *Present Limits of Neurosurgery*, Avicenum, Czechoslov. Med. Press, 323-328. Prague 1972.
32. POOL, J. L.: Treatment of arteriovenous malformations of the cerebral hemisphere. *J. Neurosurg.* 19, 136-141 (1962).
33. POTTER, J. M.: Angiomatous malformations of the brain: their nature and prognosis. *Ann. roy. Coll. Surg. (Engl.)* 16, 227-243 (1955).
34. RIECHERT, T.: Die operative Versorgung von Blutungen im Bereich der Stammganglien. *Therapiewoche* 22, 1152-1153 (1972).
35. RIECHERT, T., MUNDINGER, F.: Combined stereotaxic operation for treatment of deep seated angiomas and aneurysms. *J. Neurosurg.* 21, 358-363 (1964).
36. SANO, K., AIBA, T., JIMBO, M.: Surgical treatment of cerebral aneurysms and arteriovenous malformations. *Neurologia med.-chir.* 7, 128-131 (1965).
37. TÖNNIS, W.: Symptomatologie und Klinik der supratentoriellen arteriovenösen Angiome. 1. Congr. Int. de Neurochirurgie. Les Editions Acta Med., 205-215. Belgica 1957.
38. TÖNNIS, W., SCHIEFER, W., WALTER, W.: Signs and symptoms of supratentorial arteriovenous aneurysms. *J. Neurosurg.* 15, 471-480 (1958).
39. TÖNNIS, W., SCHIEFER, W.: Zirkulationsstörungen des Gehirns im Serienangiogramm. Berlin-Göttingen-Heidelberg: Springer 1959.
40. TÖNNIS, W., WALTER, W.: Die Indikation zur Totalexstirpation der intracraniellen arteriovenösen Angiome. *Dtsch. Z. Nervenheilk.* 186, 279-298 (1964).
41. TRUPP, M., SACHS, E.: Vascular tumors of the brain and spinal cord and their treatment. *J. Neurosurg.* 5, 354-371 (1948).
42. VERBIEST, H.: Arteriovenous aneurysms of the posterior fossa. In *Progr. Brain Research* 30, 383-396. Amsterdam-London-New York: Elsevier 1968.
43. WALDER, HAD.: Die Behandlung arteriovenöser Anomalien im Gehirn mit Hilfe von Kryotherapie. *Therapiewoche* 22, 1542-1543 (1972).
44. WALDER, HAD.: Freezing Arteriovenous Anomalies in Brain. In *Avicenum, Czechoslov. Med. Press*, 303-308. Prague 1972.