

MATHEMATICAL NOTES

Complex Ball Quotients
and Line Arrangements in
the Projective Plane

Paula Tretkoff


PRINCETON UNIVERSITY PRESS

Complex Ball Quotients and Line Arrangements in the Projective Plane

Paula Tretkoff

With an appendix by Hans-Christoph Im Hof

Mathematical Notes 51



PRINCETON UNIVERSITY PRESS
PRINCETON AND OXFORD

Copyright © 2016 by Princeton University Press
Published by Princeton University Press,
41 William Street, Princeton, New Jersey 08540
In the United Kingdom: Princeton University Press,
6 Oxford Street, Woodstock, Oxfordshire, OX20 1TW

All Rights Reserved

Library of Congress Cataloging-in-Publication Data

Tretkoff, Paula, 1957–

Complex ball quotients and line arrangements in the projective plane /
Paula Tretkoff.

pages cm. – (Mathematical notes ; 51)

Includes bibliographical references and index.

ISBN 978-0-691-14477-1 (pbk. : alk. paper) 1. Curves, Elliptic. 2. Geometry,
Algebraic. 3. Projective planes. 4. Unit ball. 5. Riemann surfaces. I. Title.

QA567.2.E44T74 2016

516.3'52–dc23

2015016120

British Library Cataloging-in-Publication Data is available

This book has been composed in Minion Pro

Printed on acid-free paper. ∞

press.princeton.edu

Typeset by S R Nova Pvt Ltd, Bangalore, India

Printed in the United States of America

1 3 5 7 9 10 8 6 4 2

Complex Ball Quotients and Line
Arrangements in the Projective Plane

To the memory of Friedrich Hirzebruch

Preface

This book is devoted to a study of quotients of the complex 2-ball yielding finite coverings of the projective plane branched along certain line arrangements. It is intended to be an introduction for graduate students and for researchers. We give a complete list of the known weighted line arrangements that can give rise to such ball quotients, and then we provide a justification for the existence of the ball quotients. The Miyaoka-Yau inequality for surfaces of general type, and its analogue for surfaces with an orbifold structure, plays a central role.

The book has its origins in a Nachdiplom course given by F. Hirzebruch at the ETH Zürich during the Spring of 1996. I (née Cohen) was at that time Directeur de Recherche au CNRS at the Université de Lille 1, and a guest of ETH Zürich. I attended the course, lectured on some of the material on hypergeometric functions, and made the original set of notes for all Hirzebruch's lectures. I also presented related material as a two-semester graduate course at Princeton University during the academic year 2001/2002 while I was Visiting Professor there. The ETH Nachdiplom course notes were subsequently developed and refined by F. Hirzebruch and me during regular visits to the Max Planck Institute in Bonn. I thank ETH Zürich, MPI Bonn, and Princeton University for their support during the preparation of this book.

After working on the book together for some years, F. Hirzebruch and I decided even more additional material seemed desirable and, at that point, F. Hirzebruch asked me to complete the book under my own name. The book entitled *Geradenkonfigurationen und Algebraische Flächen*, Vieweg, 1987, by G. Barthel, F. Hirzebruch, and T. Höfer, served as a valuable resource and reference. This book is in German, and we have mined some of its contents as needed. The present book assumes less background than the earlier book, and we revisit in more detail several of its important subjects. We have also added material not found in that book. For example, we present a more organized list of the possible weights on line arrangements that yield finite covers that are ball quotients, and we devote a whole chapter, rather than just a few pages, to the question of the existence of such ball quotients. Also, we include material on hypergeometric functions.

The book is dedicated to the memory of F. Hirzebruch. I am grateful beyond words for the time and attention he paid to this project amid the many commitments of his important and busy life.

Complex Ball Quotients and Line
Arrangements in the Projective Plane

Contents

Preface	ix
Introduction	1
1 Topological Invariants and Differential Geometry	6
1.1 Topological Invariants	7
1.2 Fundamental Groups and Covering Spaces	10
1.3 Complex Manifolds and Metrics	13
1.4 Divisors, Line Bundles, the First Chern Class	16
2 Riemann Surfaces, Coverings, and Hypergeometric Functions	23
2.1 Genus and Euler Number	23
2.2 Möbius Transformations	25
2.3 Metric and Curvature	29
2.4 Behavior of the Euler Number under Finite Covering	33
2.5 Finite Subgroups of $\mathrm{PSL}(2, \mathbb{C})$	34
2.6 Gauss Hypergeometric Functions	36
2.7 Triangle Groups	41
2.8 The Hypergeometric Monodromy Group	45
3 Complex Surfaces and Coverings	47
3.1 Coverings Branched over Subvarieties with Transverse Intersections	47
3.2 Divisor Class Group and Canonical Class	49
3.3 Proportionality	54
3.4 Signature	59
3.5 Blowing Up Points	61
4 Algebraic Surfaces and the Miyaoka-Yau Inequality	65
4.1 Rough Classification of Algebraic Surfaces	65
4.2 The Miyaoka-Yau Inequality, I	70
4.3 The Miyaoka-Yau Inequality, II	73
5 Line Arrangements in $\mathbb{P}_2(\mathbb{C})$ and Their Finite Covers	85
5.1 Blowing Up Line Arrangements	87
5.2 Höfer's Formula	88

5.3 Arrangements Annihilating R and Having Equal Ramification along All Lines	92
5.4 Blow-Up of a Singular Intersection Point	99
5.5 Possibilities for the Assigned Weights	103
5.6 Blowing Down Rational Curves and Removing Elliptic Curves	115
5.7 Tables of the Weights Giving $\text{Prop} = 0$	122
6 Existence of Ball Quotients Covering Line Arrangements	126
6.1 Existence of Finite Covers by Ball Quotients of Weighted Configurations: The General Case	128
6.2 Remarks on Orbifolds and b -Spaces	133
6.3 $K_{X''} + D''$ for Weighted Line Arrangements	135
6.4 Existence Question	139
6.5 Ampleness of $K_{X''} + D''$	140
6.6 Log-Terminal Singularities and LCS	145
6.7 Existence Theorem for Line Arrangements	148
6.8 Isotropy Subgroups of the Covering Group	164
7 Appell Hypergeometric Functions	167
7.1 The Action of S_5 on the Blown-Up Projective Plane	168
7.2 Appell Hypergeometric Functions	173
7.3 Arithmetic Monodromy Groups	181
7.4 Some Remarks about the Signature	186
A Torsion-Free Subgroups of Finite Index	189
A.1 Fuchsian Groups	190
A.2 Fenchel's Conjecture	191
A.3 Reduction to Triangle Groups	192
A.4 Triangle Groups	193
B Kummer Coverings	197
Bibliography	205
Index	213

Introduction

Historical precedent for the results in this book can be found in the theory of Riemann surfaces. Every compact Riemann surface of genus $g \geq 2$ has representations both as a plane algebraic curve, and so as a branched covering of the complex projective line, and as a quotient of the complex 1-ball, or unit disk, by a freely acting cocompact discrete subgroup of the automorphisms of the 1-ball. The latter result is a direct consequence of the uniformization theorem, which states that the simply-connected Riemann surfaces are the Riemann sphere, the complex plane, and the complex 1-ball.

In complex dimension 2, Hirzebruch (1956) and Yau (1978) showed that the smooth compact connected complex algebraic surfaces, representable as quotients of the complex 2-ball B_2 by freely acting cocompact discrete subgroups of the automorphisms of the 2-ball, are precisely the surfaces of general type whose Chern numbers satisfy $c_1^2 = 3c_2$. Here, c_2 is the Euler characteristic and c_1^2 is the self-intersection number of the canonical divisor. Accordingly, the *proportionality deviation* of a complex surface is defined by the expression $3c_2 - c_1^2$. This book examines the explicit computation of this proportionality deviation for finite covers of the complex projective plane ramified along certain line arrangements. Candidates for ball quotients among these finite covers arise by choosing weights on the line arrangements such that the proportionality deviation vanishes. We then show that these ball quotients actually exist.

The intention of F. Hirzebruch in the original notes was for the material to be presented in a nontechnical way, assuming a minimum of prerequisites, with definitions and results being introduced only as needed. The desire was for the reader to be exposed to the theory of complex surfaces—and of complex surfaces with an orbifold structure—through the examples provided by weighted line arrangements in the projective plane, with an emphasis on their finite covers which are ball quotients. There was no desire to develop a complete theory of surfaces and orbifolds although relevant references are provided, or to treat the latest developments, or to also survey results in higher dimensions. Instead, the goal was that, on reading the book, the student or researcher should be better equipped to go into more technical or more modern territory if interested. It was felt that the topic of the notes was important enough historically to warrant their appearance. F. Hirzebruch wanted the style of the original notes to be like that of his series of lectures,

and for them to remain readable and conversational; and, even though a lot of material has been added since by me, I have aimed to retain those qualities.

The plan of the book is as follows. In Chapter 1, we collect in one place the main prerequisites from topology and differential geometry needed for subsequent chapters, although, for convenience, pieces of this material are at times repeated or expanded upon later in the book. Even if the reader is familiar with these topics, it is worth his or her while to leaf through Chapter 1 to see what notions and notations are used later.

In Chapter 2, we apply some of the material from Chapter 1 to Riemann surfaces—which are of real dimension 2—by way of a historical and conceptual motivation for the material on complex surfaces. We also discuss in this chapter the classical Gauss hypergeometric functions of one complex variable and the triangle groups that serve as their monodromy groups. When these triangle groups are hyperbolic, they act on the unit disk, that is, the complex 1-ball. Infinitely many such groups act discontinuously, and their torsion-free normal subgroups of finite index define quotients of the complex 1-ball that are finite coverings of the projective line branched over three points.

In Chapter 3, we study complex surfaces and their coverings branched along divisors, that is, subvarieties of codimension 1. In particular, we discuss coverings branched over transversally intersecting divisors. Applying this to line arrangements in the complex projective plane, we first blow up the projective plane at non-transverse intersection points, that is, at those points of the arrangement where more than two lines intersect. These points are called singular points of the arrangement. This results in a complex surface and transversely intersecting divisors that contain the proper transforms of the original lines. Next, we introduce the group of divisor classes, their intersection numbers, and the canonical divisor class. The Chern numbers c_1^2 and c_2 , as well as the proportionality deviation, $\text{Prop} := 3c_2 - c_1^2$, are then defined.

In Chapter 4, we give an overview of the rough classification of (smooth complex connected compact algebraic) surfaces. We present two approaches that, in dimension 2, give the Miyaoka-Yau inequality; $c_1^2 \leq 3c_2$, for surfaces of general type. We only make a few remarks about the first of these, due to Miyaoka, which uses algebraic geometry. The second, due to Aubin and Yau, uses analysis and differential geometry. Here we give more details, since we use the analogous approach to the Miyaoka-Yau inequality for surface orbifolds in Chapter 6. We also discuss why equality in the Miyaoka-Yau inequality characterizes surfaces of general type that are free quotients of the complex 2-ball B_2 , that is, orbit spaces for the action of discrete cocompact subgroups of the automorphisms of B_2 that have no fixed points.

In Chapter 5 we arrive at the main topic of the book: the free 2-ball quotients arising as finite covers of the projective plane branched along line arrangements. The material of this chapter is self-contained and largely combinatorial. Let X be the surface obtained by blowing up the singular

intersection points of a line arrangement in the complex projective plane. Let Y be a smooth compact complex surface given by a finite cover of X branched along the divisors on X defined by the lines of the arrangement, that is, by their proper transforms and by the blown-up points, also known as exceptional divisors. If Y is of general type with vanishing proportionality deviation Prop, then it is a free 2-ball quotient. In order to find such a Y , we use a formula for Prop due to T. Höfer. This formula expresses Prop as the sum of, first, a quadratic form evaluated at numbers related only to the ramification indices along the proper transforms of the lines, and, second, nonnegative contributions from each blown-up point. The quadratic form itself depends only on the original line arrangement. The contributions from the blown-up points vanish when we impose certain diophantine conditions on the choice of ramification indices. Next, we seek line arrangements and ramification indices that make the quadratic form vanish. Initially, we ask of the arrangements that they have equal ramification indices along each of the proper transforms of the original lines. For these arrangements, the number of intersection points on each line is $(k+3)/3$, where k is the number of lines. We then list all known line arrangements with this property and restrict our attention to them. Next, we replace the ramification indices by weights that are positive or negative integers or infinity, and we list all possible weights that satisfy the diophantine conditions at the blown-up points and also annihilate the quadratic form. Finally, we enumerate all possibilities for the assigned weights of the arrangements, under the assumption that divisors of negative or infinite weight on the blown-up line arrangements do not intersect. When the weight of a divisor on a blown-up line arrangement is negative, the curves above it on Y are exceptional and can be blown down. Blowing down all curves on Y arising from such negative weights, we obtain a smooth surface Y' . When the weight of a divisor on a blown-up line arrangement is infinite, the curves above it on Y are elliptic. Letting C be the union on Y' of all elliptic curves arising in this way, we derive the appropriate expression for the Prop of the possibly non-compact smooth surface $Y' \setminus C$. It is known that we cannot have rational or elliptic curves on a free 2-ball quotient, so this construction is quite natural. For the line arrangements of Chapter 5, we give the complete list of weights such that the Prop of $Y' \setminus C$ vanishes, meaning that $Y' \setminus C$ is a free 2-ball quotient. These weights are all presented in a series of tables in §5.7 at the end of Chapter 5, except for a few extra cases that are listed in §5.6.1. Throughout this chapter, we assume that the finite covers of the line arrangements with vanishing proportionality actually exist. The expression for Prop of $Y' \setminus C$ depends only on the original weighted line arrangement, so we can work with this assumption; but we still need to show that there are such covers.

In Chapter 6 we justify the existence assumption of Chapter 5. Let X be the blow-up of the projective plane at the non-transverse intersection points of a line arrangement. Assign weights—allowed to be positive or negative

integers or infinity, denoted by n_i for the proper transforms D_i of the lines, and denoted by m_j for the exceptional divisors E_j given by the blown-up points—in such a way that the divisors with negative or infinite weights are distinct. Let X' be the possibly singular surface obtained by blowing down the D_i and E_j with negative weight, and X'' the possibly singular surface obtained by contracting the images of the D_i and E_j on X' with infinite weight. Let D_i'' be the image in X'' of D_i with weight $n_i > 0$, and E_j'' the image in X'' of E_j with weight $m_j > 0$. The central question of Chapter 6 is the following. When is X'' a (possibly compactified by points) ball quotient $X'' = \overline{\Gamma \backslash B_2}$ for a discrete subgroup Γ of the automorphisms of the ball with natural map $B_2 \rightarrow X''$ ramified of order n_i along D_i'' and m_j along E_j'' ? Any normal subgroup Γ' of Γ of finite index N in Γ , acting on B_2 without fixed points, gives rise to a free ball quotient $\Gamma' \backslash B_2 = Y' \setminus C$, as described in Chapter 5, that is a finite cover of $X_0'' = \Gamma \backslash B_2$ of degree N , ramified of order $n_i > 0$ along $D_i'' \cap X_0''$ and $m_j > 0$ along $E_j'' \cap X_0''$. Such normal subgroups exist due to the work of Borel and Selberg.

Therefore, we approach the proof of the existence of $Y' \setminus C$ by first showing the existence of Γ . The group Γ acts with fixed points, so, in answering this central question, the language of orbifolds is appropriate. Given the simplicity and explicit nature of the orbifolds we study, we use instead the related notion of b -space due to Kato. For the reader seeking a more sophisticated and general treatment, it is worth learning the basics of complex orbifolds that we exclude. Roughly speaking, we show that if all the diophantine conditions on the weights, derived in Chapter 5 to ensure that Prop vanishes, are satisfied, then there is equality in an orbifold, or b -space, version of the Miyaoka-Yau inequality, and, by arguments analogous to those of Chapter 4, the space X_0'' is of the form $\Gamma \backslash B_2$ and the map $B_2 \rightarrow X_0''$ has the desired ramification properties described above. To do this, we use the work of R. Kobayashi, S. Nakamura, and F. Sakai [83], which generalizes the work of Aubin-Yau on the Miyaoka-Yau inequality to surfaces with an orbifold structure.

There are both related and alternative approaches dating from about the same time, for example, the approach in the independent work of S. Y. Cheng and S. T. Yau [27], as well as more modern and more general approaches, such as in the work of A. Langer [91], [92], to mention just a few important instances. We choose to base our discussion on the work of R. Kobayashi, S. Nakamura, and F. Sakai, as it was part of the original notes made with F. Hirzebruch and it fits well with the presentation of the work of Aubin-Yau in Chapter 4 as well as with our approach to the material of Chapter 5. The main point of Chapter 6 is to flesh out the details of the orbifold version of the Miyaoka-Yau inequality for our particular situation of line arrangements in the projective plane.

Chapter 7 focuses on the complete quadrilateral line arrangement, and especially on its relationship with the space of regular points of the system