

LECTURE NOTES ON
CLINICAL ONCOLOGY

Barry W. Hancock

J. David Bradshaw

SECOND EDITION

43
10.2

05985

R73-43

B812 no.2

LECTURE NOTES ON CLINICAL ONCOLOGY

Barry W. Hancock

MD, FRCP, DCH

Senior Lecturer in Medicine

University of Sheffield

Honorary Consultant Physician

Royal Hallamshire Hospital and

Weston Park Hospital, Sheffield

J. David Bradshaw

MB, ChB, FRCR, DMRT

Consultant Radiotherapist and Oncologist

Weston Park Hospital, Sheffield

Honorary Clinical Lecturer in Radiotherapy

and Oncology, University of Sheffield

SECOND EDITION

馆藏专用章

BLACKWELL SCIENTIFIC PUBLICATIONS

OXFORD LONDON EDINBURGH

BOSTON PALO ALTO MELBOURNE

TO HELEN AND ELSA

© 1981, 1986 by
Blackwell Scientific Publications
Editorial offices:
Osney Mead, Oxford, OX2 0EL
8 John Street, London, WC1N 2ES
23 Ainslie Place, Edinburgh, EH3 6AJ
52 Beacon Street, Boston
Massachusetts 02108, USA
667 Lytton Avenue, Palo Alto
California 94301, USA
107 Barry Street, Carlton
Victoria 3053, Australia

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without the prior permission of the copyright owner

First published 1981
Second Edition 1986

Set by Downdell Ltd, Oxford
Printed by Biddles Ltd, Guildford
and King's Lynn

DISTRIBUTORS

USA

Blackwell Mosby Book Distributors
11830 Westline Industrial Drive
St Louis, Missouri 63141

Canada

The C.V. Mosby Company
5240 Finch Avenue East
Scarborough, Ontario

Australia

Blackwell Scientific Publications
(Australia) Pty Ltd
107 Barry Street
Carlton, Victoria 3053

British Library
Cataloguing in Publication Data

Hancock, Barry W.

Lecture notes on clinical
oncology.—2nd ed.

I. Cancer

I. Title

II. Bradshaw, J. David

616.99'4

RC261

ISBN 0-632-01548-9

Preface to Second Edition

Our thanks to colleagues who have pointed out errors and omissions in the first edition. The second edition has been updated to encompass recent advances in technology and therapy. An extra chapter on oncological emergencies has been included.

Barry W. Hancock

J. David Bradshaw

Preface to First Edition

This new book in the 'Lecture Notes' series is aimed at senior undergraduate students and recently qualified practitioners in all specialities; it is intended as an up-to-date guide to the theory and practice of all aspects of clinical oncology and should provide insight into the diverse nature of the subject, emphasising the multidisciplinary approach necessary for the successful management of the patient with cancer. We hope that our attempt to cover such a large field concisely will not contain too many omissions or dogmatisms. We are grateful to Professor J. Richmond, Dr F. E. Neal and Dr E. M. Pickering for helpful criticism and to the Department of Medical Illustration, Royal Hallamshire Hospital, Sheffield, for their help with many of the figures.

Contents

- Preface to Second Edition, v
Preface to First Edition, vi
- 1 Cancer Overview, 1
 - 2 Investigation and Staging, 23
 - 3 Radiotherapy—Principles and Practice, 33
 - 4 Cytotoxic Chemotherapy—Principles and Practice, 49
 - 5 Other Treatments, 66
 - 6 Cancer of the Lung and Mediastinum, 74
 - 7 Breast Cancer, 81
 - 8 Alimentary Tract Cancer, 89
 - 9 Head and Neck Cancer, 97
 - 10 Cancer of the Genitourinary Tract, 115
 - 11 Haemopoietic and Lymphoreticular Malignancies, 133
 - 12 Childhood Tumours, 152
 - 13 Tumours of the Central Nervous System, 159
 - 14 Tumours of Bone and Soft Tissue, 170
 - 15 Skin Cancer, 178
 - 16 Paraneoplastic (Non-Metastatic) and Hormonal Syndromes Associated with Tumours, 182
 - 17 Surveillance, Palliative Support and Terminal Care, 193
 - 18 Education, Screening and Prevention, 199
 - 19 Oncological Emergencies, 204
- Further Reading, 208
Index, 211

Chapter 1

Cancer Overview

Incidence and epidemiology

Impressions can be very misleading, and this is especially true in respect of the incidence of malignant disease. A worker in general practice might well regard the overall incidence as low; one in a general hospital is likely to regard it as higher; one in a specialist oncological hospital might get the impression that it is very common.

In fact, in the United Kingdom about three new cases are diagnosed each year for every 1000 of the population. One in every five persons born is likely to develop some form of the disease at some time during life. The average general practitioner will see seven new cases each year, but will see some types of malignancy very rarely. Cancer is not the commonest cause of death; the annual death rate from cardiac diseases is approximately three times that from malignant disease.

Cancer registration

With the development of cancer registration schemes, accurate estimations of the incidence of the disease in general and of its different types have become possible. It has become evident that overall the incidence has changed little over the past 50 years.

The registration of new cases on presentation provides a much more accurate estimation of incidence than do mortality statistics. Successfully treated malignant disease may not contribute to an individual's death, and therefore may not feature in the certified cause of death. Nevertheless, the incidence of death from the disease must rank high in the emotional reactions to the disease, of people in general.

Changing incidence of diseases

Since the beginning of the present century, there has been a steady reduction in the infant mortality rate, due in large part to improved neonatal care, and more recently to the availability of antibiotics. Whereas the principal causes of death in early life were infectious diseases, tuberculosis and other lung diseases, malignant disease now shows a relatively increased incidence in childhood.

Age and sex incidence of malignancy

The incidence of malignancy increases with age for most types of the disease. However, it is higher in the first five years of life than in the next two 5-year periods, principally due to leukaemia, to tumours of the central nervous system and to embryonal tumours.

In an increasing and ageing population the number of patients developing the disease will increase, even though the incidence at any age remains the same, and despite the fact that the chance of dying of the disease at any one age is gradually decreasing.

The overall incidence is the same in males and in females, but the relative rates vary with age. Below the age of ten years, it is higher in males than in females; over the period 20–60 years it is somewhat higher in females, especially in the period 35–50 years due to the relatively high incidence of malignancy of the breast and uterine cervix; over the age of 60 years, the incidence in males is markedly higher than that in females. Fig. 1.1 reflects approximately the overall incidence of the disease by age.

In western countries, the order of overall incidence of malignant disease for each sex and site is shown approximately in Table 1.1 and Fig. 1.2.

The predominating type of malignancy also varies with age. In children of both sexes up to the age of ten years, brain tumours and the leukaemias are commonest. In males, testicular tumours are commonest in the age





Fig. 1.1. Malignant disease; overall mortality by decades, England and Wales 1976. (Based on Government Statistical Service data from *Mortality Statistics—Cause.*)

Table 1.1. Incidence by site.

Males (%)	Females (%)
Bronchus (30)	Breast (25)
Digestive tract (20)	Digestive tract (20)
Urinary tract and prostate (15)	Uterus (10)
Skin (10)	Skin (10)
	Bronchus (5)

**Fig. 1.2.** Incidence of malignant disease by site. (Based on Trent Regional Health Authority data from *Radiotherapy Statistical Tables for 1977*.)

range 20-30 years, carcinoma of the bronchus in the age range 45-65 years, and adenocarcinoma of the prostate over the age of 70 years. In females, carcinomas of the breast and uterine cervix are commonest in the age range 25-65 years, and the predominance of carcinoma of the breast persists at all ages thereafter.

Geographical factors in incidence

The incidence of malignant disease can show marked variations between different countries and between different races. An outstanding example is the very low incidence of carcinoma of the uterine cervix in Jewish women. Other examples include the very high incidence of carcinoma of the postnasal space in the Chinese, the high incidence of malignancy of the

uterus in Indian women, of primary liver malignancy in South and West Africans, of bladder malignancy in Egyptians, of stomach tumours in Japanese and Scandinavians, and the very low incidence of breast malignancy in the Japanese.

It is of great interest, however, to note that, in general, immigrant populations tend to assume the pattern of incidence appropriate to their adopted country, suggesting that environmental factors have a major role in aetiology.

Social factors in incidence (Table 1.2)

These environmental factors may be related to local geographical conditions, or to different life styles and habits. In females the age of marriage, the number of pregnancies and the attitude to breast feeding may be relevant. To some extent, these factors may be determined by the degree of economic development of the country, and the associated social and economic status of the population. It is becoming increasingly evident that environmental factors are responsible for many forms of malignant disease.

Industrial factors in incidence

As will be seen in more detail later the increased risk of workers in some industries developing malignant disease is well recognised. As long ago as 1775, Percival Pott noted an association between carcinoma of the scrotal skin and chimney sweeping as an occupation. Earlier this century, the tendency to develop similar tumours in mule-spinners became evident, this being related to the period when certain mineral oils were used to lubricate the spindles of the mules.

More recently, an increased incidence of bladder carcinomas in workers in the azo dye industry and in the rubber and cable industries has been

Table 1.2. Standardised mortality ratios by occupation, England and Wales 1970-72. (Based on Government Statistical Service data from *Social Trends*.)

Site	Professional and other		Inter-mediate		Skilled				Partly skilled		Unskilled	
	M	F	M	F	Non-manual		Manual		M	F	M	F
					M	F	M	F				
Trachea, bronchus and lung	53	73	68	82	84	89	118	118	123	125	143	134
Uterine cervix	—	<20	—	66	—	69	—	120	—	140	—	161
Breast	—	117	—	121	—	110	—	109	—	103	—	92
Prostate	91	—	89	—	99	—	115	—	106	—	115	—

recognised. Workers having prolonged contact with tar and pitch show a tendency to develop warty lesions in the exposed skin, and these can progress to become carcinomas. Prolonged contact with arsenical compounds can have similar effects.

Inhalation of dust containing chromates or dichromates can lead to lung malignancy; even more active is asbestos dust, which can result in tumours of the pleura (mesotheliomas) as well as of the bronchial mucosa. Long-term inhalation of benzol vapour can result in bone marrow changes ranging from anaemia and pancytopenia to leukaemia.

Prolonged exposure of the skin to strong sunlight will result in a higher than average incidence of keratotic lesions which have a marked tendency to undergo malignant change. This is seen, for example, in Indian tea planters and in Australian sheep farmers. The incidence of malignant melanoma of the skin increases with increasing exposure to ultraviolet light, natural or artificial.

Exposure over long periods to ionising radiations also can induce skin malignancies; this was seen particularly in early radiation workers, some of whom accumulated relatively large doses to the hands before the dangers were recognised.

All these forms of malignant disease are related to identifiable chemical carcinogens or to physical agents which can have carcinogenic effects. Once recognised, steps can be taken to limit or prevent exposure, or to introduce alternative non-carcinogenic agents in industry.

Mortality from malignancy

For many forms of the disease, mortality rates are showing a gradual fall; this is true particularly for malignancy of the stomach, uterus, bones and tongue. Malignancy of the pharynx is showing a fall in females but not in males. A rise in mortality is evident for malignancy of the ovary, pancreas, bladder, kidney and lung, and for the leukaemias and the lymphomas. The increasing mortality from carcinoma of the lung (bronchus) is most striking.

The overall mortality for males is higher than that for females. This is due to the higher incidence of malignancies of lower curability in males, especially bronchial and gastric carcinomas.

Changing mortality patterns therefore reflect firstly the changes in incidence and detection and secondly the effects of therapy on different tumours.

Aetiology

The precise cause of cancers is still unknown but it is likely that many, if not all, have a multifactorial aetiology. Genetic and environmental factors are important (Table 1.3).

Table 1.3. Factors related to the cause of cancer.**GENETIC**

Familial predisposition to cancer

Chromosomal abnormality associated with increased incidence of cancer

Inherited syndromes associated with increased incidence of cancer

Histocompatibility antigen status predisposing to cancer

ENVIRONMENTAL

Irradiation

Chemicals

Viruses

Hormonal

Immunological

Chronic irritation

Genetic factors

There is evidence that individuals with certain genetic make-ups are more susceptible to cancer, though there is rarely a predictable mode of inheritance.

Familial predisposition

Some tumours occurring in children have a strong hereditary predisposition. One form of childhood retinoblastoma appears to be inherited as a dominant trait. Nephroblastoma, neuroblastoma, pheochromocytoma and neurofibromatosis also have a strong familial element.

Chromosomal abnormality syndromes

In certain types of chromosomal abnormality disorders the incidence of neoplasia is increased. In Down's syndrome (trisomy 21) acute leukaemia is a well-known complication. This complication is also found in Klinefelter's syndrome (XXY), a condition in which breast carcinoma is also more common.

Associated inherited syndromes

Certain immunodeficiency syndromes are associated with an increased incidence of cancer. Ten per cent of patients with ataxia telangiectasia (an autosomal recessive disorder with telangiectasia, progressive ataxia and variable mixed immunodeficiency) and Wiskott-Aldrich syndrome (X-linked recessive immunodeficiency with eczema and thrombocytopenia)

die of malignant disease, and the incidence may well be higher in patients with other variable immunodeficiency states. Lymphoreticular malignancy seems to be the commonest complication but the incidence of epithelial malignancy is also unexpectedly high.

Other 'non-immunodeficiency' disorders seem to predispose to cancer. Xeroderma pigmentosum, a skin condition with enhanced sun light sensitivity and increased incidence of skin and subcutaneous cancers, and polyposis coli predisposing to colonic carcinoma, are examples of these.

Histocompatibility antigen status

The histocompatibility antigen (human leucocyte antigen, HLA) system is a genetically determined single major transplantation antigen system, located within a region termed the major histocompatibility complex (MHC) on the 6th chromosome, which plays an important part in determining susceptibility and resistance to disease. Four of potentially many HLA gene loci have been identified—A, B, C and D (or D-related, DR). Particular HLA types are associated with malignancy, for example with Asiatic nasopharyngeal carcinoma (A2-Bw46) and long survivors with Hodgkin's disease (B8), acute lymphatic leukaemia (B2 and B9) and myelogenous leukaemia (B12).

An apparent increase of common cancers (such as breast, ovary and colon) is often seen in family groups; these could arise from some shared genetic characteristic, possibly HLA-mediated susceptibility, or equally from the action of similar environmental factors in a close-contact group or indeed from a combination of both.

Environmental factors

Irradiation

Ionising radiations have the capacity to displace electrons from atoms thus converting them to ions. They can therefore cause chemical changes within living cell molecules, the most important being the damage to the susceptible nuclear DNA with consequent changes in the structure and linkage of spiral strands. If these changes are severe enough cell death will result; less severe changes may cause the cells to become permanently altered in such a way as to escape normal control mechanisms—i.e. to become neoplastic.

Several examples of radiation induced tumours can be found. The high incidence of skin cancers in early X-ray workers, of lung cancer in the miners of radioactive ores, of bone tumours in girls who painted luminous

watch dials (with radioactive radium), of thyroid cancer in people who survived the atomic bomb blasts or who had neck irradiation in childhood for some reason, and of leukaemia in patients with ankylosing spondylitis treated by radiotherapy, all suggest that radiation induces mutations in the genetic material of cells in the irradiated tissues.

Sunlight, by virtue of its ultraviolet irradiation, may over the years cause skin cancer in fair skinned people, presumably by damaging DNA in skin cells.

Chemicals

A list of the better known of the chemical carcinogens is given in Table 1.4. Many of them (probably more than 10%) are encountered during occupational exposure. Chemical carcinogens may be active in their primary form (direct acting) or may need to be modified in the body before becoming active (procarcinogens). Interference with nuclear DNA is again the main mechanism of oncogenesis.

Table 1.4. Chemical carcinogens and cancer.

OCCUPATIONAL EXPOSURE

Asbestos (lung)
 Arsenic (skin, lung)
 Chromium (lung)
 Nickel (lung, paranasal sinuses)
 PVC (liver)
 Organic chemicals (lung, skin, bladder)
 petroleum fractions
 aromatic amines
 benzene

ENVIRONMENTAL AND FOOD

Aromatic hydrocarbons	}	atmospheric (? lung)
Asbestos		
Arsenic—drinking water (skin)		
Aflatoxin—moulds (liver)		
Preservatives (?)		

SOCIAL CUSTOMS

Tobacco smoking (lung, oesophagus)

IATROGENIC

Arsenic (skin)
 Cytotoxic drugs (various)
 Immunosuppressive drugs (various)
 Exogenous hormones (various)

Occupational. The most important occupational carcinogens are asbestos, arsenic, benzene, chromium, nickel and petroleum fractions, and it is important to remember that exposure to several of these chemicals can occur in one man's working lifetime with possible carcinogenesis of a number of tumours (e.g. lung carcinoma, mesothelioma, head and neck cancers). As has been seen, historically the soot induced scrotal carcinoma of chimney sweeps and the aniline dye induced bladder cancers achieved notoriety. More recently the effects of asbestos, particularly on the lung, of industrial mineral oils on the skin and of polyvinyl chloride (PVC) on the liver have provided the main debating points on chemical carcinogenesis.

Environmental. The importance of urban atmospheric pollution (e.g. with aromatic hydrocarbons and asbestos particles) in carcinogenesis is still uncertain. The roles of arsenic in drinking water of certain populations as a cause of skin cancer, and of aflatoxins from *Aspergillus flavus* in the staple foodstuffs of certain tropical peasants as a cause of liver cancer seem undeniable. There is also a causative association between pentose-rich fibres and lower gastrointestinal malignancy, and between poor nutrition and upper gastrointestinal cancer. In fact differences in diet probably account for more variation in the incidence of cancer than any other factor.

There are undoubtedly substances (e.g. nitrosamine) used as preservatives or colouring reagents in everyday foods, which if present in large enough quantities could be carcinogenic. The amounts present in food however are minute and the hazards which they present uncertain.

Social customs. Indisputably the main offender in this category is the habit of cigarette smoking—the carcinogenic effects of which are well known. The chewing of quids (beetle nuts, tobacco, burnt lime) is an important cause of oral cancer. Alcohol is thought to increase the incidence of mouth and throat cancer, particularly in smokers. It is estimated that tobacco has an aetiological role in 25–30% of all cancers.

Iatrogenic. Iatrogenic (doctor induced) chemical carcinogenesis is of uncertain importance but the effects of our new medications on the population may take years to evaluate. The carcinogenic problems seen at the present time are mainly related to the use of immunosuppressive and cytotoxic drugs in cancer, transplantation and autoimmune disease, and to the use of exogenous hormones (e.g. anabolic steroids and liver cancer, prenatal oestrogens and vaginal carcinoma).

Viruses

Several types of virus have been implicated in various animal tumours both naturally occurring and experimentally induced. Perhaps the most

important are the Polyoma and SV40 (Papova) viruses, which have been very useful in experimental cancer research, and the herpes type viruses, which have been incriminated in Marek's disease—a lymphoproliferative disorder of chickens.

There is little conclusive evidence of viral oncogenesis in man but circumstantial evidence suggests that infection may be an important cause in certain tumours. The occurrence of time-space clusters of patients with Hodgkin's disease and leukaemia is very suggestive of an infective aetiology. In carcinoma of the cervix an aetiological association with herpesvirus and/or human Papillomavirus type 16, is now apparent. In these conditions however, other environmental factors may be equally important. The hepatitis B virus also has a part to play in hepatoma.

No discussion of viral oncogenesis would be complete without mention of Burkitt's lymphoma. Patients with this type of lymphoma occur in clusters and invariably have antibodies to the Epstein-Barr virus; the virus has been isolated from cultured Burkitt's lymphoma cell lines. However the virus does in fact also cause infectious mononucleosis, a benign self-limiting infection, and it is difficult to see how the same virus could cause two such different disorders without some major host-virus adaptations. It now seems likely that chronic severe malaria depresses the host immunity in such a way as to allow continued infection with the virus and subsequent oncogenesis. The Epstein-Barr virus is also causally associated with nasopharyngeal carcinoma in the Chinese.

Another, albeit rare, neoplasm seen with viral infection is that caused by the human T cell leukaemia/lymphoma virus (HTLV) type I. It is also possible that this, or a similar virus, is involved in a T cell lymphoma of the skin (mycosis fungoides).

The acquired immune deficiency syndrome (AIDS) was first recognised as a new disease in 1981. Patients with this syndrome had persistent lymphadenopathy and were very susceptible to infections, as well as suffering from generalised ill health. These patients were found to have low helper/suppressor peripheral T cell ratios. The disease was noted to be more prevalent in homosexuals, drug abusers and recipients of blood products (particularly haemophiliacs), and patients were prone to develop rare opportunistic infections such as *Pneumocystis carinii* pneumonia. They also tended to develop malignancies such as Kaposi's sarcoma. This is normally an indolent malignancy, responsive to radio and/or chemotherapy; in AIDS however, it is very aggressive in its behaviour and may be rapidly lethal. It is now thought that a virus, HTLV type III, is causally associated with the development of AIDS. It is possible that some or all of the associated malignancies are caused by HTLV, or that the resulting immuno-

suppression allows other viruses (e.g. cytomegalovirus or Epstein-Barr virus) to become oncogenic.

Hormonal

Perhaps the most striking example of hormonal influence in cancer is in breast carcinoma. Some breast tumours are hormone responsive and this seems to correlate with the presence or absence of steroid receptor proteins in breast tissue cells. This has considerable therapeutic implications as we shall see later. It is likely that imbalance of endogenous hormones, rather than any direct oncogenic effects of these hormones, predisposes to cancer. Other examples of hormone dependent tumours are prostatic carcinoma and carcinoma of the body of the uterus.

Immunological

The role of immunology in the aetiology of tumours will be discussed more fully later. Suffice to say, that one theory of oncogenesis is that potentially neoplastic cells are normally eradicated by a competent immunosurveillance system; should this system become incompetent for any reasons (genetic or environmental) the cells escape control and proliferate, thus forming the cancer.

Chronic irritation

Cancers can arise at sites of chronic irritation, and in relation to scars, foreign bodies and chronic inflammation. Presumably cell damage at these sites gives rise to abnormal tissue differentiation. Oral cancer is probably the best example of this type of carcinogenesis—factors such as pipe smoking, ill-fitting dentures, poor dental hygiene and chronic infection (e.g. syphilis) are all recognised predisposing causes. In carcinoma of the vagina and cervix uteri there is an aetiological association with early coitus and/or poor genital hygiene. Certain chronic diseases also predispose to human cancer. Good examples of this are achlorhydria and pernicious anaemia and stomach cancer, Paterson-Kelly syndrome and post cricoid carcinoma, ulcerative colitis and colonic carcinoma, cirrhosis and hepatoma, Paget's disease and bone sarcoma, bilharzia and bladder cancer.

Oncogenes

A key problem in the fight against cancer is to find differences between normal and malignant cells. One such difference is now known to be