

American
DRUG
INDEX

1988

American
DRUG INDEX
1988

32nd Edition

NORMAN F. BILLUPS, R.Ph., M.S., Ph.D.

Dean and Professor of Pharmacy
College of Pharmacy
The University of Toledo

Associate Editor

SHIRLEY M. BILLUPS, R.N., B.Ed.

Oncology Nurse
The Toledo Hospital



J. B. LIPPINCOTT COMPANY
Philadelphia

London
Mexico City

New York
St. Louis

São Paulo
Sydney

Copyright © 1988, by J. B. Lippincott Company

Copyright © 1956, 1958, 1959, 1960, 1961, 1962, 1963, 1964, 1965, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983, 1984, 1985, 1986, 1987 by J. B. Lippincott Company

This book is fully protected by copyright and, with the exception of brief excerpts for review, no part of it may be reproduced in any form by print, photoprint, microfilm or any other means, without the written permission of the publishers.

ISBN 0-932686-25-7

Library of Congress Catalog Card Number 55-6286

Printed in the United States of America

Preface

The 32nd Edition of the *American Drug Index* (ADI) has been prepared for the identification, explanation, and correlation of the many pharmaceuticals available to the medical, pharmaceutical, and allied health professions. The need for this index has become acute as the variety and number of drugs and drug products have continued to multiply. For this reason the *ADI* should be most useful to pharmacists, nurses, physicians, medical transcriptionists, dentists, sales personnel, students, and teachers in the fields incorporating pharmaceuticals.

Special note to medical transcriptionists: Although all names in the *ADI* appear in bold capitalized type, trade products are easily identifiable by the manufacturer's name in parentheses immediately following the trade name. The names for official products are preceded by a bullet (•) and should appear in lower case in transcription. The therapeutic and pharmaceutical classes are denoted by a box (■) preceding the name and should be a helpful cross-reference guide.

The organization of the *ADI* falls into the following 11 major sections: Monographs of Drug Products, Common Abbreviations Used in Medical Orders, Approximate Practical Equivalents, Common Systems of Weight and Measure, Glossary, New U.S.P. XXI and N.F. XVI Monographs, Container Requirements for U.S.P. XXI Drugs, Container and Storage Requirements for Sterile U.S.P. XXI Drugs, Oral Dosage Forms That Should Not Be Crushed, Pharmaceutical Company Labeler Code Index Numbers, and Pharmaceutical Manufacturer and/or Drug Distributor Addresses. The New U.S.P. XXI and N.F. XVI Monographs section has been added to make the *ADI* an even more complete and valuable reference tool.

MONOGRAPHS: The organization of the monograph section of the *ADI* is fundamentally alphabetical with extensive cross-indexing. Names listed are generic (also called nonproprietary, public name, or common name); brand (also called trademark, proprietary, or specialty); and chemical. Synonyms that are in general use also are included. All names used for a pharmaceutical appear in alphabetical order with the pertinent data given under the brand name by which it is made available.

The monograph for a typical brand name product consists of the manufacturer, generic and/or chemical names, composition and strength, pharmaceutical forms available, package size, dosage, and use.

Generic names appear in alphabetical order, followed by the corresponding recognition of the drug to the U.S.P. (United States Pharmacopeia), N.F. (National Formulary), and USAN (United States Adopted Names). Each of these officially recognized generic names is preceded by a bullet (•) at the beginning of each entry. The information is in accord with the U.S.P. XXI and N.F. XVI and through

Supplement 5, which became official on May 15, 1987; and the USAN-1988.

British Approved Names, indicated by B.A.N., are included in this edition with the permission of the Controller of Her Britannic Majesty's Stationery Office.

Because of the multiplicity of specialty names (brand names) used for the same therapeutic agent or the same combination of therapeutic agents, it was apparent that some correlation could be done. As an example of this, please turn to tetracycline HCl. Here under the generic name are listed the various brand names. Following are combinations of tetracycline HCl organized in a manner to point out relationships among the many products. Reference then is made to the specialty name or names having the indicated composition. Under the specialty name are given manufacturer, composition, available forms, sizes, dosage, and use.

The multiplicity of generic names for the same therapeutic agent has complicated the nomenclature of these agents. Examples of multiple generic names for the same chemical substance are (1) parabromdylamine, brompheniramine; (2) acetaminophen, p-hydroxy acetanilid, N-acetyl-p-aminophenol; (3) guaifenesin, glyceryl guaiacolate, glyceryl guaiacol ether, guaianesin, guaifylline, guaiphenesin, guayanesin, methphenoxdyol; (4) pyrilamine, pyranisamine, pyranilamine, pyraminyl, anisopyradamine.

The cross-indexing feature of the *ADI* permits the finding of drugs or drug combinations when only one major ingredient is known. For example, a combination of aluminum hydroxide gel and magnesium trisilicate is available. This combination can be found by looking under the name of either of the two ingredients, and in each case the specialty names are given. A second form of cross-indexing involves the listing of drugs under various therapeutic and pharmaceutical classes (i.e., antacids, antihistamines, diuretics, laxatives, etc.). Each of these classes is preceded by a box (■) at the beginning of each entry.

ABBREVIATIONS: A listing of Common Abbreviations Used in Medical Orders is included as an aid in interpreting medical orders. The Latin or Greek word and abbreviation are both given with the meaning.

CONVERSION FACTORS: The listing of Approximate Practical Equivalents is added as an aid in calculating and converting dosages among the metric, apothecary, and avoirdupois systems.

WEIGHT AND MEASURE: Tables containing the Common Systems of Weight and Measure are included to aid the practitioner in calculating dosages in the metric, apothecary, and avoirdupois systems.

GLOSSARY: A listing of commonly used terms, with brief definitions, is included as an aid in interpreting the use given for drug monographs listed in the *ADI*.

NEW TITLES FOR U.S.P. XXI AND N.F. XVI MONOGRAPHS: 252

new official monographs appeared through Supplement 5 to the 1986 U.S.P. XXI and N.F. XVI, and an additional 535 new monographs have appeared since the 1980 U.S.P. XX and N.F. XV. Since many of these new monographs apply to commonly used drug products, this section has been added to aid the practitioner in identifying drug products.

CONTAINER AND STORAGE REQUIREMENTS FOR U.S.P. XXI COMPENDIAL DRUGS: These sections have been added on container and storage requirements specified by the U.S.P. XXI for compendial drugs, to aid the practitioner in storing and dispensing.

ORAL DOSAGE FORMS THAT SHOULD NOT BE CRUSHED: This section has been added to alert the health care practitioner about oral dosage forms that should not be crushed, and to serve as an aid in consulting with patients. Examples of products falling into the "non-crush" category are extended-release, enteric-coated, encapsulated beads, wax matrix, sublingual, and encapsulated liquid formulations.

LABELER CODE INDEX: The Pharmaceutical Labeler Code Index is presented to aid the practitioner in the identification of drug products. In this section, the codes are listed in numerical order followed by the name of the manufacturer.

MANUFACTURER ADDRESSES: The name, address, and zip code of virtually every American Pharmaceutical Manufacturer and/or Drug Distributor are listed in alphabetical order in this section. Additionally, a pharmaceutical labeler code number appears before the address of each company as a further aid in the identification of drug products.

A special word of appreciation and acknowledgment is given to my wife, Shirley, who served again this year as my Associate Editor—and helped to compile much of the information that appears in the monograph section of this volume. Special thanks are also extended to the manufacturers who supplied product information and made helpful comments and suggestions for this index, and to Dr. Kenneth S. Alexander, who organized the information on Container and Storage Requirements for U.S.P. XXI Drugs.

Correspondence or communication with reference to a drug or drug products listed in the *American Drug Index* should be directed to Norman F. Billups, Ph.D., Dean, College of Pharmacy; The University of Toledo; Toledo, Ohio 43606, or call 419-537-4235.

Norman F. Billups, R.Ph., M.S., Ph.D.

Contents

1. Monographs	1-661
2. Common Abbreviations Used In Medical Orders	662-663
3. Approximate Practical Equivalents	664
4. Common Systems of Weight and Measure	665
5. Glossary	666-674
6. New U.S.P. XXI and the N.F. XVI Monographs	675-678
7. Container Requirements for U.S.P. XXI Drugs	679-691
8. Container and Storage Requirements for Sterile U.S.P. XXI Drugs	692-701
9. Oral Dosage Forms That Should Not Be Crushed	702-705
10. Pharmaceutical Company Labeler Code Index	706-711
11. Pharmaceutical Manufacturers and/or Drug Distributors	712-726

[•] Denotes official name: Generic name or chemical name approved by F.D.A. or recognized by the U.S.P., N.F., or USAN.

[■] Denotes therapeutic and pharmaceutical classes.

A

A-200 PYRINATE. (Norcliff Thayer) **Liquid:** Pyrethrins 0.17%, piperonyl butoxide technical 2.00%, petroleum distillate 5.00%. Bot. 2 fl. oz., 4 fl. oz.

Gel: Pyrethrins 0.33%, piperonyl butoxide technical 4.00%, petroleum distillate 5.33%. Tube 1 oz.

Use: Topically, pediculosis.

A.A.A. OINTMENT. (Jenkins) Ammoniated mercury 2%, acid salicylic 1.25%, acid boric 1%, zinc oxide 15%. Jar 1 oz., 1 lb.

Use: Impetigo, nonspecific eczema & minor skin irritations.

A AND D HAND CREAM. (Schering) Emollients isopropyl myristate & squalane. Tube 2 oz.

Use: Dry skin, sunburn, irritations.

A AND D OINTMENT. (Kenyon) Vits. A & D in lanolin-petrolatum base. Jar 1 lb.

A AND D OINTMENT. (Schering) Petrolatum-lanolin. Tube 1.5 oz., 4 oz., Jar 1 lb.

A & D TABLETS. (Barth's) Vit. A 10,000 I.U., D 400 I.U./Tab. Bot. 100s, 500s.

A AND D VITAMIN CAPSULES. (Lannett) Vit. A 5000 I.U., D 400 I.U./Cap. Bot. 500s, 1000s.

Use: Vit. supplement.

A & D VITAMIN OINTMENT. (Quality Generics) Vit. A & D in lanolin-petroleum base Tube 2 oz., Jar 1 lb. Use: Emollient.

A AND D VITAMIN OINTMENT. (Lannett) In a petrolatum-lanolin base. Jar 1, 5 lb.

A-A PLUS. (Galen) Al hydroxide gel 400 mg., Mg oxide 175 mg., Na lauryl sulfate 25 mg., Me cellulose 100 mg., hyoscyamine sulfate 0.1037 mg., atropine sulfate 0.0194 mg., hyoscine HBr 0.0065 mg./Tab. Bot. 100s, 1000s. Susp. 1 pt.

AAS INFANTIL W/VITAMIN C. (Winthrop Products) Acetylsalicylic acid, Vit. C.

Use: Analgesic, antipyretic.

AAS TABLETS. (Winthrop Products) Acetylsalicylic acid.

Use: Analgesic, antipyretic.

ABBOCIN. Oxytetracycline, B.A.N.

ABBOKINASE. (Abbott) Urokinase 250,000 I.U./5 ml. Lyophilized powder. Vial 5 ml.

Use: Plasminogen activator, thrombolytic agent.

ABBOKINASE OPEN-CATH. (Abbott) Urokinase for catheter clearance 5,000 I.U./ml. Unival.

Use: Clearance of fibrin occluded catheters.

ABBOTT AFP-EIA. (Abbott Diagnostics) Test Kits 100s.

Use: Enzyme immunoassay for the quantitative measurement of alpha-fetoprotein in human serum and amniotic fluid.

ABBOTT AFP-EIA MONOCLONAL. (Abbott Diagnostics)

Use: Enzyme immunoassay for the quantitative measurement of Alpha-fetoprotein (AFP) in human serum and amniotic fluid.

ABBOTT ANTI-DELTA. (Abbott Diagnostics) Radioimmunoassay for the detection of antibody to delta antigen (HDag) in human serum or plasma.

For research only. Not for use in diagnostic procedures.

ABBOTT ANTI-DELTA EIA. (Abbott Diagnostics) Enzyme immunoassay for the detection of antibody to hepatitis delta antigen in human serum or plasma. For research only. Not for use in diagnostic procedures.

ABBOTT β -HCG 15/15. (Abbott Diagnostics) Enzyme immunoassay for the quantitative determination of human chorionic gonadotropin in human serum.

ABBOTT CA125-EIA. (Abbott Diagnostics) Enzyme immunoassay for the quantitative measurement of cancer antigen (CA) 125 in human serum. For research use only. Not for diagnostic procedures.

ABBOTT CEA-EIA MONOCLONAL. (Abbott Diagnostics) Enzyme immunoassay for the quantitative measurement of carcinoembryonic antigen (CEA) in human serum or plasma to aid in the management of cancer patients and assessing prognosis.

ABBOTT CEA-RIA. (Abbott Diagnostics) Solid phase radioimmunoassay for the quantitative measurement of carcinoembryonic antigen (CEA) in human serum or plasma to aid in the management of cancer patients and assessing prognosis.

ABBOTT CMV TOTAL AB EIA. (Abbott Diagnostics) Test kits 100s.

Use: Enzyme immunoassay for the detection of antibody to cytomegalovirus in human serum, plasma, and whole blood.

ABBOTT DIAGNOSTIC REAGENTS. (Abbott Diagnostics) A series of diagnostic tests for cancer, cardiovascular, hepatitis, infectious disease and immunology, metabolic and digestive disease, OB/GYN, rubella, and thyroid.

ABBOTT ER-EIA MONOCLONAL. (Abbott Diagnostics) Enzyme immunoassay for the quantitative measurement of human estrogen receptor in tissue cytosol. For research use only. Not for diagnostic purposes.

ABBOTT ER-ICA MONOCLONAL. (Abbott Diagnostics) Immunoassay for the detection of estrogen receptor. For research use only. Not for diagnostic procedures.

ABBOTT-HB EIA. (Abbott Diagnostics) Enzyme immunoassay for the detection of hepatitis Be antigen and/or antibody to hepatitis Be antigen.

ABBOTT-HB_e TEST. (Abbott Diagnostics) Test kits 100s.

Use: Radioimmunoassay or enzyme immunoassay for detection of hepatitis Be antigen and/or antibody to hepatitis Be antigen.

ABBOTT HTLV III ANTIGEN EIA. (Abbott Diagnostics) Enzyme immunoassay for the detection of Human T-Lymphotropic Virus Type III (HTLV III) antigens. For research use only.

ABBOTT HTLV III EIA. (Abbott Diagnostics) Enzyme immunoassay for the detection of antibody to human T-lymphotropic virus type III (HTLV III) in human serum or plasma.

ABBOTT IgE EIA. (Abbott Diagnostics) Test kits 100s.

Use: Enzyme immunoassay for quantitative determination of IgE in human serum and plasma.

ABBOTT PAP-EIA. (Abbott Diagnostics) Enzyme immunoassay for the measurement of prostatic acid phosphatase (PAP) in serum or plasma.

ABBOTT RSV-EIA. (Abbott Diagnostics) Enzyme immunoassay for the detection of respiratory syncytial virus (RSV) in nasopharyngeal washes and aspirates.

ABBOTT SCC-RIA. (Abbott Diagnostics) Radioimmunoassay for the quantitative measurement of squamous cell carcinoma associated antigen in human serum. For research use only. Not for diagnostic procedures.

ABBOTT TdT EIA. (Abbott Diagnostics) Enzyme immunoassay for the quantitative measurement of terminal deoxynucleotidyl transferase (TdT), in extracts of human whole blood or isolated mononuclear cells.

ABBOTT TESTPACK HCG-SERUM. (Abbott Diagnostics) Monoclonal antibody, enzyme immunoassay for the qualitative determination of human chorionic gonadotropin (hCG) in serum. No instrumentation required.

ABBOTT TESTPAK HCG-URINE. (Abbott Diagnostics) Monoclonal antibody, enzyme immunoassay for the qualitative determination of human chorionic gonadotropin (HCG) in urine. No instrumentation required.

ABBOTT TESTPACK-STREP A. (Abbott Diagnostics) A rapid screening and confirmatory test for the detection of Group A beta-hemolytic streptococci from throat swabs. No instrumentation required.

ABBOTT TOXO-G EIA. (Abbott Diagnostics) Enzyme immunoassay for the qualitative and quantitative determination of IgG antibody to toxoplasma gondii in human serum and plasma.

ABBOTT TOXO-M EIA. (Abbott Diagnostics) Enzyme immunoassay for the qualitative determination of IgM antibody to toxoplasma gondii in human serum.

ABCDG VITAMIN CAPSULES. (Lannett) Vit. A 5000 u., D 400 u., B-1 1 mg., B-2 2 mg., C 30 mg./Cap. Bot. 1000s.

Use: Vitamin supplement.

ABDEC BABY VITAMIN DROPS. (Parke-Davis) Vit. A 1500 I.U., D 400 I.U., E 5 I.U., B-1 0.5 mg., B-2 0.6 mg., B-6 0.4 mg., B-12 2 mcg., biotin 0.05 mg., pantothenic acid 3 mg., niacin 8 mg., C 35 mg./ml. Bot. 1½ oz. w/dropper.

Use: Vitamin deficiencies.

ABDEC WITH FLUORIDE BABY VITAMIN DROPS. (Parke-Davis) Vit. A 1500 I.U., D 400 I.U., C 35 mg., E 5 I.U., B-1 0.5 mg., B-2 0.6 mg., B-6 0.4 mg., niacin 8 mg., B-12 2 mcg., biotin 0.05 mg., pantothenic acid 3 mg./ml. w/fluoride 0.25 mg. Bot. 50 ml.

Use: Vitamin therapy with fluoride.

ABDO. (Eastwood) Cap. Bot. 100s.

ABSORBABLE CELLULOSE COTTON OR GAUZE.

See: Oxidized Cellulose (Various Mfr.)

• **ABSORBABLE DUSTING POWDER,** U.S.P. XXI.

• **ABSORBABLE GELATIN FILM,** U.S.P. XXI. Sterile nonantigenic, absorbable, water-insoluble, gelatin film. See: Gelfilm (Upjohn)

• **ABSORBABLE GELATIN SPONGE,** U.S.P. XXI. Gelatin sponge.

Use: Surgical aid.

See: Gelfoam, Preps. (Upjohn)

• **ABSORBABLE SURGICAL SUTURE,** U.S.P. XXI. (Various Mfr.) Surgical Gut, Surgical Catgut, Catgut suture.

Use: Surgical aid.

• **ABSORBENT GAUZE,** U.S.P. XXI.

Use: Surgical aid.

ABSORBENT RUB RELIEF FORMULA. (DeWitt)

Green soap 11.64%, camphor 1.63%, menthol 1.63%, pine tar soap 0.87%, oil wintergreen 0.71%, oil sassafras 0.54%, benzocaine 0.48%, capsicum 0.03%, oil wormwood 0.60%, iso alcohol 75%. Bot. 2 oz.

Use: Sore muscle and joint pain.

ABSORBINE ARTHRITIC PAIN LOTION. (W.F. Young) Bot. 2 oz., 4 oz.

ABSORBINE FOOT POWDER. (W.F. Young) Zinc stearate, parachloroxylenol, aluminum chlorhydroxy, allantoinate, benzethonium chloride, menthol.

Plastic bot. 3 oz. w/shaker top.

Use: Athlete's foot treatment.

ABSORBINE, JR. (W.F. Young) Wormwood, thymol, chloroxylenol, menthol, acetone, zinc stearate, parachloroxylenol, aluminum chlorhydroxy, allantoinate, benzethonium chloride, menthol. Liq. Bot. 1 oz., 2 oz., 4 oz., 12 oz. w/applicator. Use: Muscle soreness, stiffness, treatment of athlete's foot.

ABUSCREEN. (Roche Diagnostics) An immunological and radiochemical assay for morphine and morphine glucuronide in nanogram levels. Utilizes I-125 labeled morphine requiring gamma scintillation equipment. 100 tests.

ACABEL. Bevonium Methylsulphate, B.A.N.

• **ACACIA,** N.F. XVI. Syrup N.F. XVI. Acacia senegal, gum arabic. (Penick) pow., 0.25 lb., 1 lb.; tears, 0.25-lb.; whole, 0.25-lb.

Use: Demulcent, emulsifier, suspending agent.

A-CAINE. (A.V.P.) Rectal ointment. Tube 1.25 oz.

Use: Rectal care, anti-hemorrhoidal.

A-CAPS. (Drug Industries) Vit. A 50,000 I.U./Cap. Bot. 100s, 500s.

Use: Vitamin A therapy.

• **ACARBOSE.** USAN.

Use: Alpha-glucosidase inhibitor.

ACCELERASE. (Organon) Pancrelipase: Lipase 4000 U., amylase 15,000 U., protease 15,000 U., other pancreatic enzymes mixed with bile salts 65 mg., cellulase 2 mg., calcium carbonate 20 mg./Cap. Bot. 60s.

Use: Functional digestive disorders.

ACCURBRON. (Merrell Dow) Theophylline, anhydrous 10 mg./ml. Bot. pt.

Use: Bronchial asthma treatment.

ACCUSENS T TASTE FUNCTION KIT. (Westport)

Test for ability to distinguish among salty, sweet,

- sour and bitter tastants. Kit contains 15 bottles (60 ml.) tastants and 30 taste record forms.
Use: Diagnostic taste function kit.
- ACCUTANE.** (Roche) Isotretinoin 10 mg., 20 mg., or 40 mg./Cap. Prescription Paks 30s.
Use: Treatment of severe recalcitrant cystic acne.
- A-C-D SOLUTION.** U.S.P. Sod. citrate, citric acid & dextrose in sterile pyrogen-free solution.
Baxter-600 cc. bot. with 70, 120 or 300 cc. of sol.; 1000 cc. with 500 cc. of sol.
Cutter-500 cc. bot. with 75 or 120 cc. of sol.; 650 cc. bot. with 80 or 130 cc. of sol.
Diamond-(Abbo-Vac) 250 cc., 500 cc.
Use: Anticoagulant for preparation of plasma or whole blood.
- A-C-D SOLUTION MODIFIED.** (Squibb) Acid citrate dextrose anticoagulant solution modified.
Use: Anticoagulant for use in radiolabeling red blood cells.
- ACDRILE.** Methyl Cysteine, B.A.N.
- **ACEBUTOLOL.** USAN. ($\pm$)-1-(2-Acetyl-4-butyramidophenoxy)-3-isopropylaminopropan-2-ol. (Ives) ($\pm$)-N-[3-Acetyl-4-[2-hydroxy-3-[(1-methylethyl)amino]propoxy]phenyl] butamide.
Use: Beta adrenergic receptor blocking agent.
See: Sectral, Cap. (Ives)
- **ACECAINIDE HYDROCHLORIDE.** USAN.
Use: Cardiac depressant.
- **ACECLIDINE.** USAN. 3-Quinuclidinol acetate (ester).
Use: Parasympathomimetic.
- **ACEDAPSONE.** USAN. 4',4''-Sulfonylbis (acetanilide).
Use: Antimalarial; antibacterial (leprostatic).
- ACEDOVAL.** (Vale) Dover's powder 15 mg., ipecac 1.5 mg., aspirin 162 mg., caffeine anhydrous 8.1 mg./Tab. Bot. 1000s, 5000s.
Use: Analgesic, antispasmodic, antiperistaltic.
- ACEFYLLINE PIPERAZINE.** Acephylline, B.A.N.
- **ACEGLUTAMIDE ALUMINUM.** USAN.
Use: Antulcerative.
- ACENOCOUMAROL.** 3-(α -Acetonyl-4-nitrobenzyl)-4-hydroxy-coumarin. Nicoumalone, B.A.N.
- ACEPHEN.** (G&W) Adult: Acetaminophen 650 mg./Supp. Box 12s, 100s. **Pediatric:** Acetaminophen 120 mg./Supp. Box 12s, 100s.
Use: Analgesic, antipyretic.
- ACEPIFYLLINE.** B.A.N. Piperazine theophyllin-7-ylacetate. Acephylline Piperazine (I.N.N.). Etophylate.
Use: Spasmodic.
- ACEPROMAZINE.** 10-(3-Dimethylaminopropyl)-3-acetylphenothiazine maleate. Plegicil. (Ayerst) 1-(10-(3-(Dimethylamino)propyl)-10H-phenothiazin-2-yl)ethanol e(Z)-2-butenedioate.
Use: Tranquilizer.
- ACEPROMAZINE.** B.A.N. 2-Acetyl-10-(3-di-methylaminopropyl)phenothiazine. Notensil maleate.
Use: Tranquilizer.
- **ACEPROMAZINE MALEATE.** USAN.
Use: Sedative.
- ACEROLA-C.** (Barth's) Vit. C 300 mg./Wafer. Bot. 30s, 90s, 180s, 360s.
Use: Vitamin C therapy.
- ACEROLA-PLEX.** (Barth's) Vit. C 100 mg., bioflavonoids 50 mg./Tab. Bot. 100s, 500s.
Use: Vitamin C therapy.
- ACETA.** (Century) Acetaminophen. Elix. (120 mg./5 cc) Bot. 4 oz., pint, gal. **Tab.** (5 gr.) Bot. 100s, 1000s, 3000s, 5000s.
Use: Analgesic.
- ACTAGEN OTC SYRUP.** (Goldline) Triprolidine HCl 1.25 mg., pseudoephedrine HCl 30 mg./5 ml. Bot. 4 oz., pt., gal.
Use: Nasal decongestant.
- ACETA-GESIC.** (Rugby)
Use: Analgesic.
- ACETA W/CODEINE.** (Century) Acetaminophen 325 mg., codeine phosphate 0.5 gr./Tab. Bot. 100s.
Use: Analgesic.
- ACETAMIDE.** N-[5-(Aminosulfonyl)-3-methyl-1,3,4-thiadiazol-2(3H)-ylidene]-. Methazolamide.
- ACETAMIDE, 2,2-DICHLORO-N-[2-HYDROXY-1-(HYDROXYMETHYL)-2-(4-NITROPHENYL)ETHYL].** Chloramphenicol, U.S.P. XXI.
Use: Antibiotic.
- ACETAMIDE, 2-(DIETHYLAMINO)-N-(2,6-DIMETHYLPHENYL)-.** Lidocaine, U.S.P. XXI.
Use: Local anesthetic.
- ACETAMIDE, N-[5-(AMINOSULFONYL)-1,3,4-THIADIAZOL-2-YL]-.** Acetazolamide, U.S.P. XXI.
Use: Carbonic anhydrase inhibitor.
- ACETAMIDE, N-(4-HYDROXYPHENYL)-.** Acetaminophen, U.S.P. XXI.
Use: Analgesic, antipyretic.
- ACETAMIDE, N-(5,6,7,9-TETRAHYDRO-1,2,3,10-TETRAMETHOXY-9-OXOBENZO[a]-HEPTALEN-7-YL)-.** Colchicine, U.S.P. XXI.
Use: Gout suppressant.
- p-ACETAMIDOBENZOIC ACID SALT** of 2-di-methylaminoethanol.
- 5-ACETAMIDO-2,4,6-TRIIODO-N-METHYLISOPHTHALIC ACID.** Iothalamic Acid, U.S.P. XXI.
Use: Diagnostic aid.
- 5-ACETAMIDO-2,4,6-TRIIODO-N-METHYLISOPHTHALIC ACID, SODIUM SALT.** Iothalamate Sodium Injection, U.S.P. XXI.
Use: Diagnostic aid.
- **ACETAMINOPHEN.** U.S.P. XXI. Elix., for Effervescent Oral Soln., Tabs, Cap. U.S.P. XXI. N-Acetyl-p-amino-phenol. Acetamide, N-(4-hydroxyphenyl)-. 4'-Hydroxyacetanilide. APAP.
Use: Analgesic, antipyretic.
See: Acephen, Supp. (G&W)
Aceta, Tab. Elix., Supp. (Century)
Actamin, Tab. (Buffington)
Acetaminophen Uniserts, Suppos. (Upsher-Smith)
Aminodyne, Elix. (Bowman)
Anapap, Tab. (Fellows)
Anelix, Elix. (Kirkman)
Anuphen, Supp. (Comatic)
Apap, Cap., Tab. (Various Mfr.)
Capital, Tab. (Carnrick)
Chlor-A-Tyl. (Jenkins)
Dapa, Tab. (Ferndale)
Datril 500, Tab. (Bristol-Myers)

Dolanex, Elix. (Lannett)
 Dorcol, Prods. (Sandoz Consumer)
 Dularin, Syr. (Donner)
 Febridol, Elix., Tab. (Amid)
 Fendon, Tab. (APC)
 G-1 (Hauck)
 Janupap, Tab. (Reid-Rowell)
 Lestemp, Elix. (Reid-Rowell)
 Liquiprin, Solu. (Norcliff Thayer)
 Lyteca, Syr., Tab. (Westerfield)
 Medigesic (Medics)
 Nebs, Tab., Liq. (Eaton)
 Nilprin 7.5, Tab. (AVP)
 Parten, Tab. (Parmed)
 Phenaphen, Cap., Tab. (Robins)
 Proval, Cap., Elix., Drop., Tab. (Reid-Rowell)
 Tapar, Tab. (Parke-Davis)
 Temetan, Elix., Tab. (Nevin)
 Temptra, Drop, Syr., (Mead Johnson)
 Tenlap, Elix. (Merrell Dow)
 Tylenol, Drops, Elix., Liq., Tab., Chew. Tab. (McNeil)
 Tylenol Extra-Strength, Tab., Cap. (McNeil)
 Valadol, Tab., Liq. (Squibb Mark)
 Valorin Tab. (Otis Clapp)

ACETAMINOPHEN W/COMBINATIONS

Aceta w/Codeine, Tab. (Century)
 Akes-N-Pain, Cap. (Edw. J. Moore)
 Al-Ay Modified, Tab. (Bowman)
 Algecol, Tab. (Winston)
 Allerest Headache Strength, Tab. (Pharmacraft)
 Allylgesic W/Ergotamine, Cap. (Elder)
 Allylvon, Cap. (Elder)
 Alumadrine, Tab. (Fleming)
 Amaphen, Cap. (Trimen)
 Aminodyne, Elix. (Bowman)
 Anaphen, Cap. (Mallard)
 Anodynos-DHC, Tab. (Berlex)
 Anodynos Forte, Tab. (Buffington)
 Apap w/Codeine, Tab. (Central)
 Apapalone, Tab. (Amfre-Grant)
 Arthol Tabs. (Towne)
 Arthralgen, Tab. (Robins)
 Bancaps (Westerfield)
 Bancaps-C (Westerfield)
 Banesin-Forte, Tab. (Westerfield)
 Bexahist, Cap. (Bexar)
 Biophens-S, Tab. (La Crosse)
 Blanex, Cap. (Edwards)
 Bowman Cold Tab. (Bowman)
 B-Pap., Liq. (Wren)
 BQ Cold Tabs (Bristol-Myers)
 Bromo-Seltzer, Gran. (Warner-Lambert)
 Butigetic, Tab. (McNeil)
 Capital and Codeine, Susp. (Carnrick)
 Cardui, Tab. (Chattem Labs.)
 Ceroase Comp., Cap. (Ives)
 Chexit, Tab. (Sandoz Consumer)
 Christodyne-DHC, Tab. (Paddock)
 Coastaladyne, Tab. (Coastal)
 Coastalgesic, Tab. (Coastal)
 Codalan, Tab. (Lannett)
 Codimal, Tab. (Central)
 Colrex Compound, Elix. (Reid-Rowell)
 Comtrex, Cap., Liq., Tab. (Bristol-Myers)
 Conar-A, Tab., Susp. (Beecham Labs)
 Conex, Preps. (Westerfield)
 Congesprin, Liq., Tab. (Bristol-Myers)
 Contac, Prods. (SmithKline Prods)
 Coricidin Sinus Headache Tab. (Schering)
 Coryban-D, Syr. (Roerig)
 Corzans, Tab. (Moore Kirk)
 Co-Tylenol, Liq., Tab. (McNeil)
 Dapase, Tab. (Ferndale)
 Darvocet-N, Tab. (Lilly)
 Darvocet-N 100, Tab. (Lilly)
 Daycare, Liq. (Vicks)
 Demerol APAP, Tab. (Winthrop Pharm)
 Dengesic, Tab. (Scott-Alison)
 Desa-Hist AT, Tab. (Pharmics)
 Dilone, Tab. (Vicks)
 Dolene AP-65, Tab. (Lederle)
 Dolopar, Liq. (O'Neal)
 Drinacet Cap. (Phillips Roxane)
 Drinophen, Cap. (Lannett)
 D-Sinus, Cap. (Laser)
 Dularin-TH, Tab. (Dooner)
 Duoprin, Tab. (Dunhall)
 Duo-3X, Tab. (Med. Spec.)
 Empracet with Codeine, Tab. (Burroughs Wellcome)
 Enz-Cold, Cap. (Edw. J. Moore)
 Esemgesic, Tab. (Spencer-Mead)
 Esgic, Tab. (Gilbert)
 Excedrin, Cap., Tab. (Bristol-Myers)
 F.C.A.H. Caps. (Scherer)
 Femicin, Tab. (Norcliff Thayer)
 Fendol, Tab. (Buffington)
 Histalets, Tab. (Edw. J. Moore)
 Histogesic, Tab. (Century)
 Histosal #2, Tab. (Ferndale)
 Hi-Temp, Tab. (Saron)
 Hycomine Comp., Tab. (Du Pont)
 Hydrocet, Cap. (Carnrick)
 Isomel, Cap. (Reid-Rowell)
 Kiddies Sialco, Tab. (Foy)
 Kleer, Syr. (Scrip)
 Koryza Tab. (Fellows-Testagar)
 Maxigesic, Cap. (W.F. Merchant)
 Midran, Tab. (Quality Generics)
 Midrin, Cap. (Carnrick)
 Myocalm, Tab. (Parmed)
 Myolate, Tab. (Panray)
 Naldetuss, Syr. (Bristol)
 N-D Gesic, Tab. (Hyrex)
 Neo-Pyranistan, Tab. (Standex Labs.)
 Nokane, Tab. (Wren)
 Novahistine Singlet, Tab. (Merrell Dow)
 Novahistine Sinus Tablets, Tab. (Merrell Dow)
 Nylorac PB, Tab. (Beard)
 Nyquil, Liq. (Vicks)
 Opasal, Cap. (Elder)
 Ornex, Cap. (SmithKline Prods)
 Pamprin, Tab. (Chattem Labs.)
 Panitol H.M.B., Tab. (Wesley)

Panritis, Tab. (Pan Amer.)
 Parafon Forte, Tab. (McNeil)
 Partuss-A, Tab. (Parmed)
 Partuss T.D., Tab. (Parmed)
 Pavadon Elix. (Coastal)
 Pedric, Elix., Wafers (Vale)
 Percogesic, Tab. (Du Pont)
 Phenahist, Cap., Elix. (Amid)
 Phenaphen #2, #3, #4 (Robins)
 Phenaphen-650, Tab. (Robins)
 Phrenilin, Tab. (Carnrick)
 Phrenilin Forte, Cap. (Carnrick)
 Phrenilin w/Codeine, Cap. (Carnrick)
 Presalin, Tab. (Mallard)
 Prodolor, Tab. (Amfre-Grant)
 Proval No. 3, Tab. (Reid-Rowell)
 Renpap, Tab. (Wren)
 Rentuss, Cap., Syr. (Wren)
 Repan, Tab. (Everett)
 Rhinex, D•Lay, Tab. (Lemmon)
 Rhinidrin, Tab. (Central)
 Rhinogesic, Tab. (Vale)
 Rhinspec, Tab. (Lemmon)
 S.A.C. Sinus, Tabs. (Towne)
 Salatin w/Codeine, Tab. (Ferndale)
 Saleto, Tab. (Mallard)
 Saleto-D, Tab. (Mallard)
 Salphenyl, Cap., Liq. (Mallard)
 Santussin, Prep. (Sandia)
 Scotgesic, Cap., Elix. (Scott/Cord)
 Scotuss Pediatric Cough Syr. (Scott/Cord)
 Sedacane, Cap. (Edw. J. Moore)
 Sedalgesic, Tab. (Table Rock)
 Sedragesic, Tab. (Lannett)
 Sialco, Tab. (Foy)
 Sinarest, Tab. (Pharmacraft)
 Sine-Aid, Tab. (McNeil)
 Sine-Off, Prods (SmithKline Prods)
 Sinoze, Tab. (Bowman)
 Sinubid, Tab. (Warner Chilcott)
 Sinulin, Tab. (Carnrick)
 Sinumal, Tab., Liq. (Cook)
 Sinus Tab., Tab. (Zenith)
 Sinutab, Prods. (Warner-Lambert)
 Spantuss, Liq. (Arco)
 Spendrisin, Tab. (Spencer-Mead)
 Stopain, Tab. (Quality Generics)
 Sunril, Cap. (Emko)
 Super-Anahist Tablets (Warner-Lambert)
 Symptomax, Liq. (Reid-Rowell)
 Talacen, Cap. (Winthrop Pharm)
 T-Caps, Cap. (Burlington)
 Tega-Code, Cap. (Ortega)
 Tegapap, Liq., Tab. (Ortega)
 Teragen, Tab. (O'Neal)
 T-Gesic, Cap. (Tennessee)
 Triaminicin, Tab. (Sandoz Consumer)
 Triaprin, Cap. (Dunhall)
 Trind, Syr. (Mead Johnson)
 Trind-DM, Syr. (Mead Johnson)
 Tussagesic, Tab., Susp. (Sandoz Consumer)
 Two-Dyne, Tab. (Hyrex)

Tylenol Preps. (McNeil)
 Tylenol with Codeine, Elixir, Tab. (McNeil)
 Tylox, Cap. (McNeil)
 Valihist, Cap. (Clapp)
 Vicks Daycare, Liq. (Vicks)
 Windolor, Tab. (Winston)
 Wygesic, Tab. (Wyeth)
 Zenex Comp., Cap. (Zenith)

• **ACETAMINOPHEN AND ASPIRIN TABLETS**, U.S.P. XXI.

Use: Analgesic.

• **ACETAMINOPHEN AND CODEINE PHOSPHATE ELIXIR**, U.S.P. XXI.

Use: Analgesic.

• **ACETAMINOPHEN AND DIPHENHYDRAMINE CITRATE TABLETS**, U.S.P. XXI.

Use: Analgesic, antihistaminic.

• **ACETAMINOPHEN ORAL SUSPENSION**, U.S.P. XXI.

Use: Analgesic.

ACETAMINOPHENOL.

See: Acetaminophen.

• **ACETAMINOPHEN ORAL SOLUTION**, U.S.P. XXI.

Use: Analgesic.

ACETAMINOPHEN UNISERTS. (Upsher-Smith)
 Acetaminophen 120 mg./Supp. Ctn. 12s, 50s; 325 mg./Supp. Ctn. 12s, 50s; 650 mg./Supp. Ctn. 12s, 50s, 500s.

Use: Analgesic, antipyretic.

ACETANILID. (Various Mfr.) (Acetylaminobenzene, acetylaniline, antifebrin) N-phenylacetamide cry.

Use: Antipyretic, analgesic.

ACETARSOL.

See: Acetarsone, Tabs.

ACETARSONE. 3-Acetamido-4-hydroxy-phenylarsonic acid. Acetarsol, Acetphenarsine, Amarsan, Dynarsan, Ehrlich 594, Limarsol, Orarsan, Osarsal, Osvarsan, Paroxyl, Stovarsol.

ACETARSONE SALT OF ARECOLINE.

See: Drocarbil.

ACETASOL HC OTIC. (Goldline) Hydrocortisone 1%, acetic acid 2%. Bot. 10 ml.

Use: Antibacterial, antifungal, hydrophilic, anti-inflammatory, anti-pruritic for auditory canal infections.

ACETASOL OTIC. (Goldline) Acetic acid—non-aqueous 2%. Bot. 5 ml.

Use: Antibacterial, antifungal for superficial auditory canal infections.

ACETAZIDE. Acetazolamide, B.A.N.

• **ACETAZOLAMIDE**, U.S.P. XXI. Tab. U.S.P. XXI. N-(5-Sulfamoyl-1,3,4-thiadiazol-2-yl) acetamide. Acetamide, N-[5-(aminosulfonyl)-1,3,4-thiadiazol-2-yl]-.

Use: Carbonic anhydrase inhibitor.

See: Diamox, Tab., Sequels (Lederle)

Hydrazol, Tab. (Softcon)

ACETAZOLAMIDE. B.A.N. 5-Acetamido-1,3,4-thiadiazole-2-sulfonamide.

Use: Treatment of glaucoma.

See: Acetazide

Diamox

• **ACETAZOLAMIDE SODIUM, STERILE**, U.S.P. XXI.

Use: Carbonic anhydrase inhibitor.

See: Diamox, Inj. (Lederle)

ACET-DIA-MER-SULFONAMIDE. Sulfacetamide, sulfadiazine & sulfamerazine.

See: Buffonamide, Susp., Tab. (Reid-Rowell)

Buf-Sul, Tabs. & Susp. (Sheryl)

Cetazine, Tab. (Bowman)

Coco-Diazine, Susp. (Lilly)

Sulfadiazine, sulfamerazine & sulfacetamide, Susp.

ACETEST REAGENT TABLETS. (Ames) Sod.

nitroprusside, disodium phosphate, aminoacetic acid, lactose. Tab. Bot. 100s.

Use: Test for acetone or diacetic acid in urine.

• **ACETIC ACID**, U.S.P. XXI, Glacial, Otic soln., U.S.P. XXI. Diluted, N.F. XVI.

Use: Pharmaceutical aid (acidifying agent).

• **ACETIC ACID IRRIGATION**, U.S.P. XXI. 0.25% solution, (Abbott) 250 ml glass cont.; 250 ml.

Use: Irrigating solution

ACETIC ACID, POTASSIUM SALT. Potassium

Acetate, U.S.P. XXI.

ACETICYL.

See: Acetylsalicylic Acid.

ACETILUM ACIDULATUM.

See: Acetylsalicylic Acid. (Various Mfr.)

• **ACETOHEXAMIDE**, U.S.P. XXI. Tab., U.S.P. XXI. N-(p-acetylphenylsulfonyl)-N'-cyclo-hexylurea. 1-((p-acetylphenyl)sulfonyl)-3-cyclohexylurea. Benzenesulfonamide, 4-acetyl-N-[[cyclohexylamino]carbonyl].

Use: Blood sugar lowering compound, antidiabetic.

See: Dymelor, Tab. (Lilly)

• **ACETOHYDROXAMIC ACID**, U.S.P. XXI. Tab., U.S.P. XXI.

Use: Enzyme inhibitor.

See: Lithostat (Mission)

ACETOJEN, JR. (Jenkins) Acetylsalicylic acid 1.5 gr., phenacetin 1 gr., hyoscyamus ext. 0.0125 gr., (total alkaloids 0.0002 gr.) gelsemium ext. 0.0125 gr./Tab. Bot. 1000s.

ACETOLAX. (Mills) Acetphenylisatin 5 mg., Vit. B-1 1 mg., sod. carboxymethyl cellulose 500 mg./Tab. Bulk Pkg. 1000s.

Use: Laxative.

ACETOL.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOMEROCTOL. Acetato [2-hydroxy-5-(1,1,3,3-tetramethylbutyl) phenyl] mercury. Under study.

Use: Topical antiseptic.

• **ACETONE**, N.F. XVI.

ACETONE or DIACETIC ACID TEST.

See: Acetest, Tab. (Ames)

3-(alpha-ACETONYLBENZYL)-4-HYDROXYCOUMARIN SODIUM SALT. Warfarin Sodium.

ACETOPHEN CAPSULE. (Jenkins) Aspirin 3.5 gr., phenacetin 2.5 gr., caffeine 0.5 gr./Cap. Bot. 1000s.

ACETOPHEN C.T. GREEN. (Jenkins) Aspirin 3.5 gr., phenacetin 2.5 gr., caffeine 0.25 gr./Tab. Bot. 1000s.

• **ACETOPHENAZINE MALEATE**, U.S.P. XXI. Tab., U.S.P. XXI. 2-Acetyl-10- [3-[4-(β-hydroxyethyl)-piperazinyl] propyl] phenothiazine dimaleate. 10-[3-[4-(2-Hydroxyethyl)-1-piperazinyl]-propyl]-phenothiazine-2-yl methyl ketone maleate (1:2).

Use: Tranquillizer.

See: Tindal Maleate, Tab. (Schering)

ACETOPHENETIDIN. Phenacetin, U.S.P. XXI.

Ethoxyacetanilide.

Use: Antipyretic, analgesic.

ACETORPHINE. B.A.N. O³-Acetyl-7,8-dihydro-7α[1(R)-hydroxy-1-methylbutyl]-O⁶-methyl-6, -14-endoethenormorphine.

Use: Narcotic analgesic.

ACETOSAL.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOSALIC ACID.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOSALIN.

See: Acetylsalicylic Acid. (Various Mfr.)

ACETOSULFONE. 4,4'-Diaminodiphenylsulfone-2-N-acetylsulfonamide sodium. (N¹-acetyl-6-sulfanilyl-metanilamido) sodium.

• **ACETOSULFONE SODIUM**. USAN.

Use: Antibacterial.

ACETOXYPHENYLMERCURY.

See: Phenylmercuric acetate.

ACETPHENARSINE.

See: Acetarson, Tab. (City Chemical)

ACETPHENOLISATIN.

See: Oxyphenisatin Acetate Prep.

W/D-Calcium pantothenate, sodium sulfosuccinate.

See: Peri-Pantyl, Tab. (McGregor)

W/Docusate sodium, sodium carboxymethylcellulose.

See: Scrip-Lax, Tab. (Scrip)

ACETRIZOATE SODIUM. 3-Acetamido-2,4,6,-triiodobenzoate sodium.

W/Polyvinylpyrrolidone.

See: Salpix, Vial (Ortho)

ACETRIZOIC ACID. 3-Acetylamino-2,4,6,-triiodobenzoic acid.

See: Acetrizoate, Sod.

ACET-THEOCIN SODIUM.

See: Theophylline Sod. acetate (Various Mfr.)

ACETYL ADALIN.

See: Acetylcarbromal. (Various Mfr.)

p-ACETYLAMINO BENZALDEHYDE THIOSEMICARBAZONE. (Amithiozone, Antib, Berculon A, Benzothiozon, Conteben, Myuizone, Neustab, Tebethion, Thiomicid, Thioparamizone, Thiacetazone).

Use: With streptomycin in the treatment of tuberculosis.

ACETYLAMINO BENZENE.

See: Acetanilid. (Various Mfr.)

N-ACETYL-p-AMINOPHENOL. Acetaminophen, U.S.P. XXI.

ACETYLANILINE.

See: Acetanilid. (Various Mfr.)

ACETYL-BETA-METHYLCHOLINE CHLORIDE.

See: Mecholyl Chloride, Amp. (Baker, J.T.)

ACETYL-BROMO-DIETHYLACETYL-CARBAMIDE.

See: Acetylcarbromal. (Various Mfr.)

ACETYL CARBROMAL. Acetyl adalin,

acetyl bromodiethylacetyl-carbamide. Pow. for mfg.

Use: Sedative.

See: Paxarel, Tab. (Circle)

W/Bromisovalum, scopolamine aminoxide HBr.

See: Tranquinal, Tab. (Barnes-Hind)

• ACETYLCHOLINE CHLORIDE, U.S.P. XXI. For

Ophth. Soln., U.S.P. XXI. Ethanaminium, 2-(acetyloxy)-

N, N, N-trimethyl-, chloride. Choline chloride acetate.

Use: S.C., I.M., I.V. Parasympathomimetic agent & vasodilator. Paroxysmal tachycardia.

See: Miochol Ophthalmic (CooperVision)

ACETYLCHOLINE-LIKE THERAPEUTIC AGENTS.

See: Cholinergic agents.

• ACETYLCYSTEINE, U.S.P. XXI. Sol. U.S.P. XXI. N-

Acetyl-L-cysteine.

Use: Mucolytic agent.

See: Mucomyst, Sol. (Bristol)

• ACETYLCYSTEINE AND ISOPROTERENOL HYDROCHLORIDE INHALATION SOLUTION, U.S.P. XXI.

Use: Mucolytic agent.

ACETYLCYSTEINE. B.A.N. N-Acetyl-L-cysteine. Air-bron

Use: Liquefaction of bronchial secretion.

ACETYLDIGITOXIN. α -Digitoxin monoacetate. Acetyl-tridigitoxose. The natural-occurring cardiac glycoside defined as the alpha acetyl ester of digitoxin.

Use: Digitoxin therapy. Cardiotonic.

ACETYLDIHYDROCODEINONE. Thebacon, B.A.N.**ACETYLIN.**

See: Acetylsalicylic Acid. (Various Mfr.)

ACETYLMETHADOL, I.N.N. Methadyl acetate, B.A.N.**ACETYLPHENYLISATIN.**

See: Oxyphenisatin Acetate.

1-(p-ACETYLPHENYL)SULFONYL-3-CYCLO-HEXYLUREA. Acetohexamide.**ACETYLRÉSOSCINOL.**

See: Resorcinol Monoacetate. (Various Mfr.)

ACETYLSAL.

See: Acetylsalicylic Acid

ACETYLSALICYLATE ALUMINUM.

See: Aluminum Aspirin

ACETYLSALICYLIC ACID. Aspirin, U.S.P. XXI.

Acidum acetylsalicylicum, 2-acetoxybenzoic acid,

Acetillum Acidulatum, Acetophen, Acetol, Acetosol,

Acetosalic Acid, Acetosalin, Acetylal, Acetylsal,

Acylpyrin, Aspro, Helicon, Rhodine, Salacetin,

Salcetogen, Saletin.

See: Aspirin Preps. (Various Mfr.)

ACETYLSALICYLIC ACID, ACETOPHENETIDIN AND CAFFEINE.

See: A.P.C., Prep. (Various Mfr.)

ACETYL SULFAMETHOXYPYRIDAZINE. 3-(N-Acetyl-sulfanilamido)-6-methoxypyridazine.**N-ACETYLSULFANILAMIDE.** (Albucid; p-Aminobenzenesulfonacetamide; Sulfacet; Sulfacetamide, N-Sulfanilylacetamide.)

Use: Sulfonamide therapy.

ACETYL SULFISOXAZOLE. Sulfisoxazole Acetyl, U.S.P. XXI.**ACETYLTANNIC ACID.** Tannic acid acetate.

Use: Antiperistaltic.

AC EYE DROPS. (Walgreen) Tetrahydrozoline HCl

0.05%, zinc sulfate 0.25%. Bot. 0.75 oz.

Use: Allergy, colds, eye relief.

ACHLORHYDRIA DETERMINATION.

See: Diagnex Blue, Prep. (Squibb)

ACHLORHYDRIA THERAPY.

See: Acidol. (Various Mfr.)

Glutamic Acid HCl (Various Mfr.)

Muripsin, Tab. (Norgine)

Normacid, Tab. (Stuart)

ACHOL. (Enzyme Process) Vit. A 4000 U., ketocholanic acids 62 mg./Tab. Bot. 100s, 250s.

Use: Vitamin A supplement and therapy.

ACHROMYCIN. (Lederle) Tetracycline HCl.

Intramuscular: (100 mg., 250 mg., sterile, buffered w/ascorbic acid) Vial. Box 1s.

Intravenous: (250 mg., 500 mg., sterile, buffered w/ascorbic acid) Vial.

Oint. (3%) 30 mg./Gm. Tube 0.5 oz., 1 oz.

Oint. (Ophth.) 10 mg./Gm. 1%. Tube 0.125 oz.

Ophth. Susp. 1%, Plastic dropper bot. 4 ml.

ACHROMYCIN-V. (Lederle) Tetracycline HCl. **Cap.:** 250

mg. Bot. 100s, 1000s, Unit dose 10 $\times$ 10s, Unit of

use 12 $\times$ 20s, 12 $\times$ 28s, 12 $\times$ 40s, 12 $\times$ 100s;

500 mg. Bot. 100s, 1000s; Unit of use 12 $\times$ 20s;

Unit dose 10 $\times$ 10s. **Oral Susp.:** 125 mg./5 ml.

w/methylparaben 0.12%, propylparaben 0.03%.

Bot. 2 oz., 1 pt.

Use: Treatment of infections caused by both gram-negative and gram-positive bacteria.

ACID ACRIFLAVINE.

See: Acriflavine Hydrochloride (Various Mfr.)

ACID CITRATE DEXTROSE SOLUTION.

See: A.C.D. Sol. (Various Mfr.)

ACID CITRATE DEXTROSE ANTICOAGULANT SOLUTION MODIFIED.

See: A-C-D Solution Modified (Squibb)

ACID HISTAMINE PHOSPHATE.

See: Histamine Phosphate. (Various Mfr.)

ACID MANTLE CREME. (Sandoz Consumer)

Aluminum acetate in specially prep. water-soluble hydrophilic cream at pH 4.2. Tube 1 oz., Jar 4 oz., 1 lb.

Use: Skin prophylactic agent in restoring normal skin acidity.

ACIDOL.

See: Betaine Hydrochloride (Various Mfr.)

ACIDOPHILUS W/PECTIN. (Barth's) *Lactobacillus acidophilus* w/natural citrus pectin 100 mg./Cap.

Bot. 100s.

Use: Aid to digestion.

ACID TRYPAFLAVINE.

See: Acriflavine Hydrochloride. (Various Mfr.)

ACIDULATED PHOSPHATE FLUORIDE. (Scherer)

Flavored soln. containing 0.31% fluoride ion in 0.1 molar phosphate. Bot. 64 oz. Office Product.

Use: Dental caries prevention.

ACIDULIN. (Lilly) Glutamic acid HCl 340 mg./Pulvule.

Bot. 100s, 1000s.

Use: Achlorhydria.

• **ACIFRAN.** USAN.

Use: Antihyperlipoproteinemic.

ACIGLUMIN.

See: Glutamic Acid Hydrochloride. (Various Mfr.)

ACI-JEL. (Ortho) Glacial acetic acid 0.921%, ricinoleic acid 0.70%, oxyquinoline sulfate 0.025%, glycerine 5%. Tube 85 Gm. w/dose applicator.

Use: Vaginal acidifier.

ACINITRAZOLE. B.A.N. 2-Acetamido-5-nitrothiazole, Aminotrazole (I.N.N.)

Use: Treatment of trichomoniasis.

See: Trichorad

Tritheon

• **ACITRETIN.** USAN.

Use: Antisporiatic.

• **ACLARUBICIN.** USAN.

Use: Antineoplastic.

ACLOBROM. Propantheline Bromide, B.A.N.

ACLOVATE CREAM. (Glaxo) Alclometasone

dipropionate 0.05%. Tube 15 Gm., 45 Gm.

Use: Topical anti-inflammatory.

ACLOVATE OINTMENT. (Glaxo) Alclometasone

dipropionate 0.05%. Tube 15 Gm., 45 Gm.

Use: Topical anti-inflammatory.

A.C.N. (Person & Covey) Vitamin A (as acetate) 25M

I.U., ascorbic acid 250 mg., niacinamide 25

mg./Tab. Bot. 100s.

Use: Dietary supplement.

ACNAVEEN. (CooperCare)

See: Aveenobar Medicated (CooperCare)

ACNA-VITE. (Cenci) Vit. A 10,000 I.U., C 250 mg., hes-

peridin 50 mg., niacinamide 25 mg./Cap. Bot. 75s.

Use: Acne vulgaris with Vit. A & C deficiency.

ACNE-5. (Goldline) Benzoyl peroxide 5%. Bot. 1 oz.

Use: Acne treatment.

ACNE-10. (Goldline) Benzoyl peroxide 10%. Bot. 1 oz.

Use: Acne treatment.

ACNE-AID CLEANSING BAR. (Stiefel) Neutral soap of

6.3% surfactant blend. Non-medicated. Bar 4.0 oz., 5.8 oz.

Use: Cleanse oily skin and acne.

ACNE-AID CREAM. (Stiefel) Benzoyl peroxide 10% in special cover-up base. Tube, 1.8 oz.

Use: Absorbs oil, hides blemishes.

ACNEDERM. (Lannett) Lotion: Dispersable sulfur

5%, zinc sulfate 1%, zinc oxide 10%, alcohol 21%.

Bot. 2 oz. Soap: Zinc oxide 2%, zinc sulfate 1%,

colloidal sulfur 5%. Cake 3.2 oz.

Use: Skin conditions.

ACNE LOTION. (Weeks & Leo) Benzoyl peroxide

10% in odorless, greaseless, vanishing lotion.

Bot. 2 oz.

Use: Treatment for acne.

ACNO CLEANSER. (Baker/Cummins) Isopropyl

alcohol 60%. Bot. 8 oz.

Use: Antiseptic astringent for the treatment of acne.

ACNO LOTION. (Baker/Cummins) Micronized sulfur

3%, salicylic acid 2%. Bot. 4 oz.

Use: Treatment of acne.

ACNOMEL. (SmithKline Prods) Resorcinol 2%, sulfur 8%, alcohol 11%.

Cream. Tube 1 oz.

Use: Acne therapy.

ACNOPHILL. (Torch) Sulfur 5%, Pot. and zinc sulfides and polysulfides 5%, zinc oxide 10% in hydrophilic vehicle. Plain or tinted. Jar 1 lb. and 2 oz.

Use: Acne treatment.

ACNOTEX. (C & M Pharmacal) Sulfur 8%, salicylic

acid 2.25% in lotion base. Bot. 2 oz.

Use: Anti-acne cover-up lotion.

• **ACODAZOLE HYDROCHLORIDE.** USAN.

Use: Antineoplastic.

ACOTUS. (Whorton) Phenylephrine HCl 5 mg.,

guaiacol glyceryl ether 100 mg., menthol 1 mg.,

alcohol, by vol. 10%/5 cc. Bot. 4 oz., 12 oz.,

gal.

Use: Colds, coughs.

ACR. (Western Research) Ammonium chloride 7.5

gr./Tab. Handicount 28s (36 bags of 28 tab.)

Use: Diuretic.

ACRIFLAVINE. Euflavine, Gonacrine, Neutroflavine

Acriflavine neutral; a mixture of

2, 8-diamino-10-methylacridinium chloride & 2,8-

diaminoacridine. (Lilly) Tab. 1.5 gr. Bot. 100s.

Use: External antiseptic.

ACRIFLAVINE HYDROCHLORIDE. (Various Mfr.)

Hydrochloride form of acriflavine. Acid acriflavine,

acid tryptaflavine, flavine, tryptaflavine.

National Aniline—Pow., Bot. (1, 5, 10, 25, & 50

Gm.). Tab. (1.5 gr.) Bot. 50s & 100s.

Use: External bacteriostatic.

• **ACRISORCIN.** U.S.P. XXI. Cream, U.S.P. XXI. 9-

Aminoacridinium 4-hexylresorcinolate.

See: Akrinol Cream (Schering)

Use: Topical, tinea versicolor. Local antifungal.

• **ACRIVASTINE.** USAN.

Use: Antihistamine.

• **ACRONINE.** USAN. 3, 12-Dihydro-6-methoxy-3,3,12-

trimethyl-7H-pyrano[2,3-c]acridin-7-one. Under

study.

Use: Antineoplastic.

A-C TABLET. (Century) Aspirin 6 gr., caffeine 0.5

gr./Tab.

Bot. 100s, 1000s

Use: Analgesic.

ACTACIN-C SYRUP. (Vangard) Codeine phosphate 10

mg., triprolidine HCl 2 mg., pseudoephedrine HCl

20 mg., guaifenesin 100 mg./ 5 ml. Bot. pt.,

gal.

Use: Antihistamine, decongestant, cough

depressant.

ACTACIN SYRUP. (Vangard) Triprolidine HCl 1.25

mg., pseudoephedrine HCl 30 mg./ 5 ml. Bot. pt.,

gal.

Use: Antihistamine, decongestant.

ACTACIN TABLETS. (Vangard) Triprolidine HCl 2.5 mg., pseudoephedrine HCl 60 mg./Tab. Bot. 100s, 1000s.

Use: Antihistamine/decongestant.

ACTAGEN-C COUGH SYRUP. (Goldline) Triprolidine HCl, pseudoephedrine HCl, codeine phosphate/Bot. 4 oz., pt., gal.

Use: Decongestant, cough suppressor.

ACTAGEN TABLETS. (Goldline) Triprolidine HCl, pseudoephedrine HCl. Bot. 100s, 1000s.

Use: Decongestant.

ACTAL SUSPENSION. (Winthrop Products) Aluminum hydroxide.

Use: Antacid.

ACTAL PLUS TABLETS. (Winthrop Products)

Aluminum hydroxide, magnesium hydroxide.

Use: Antacid, demulcent.

ACTAL TABLETS. (Winthrop Products) Aluminum hydroxide.

Use: Antacid.

ACTAMIN. (Buffington) Acetaminophen 325 mg./Tab.

Sugar, caffeine, lactose, and salt free. Dispens-A-Kit 500s, Aidpaks 100s. Unit boxes 20s. Medipaks 200s.

Use: Analgesic, anti-pyretic.

ACTAMIN EXTRA. (Buffington) Acetaminophen 500

mg./Tab. Sugar, caffeine, lactose, and salt free.

Dispens-A-Kit 500s. Medipaks 200s.

Use: Extra strength nonaspirin analgesic and antipyretic.

ACTAMIN SUPER. (Buffington) Acetaminophen 500

mg. Tab. w/caffeine. Sugar, salt, and lactose free.

Dispens-A-Kits 500s, Medipaks 200s.

Use: Maximum strength analgesic, antipyretic.

• **ACTAPLANIN.** USAN.

Use: Growth stimulant.

ACTH-ACTEST GEL. (Fellows) Repository corticotropin 40 or 80 u./cc. Vial 5 cc.

ACTH. (Adrenocorticotrophic hormone.

Adrenocorticotrop(h)in).

See: Corticotropin, U.S.P.

Mallard—(40 u./ml., 5 ml.)

Forest—(40 u. or 80 u./ml., 5 ml.)

Parke, Davis—25 units/Vial; 40 units/Vial.

Pharmex—(40 u. or 80 u./ml., 5 ml.)

ACTX GEL, PURIFIED. Arcum—40 or 80 units/ml.,

vial 5 ml. Conal.—40 or 80 units/ml., vial 5 ml. Hart

Labs.—40 units/ml., vial 5 ml. Bowman—

Adrenocorticotrophic hormone 40 u., aqueous gelatin

16%, phenol 0.5%/cc. Vial 5 cc.

Use: Repository corticotropin.

Arcum—(40 u. or 80 u./ml., 5 ml.)

Bell—(40 u. or 80 u./ml., 5 ml.)

Bowman—(40 u./ml., 5 ml.)

Hyrex—(40 u. or 80 u./ml., 5 ml.)

Jenkins—(40 u. or 80 u./ml., 5 ml.)

Vortech—40 units/ml., 80 units/ml., Vial 5 ml.

Rocky Mtn. Pharmacal—40 units/ml., 80 units/ml. Vial 5 ml.

Wesley Pharm.—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.

Wyeth—(40 u. or 80 u./ml. or Tubex)

ACTHAR. (Armour) Corticotropin for Injection. Vial 25, 40 u./Vial. (Lyophilized w/gelatin). Pkg. 10s.

Use: I.M., S.C. or I.V. stimulation of adrenal cortex and diseases related to it.

ACTICORT LOTION 100. (Baker/Cummins)

Hydrocortisone alcohol 1% in a specially formulated vehicle. Bot. 2 oz.

Use: Anti-inflammatory & antipruritic.

ACTIDIL. (Burroughs Wellcome) Triprolidine HCl Tab.

2.5 mg./Tab. Bot. 100s. **Syr.** 1.25 mg./5 ml. Bot. 1 pt.

Use: Antihistamine.

ACTIDOSE. (Paddock) Activated charcoal solution 25

Gm./120 ml. or 50 Gm./240 ml.

Use: Orally or through gastric tube to adsorb toxins in GI tract in drug overdose or poisoning emergency.

ACTIDOSE-AQUA. (Paddock) Activated charcoal aqueous

suspension 25 Gm./120 ml. or 50 Gm./240 ml.

Use: Orally or through gastric tube to adsorb toxins in GI tract in drug overdose or poisoning.

ACTIFED. (Burroughs Wellcome) Tab.: Triprolidine

HCl. 2.5 mg., pseudoephedrine HCl. 60 mg./Tab.

Bot. 100s, 1000s. Boxes 12s, 24s, 48s. Unit dose

pack 100s. **Cap.:** Triprolidine HCl 2.5 mg.,

pseudoephedrine HCl 60 mg./Cap. Box 10s, 20s.

Syr.: Triprolidine HCl 1.25 mg., pseudoephedrine

HCl 30 mg./5 ml. Bot. 4 oz., 1 pt.

Use: Common cold, allergic rhinitis.

ACTIFED 12-HOUR CAPSULES. (Burroughs

Wellcome) Triprolidine HCl 5 mg., pseudoephedrine

HCl 120 mg./Cap. Boxes 10s, 20s.

Use: Antihistamine/decongestant.

ACTIFED CAPSULES. (Burroughs Wellcome)

Triprolidine HCl 2.5 mg., pseudoephedrine HCl 60

mg./Cap. Box 10s, 20s.

Use: Nasal decongestant, antihistamine.

ACTIFED WITH CODEINE COUGH SYRUP.

(Burroughs Wellcome) Codeine phosphate 10 mg.,

triproloidine HCl 1.25 mg., pseudoephedrine HCl 30

mg./5 ml. w/alcohol 4.3%. Bot. pt.,

gal.

Use: Treatment of the common cold.

ACTINOMYCIN. B.A.N. Antimicrobial substances with

antitumor activity produced by Streptomyces

antibioticus and Streptomyces chrysomallus.

Sanamycin is Actinomycin C.

ACTINOMYCIN C. Name previously used for

Cactinomycin.

ACTINOMYCIN D. Dactinomycin, U.S.P. XXI.

See: Cosmegen (Merck Sharp & Dohme)

• **ACTINOQUINOL SODIUM.** USAN.

Use: Ultraviolet screen.

ACTINOSPECTOCIN. Name previously used for

Spectinomycin.

ACTIVATED ATTAPULGITE.

W/Aluminum hydroxide, magnesium carbonate

coprecipitate, compressed gel.

See: Hykasil, Cream (Philips Roxane)

W/Polysorbate 80, colloidal sulfur, salicylic acid,

propylene glycol.

See: Sebasorb Lotion (Summers Labs.)

ACTIVATED CHARCOAL TABLETS. (Cowley) 5 gr./Tab. Bot. 1000s.

ACTIVATED 7-DEHYDROCHOLESTEROL.

See: Vitamin D-3 (Various Mfr.)

ACTIVATED ERGOSTEROL.

See: Calciferol.

ACTOMOL. Mebanazine, B.A.N.

ACTOQUINOL SODIUM. 8-Ethoxy-5-quinoline-sulfonic acid sodium salt.

Use: Ultraviolet screen.

ACTOS 50. Corticotrophin, B.A.N.

ACTRAPID HUMAN INSULIN. (Squibb/Novo) Human insulin 100 Units/ml. Vial 10 ml.

Use: Human insulin (rapid action).

ACTRIOL. Epioestril, B.A.N.

ACTYLATE. (Kinney) Ammonium salicylate 80 mg., pot. salicylate 80 mg., strontium salicylate 80 mg., pot. para-aminobenzoate 0.32 Gm., ascorbic acid 20 mg./Tab. Bot. 100s.

Use: Rheumatoid arthritis.

ACUCRON. (Seatrice) Acetaminophen 300 mg., salicylamide 200 mg., phenyltoloxamine 20 mg./Tab. Bot. 100s, 1000s, 5000s.

Use: Antihistaminic, analgesic.

• **ACYCLOVIR.** USAN. 9-(2-hydroxyethoxymethyl)-guanine.

Use: Management of initial herpes genitalis and mucocutaneous herpes simplex virus infections.

See: Zovirax Caps., Oint. (Burroughs Wellcome)

• **ACYCOLOVIR SODIUM.** USAN.

Use: Antiviral.

ACYLPYRIN.

See: Aspirin

ADALAN LANATABS. (Lannett) Amobarbital 50 mg., homatropine methylbromide 7.5 mg., methamphetamine HCl 10 mg./T.D. Tab. Bot. 100s, 1000s.

ADALAT. (Miles Pharm) Nifedipine 10 mg. or 20 mg./Cap. Bot. 100s, 300s. Unit-dose 100s.

Use: Calcium channel blocker for angina pectoris.

ADAMANTANAMINE HCl. Amantadine HCl. Anti-flu capsule. This drug is thought to protect cells against entry of the flu virus without actually destroying the virus. Also used in the treatment of Parkinson's disease.

See: Symmetrel, Cap., Syr. (Du Pont)

ADANON. Methadone, B.A.N.

ADAPETTES FOR SENSITIVE EYES. (Alcon Surgical) EDTA 0.1%, sorbic acid 0.1%. Pkg. 15 ml.

Use: Contact lens rewetting and lubricating solution.

ADAPIN. (Pennwalt) Doxepin HCl 10 mg., 25 mg. 50 mg., 75 mg., 100 mg./Cap. Bot. 30s, 100s, 1000s. Strip-pack 100s. 25 mg., 50 mg./Cap. also available in 5000s.

Use: Antianxiety, antidepressant.

ADAPT. (Alcon Surgical) Povidone, EDTA 0.1%, thimerosal 0.004%. Bot. 15 ml.

Use: Contact lens cushioning solution.

ADATUSS. (Merchant) Hydrocodone bitartrate 5 mg., guaifenesin 100 mg., alcohol 5%/5 ml. Syr. Bot. pt., gal.

Use: Cough relief.

A.D.C. DROPS. (CMC) Vit. A 3000 I.U., D 400 I.U., C 60 mg./0.6 cc. Bot. 60 cc.

ADCORTYL. Triamcinolone, B.A.N.

ADEECON. (CMC) Vit. A 5000 I.U., D 1000 I.U./Cap. Bot. 1000s.

Use: Supplement.

ADEFLOL CHEWABLE. (Upjohn) Fluoride 0.5 mg. or 1 mg., Vitamin A 4000 I.U., D 400 I.U., C 75 mg., B-1 2 mg., B-2 2 mg., niacinamide 18 mg., B-6 1 mg., calcium pantothenate 5 mg., B-12 2 mcg./Chewable Tab. **0.5 mg.** Bot. 100s. **1 mg.** Bot. 100s.

Use: Prevention of dental caries.

ADEFLOL DROPS. (Upjohn) Vit. A 2000 I.U., D 400 I.U., C 50 mg., B-6 1 mg., fluoride (as sodium fluoride) 0.5 mg./0.6 ml. Bot. 50 ml. w/dropper.

Use: Prevention of dental caries.

ADEFLOL M TABLETS. (Upjohn) Vit. A 6000 I.U., D 400 I.U., B-1 1.5 mg., B-2 2.5 mg., C 100 mg., niacinamide 20 mg., Vit. B-6 10 mg., calcium pantothenate 10 mg., Vit. B-12 2 mcg., folic acid 0.1 mg., fluoride (as sodium fluoride) 1 mg., calcium (as calcium carbonate) 250 mg., iron 30 mg./Tab. Bot. 100s.

Use: Dietary supplement.

• **ADENINE.** U.S.P. XXI. 6-Aminopurine.

ADENO TWELVE GEL INJECTION. (Fellows)

Adenosine-5-monophosphate 25 mg., methionine 25 mg., niacin 10 mg./cc. Vial 10 cc.

ADENOCREST. (Nutrition) Adenosine-5-monophosphate sod. 25 mg., nicotinic acid as sod. salt 25 mg., gelatin 100 mg. w/benzyl alcohol 1.5%/cc. Vials 10 cc.

Use: Capillary and arterial vasodilation.

ADENOLIN FORTE. (Lincoln) Adenosine-5-monophosphate, 25 mg., methionine 25 mg., niacin 10 mg./cc. Vial 15 cc.

Use: Arthritis, bursitis, tendinitis & other degenerative diseases.

• **ADENOSINE.** F.D.A. 6-Amino-9-beta-D-Ribofuranosyl-9H-purine.

ADENOSINE IN GELATIN. (Fellows) **Forte:**

Adenosine-5-monophosphate 50 mg./cc. **Super:**

Adenosine-5-monophosphate 100 mg./cc.

ADENOSINE-5-MONOPHOSPHATE AS THE SODIUM SALT.

See: Adenocrest, Amp. (Nutrition)

• **ADENOSINE PHOSPHATE.** USAN. 5-Adenylic acid. B.A.N. Adenosine-5'-(dihydrogen phosphate). Adenosine monophosphate, AMP.

Use: Nutrient.

See: Adenyl

Cobalazine, Inj. (Keene)

My-B-Den (Miles Pharm)

ADENYL. Adenosine Phosphate, B.A.N.

ADEPSINE OIL.

See: Petrolatum Liquid (Various Mfr.)

ADICILLIN. B.A.N. 6-[p(+)-5-Amino-5-carboxy-valeramido]penicillanic acid. Penicillin N.

Use: Antibiotic.