

Wiley Classics Library

GRIFFITHS & HARRIS

代数几何原理

Principles of
Algebraic Geometry



www.wiley.com

世界图书出版公司
www.wpcbj.com.cn

PRINCIPLES OF ALGEBRAIC GEOMETRY

PHILLIP GRIFFITHS and JOSEPH HARRIS

Harvard University

Wiley Classics Library Edition Published 1994



A WILEY-INTERSCIENCE PUBLICATION

JOHN WILEY & SONS, INC.

New York • Chichester • Brisbane • Toronto • Singapore

图书在版编目 (CIP) 数据

代数几何原理=Principles of Algebraic Geometry:
英文 / (英) 格里菲思 (Griffiths, P.), (英) 哈里
斯 (Harris, J.) 著. —北京: 世界图书出版公司北京公
司, 2007.5

ISBN 978-7-5062-8277-2

I.代… II.①格… ②哈… III.代数几何—英文 IV.
0187

中国版本图书馆CIP数据核字 (2007) 第064418号

书 名: Principles of Algebraic Geometry

作 者: P. Griffiths, J. Harris

中 译 名: 代数几何原理

责任编辑: 焦小浣

出 版 者: 世界图书出版公司北京公司

印 刷 者: 北京世图印刷厂

发 行: 世界图书出版公司北京公司 (北京朝内大街 137 号 100010)

联系电话: 010-64015659, 64038348

电子信箱: kjsk@vip.sina.com

开 本: 24 开

印 张: 35

版 次: 2007 年 5 月第 1 次印刷

版权登记: 图字:01-2007-1464

书 号: 978-7-5062-8277-2/O · 560

定 价: 78.00 元

世界图书出版公司北京公司已获得 Wiley 授权在中国大陆独家重印发行

PREFACE

Algebraic geometry is among the oldest and most highly developed subjects in mathematics. It is intimately connected with projective geometry, complex analysis, topology, number theory, and many other areas of current mathematical activity. Moreover, in recent years algebraic geometry has undergone vast changes in style and language. For these reasons there has arisen about the subject a reputation of inaccessibility. This book gives a presentation of some of the main general results of the theory accompanied by—and indeed with special emphasis on—the applications to the study of interesting examples and the development of computational tools.

A number of principles guided the preparation of the book. One was to develop only that general machinery necessary to study the concrete geometric questions and special classes of algebraic varieties around which the presentation was centered.

A second was that there should be an alternation between the general theory and study of examples, as illustrated by the table of contents. The subject of algebraic geometry is especially notable for the balance provided on the one hand by the intricacy of its examples and on the other by the symmetry of its general patterns; we have tried to reflect this relationship in our choice of topics and order of presentation.

A third general principle was that this volume should be self-contained. In particular any “hard” result that would be utilized should be fully proved. A difficulty a student often faces in a subject as diverse as algebraic geometry is the profusion of cross-references, and this is one reason for attempting to be self-contained. Similarly, we have attempted to avoid allusions to, or statements without proofs of, related results. This book is in no way meant to be a survey of algebraic geometry, but rather is designed to develop a working facility with specific geometric questions. Our approach to the subject is initially analytic: Chapters 0 and 1 treat the basic techniques and results of complex manifold theory, with some emphasis on results applicable to projective varieties. Beginning in Chapter 2 with the theory of Riemann surfaces and algebraic curves, and continuing in Chapters 4 and 6 on algebraic surfaces and the quadric line

PREFACE

complex, our treatment becomes increasingly geometric along classical lines. Chapters 3 and 5 continue the analytic approach, progressing to more special topics in complex manifolds.

Several important topics have been entirely omitted. The most glaring are the arithmetic theory of algebraic varieties, moduli questions, and singularities. In these cases the necessary techniques are not fully developed here. Other topics, such as uniformization and automorphic forms or monodromy and mixed Hodge structures have been omitted, although the necessary techniques are for the most part available.

We would like to thank Giuseppe Canuto, S. S. Chern, Maurizio Cornalba, Ran Donagi, Robin Hartshorne, Bill Hoffman, David Morrison, David Mumford, Arthur Ogus, Ted Shifrin, and Loring Tu for many fruitful discussions; Ruth Suzuki for her wonderful typing; and the staff of John Wiley, especially Beatrice Shube, for enormous patience and skill in converting a very rough manuscript into book form.

PHILLIP GRIFFITHS
JOSEPH HARRIS

May 1978
Cambridge, Massachusetts

目 录

第 0 章 基础知识	1
1. 多复变初步	
柯西公式及应用	2
多变量	6
魏尔斯特拉斯定理及其推论	7
解析簇	12
2. 复流形	
复流形	14
子流形与子簇	18
De Rham 和 Dolbeault 上同调	23
复流形上的积分	27
3. 层和上同调	
起源：米塔—列夫勒问题	34
层	35
层的上同调	38
De Rham 定理	43
Colbeault 定理	45
4. 流形的拓扑	
闭链的相交	49
庞加莱对偶	53
解析闭链的相交	60
5. 向量丛、联络和曲率	
全纯复向量丛	66
度量、联络和曲率	71
6. 紧致复流形的调和理论	

霍奇定理	80
霍奇定理 I 的证明——局部理论	84
霍奇定理 II 的证明——全局理论	92
霍奇定理的应用	100
7. Kähler 流形	
Kähler 条件	106
霍奇恒等式和霍奇分解	111
Lefschetz 分解	118
第 1 章 复代数簇	128
1. 除子与线丛	
除子	129
线丛	132
线丛的陈类	139
2. 消灭定理及推论	
小平消灭定理	148
超平面截面的 Lefschetz 定理	156
定理	159
$(1, 1)$ 类的 Lefschetz 定理	161
3. 代数簇	
解析簇和代数簇	164
簇的次数	171
代数簇的切空间	175
4. 小平嵌入定理	
线丛和到投影空间的映射	176
胀开	182
小平定理的证明	189
5. 格拉斯曼理论	
定义	193
胞腔分解	194
Schubert 微积分	197
万有丛	207
Plücker 嵌入	209

第 2 章 Riemann 曲面和代数曲线	212
1. 预备知识	
Riemann 曲面的嵌入	213
Riemann—Hurwitz 公式	216
亏格公式	219
$G=0, 1$ 的情况	222
2. 阿贝尔定理	
阿贝尔定理——第一种描述	224
第一互反定律及推论	229
阿贝尔定理——第二种描述	232
雅可比反演问题	235
3. 曲线的线性系统	
互反定律 II	240
Riemann—Roch 公式	243
典范曲线	246
特殊线性系统 I	249
超椭圆曲线与黎曼点数	253
特殊线性系统 II	259
4. Plücker 公式	
伴随曲线	263
分歧	264
广义 Plücker 公式 I	268
广义 Plücker 公式 II	271
Weierstrass 点	273
平面曲线的 Plücker 公式	277
5. 对应	
定义和公式	282
空间曲线的几何性	290
特殊线性系统 III	298
6. 复环面和 Abel 簇	
黎曼条件	300
复环面上的线丛	307

函数	317
Abel 簇上的群结构	324
固有公式	326
7. 曲线及曲线的行列式	
初步知识	333
黎曼定理	338
黎曼奇异定理	341
特殊线性系统 IV	349
Torelli 定理	359
第 3 章 深入技巧	364
1. 分布与流	
定义; 幂公式	366
平滑与整齐	373
流的上同调	382
2. 流在复分析上的应用	
解析簇相关的流	385
解析簇的相交数	392
莱维扩展与常态映射定理	395
3. 陈类	
定义	400
高斯博内公式	409
关于全纯向量丛陈类讨论	416
4. 不动点与剩余公式	
莱夫谢茨不动点公式	419
全纯莱夫谢茨不动点公式	422
博特剩余公式	426
广义 Hirzebruch—Riemann—Roch 公式	435
5. 谱序列及其应用	
滤子化双重复形的谱序列	438
超上同调	445
二类微分	454
勒雷谱序列	462

第 4 章 曲面	469
1. 初步知识	
相交数、从属公式与 Riemann—Roch	470
胀开与收缩	473
二次曲面	478
三次曲面	480
2. 有理映射	
有理和双有理映射	489
曲线与代数面	498
面之间双有理映射的结构	510
3. 有理曲面 I	
诺特引理	513
有理直纹面	514
广义有理曲面	520
极小度曲面	522
最大类曲线	527
施泰纳构造	528
Enriques—Petri 定理	533
4. 有理曲面 II	
Castelnuovo—Enriques 定理	536
Enriques 曲面	541
修正的三次曲面	545
中两个二次曲面的相交	550
5. 无理曲面	
阿尔巴内塞映射	552
无理直纹曲面	553
椭圆曲面简介	564
小平数和分类定理 I	572
分类定理 II	582
K—3 曲面	590
诺特曲面	594
6. 诺特公式	

平滑超平面的诺特公式	600
胀开子流形	602
曲面的寻常奇点	611
一般曲面的诺特公式	618
几个例子	628
曲面的孤立奇点	636
第 5 章 留数 (残数)	647
1. 留数的基本性质	
定义和上同调解释	649
整体留数定理	655
变换法则与局部对偶性	656
2. 留数的应用	
相交数	662
有限全纯映射	667
平面投影几何中的应用	670
3. 交换同调代数应用初步	
交换代数	678
同调代数	682
科斯居尔复形及其应用	687
凝聚层的简短游程	695
4. 整体对偶	
整体扩展	705
广义整体对偶定理解释	707
整体扩展和带孤立零点的向量场	708
整体对偶和曲面上点的剩余	712
模的扩张	722
曲面上的点和秩 2 向量丛	726
留数和向量丛	729
第 6 章 二次线丛	733
1. 二次曲面初步	
二次曲面的秩	734

二次曲面中的线性空间	735
二次曲面的线性系统	741
五个锥线论问题	749
2. 二次线丛介绍	
格拉斯曼 $G(2, 4)$ 几何	756
线丛	759
二次线丛和伴随库默尔曲面 I	762
二次线丛的奇异线	767
两个构形	773
3. 二次线丛的线	
二次线丛的线簇	778
线簇上的曲线	780
两个修正构形	784
群法则	787
4. 二次线丛: Reprise	
二次线丛和伴随库默尔曲面 II	791
二次线丛的有理性	796
索引	805

0

基础知识

In this chapter we sketch the foundational material from several complex variables, complex manifold theory, topology, and differential geometry that will be used in our study of algebraic geometry. While our treatment is for the most part self-contained, it is tacitly assumed that the reader has some familiarity with the basic objects discussed. The primary purpose of this chapter is to establish our viewpoint and to present those results needed in the form in which they will be used later on. There are, broadly speaking, four main points:

1. *The Weierstrass theorems and corollaries*, discussed in Sections 1 and 2. These give us our basic picture of the local character of analytic varieties. The theorems themselves will not be quoted directly later, but the picture—for example, the local representation of an analytic variety as a branched covering of a polydisc—is fundamental. The foundations of local analytic geometry are further discussed in Chapter 5.

2. *Sheaf theory*, discussed in Section 3, is an important tool for relating the analytic, topological, and geometric aspects of an algebraic variety. A good example is the *exponential sheaf sequence*, whose individual terms \mathbb{Z} , \mathcal{O} , and \mathcal{O}^* reflect the topological, analytic, and geometric structures of the underlying variety, respectively.

3. *Intersection theory*, discussed in Section 4, is a cornerstone of classical algebraic geometry. It allows us to treat the incidence properties of algebraic varieties, a priori a geometric question, in topological terms.

4. *Hodge theory*, discussed in Sections 6 and 7. By far the most sophisticated technique introduced in this chapter, Hodge theory has, in the present context, two principal applications: first, it gives us the *Hodge decomposition* of the cohomology of a Kähler manifold; then, together with the formalism introduced in Section 5, it gives the vanishing theorems of the next chapter.

1. 多复变初步

Cauchy's Formula and Applications

NOTATION. We will write $z = (z_1, \dots, z_n)$ for a point in \mathbb{C}^n , with

$$z_i = x_i + \sqrt{-1} y_i;$$

$$\|z\|^2 = (z, z) = \sum_{i=1}^n |z_i|^2.$$

For U an open set in \mathbb{C}^n , write $C^\infty(U)$ for the set of C^∞ functions defined on U ; $C^\infty(\bar{U})$ for the set of C^∞ functions defined in some neighborhood of the closure \bar{U} of U .

The cotangent space to a point in $\mathbb{C}^n \cong \mathbb{R}^{2n}$ is spanned by $\{dx_i, dy_i\}$; it will often be more convenient, however, to work with the complex basis

$$dz_i = dx_i + \sqrt{-1} dy_i, \quad d\bar{z}_i = dx_i - \sqrt{-1} dy_i$$

and the dual basis in the tangent space

$$\frac{\partial}{\partial z_i} = \frac{1}{2} \left(\frac{\partial}{\partial x_i} - \sqrt{-1} \frac{\partial}{\partial y_i} \right), \quad \frac{\partial}{\partial \bar{z}_i} = \frac{1}{2} \left(\frac{\partial}{\partial x_i} + \sqrt{-1} \frac{\partial}{\partial y_i} \right).$$

With this notation, the formula for the total differential is

$$df = \sum_i \frac{\partial f}{\partial z_i} dz_i + \sum_j \frac{\partial f}{\partial \bar{z}_j} d\bar{z}_j.$$

In one variable, we say a C^∞ function f on an open set $U \subset \mathbb{C}$ is *holomorphic* if f satisfies the Cauchy-Riemann equations $\partial f / \partial \bar{z} = 0$. Writing $f(z) = u(z) + \sqrt{-1} v(z)$, this amounts to

$$\operatorname{Re} \left(\frac{\partial f}{\partial \bar{z}} \right) = \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} = 0,$$

$$\operatorname{Im} \left(\frac{\partial f}{\partial \bar{z}} \right) = \frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} = 0.$$

We say f is *analytic* if, for all $z_0 \in U$, f has a local series expansion in $z - z_0$, i.e.,

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n$$

in some disc $\Delta(z_0, \epsilon) = \{z : |z - z_0| < \epsilon\}$, where the sum converges absolutely and uniformly. The first result is that f is analytic if and only if it is holomorphic; to show this, we use the

Cauchy Integral Formula. For Δ a disc in \mathbb{C} , $f \in C^\infty(\bar{\Delta})$, $z \in \Delta$,

$$f(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w) dw}{w - z} + \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} \frac{\partial f(w)}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w - z},$$

where the line integrals are taken in the counterclockwise direction (the fact that the last integral is defined will come out in the proof).

Proof. The proof is based on Stokes' formula for a differential form with singularities, a method which will be formalized in Chapter 3. Consider the differential form

$$\eta = \frac{1}{2\pi\sqrt{-1}} \frac{f(w)dw}{w-z};$$

we have for $z \neq w$

$$\frac{\partial}{\partial \bar{w}} \left(\frac{1}{w-z} \right) = 0$$

and so

$$d\eta = -\frac{1}{2\pi\sqrt{-1}} \frac{\partial f(w)}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w-z}.$$

Let $\Delta_\varepsilon = \Delta(z, \varepsilon)$ be the disc of radius ε around z . The form η is C^∞ in $\Delta - \Delta_\varepsilon$, and applying Stokes' theorem we obtain

$$\begin{aligned} \frac{1}{2\pi\sqrt{-1}} \int_{\partial \Delta_\varepsilon} \frac{f(w)dw}{w-z} &= \frac{1}{2\pi\sqrt{-1}} \int_{\partial \Delta} \frac{f(w)dw}{w-z} \\ &\quad + \frac{1}{2\pi\sqrt{-1}} \int_{\Delta - \Delta_\varepsilon} \frac{\partial f}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w-z}. \end{aligned}$$

Setting $w - z = re^{i\theta}$,

$$\frac{1}{2\pi\sqrt{-1}} \int_{\partial \Delta_\varepsilon} \frac{f(w)dw}{w-z} = \frac{1}{2\pi} \int_0^{2\pi} f(z + \varepsilon e^{i\theta}) d\theta,$$

which tends to $f(z)$ as $\varepsilon \rightarrow 0$; moreover,

$$dw \wedge d\bar{w} = -2\sqrt{-1} dx \wedge dy = -2\sqrt{-1} r dr \wedge d\theta$$

so

$$\left| \frac{\partial f(w)}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w-z} \right| = 2 \left| \frac{\partial f}{\partial \bar{w}} dr \wedge d\theta \right| \leq c |dr \wedge d\theta|.$$

Thus $(\partial f / \partial \bar{w})(dw \wedge d\bar{w}) / (w - z)$ is absolutely integrable over Δ , and

$$\int_{\Delta_\varepsilon} \frac{\partial f}{\partial \bar{w}} \frac{dw \wedge d\bar{w}}{w-z} \rightarrow 0$$

as $\varepsilon \rightarrow 0$; the result follows.

Q.E.D.

Now we can prove the

Proposition. For U an open set in \mathbb{C} and $f \in C^\infty(U)$, f is holomorphic if and only if f is analytic.

Proof. Suppose first that $\partial f/\partial \bar{z}=0$. Then for $z_0 \in U$, ε sufficiently small, and z in the disc $\Delta=\Delta(z_0, \varepsilon)$ of radius ε around z_0 ,

$$\begin{aligned} f(z) &= \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{w-z} \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{(w-z_0)-(z-z_0)} \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{(w-z_0)\left(1-\frac{z-z_0}{w-z_0}\right)} \\ &= \sum_{n=0}^{\infty} \left(\frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{(w-z_0)^{n+1}} \right) (z-z_0)^n; \end{aligned}$$

so, setting

$$a_n = \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{(w-z_0)^{n+1}},$$

we have

$$f(z) = \sum_{n=0}^{\infty} a_n (z-z_0)^n$$

for $z \in \Delta$, where the sum converges absolutely and uniformly in any smaller disc.

Suppose conversely that $f(z)$ has a power series expansion

$$f(z) = \sum_{n=0}^{\infty} a_n (z-z_0)^n$$

for $z \in \Delta=\Delta(z_0, \varepsilon)$. Since $(\partial/\partial \bar{z})(z-z_0)^n=0$, the partial sums of the expansion satisfy Cauchy's formula without the area integral, and by the uniform convergence of the sum in a neighborhood of z_0 the same is true of f , i.e.,

$$f(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{f(w)dw}{w-z}.$$

We can then differentiate under the integral sign to obtain

$$\frac{\partial}{\partial \bar{z}} f(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\partial\Delta} \frac{\partial}{\partial \bar{z}} \left(\frac{f(w)}{w-z} \right) dw = 0,$$

since for $z \neq w$

$$\frac{\partial}{\partial \bar{z}} \left(\frac{1}{w-z} \right) = 0.$$

Q.E.D.

We prove a final result in one variable, that given a C^∞ function g on a disc Δ the equation

$$\frac{\partial f}{\partial \bar{z}} = g$$

can always be solved on a slightly smaller disc; this is the

$\bar{\partial}$ -Poincaré Lemma in One Variable. Given $g(z) \in C^\infty(\bar{\Delta})$, the function

$$f(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} \frac{g(w)}{w-z} dw \wedge d\bar{w}$$

is defined and C^∞ in Δ and satisfies

$$\frac{\partial f}{\partial \bar{z}} = g.$$

Proof. For $z_0 \in \Delta$ choose ε such that the disc $\Delta(z_0, 2\varepsilon) \subset \Delta$ and write

$$g(z) = g_1(z) + g_2(z),$$

where $g_1(z)$ vanishes outside $\Delta(z_0, 2\varepsilon)$ and $g_2(z)$ vanishes inside $\Delta(z_0, \varepsilon)$. The integral

$$f_2(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} g_2(w) \frac{dw \wedge d\bar{w}}{w-z}$$

is well-defined and C^∞ for $z \in \Delta(z_0, \varepsilon)$; there we have

$$\frac{\partial}{\partial \bar{z}} f_2(z) = \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} \frac{\partial}{\partial \bar{z}} \left(\frac{g_2(w)}{w-z} \right) dw \wedge d\bar{w} = 0.$$

Since $g_1(z)$ has compact support, we can write

$$\begin{aligned} \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} g_1(w) \frac{dw \wedge d\bar{w}}{w-z} &= \frac{1}{2\pi\sqrt{-1}} \int_{\mathbb{C}} g_1(w) \frac{dw \wedge d\bar{w}}{w-z} \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\mathbb{C}} g_1(u+z) \frac{du \wedge d\bar{u}}{u}, \end{aligned}$$

where $u = w - z$. Changing to polar coordinates $u = re^{i\theta}$ this integral becomes

$$f_1(z) = -\frac{1}{\pi} \int_{\mathbb{C}} g_1(z + re^{i\theta}) e^{-i\theta} dr \wedge d\theta,$$

which is clearly defined and C^∞ in z . Then

$$\begin{aligned} \frac{\partial f_1(z)}{\partial \bar{z}} &= -\frac{1}{\pi} \int_{\mathbb{C}} \frac{\partial g_1}{\partial \bar{z}}(z + re^{i\theta}) e^{-i\theta} dr \wedge d\theta \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{\Delta} \frac{\partial g_1}{\partial \bar{w}}(w) \frac{dw \wedge d\bar{w}}{w-z}; \end{aligned}$$