

AMERICAN DRUG INDEX

1958

CHARLES O. WILSON, Ph.D.

*Professor of Pharmaceutical Chemistry, College of Pharmacy,
University of Texas*

AND

TONY EVERETT JONES, Ph.D.

*Assistant Professor of Pharmaceutical Chemistry, College of Pharmacy,
University of Colorado; Director of
Pharmaceutical Research, Carbisulphoil Company*

J. B. LIPPINCOTT COMPANY

Philadelphia and Montreal

PREFACE

The American Drug Index has been prepared for the identification and correlation of the many pharmaceuticals available to the medical and allied professions. The need for this index has become acute as the number of drugs and drug products in all their tremendous variety has multiplied.

The organization of the index is fundamentally alphabetical with extensive cross-indexing. Names listed are generic; brand (trade-mark, proprietary or specialty), chemical, U.S.P., N.F., N.N.R. and A.D.R. Synonyms that are in general use also are included. All names used for a pharmaceutical occur in alphabetical order with the pertinent data given under the brand name and the name by which it is made available. Data included are generic name, chemical name, manufacturer, pharmaceutical forms, size, dosage and use. Generic names occur in alphabetical order and here the recognition of the drug is indicated by U.S.P. (United States Pharmacopeia), N.F. (National Formulary), N.N.R. (New and Nonofficial Remedies) and A.D.R. (Accepted Dental Remedies). The information is in accord with the latest editions of these books: U.S.P. XV, N.F. X, N.N.R. 1957, A.D.R. 1957.

Because of the multiplicity of specialty names (brand names) used for the same therapeutic agent or the same combination of therapeutic agents, it was apparent that some correlation could be done. As an example of this, please turn to Aluminum Hydroxide Gel. Here under the generic name are given the brand names. Following are combinations of aluminum hydroxide gel organized in a manner to point out relationships among the many products. Reference then is made to the specialty name or names having the indicated composition. Under the specialty name are given manufacturer, composition, available forms, sizes, dosage and use.

The American Drug Index should be most useful to physicians, pharmacists, dentists, nurses, and to sales personnel, students and teachers in the fields incorporating pharmaceuticals. The cross-indexing feature permits the finding of drugs or drug combinations when only one major ingredient is known. For example, a combination of aluminum hydroxide gel and magnesium trisilicate is available. This combination can be found by looking under the name of either of the two ingredients, and in each case the specialty names are given.

CHARLES O. WILSON
TONY E. JONES

ABSORBENT OINTMENT BASE

A.A.S. COMPOUND. (Tilden) Aspirin 5 gr., nux vomica ext. 1/10 gr., hyoscyamus ext. 1/6 gr., gelatinum tr. 3 min., cascara sagrada ext. 1 gr./Tab. Bot. 100s, 1000s.
Use: Analgesic.

ABAC. (Massengill) Aspirin 5 gr., caffeine 1/4 gr., Pot. bromide 1/4 gr., Sod. bromide 1/4 gr./Tab. or Cap. Bot. 100s, 1000s.
Use: (1-2 q. 3 h.) Analgesic sedative.

ABAC FOR CHILDREN. (Massengill) Aspirin 1 gr., caffeine 1/10 gr., Pot. bromide 1/20 gr., Sod. bromide 1/20 gr., aromatics q.s./Tab. Bot. 500s, 1000s.
Use: (1 q. 4 h.) Analgesic.

ABACTAL. (Massengill) Aspirin 5 gr., caffeine 1/4 gr., Pot. bromide 1/4 gr., Sod. bromide 1/4 gr., phenobarbital 1/2 gr./Tab. Bot. 100s, 1000s.
Use: (1-2 t.i.d.) Analgesic sedative.

ABASIN. (Winthrop Labs.) Acetylcarbamol, N-acetyl-N-bromodiethylacetylcarbamide. Tab. (4 gr.) Bot. 100s.
Use: (1-2 t.i.d.) Sedative.

ABBOCILLIN 800 M., A.D.R. (Abbott) Procaine penicillin G 600,000 u., buffered Pot. penicillin G 200,000 u. for aq. inj. Vial, 1 cc., Box 25s, 5 cc., 1s.
Use: I.M.; bacterial infections.

ABBOCILLIN-DC., A.D.R. (Abbott) Procaine penicillin G in aq. susp. 600,000 u. 1 cc. Abboject disposable syr. w/needle, 1s, 25s.
Use: I.M.; penicillin therapy.

ABBO-VAC. See: A.C.D. Solution. (Abbott).
A-B-C-D-G VITAMINS. (C. D. Smith) Vit. A 2500 u., ascorbic acid 25 mg., thiamine HCl 1 mg., riboflavin 1 mg., Vit. D 250 u., niacinamide 10 mg., Cal. pantothenate 0.5 mg./Tab. Bot. 100s, 1000s.
Use: Multivitamin therapy.

ABDEC DROPS. (Parke, Davis) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 0.4 mg., B-6 1 mg., Sod. pantothenate 2 mg., niacinamide 5 mg., Vit. C 50 mg./0.6 cc. Bot. w/ dropper 15 cc., 50 cc.
Use: Vitamin deficiencies.

ABDEC KAPSEALS. (Parke, Davis) Vit. A 10,000 u., D 1000 u., B-1 5 mg., B-2 3 mg., B-6 1.5 mg., B-12 2 mcg., Sod. pantothenate 5 mg., niacinamide 25 mg., Vit. C 75 mg., mixed tocopherols 5 mg./Cap. Bot. 50s, 100s, 250s, 1000s.
Use: (1 daily) Multivitamin therapy.

A-B-D-O. (Blue Line) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 2 mg./Cap. Bot. 100s.
Use: (1 daily) Dietary supplement.

A-B-D-G VITAMIN. (Horton & Converse) Vit. A 5000 u., D 1000 u., B-1 333 u., riboflavin 2000 mcg./Tab. Bot. 100s, 250s, 500s, 1000s.

Use: (1 daily) Multivitamin therapy.

A-B-D-G VITAMINS. (Upjohn) Vit. A 2500 u., D 250 u., B-1 1.5 mg., B-2 1 mg., niacinamide 10 mg./Cap. Bot. 50s, 100s, 250s, 1000s.
Use: (1 daily) Multivitamin therapy.

ABDOL W/Minerals. (Parke, Davis) Vit. C 50 mg., B-2 2.5 mg., B-12 1 mcg., B-6 0.5 mg., B-1 2.5 mg., niacinamide 20 mg., Vit. A 5000 u., D 1000 u., Cal. pantothenate 2.5 mg., folic acid 0.1 mg., iodine 0.15 mg., Mang. 1 mg., cobalt 0.1 mg., Pot. 5.0 mg., molybdenum 0.2 mg., iron 15 mg., copper 1 mg., zinc 0.5 mg., Mag. 1 mg., Cal. 44.0 mg., Phos. 34 mg./Cap. Bot. 100s, 250s.
Use: (1 daily) Vitamin-mineral deficiency states.

ABDOL W/Vit. C Cap. (Parke, Davis) Vit. A 5000 u., D 1000 u., B-1 2.5 mg., B-2 2.5 mg., B-6 0.5 mg., B-12 1 mcg., niacinamide 20 mg., Vit. C 50 mg., pantothenic acid (as Cal. salt) 5 mg./Cap. Bot. 25s, 100s, 250s, 500s.
Use: (1 daily) Vitamin deficiency.

A-B-M-C OINTMENT. See: Lembrose, Oint. (Wyeth).

ABRODIL. See Methiodal Sodium. (Winthrop-Stearns).

A-B-S & C. (Horton & Converse) Aloin 1/5 gr., strychnine 1/120 gr., belladonna lvs ext 1/6 gr., cascara sagrada 1/2 gr./Tab. Bot. 100s, 1000s.

Use: (1 h.s.) Laxative.

ABSORBABLE CELLULOSE COTTON OR GAUZE. See: Oxidized Cellulose. (Various Mfr.).

ABSORBABLE DUSTING POWDER, U.S.P., N.N.R. See: Bio-Sorb, Pow. (Ethicon).

ABSORBABLE GELATIN FILM, N.N.R. Sterile nonantigenic, absorbable, water-insoluble, gelatin film. See: Gelfilm. (Upjohn).

ABSORBABLE GELATIN SPONGE, U.S.P., N.N.R., A.D.R. Gelatin sponge. See: Gel-foam, Preps. (Upjohn)

ABSORBABLE SURGICAL DRESSINGS. See: Hemo-Pak. (Johnson & Johnson).

ABSORBABLE SURGICAL SUTURE, U.S.P., A.D.R. (Various Mfr.) Surgical Gut, Surgical Catgut, Catgut suture.

ABSORBENT BASE. (Upsher-Smith) Oint. Base: Cetyl alcohol, stearyl alcohol, lauryl sulfate, methyl p-hydroxybenzoate & hydrocarbons. Jar 1.5, 6, 25 lb.
Use: Ointment base.

ABSORBENT OINTMENT BASE.
See: Plastibase Hydrophilic (Squibb).
Polysorb (Fougera).

ABTEN. (Paul Maney) *Rauwolfia serpentina* alkaloids ext. Tab. (1 mg. & 2 mg.) Bot. 100s, 1000s.

Use: (2 daily) Hypotensive and tranquilizing agent.

ABUPHENINE CITRATE. (Chemo Puro) Diethylaminoethyl phenylethylacetate citrate. Pow. for mfg.

Use: Antispasmodic.

ABUPHENINE HYDROCHLORIDE. (Chemo Puro) Diethylaminoethyl phenylethylacetate hydrochloride. Pow. for Mfg.

Use: Antispasmodic.

A-C TROCLES. (Abbott) Anesthesin 1/4 gr., calcidin & licorice. Bot. 24s, 100s, 1000s.

Use: Throat & mouth infections.

ACACIA GUM, U.S.P., A.D.R. (Penick) *Acacia senegal*, gum arabic. Mucilage U.S.P.; pow., 1/4 lb., tears, 1/4-1 lb., syr. U.S.P.; whole, 1/4-1 lb.

Use: Demulcent, emulsifier.

W/Mag. oxide & veg. mucinoid. See: Acarosil, Pow. (Sherman).

W/Veg. mucinoid. See: Acarojel, Pow. (Sherman).

A.C.A. COMPOUND. (Rowell) Aspirin 3 1/2 gr., phenacetin 2 1/2 gr., caffeine 1/2 gr./Cap. or Tab. Bot. 100s, 1000s. Also with codeine phos. 1/8, 1/4, 1/2 and 1 gr. See: A.P.C.

Use: (1-2 q. 3 h.) Analgesic & antipyretic.

ACAPNOID. (Chemico) Belladonna pow. 1/10 gr., phenobarbital 1/2 gr., stramonium pow. 1 gr., ephedrine sulfate 1/4 gr./Tab. Bot. 100s, 500s, 1000s.

Use: Sedative & antispasmodic.

ACARALUM. (Sherman) Carbohydrate uronides (veg. mucinoid) 50%, acacia 20%, aluminum hydroxide gel 25%, Mag. oxide 5%. Pow. Can 8 oz.

Use: (2 dr. p.r.n.) Gastric antacid.

ACAROJEL. (Sherman) Carbohydrate uronides (veg. mucinoid) 50%, acacia 50%. Pow. Can 8 oz.

Use: Laxative.

ACAROSIL. (Sherman) Carbohydrate uronides (veg. mucinoid) 50%, acacia 25%, Mag. oxide 25%. Pow. Can 8 oz.

Use: (1 dr. p.r.n.) Gastric antacid.

ACCESSORONE. (Upjohn) Vit. A 22,600 u., D 267 u., red bone marrow ext. 0.62 cc., iron & ammonium citrate 0.259 Gm., malt ext., yeast ext., citrus conc. q.s./fl. oz. Bot. 10 fl. oz., 24 fl. oz.

Use: Dietary supplement.

ACCO. (Assoc. Concentrates) Protein, 32%, carbohydrate 54%, ash 5.75%, fat 0.75%, crude fiber 1.61%, Vit. B-1 2 mg., B-2 .3 mg., nicotinic acid 15 mg. Jar 15 oz.

Use: Nutritional supplement.

A-C-D SOLUTION, U.S.P. Sod. citrate, citric acid & dextrose in sterile pyrogen-free solution.

Abbott—(Abbo-Vac & non-Vac) 250 cc., (Piapab) w/Blood collection set 500 cc.

Baxter—600 cc. bot. with 70, 120 or 300 cc. of sol.; 1000 cc. with 500 cc. of sol.

Cutter—500 cc. bot. with 75 or 120 cc. of sol.; 650 cc. bot. with 80 or 130 cc. of sol.

Use: Anticoagulant for preparation of plasma or whole blood.

ACE-CAF-EDINE, A.D.R. 1955. (Paul Maney)

Aspirin 3 1/2 gr., acetophenetidin 2 1/2 gr., caffeine 1/2 gr./cap. or tab. Cap., Bot. 100s; Tab., No. 1, Bot. 1000s; also w/phenobarbital, Bot. 1000s. See: A.P.C.

Use: (1-2 q. 3 h.) Analgesic & antipyretic.

ACECOLINE. (Anglo-French) Acetylcholine chloride 20, 50, 100 or 200 mg. w/saligenin 40 mg./cc. in propylene glycol. Amp., 1 cc. Box 6s.

Use: S.C., I.M.; parasympathetic agent.

ACEDICON. See: Dihydrocodeinone Enol Acetate. (No Mfr. currently lists.)

ACE-JEL. (Ortho) Acid vaginal jelly buffered to pH of 4.0, containing ricinoleic acid, boric acid, oxyquinoline sulfate & propyl ester parahydroxybenzoic acid in water-dispersible base. Tube with or without applicator.

ACENOCOUMARIN. See: Sintrom, Tab. (Geigy)

ACETABAR. (Columbus) Phenobarbital 1/4 gr., acetophenetidin 2 1/2 gr., acetylsalicylic acid 3 1/2 grs., caffeine 1/4 gr./Tab. or wafer. Bot. 100s, 1000s.

Use: (1-2 q. 4 h.) Analgesic-sedative.

ACETALAX. (G. F. Harvey) Diacetylhydroxyphenolisatin. Cap. 5 mg., Bot. 100s, 1000s. Use: (1-2 h.s.) Laxative.

3 - ACETAMIDO-2, 4, 6 - TRIIODOBENZOIC ACID. See: Acetizicoic acid, U.S.P. (Mallinckrodt).

2 - ACETAMIDO-5 - NITROTHIAZOLE. See: Tritheon (Ortho)

ACETAMINOPHEN. See: N-Acetyl-p-aminophenol.

ACETANILID, N.F., A.D.R.; Tablets, N.F. (Major Mfr.). (Acetylaminobenzene, acetyl-aniline, antifebrin) N-Phenylacetamide Cry. Bot. 1 oz., 1/4 lb., 1 lb.

Use: Antipyretic, analgesic.

Powder Compound, N.F. (Major Mfr.).

W/Acetophenetidin, salol & caffeine. See: Ferbrisol, liq. (Tilden).

W/Anesthesin, ephedrine alkaloid, bismuth subnitrate, petrolatum, menthol, eucalyptol. See: Nasalene, No. 1, Liq. (Abbott).

W/Quinine sulfate, podophyllin, aloin, aco-

ACETOPHENETIDIN PHENYL SALICYLATE TABLETS

nite root, capsicum, atropine sulfate, and camphor. See: C.C.G., Tab. (Noyes).

ACETANNIN, N.N.R. 1941. See: Acetylannic Acid. (No Mfr. currently lists.)

ACETARSOL. See: Acelarsone, Tabs. (Abbott)

ACETARSONE, N.F., N.N.R. 1950; Tablets, N.F. (Abbott) 3-Acetamido-4-hydroxy-phenylarsonic acid (Acetarsol, Acetphenarsine, Amarsan, Dynarsan, Ehrlich 594, Limarsol, Orarsan, Osarsal, Paroxyl, Stovarsol.) Tab. (0.25 Gm.). Bot. 100s, 500s.

Use: Amebiasis & trichomoniasis.

W/Hydrolyzed carbohydrates. See: Devegan, Tabs. (Winthrop-Stearns).

ACETARSONE SALT OF ARECOLINE. See: Drocabil, N.F. (Winthrop Labs.)

ACETASOL. See: Antipyrine acetylsalicylate. (No Mfr. currently lists.)

ACETAZOLAMIDE, N.N.R. See: Diamox, Tabs. (Lederle).

ACET-DIA-MER-SULFONAMIDE, N.F., N.N.R. 1955. Sulfaclamide, Sulfadiazine & Sulfa-merazine.

See: Buffonamide, Susp., N.N.R. (Tutag) Cetazine, Tab., N.N.R. (Bowman Bros.). Dorsulfas, Susp. & Tabs., N.N.R. (Smith-Dorsey).

Incorposul, Susp. & Tabs., N.N.R. (Blue Line). Pansulfa, Susp., Tabs. (Merrell).

Sulfadiazine, sulfamerazine & sulfacetamide, Susp., N.F.; Tab., N.F. Tilamide, Liq. (Tilden).

Tricombisul, Susp. & Tabs., N.N.R. (Schering). Trizyl, Susp. & Tabs. (Central).

W/Gantrisin & Pot. penicillin G. See: Triocinil, Tab. (Flint, Eaton)

W/Hyoscyamus Ext. See: Suromate, Tab. (Patch)

W/Sulfamethazine. See: Delomets, Tab. (Durst). Deltamide, Tab. or Susp. (Armour)

Katrasul, Tabs., Susp. (Chicago). Quad-Sulfa, Tab. (Zimmer).

W/Sulfamethazine, Pot. penicillin G. See: Deltamide w/Penicillin, Tab. or Supp. (Armour)

W/Urea, neocalamine, talc, glycerin, camphor, menthol. See: Neofonamide, Liq. (McNeil)

ACETEST. (Ames). Soda, nitroprusside, disodium phosphate, aminoacetic acid. Tab. Bot. 100s, 250s.

Use: Test for acetone or diacetic acid in urine.

ACETICYL. See: Acetylsalicylic Acid.

ACETIDINE, A.D.R. (Merck Sharp & Dohme)

Aspirin 6 pt., phenacetin 4 pt., caffeine 1 pt. Caps., pink, 5 gr., Bot. 100s, 1000s;

white, 5 gr., Bot. 100s, 1000s; Tab., pink or white, 5 gr., Box 12s, 36s, Bot. 100s, 1000s. See: A.P.C.

Use: (1-2 q. 3 h.) Analgesic, antipyretic. W/Codeine phosphate $\frac{1}{4}$ gr., A.D.R. Tab. 100s, 500s.

Use: (1-2 q. 3 h.) Analgesic, antipyretic. W/Codeine phosphate $\frac{1}{2}$ gr., A.D.R. Tab. 100s, 500s.

Use: (1-2 q. 3 h.) Analgesic, antipyretic. W/Codeine phosphate $\frac{1}{4}$ gr., A.D.R. Tab. 100s, 500s.

Use: (1-2 q. 3 h.) Analgesic, antipyretic.

ACETILUM ACIDULATUM. See: Acetylsalicylic Acid. (Various Mfr.).

ACETOBRO, PEDIATAB No. 1. (Columbus Pharm.) Bromisovalum $\frac{1}{2}$ gr., ethyl salicylate carbonate 1 gr., acetophenetidin $\frac{1}{2}$ gr., caffeine 1/16 gr./tab. Bot. 100s, 1000s.

Use: (1-2 q. 3 hr.) Sedative. **ACETOL**. See: Acetylsalicylic Acid. (Various Mfr.).

ACETOLACTIN. (G. F. Harvey) Acetophenol-satin 5 mg./Cap. Bot. 100s, 1000s

Use: Laxative or purgative. **ACETOMEROCTOL**, N.N.R. See: Merbak, Prep. (Schieffelin)

ACETOMIN. (Central) Salicylamide 60 mg., acetophenetidin 30 mg., atropine sulfate 0.03 mg., camphor monobromated 8 mg., phenolphthalein 6 mg./Palatab. Bot. 100s, 1000s

Use: Analgesic, antipyretic. **ACETONE or DIACETIC ACID TEST**. See: Acetest, Tab. (Ames).

ACETONYL. (Upjohn) Granular efferv. salt. Aspirin as sod. salt $7\frac{1}{2}$ gr. w/Pot. & Soda citrates & tartrates, calcium lactate, Soda phosphate & magnesium sulfate/2 oz. Bot. 4 oz.

Use: (1 tps.) Analgesic & alkalizer. **3-(α -ACETONYLBENZYL)-4-HYDROXY-COUMARIN**. (Warfarin sod. N.N.R.)

See: Coumadin, Tab. (Endo) Prothromadin, Tab. (G. F. Harvey)

3-(α -ACETONYL-4-NITRO-BENZYL)-4-HYDROXYCOUMARIN. See: Sintrom, Tab. (Geigy)

ACETOPHEN. (Smith-Dorsey) Phenobarbital $\frac{1}{4}$ gr., aspirin 3 gr., acetophenetidin 2 gr., hyoscyamus ext. $\frac{1}{2}$ gr./Cap. Bot. 100s, 1000s.

Use: (1 t.i.d.) Sedative—analgesic. **ACETOPHENETIDIN**, U.S.P., N.N.R. 1941, A.D.R.; Tablets, U.S.P. (Various Mfr.) (Phenacetin, p. Ethoxyacetanilide.) Cry. or Pow. 1 oz., $\frac{1}{4}$ lb., 1 lb.

Use: Antipyretic, analgesic. W/Aspirin, caffeine. See: A.P.C.

Acetophenetidin Combinations on pp. 4 to 8.

ACETOPHENETIDIN PHENYL SALICYLATE TABLETS, N.F.

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
16 mg. 120 mg.	Acetanilid 10 mg., caffeine 2 gr., salol 4 gr. Achromycin HCl 125 mg., caffeine 30 mg., salicylamide 150 mg., chlorothen citrate 25 mg.	Tilden Lederle	Ferbrisol, Tab. Achrocidin, Tab.
120 mg.	Achromycin HCl 125 mg., salicylamide 150 mg., Vit. C 25 mg., pyrilamine maleate 15 mg.	Lederle	Achrocidin, Syr./5 cc.
3 gr. 1 gr. 1 1/2 gr. 1 1/2 gr.	Alurate 1 gr. Amidopyrine 3 1/2 gr., caffeine 1/2 gr., hyoscyamus ext. 1/4 gr. Aminopyrine 3 1/2 gr., caffeine 1/2 gr., hyoscyamus ext. 1/4 gr. Amidopyrine 2 gr., caffeine 3/4 gr., phenobarbital 1/2 gr. Amidopyrine 3 gr., caffeine 3/4 gr., phenobarbital 1/6 gr., Mag. oxide	Roche Lilly Massengill Chicago Harco	Allonal, Tab. Amidophen, Puy. Andofive, Cap. Nalgised, Tab. Compirin, Tab.
2 1/2 gr. 2 1/2 gr. 2 1/2 gr. 2 gr. 2 1/2 gr. 2 1/2 gr. 2 1/2 gr. 1 1/2 gr.	Amobarbital 1/2 gr., d-amphetamine sulf. 5 mg., aspirin 2 1/2 gr. Amobarbital 1/2 gr., d-amphetamine 5 mg., aspirin 2 1/2 gr. d-Amphetamine phosphate 1/24 gr., aspirin 2 1/2 gr. rac-Amphetamine 1/26 gr., aspirin 4 gr. Amphetamine sulfate 2.5 mg., aspirin 2 1/2 gr. d-Amphetamine sulfate 2.5 mg., aspirin 3 1/2 gr. d-Amphetamine sulfate 2.5 mg., pentobarbital sod. 1/4 gr., aspirin 3 1/2 gr. Aspirin 3 gr., opium pow. 1/20 gr., ipecac pow. 1/20 gr., atropine sulf. 1/500 gr., camphor 1/2 gr.	Smith; Kline & French Smith, Kline & French Neisler Harvey Smith, Kline & French Pitman-Moore Arnar-Stone	Deprisal, Tab. Deprisal, Tab. Threaze, Tab. Harodyne # 4, Tab. Edrisal, Tab. Aletin, Tab. Polygesic, Tab.
194 mg.	Aspirin 162 mg., phenobarbital 16.2 mg., hyoscymine sulf. 0.031 mg., propenopyridamine maleate 12.5 mg., phenylephrine HCl 10 mg.	Robins	Dophenco, Cap. Phenaphen Plus, Tab.
2.5 gr.	Aspirin 3 gr., promethazine HCl 6.25 mg., mephenetermine sulf. 7.5 mg.	Ives-Cameron	Synalgos, Cap.
2 gr.	Aspirin 4 gr., racetephedrine HCl 1/12 gr., pyrilamine maleate 1/5 gr.	Ulmer	Hiscaphen, Tab.
2 gr. 120 mg.	Barbital 1 1/2 gr., aspirin 3 1/2 gr. Belladonna alkaloids 0.0625 Gm., aspirin 180 mg., phenobarbital 15 mg.	Warner Rorer	Peralga, Tab. Paarden, Cap.
1/2 gr. 2 gr.	Belladonna lr. 3.8 min., aspirin 3 1/2 gr., caffeine 1/2 gr. Belladonna ext. 1/5 gr., Cal. iodide 1/8 gr., Soc. salicylate 2 gr., camphor 1/8 gr., capsicum 1/10 gr.	Vanpelt & Brown Bellaspro, Tab. Cephbe, Tab.	Superior Pharmacal

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
2½ gr.	Benzedrine sulfate 2.5 mg., aspirin 2½ gr. Codeine sulfate ½ gr.	Smith, Kline & French Smith, Kline & French Smith, Kline & French Columbus Pharmacal	Edisal, Tab. w/Codeine w/Codeine Acetobro, Pediatab No. 1
¾ gr.	Codeine sulfate ¼ gr., ½ gr. Bromosalvolum ½ gr., ethyl salicylate carbonate 1 gr., caffeine 1/16 gr.		
½ gr.	Caffeine 1/10 gr., sod. bicarbonate 1/5 gr., amm. carbonate 1/32 gr., licorice ext.	Jenkins Labs.	Anodyne Jr., Tabs.
1 gr.	Colchicine salicylate 1/700 gr., aspirin 5 gr., caffeine ½ gr. 320 mg. Chlorothen citrate 25 mg., caffeine 32 mg.	First Texas Whittier	For-Dyne, Cap. & Tab.
320 mg.	Chlorothenpyramine 25 mg., caffeine 32 mg.		Caribren Comp., Tab.
16 mg.	Chloro-Trimeton maleate .75 mg., aspirin 80 mg.	Breon	Fenoxydine Comp., Tab.
130 mg.	Chloro-Trimeton maleate 4 mg., salicylamide 190 mg., caffeine 30 mg., ascorbic acid 50 mg., methamphetamine HCl 1.25 mg.	Schering	Coricidin Pediatric Medilets
2 gr.	d-Desoxyephedrine HCl 1/24 gr., aspirin 4 gr., butabarbit ¼ gr.	Central	Asphedex, Tab.
200 mg.	di-Desoxyephedrine HCl 1 mg., Sod. gentisate 200 mg.	Crookes	Gentexin, Tab.
2 gr.	Diallybarbital 2/3 gr., aspirin 3½ gr.	Ulmer	Alprine, Tab.
160 mg.	Dihydrohydroxycodeinone HCl 4.5 mg., dihydrohydroxycodeinone Terephthalate 0.38 mg., hexobarbital 100 mg., homatropine Terephthalate 0.38 mg., aspirin 224 mg., caffeine 32 mg.	Endo	Percobarb, Cap.
1 gr.	Dover's pow. 3/16 gr., aspirin 3 gr., caffeine ⅛ gr.	C. D. Smith	Acetyl-Cat-Phen, Cap.
1½ gr.	Dover's pow. ¼ gr., aspirin 3½ gr., caffeine ½ gr., camph. ½ gr.	Harvey	Harvadine # 3, Tab.
1½ gr.	Dover's pow. ¼ gr., aspirin 2 gr., caffeine ⅛ gr., camphor mono bromide ¼ gr.	Warren-Teed	Oveen, Tab.
1½ gr.	Dover's pow. ¼ gr., aspirin 2 gr., caffeine ⅛ gr., atropine sulfate 1/500 gr.		
1½ gr.	Dover's pow. ¼ gr., camphor ¼ gr.		Ovofen, Cap. & Tab.
1½ gr.	Dover's pow. ¼ gr., aspirin 2 gr., atropine sulfate 1/500 gr.		
1½ gr.	camphor monobromide ¼ gr.		
1½ gr.	Dover's pow. ¼ gr., aspirin 2 gr., caffeine ⅛ gr., atropine sulfate 1/500 gr.	Hance	
3 gr.	Ephedrine HCl ½ gr.	Abbott	
160 mg.	Ephedrine HCl ½ gr., phenobarbital ¼ gr. 25 mg., caffeine 50 mg.	Mallard	
		Field Labs.	

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
130 mg. 100 mg. 2 gr.	Ergotamine tartrate 1 mg., belladonna alkaloids 0.1 mg., caffeine 1/2 gr.	Organan	Wigraine, Tab.
2 gr.	Gelsemium 1/10 gr., aspirin 3 gr., caffeine 1/2 gr.	Smith-Dorsey	Phen-Acetyl, Cap.
2 gr.	Gelsemium 3/10 gr., aspirin 4 gr., caffeine 1/2 gr.	Smith-Dorsey	Salsiphen, Tab.
2 gr.	Gelsemium 1/3 gr., aspirin 4 gr., caffeine 1/2 gr., strychnine sulfate 1/240 gr.	Elder	Narelic, Tab.
2 1/2 gr.	Gelsemium 1/5 gr., aspirin 2 1/2 gr., colchicine salicylate, camphor	First Texas	Formosal & Camphor Comp., Tab.
130 mg.	Gelsemium ext. 2.2 mg., aspirin 130 mg., caffeine 16 mg., hyoscyamus 11 mg.	Drug Products	Henasphen, Cap.
2 1/2 gr.	Gelsemium tr. 2 min., aspirin 3 1/2 gr., colchicine salicylate 1/4 gr.	Strasenburgh	Semapirin, Tab.
2 gr.	Gelsemium tr. 3 min., aspirin 4 gr., colchicine salicylate 1/4 gr.	Central	Asphac-G, Tab.
2 gr.	Gelsemium tr. 2 min., aspirin 3 gr., colchicine salicylate 1/4 gr., methyl salicylate	Noyes	Aidental, Tab.
2 1/2 gr.	Gelsemium tr. 1 min., aspirin 3 1/2 gr., colchicine salicylate 1/4 gr., diazepam 1/4 gr.	Buffington	Doradol, Tab.
120 mg.	Heperidin 100 mg., vit. C 50 mg., phenylephrine HCl 5 mg., prophenylpyridamine maleate 6.25 mg., methapyrilene HCl 8.33 mg., pyrilamine maleate 8.33 mg., salicylamide 200 mg., caffeine 30 mg.	Boyle	Boyle Citra, Cap.
1 gr. 2 gr.	Hyoscynamus ext. 1/4 gr., amidopyrine 3 1/2 gr., caffeine 1/2 gr., Methapyrilene fumarate 8.33 mg., phenyltoloxamine dihydrogen citrate 8.3 mg., pyrilamine maleate 8.3 mg., aspirin 3 gr., caffeine 1/2 gr.	Lilly Walker Corp.	Amidophen, Pulv. Tris-A-Lin, Tab.
100 mg. 130 mg.	Panabrom 50 mg., Penicillin V 100,000 u., salicylamide 194 mg., promethazine HCl 6.25 mg., mephenetermine sulf. 3 mg.	Whittier Wyeth	M-Minus-5, Tab. Pen-Vee-Cidin, Cap.
2 1/2 gr. 2 1/2 gr.	Pentobarbital 1/4 gr., aspirin 4 gr. Pentobarbital 1/4 gr., dextroamphetamine sulfate 2.5 mg., aspirin 3 1/2 gr.	Smith-Dorsey U.S. Standard Prods.	Angesal, Tab. Damital, Tab.
2 gr. 2 gr.	Phenobarbital 1/4 gr., aspirin 4 gr. Phenobarbital 1/4 gr., aspirin 4 gr.	Warren-Teed Harvard, Tab.	Axotol, Tab.
2 gr.	Phenobarbital 1/4 gr., aspirin 4 gr., atropine sulfate 1/500 gr.	Warren-Teed Rorer	Axopin, Tab.
120 mg.	Phenobarbital 15 mg., aspirin 180 mg., belladonna alkaloids 0.0625 Gm.	Paepon, Cap. & Tab.	Paepon, Cap. & Tab.

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
1½ gr.	Phenobarbital ½ gr., amidopyrine 2 gr., caffeine ½ gr.	Chicago	Nalgesid, Tab.
1½ gr.	Phenobarbital 1/6 gr., amidopyrine 3 gr., caffeine ¼ gr., Mag.	Harcos	Compirin, Tab.
oxide			
3 gr.	Phenobarbital ¼ gr., ephedrine HCl ¾ gr.	Mallard	Phenodrin, Cap.
100 mg.	Pot. Penicillin G 250,000 u., salicylamide 100 mg., phenyltolox-	Patch	Elipatec, Tab.
amine dihydrogen citrate 25 mg.	amine dihydrogen citrate 25 mg.	Whitier	M-Minus-4, Tab.
100 mg.	Pyranisamine maleate 25 mg., aspirin 2½ gr., gelsemium 1/5 gr.	First Texas	Formasol & Camphor Comp.
2½ gr.	colchicine salicylate camphor		w/Antihistamine
2½ gr.	Pyriaramine maleate 25 mg., aspirin 3½ gr., caffeine ½ gr.,	Horton & Converse	Hisia-Pacq., Tab.
	quinine sulfate ½ gr.	Hance	Histo-Pac., Tab. & Cap.
2½ gr.	Pyriaramine maleate 25 mg., aspirin 3½ gr., caffeine ½ gr.	Rexall	Monistatin A.P.C., Tab.
1½ gr.	Pyriaramine maleate 25 mg., aspirin 3½ gr., caffeine ½ gr.	Gold Leaf	Parisamine, Tab.
2½ gr.	Pyriaramine maleate 10 mg., aspirin 3½ gr., PABA ½ gr.	Brewer	Pyratist Comp. w/Vit. C, Cap.
2½ gr.	Pyriaramine maleate 5/12 gr., aspirin 3 gr., caffeine 1 gr., Vit. C	Taylor	Almegesic, Tab.
2½ gr.	Pyriaramine maleate 25 mg., aspirin 3½ gr., caffeine ½ gr.,	Rowell	Histaquin, Cap.
	hydrocyanine HBr 0.4 mg., hyoscine HBr .008 mg.	Bryant	
2 gr.	Pyriaramine maleate 25 mg., salicylamide 3 gr., caffeine ½ gr.,	Wampole	
	atropine sulf. 1/600 gr., quinine sulf. ½ gr., ascorbic acid	Mallard	
	25 mg.		
2½ gr.	Salicylamide 3½ gr., caffeine ½ gr.		
2.5 gr.	Salicylamide 3.5 gr., caffeine 0.5 gr., ascorbic acid 50 mg., pyril-		
	amine maleate 25 mg., hesperidin 25 mg.		
2½ gr.	Salicylamide 3½ gr., di-desoxyephedrine HCl 1/30 gr.	Tryads, Tab.	
2 gr.	Salicylamide 1 gr., di-desoxyephedrine HCl 5 mg., Mag. glu-	Sedonate, Tab.	
	conate 3 gr.		
120 mg.	Salicylamide 230 mg., prophenylpyridamine maleate 10 mg., caf-	Roerig	
	feine 30 mg., ascorbic acid 20 mg., hesperidin purified 20 mg.		
120 mg.	Salicylamide 250 mg., racophedrine HCl 5 mg., pyramine male-	Central	
	late 12 mg., ascorbic acid 50 mg., Dover's powder 24 mg.		
65 mg.	Salicylate sod. 320 mg., salicylic ethyl ester carbonate 32 mg.,	Endo	
	colchicum root pow. ext. 1.6 mg.		

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
1/4 gr.	Salicylic acid 1/2 gr., sod. bicarbonate 3 gr., mag. salicylate 2 gr., strontium salicylate 2 gr., methyl salicylate, pancreatic, diastase	Jenkins Labs.	Foursalco, Tab.
2 1/2 gr.	Salsal 1 1/2 gr., caffeine 1/2 gr.	Rare-Galen	Salphene, Tab.
2 gr.	Sandoval 3/4 gr., aspirin 3 gr., caffeine 1/2 gr.	Sandoz	Fiorinal, Tab.
200 mg.	Sod. gentisate 200 mg., di-desoxyephedrine HCl 1 mg.	Crookes	Gentexin, Tab.
2 1/2 gr.	Sod. salicylate, PABA, Sod. bicarb.	Anglo-French	Encynex, Tab.
120 mg.	Tetracycline HCl 125 mg., caffeine 30 mg., salicylamide 150 mg., buclizine HCl 15 mg.	Pfizer	Tetracydin, Tab.
2 1/2 gr.	Thiényramine HCl 25 mg., aspirin 3 3/4 gr., d-amphetamine sulfate 1 mg.	Pitman-Moore	Cohisine, Tab.
2 1/2 gr.	Vit. C 25 mg., Cal. glutamate 2 gr., aspirin 5 gr., caffeine 1 gr.	Testagar	Nuprin Comp., Tab.

ACETOPHENONE. (No Mfr. currently lists.) Hypnone, phenyl methyl ketone. Use: Hypnotic.

ACETOSAL. See: Acetylsalicylic Acid. (Various Mfr.).

ACETOSALIC ACID. See: Acetylsalicylic Acid. (Various Mfr.).

ACETOSALIN. See: Acetylsalicylic Acid. (Various Mfr.).

ACETO-VER. (Mallard) Aspirin 5 gr., veratrum viride Pow. 1 gr., rutin 20 mg., Vit. C 30 mg./Tab. Bot. 100s, 500s, 1000s. Use: Capillary diseases.

21-ACETOXYPREGNENOLONE. Artisone acetate (Not listed).

Use: Rheumatoid arthritis.

See: Acetoxanon. [Organon (Not listed)]. Aceloxy-Prenolon [Schering (Not listed)].

ACETOZONE, N.N.R. 1918. (Parke, Davis). Equal parts of acetylbenzoyl peroxide and inert pow.

ACETPHEN-ACETYL COMPOUND. (First Texas) Aspirin 3 gr., acetophenetidin 2 gr., caffeine 1/2 gr., gelsemium tr. 1 min./Cap. Bot. 100s, 500s, 1000s. See A.P.C. w/gelsemium

Use: (2-3 cap. daily) Analgesic & antipyretic.

ACETPHENARSINE. See: Acetarsone, Tab. (Abbott).

ACETPHENOLISATIN. (Chemo Puro) Diacetyl-dihydroxy-diphenylisatin. Pow. Pkg. for mfg.

See: Diacetylhydroxyphenylisatin, Prep. (Various Mfr.).

Use: Laxative.

ACETPYROGALL, N.N.R. 1951. Pyrogallol triacetate.

ACETRIZOATE SODIUM, N.N.R. Injection, U.S.P. 3-Acethylamino-2,4,6-triiodobenzoate sodium. See: Urokon Sodium, Amp. (Mallinckrodt)

W/Polyvinylpyrrolidone. See: Salpix, Vial (Ortho).

ACETRIZOIC ACID, U.S.P., N.N.R. 1951. 3-Acethylamino-2,4,6-triiodobenzoic acid. See: Acetrizoate, Sod. (Mallinckrodt).

ACET-THEOCIN SODIUM. See: Theophylline Sod. Acetate. (Various Mfr.).

ACETYCOL. (Warner-Chilcott) Colchicine salicylated 0.25 mg., p-aminobenzoic acid 162 mg., aspirin 325 mg., thiamine 5 mg., ascorbic acid 20 mg., niacin 15 mg./Tab. Bot. 100s, 500s, 1000s.

Use: (1 t.i.d.) Analgesic in arthritis, gout & neuritis.

ACETYL ADALIN. See: Acetylcarbromal. (Various Mfr.).

p-ACETYLAMINOBENZALDEHYDE THIO-SEMICARBAZONE. (Antib, Berculan A, Benzothiozon, Conteben, Myuizone, Neustab, Tebethion, Thiomicid, Thioparamizone, Thiacetazone)

See: Panrone, Tab. (Panray).

Teebazone, Tab. (Consol. Midland)

Tibione, Tab. (Schenley).

Use: With streptomycin in the treatment of tuberculosis.

ACETYLMAMINOBENZENE. See: Acetanilid. (Various Mfr.).

2-ACETYLAMINO-5-NITROTHIAZOLE. See: Tritheon, Tab. (Ortho)

N-ACETYL-p-AMINOPHENOL. Acetaminophen.

See: Apamide, Tab. (Ames)

Metalid, Tab. (Columbus)

Nebs, Tab. (Norwich Pharmacal)

Tempra (Mead)

Tylenol, Elix. (McNeil)

BUFFERED TABLETS.

See Apamide-Ves, Tab. (Ames)

W/Acetylcarbromal. See: Apromal, Tab. (Ames).

W/A.P.C. See: Doloro Compound, Tab. (Buffington)

W/Aspirin & caffeine.

See: Acetyl duo, Tab. (Chicago Pharm.).

Trigesic, Tab. (Squibb).

W/Aspirin & caffeine with codeine. See: Trigesic w/Codeine, Tab. (Squibb).

W/Butabarbital Sod. & methamphetamine HCl. See: Algoson, Tab. (McNeil).

W/Demeral. See: Demeral APAP (Breon)

W/Flexin. See: Flexilon, Tab. (McNeil)

W/Flexin, hydrocortisone. See: Flexilon-HC, Tab. (McNeil)

W/Methyliso-octenylamine mucate, salicylamide, and caffeine. See: Nygram, Cap. (Dumas-Wilson)

W/Phenyltoloxamine dihydrogen citrate, salicylamide, dl-amphetamine phosphate, metropine. See: Acogesic, Cap. (Strasenburgh)

W/Phenyltoloxamine dihydrogen citrate, salicylamide, Vit. C, citrus bioflavonoids, caffeine, racemic amphetamine sulfate. See: Euphenex, Tab. (Lloyd, Dabney & Westerfield)

W/Salicylamide, codeine phosp., caffeine. See: Cylamide, Tab. (Chicago Pharmacal Co.)

W/Salicylamide, phenobarbital, caffeine. See: Cylamide w/Phenobarbital, Tab. (Chicago Pharmacal Co.)

3-ACETYLAMINO-2,4,6-TRIODOBENZOATE SODIUM. See: Acetrizoate Sod. (Mallinckrodt).

3-ACETYLAMINO-2,4,6-TRIODOBENZOIC ACID. See: Acetrizoic Acid, U.S.P. (Mallinckrodt).

ACETYLANILINE. See: Acetanilid. (Various Mfr.).

ACETYLARSAN. (No Mfr. currently lists) Diethylamine acetarsone

Use: Antisyphilitic

ACETYL-BETA-METHYLCHOLINE CHLORIDE, U.S.P., N.N.R. 1947. See: Mecholyl Chloride, Amp. (Merck Sharp & Dohme)

ACETYL-BROMO-DIETHYLACETYL-CARBAMIDE. See: Acetylcarbomal. (Various Mfr.).

ACETYL-CAF-PHEN. (C. D. Smith) Aspirin 3 gr., acetophenetidin 1 gr., caffeine $\frac{1}{8}$ gr., Dover's Pow. 3/16 gr./Cap. Bot. 100s, 1000s.

Use: (1 t.i.d.) Analgesic.

ACETYL-CARB. (Sherman) Aspirin & Sod. bicarb. Tab. 500s, 1000s.

Use: (1-2 q. 3 h.) Analgesic.

ACETYLCARBROMAL. (Chemo Puro, Meyer)

Acetyl adalin, acetyl-bromo-diethylacetyl-carbamide. Pow. for mfg.

See: Abasin. (Winthrop Labs.)

Sedamyl. (Schenley).

Use: Sedative.

W/N-acetyl-p-aminophenol. See: Apromal, Tab. (Ames)

W/Apamide. See: Apromal, Tab. (Ames).

W/Bromisovalum.

See: Daysed, Tab. (Tailby-Nason).

Sedayson, Tab. (Tailby-Nason).

W/d-Desoxyephedrine HCl.

See: Euphased-5, Tab. (Schenley)

W/Hyoscamine sulfate, atropine sulfate & hyoscine HBr. See: Disperdon, Tab. (Standard).

W/Mephenesin & resperine. See: Amril, Tab. (Leeming)

ACETYLCHOLINE CHLORIDE. (Various Mfr.).

Leader—Amp. (0.1 Gm., 0.2 Gm.) 1 cc., 12s.

Merck—Amp. (0.1 Gm.). Vial 10 Gm.

Use: S.C., I.M., I.V. Parasympathomimetic agent & vasodilator.

W/Saligenin. See: Acecoline, Amp. (Anglo-French).

ACETYLCHOLINE-LIKE THERAPEUTIC AGENTS. See: Cholinergic agents.

ACETYLDIGITOXIN. See: Acylanid, Tab. (Sandoz).

N¹-(ACETYL-3,4-DIMETHYL-5-ISOOXAZOLYL) SULFANILAMIDE. See: Gantrisin Acetyl, Prep. (Hoffmann-La Roche).

ACETYLDUO. (Chicago Pharm.) Acetyl-p-aminophenol 2.5 gr., aspirin 3.5 gr., caffeine 0.5 gr./Tab. Bot. 100s, 1000s.

Use: (1 q. 3-4 h.) Analgesic.

ACETYLYC COMPOUND. (Central) Salicylamicde 30 mg., acetophenetidin 30 mg., atropine sulfate 0.03 mg.,aconite root 1.5 mg., camphor monobromated 8.0 mg., phenolphthalein 6.0 mg./Tab. Bot. 100s, 1000s
Use: Analgesic, antipyretic

ACETYLIN. See: Acetylsalicylic Acid. (Various Mfr.).

ACETYLMETHIONINE. (Chemo Puro) Methionine 78%. Pow. for mfg.

Use: Nutrient.

ACETYLMETHIONINE CALCIUM. (Chemo Puro) Methionine 71%, calcium 9.5%. Pow. for mfg.

Use: Nutrient.

ACETYLPHENYLSALICYLATE. See: Phenyl Acetysalicylate. (No Mfr. currently lists.).

ACETYRESORCINOL. See: Resorcinol Monoacetate. (Various Mfr.).

ACETYSAL. See: Acetylsalicylic Acid. (Various Mfr.).

ACETYSALICYLIC ACID, U.S.P., A.D.R., N.N.R. 1947; Capsules, N.F.; Tablets, U.S.P. (Various Mfr.) Acidum acetyl salicylicum, 2-acetoxybenzoic acid, Acetilum Acidulatum, Acetophen, Acetol, Acetosal, Acetosalic Acid, Acetosalin, Acetyl, Acetylin, Acetyl-sal, Acylpyrin, Aspro, Helicon, Rhodine, Salacetin, Salcetogen, Salatin.
See: Aspirin Preps. (Various Mfr.).

Empirin (Burroughs Wellcome).

Salts.

See: Aluminum Acetysalicylate.

Calcium Acetysalicylate.

Use: Analgesic & antipyretic.

W/Acetophenetidin & caffeine. See: A.P.C. preparations. (Various Mfr.).

W/Mixtures. See: Aspirin preparations. (Various Mfr.).

ACETYSALICYLIC ACID, ACETOPHENETIDIN AND CAFFEINE, N.F. See: A.P.C., Prep. (Various Mfr.).

ACETYSALICYLIC ACID ENTERIC COATED. See: Asteric, Tab. (Brewer).

ACETYLSTROBELINE. See: Asmatone, Cap. (National Research).

N¹-ACETYSULFANILAMIDE. (Albucid; p-Aminobenzenesulfonacetamide; Sulfacet; Sulacetamide, incorrectly called A.P.C.; N-Sulfanilylacetamide.) See: Sulamyd. (Schering).
Use: Sulfonamide therapy.

ACETYL SULFOXAZOLE, N.N.R. N₁-acetyl-3,4-dimethyl-5-sulfanilamidoisoxazole. See: Gantrisin Acetyl. (Hoffmann-La Roche)

ACETYLTANNIC ACID, N.N.R. 1941. (Acetannin, diacetyl tannic acid, tannyl acetate.) See: Tannigen, Pow. (Winthrop Labs.)

Use: Symptomatic treatment of diarrhea.

ACETYL-VESS. (Ames) (Not listed) Aspirin 7½ gr., monocalcium phosphate & citrate carbonate effervescent base/Tab. Bot. 25s Use: (1 q. 3 h.) Antipyretic & analgesic.

ACHLORHYDRIA DETERMINATION. See: Diagnex.

ACHLORHYDRIA THERAPY.

See: Acidol. (Various Mfr.)

Acidol-Pepsin, Cap. (Winthrop Labs.)

Acid-V, Cap. (Merck Sharp & Dohme)

Aclor, Cap. (Cale Chem. Co.)

Chlor-Ethamine, Tab. (Pitman-Moore)

ACHROCIDIN. (Lederle) Tabs. Achromycin HCl 125 mg., phenacetin 120 mg., caffeine 30 mg., salicylamide 150 mg., chlorothen citrate 25 mg./tab. Bot. 24s.

Syr.: Achromycin HCl 125 mg., phenacetin 120 mg., salicylamide 150 mg., Vit. C 25 mg., pyrilamine maleate 15 mg., methylparaben 4 mg., propylparaben 1 mg./5 cc. Syr. Bot. 4 fl. oz.

Use: Antibiotic, antihistamine, analgesic therapy.

ACHROCORT. (Lederle) Tetracycline equiv. to tetracycline HCl 30 mg., hydrocortisone acetate 10 mg./Gm. Cream. Tube 5 Gm. Use: Treatment of superficial pyogenic infections of the skin

ACHROMYCIN, U.S.P., N.N.R. (Lederle) Tetracycline HCl.

Cap.: (50 mg.) Bot. 25s & 100s; (100 & 250 mg.) Bot. 16s & 100s.

Ear Solution: (50 mg. vial and a 10 cc. vial of 5% benzocaine in propylene glycol).

Intravenous: (100, 250 & 500 mg.) Vial.

Intramuscular: (100 mg.) Vial.

Oral Susp.: (250 mg./5 cc.) Bot. 1.5 Gm., dry pow. 1 oz.

Oint.: (3%) Tube ½ & 1 oz.

Ophth. Oil Susp.: 1%, Bot. w/dropper 4 cc.

Ophth. Oint.: (1%) Tube ½ oz.

Ophth. Sol.: Vial (25 mg.) w/dropper.

Ophth. Sterilized: Vial 25 mg.

Pediatric Drops, Liquid: (100 mg./cc.) Bot. w/dropper 10 cc.

Pharyngets: (15 mg.) Pkg. 10s.

Speroids: Choc. flavor (50 mg./isp.) Bot. 12 & 25 doses.

Surgical Powder: Vial (200 mg./Gm.) 5 Gm.

Tab.: (50 & 100 mg.) Bot. 25s & 100s; (250 mg.) Bot. 50s, 250s & 1000s.

Tab. Soluble: (50 mg.) Tube 40s, 100s.

Topical Spray: 710 mg.

Troches: (15 mg.) Pkg. 25s, 250s.

W/Hydrocortisone Oint. (Lederle). Tetra-cycline HCl 3%, Hydrocortisone 2%. Tube 5 Gm.

Use: Pyogenic infections of the skin.
W/Hydrocortisone (Lederle) Ophth. Oint. Tetracycline HCl 1%, Hydrocortisone 1.5%. Tube 1/8 oz. Topical: Tetracycline HCl 3%, Hydrocortisone 2%. Tube 5 Gm.
Use: Eye infections and skin infections
W/Phenacetin, caffeine, salicylamide, chlorothen citrate. See: Achromycin, Syr., Tab. (Lederle)

ACHROMYCIN NASAL SUSPENSION. (Lederle) Tetracycline HCl 3.75 mg., hydrocortisone acetate 2 mg., phenylephrine HCl 0.125%/cc. Bot. squeeze 15 cc.

Use: Upper respiratory infections.

ACHROMYCIN SF CAPSULES. (Lederle) Achromycin 250 mg., Vit. C 75 mg., B-1 2.5 mg., B-2 2.5 mg., niacinamide 25 mg., Vit. B-6 0.5 mg., Cal. pantothenate 5 mg., Vit. B-12 1 mcg., folic acid 0.375 mg., Vit. K 0.5 mg./Cap. Bot. 16s, 100s.

Use: Antibiotic with stress formula vitamins.

ACHROMYCIN SF ORAL SUSPENSION. (Lederle) Achromycin 125 mg., Vit. C 37.5 mg., B-1 1.25 mg., B-2 1.25 mg., niacinamide 12.5 mg., Vit. B-6 0.25 mg., Cal. pantothenate 2.5 mg., Vit. B-12 0.5 mcg., folic acid 0.1875 mg., Vit. K 0.25 mg./5 cc. Bot. 2 fl. oz.

Use: Antibiotic with stress formula vitamins.

ACHROMYCIN SYRUP. (Lederle) Tetracycline HCl 125 mg., methylparaben 0.08%, propylparaben 0.02%, alcohol 1%/5 cc. Bot. 2 oz., 1 pt.

Use: Antibiotic therapy.

ACHROMYCIN-V. (Lederle) Achromycin 250 mg. buffered w/sod. metaphosphate 380 mg./Cap. Bot. 16s, 100s. Pink Caps.: Achromycin 100 mg. w/sod. metaphosphate 152 mg./Cap. Bot. 25s, 100s. **Syr.:** Achromycin 125 mg. buffered w/sod. metaphosphate/5 cc. Bot. 2 oz., 1 pt. **Pediatric Drops:** Achromycin 100 mg. w/ sod. metaphos./cc. Dropper Bot. 10 cc.

Use: Treatment of infections caused by both gram-negative and gram-positive bacteria

ACROSTATIN-V. (Lederle) Tetracycline phosphate buffered 250 mg., nystatin 250M u./Cap. Bot. 16s, 100s. Oral Susp. tetracycline phosp. buf. 125 mg. & nystatin 125M u./5 cc. Bot. 60 cc.

Use: Broad-spectrum antibiotic w/antifungal therapy

ACIBAN. (Lederle) Calcium caseinate 0.4 Gm., calcium carb. 0.1 Gm./Tab. Bot. 50s, 250s, 1000s.

Use: (1-2 t.i.d.) Antacid.

ACICHEK. (Valentine) Dihydroxy aluminum aminoacetate 250 mg., Sod. carboxymethyl-

cellulose 150 mg., glycine 30 mg./Tab. Bot. 100s, 1000s.

Use: Gastric hyperacidity in peptic ulcer.

ACID ACRIFLAVINE. See: Acriflavine Hydrochloride. (Various Mfr.).

ACID CARBOLIC, NUTGALL, AND ATROPINE COMPOUND. (Upjohn) Phenol 6 gr., nutgall (equiv.) 36 gr., atropine 1/4 gr., benzocaine 10 1/10 gr./oz. Oint. Tube 1 oz. Jar 1 lb.

Use: Rectally, local anesthetic, analgesic & antipruritic.

ACID CITRATE DEXTROSE SOLUTION. See: A.C.D. Sol.

ACI-DERMA. (Princer) An aqueous emulsion of glyceryl monostearate, cetyl alcohol, sodium lauryl sulfate, propyl and benzyl p-hydroxy benzoates and propylene glycol, buffered to pH 2-3. Bot. 4 oz.

Use: Restore the acid mantle of the skin

ACID HISTAMINE PHOSPHATE. See: Histamine Phosphate. (Various Mfr.).

ACIDIIRON. (Walker Labs.) Ferrous sulfate exsic. 3 gr., hydrochloric acid diluted 125 mg./tab. Bot. 100s.

Use: Anemias.

ACID MANTLE CREME. (Dome) Aluminum acetate in specially prep. water-soluble hydrophilic cream at pH 4.2 Tube 1 oz. Jar 4 oz., 1 lb.

Use: Skin prophylactic agent in restoring normal skin acidity.

ACID MANTLE LOTION. (Dome) Aluminum acetate lotion. Bot. plastic, 1 oz. Bot. 4 oz. squeeze, 1 pt.

Use: Dermatitis.

ACIDOL. See: Betaine Hydrochloride. (Various Mfr.).

ACIDOLATE. (White) (Hypoallergenic liquid skin cleanser.) Sulfated vegetable oils, liquid petrolatum & water. Bot. 8 oz., & gal. Use: Cleanses skin surfaces without irritation.

ACIDOL-PEPSIN. (Winthrop Labs.) Betaine HCl 4 gr., pepsin 3.5 gr./Cap. Bot. 50s. Use: (1-2 t.i.d.) Achlorhydria.

ACIDORIDE. (Abbott) d-Glutamic acid HCl. Cap. (5 gr.), Bot. 100s, 500s, 1000s. Use: Achlorhydria,

ACIDOTHYN. (Flint-Eaton) Glutamic acid HCl. (Each cap. is equivalent to 10 min. of Dilute Hydrochloric Acid U.S.P.) Bot. 100s, 500s, 1000s. Use: (1-2) Achlorhydria.

ACID TRYPAFLAVINE. See: Acriflavine Hydrochloride. (Various Mfr.).

ACIDULIN. (Lilly) Glutamic acid HCl. (Equivalent to 0.62 cc. [10 min.] of diluted hydro-

chloric acid.) Pulvules. Bot. 100s, 500s, 1000s.

Use: (1-2) Achlorhydria.

ACIDUSOL. (Texas) Oxyquinoline sulfate, lactic acid, zinc sulfocarboilate, citric acid, dextrose, buffered solution. Bot. 8 oz.

Use: Vaginal cleanser.

ACID-V. (Merck Sharp & Dohme) Glutamic acid HCl 300 mg., vit. B-1 2 mg., riboflavin 1 mg., niacinamide 10 mg./cap. Bot. 100s, 500s, 1000s.

Use: (1 t.i.d.) Achlorhydria.

ACIGLUMIN. See: Glutamic Acid Hydrochloride. (Various Mfr.).

ACI-JEL. (Ortho) Acetic acid, ricinoleic acid, oxyquinoline sulfate, and glycerine in a vegetable gum base. Jelly, Tube with applicator, 3 oz.

Use: Vaginal antiseptic.

ACINITRAZOLE. See: Tritheon (Ortho)

ACITAMIN. (Massengill) Glutamic acid HCl 0.324 Gm., thiamine HCl 1 mg., riboflavin 1 mg., niacinamide 5 mg./cap. Bot. 100s, 500s, 1000s.

Use: (1-2) Achlorhydria.

A-C-K. (G. F. Harvey) Aspirin 5 gr., menadione 1/200 gr., ascorbic acid 1/2 gr./Tab. Bot. 100s, 1000s

Use: Salicylate therapy.

A-C-K BUFFERED. (G. F. Harvey) Aspirin 333 mg., ascorbic acid 33.3 mg., menadione 0.33 mg., calcium carbonate 60 mg./tab. Bot. 100s, 1000s

Use: (1-2 q. 3 h.) Analgesic.

ACLETIN. (Associated Concentrates) Lecithin, soya 8 gr. with vitamins/Cap. Bot. 100s, 250s.

Use: (1 to 2 cap. daily) Fat metabolism.

ACLOR. (Cole) Glutamic acid HCl. Cap. (5 gr.) Bot. 100s

Use: (1-2) Achlorhydria.

ACNAVEEN. (Aveeno Corp.) Colloidal oatmeal, kaolin, sulfur 3%, and hexachlorophen 0.5%. Lot., Pow., Pkg. 3 1/4 oz.

Use: Topically, cleansing agent in the treatment of acne

ACNEDERM. (Lannett) 5% Dispersible sulfur, 3% zinc sulfate. Bot. 2 oz.

Use: Skin conditions.

ACNE MIXED "U.B.A." (Lilly) Undenatured antigens of bacillus acne & staphylococci standardized to 0.05 mg. nitrogen/cc. Vial 5 cc., 20 cc.

Use: Pustular acne.

ACNE SEROBACTERINE MIXED VACCINE. (Merck Sharp & Dohme) Acne bacillus 800 million, Staphylococcus aureus 2000 million, Staphylococcus albus 2000 million/cc. Vial 5 cc.

Use: Certain types of acne.

ACNE VACCINES. (Sherman) Acne bacillus 100 million, Colon bacillus 300 million, Staphylococcus albus 1000 million, Staphylococcus aureus 1000 million/cc. Vial 12.5 cc., 20 cc.

Use: S.C. with increasing dosage, acute & chronic acne infections.

ACNOMEL. (Smith, Kline & French) Resorcinol 2%, sulfur 8%, hexachlorophene 0.25% w/bentonite, propylene glycol, alcohol & titanium dioxide. Cream. Tube 1 1/2 oz., also available in cakes.

Use: Acne therapy.

ACOBRYONNA. (Lemonon) Aconite pow. 1.33 mg., byronia pow. 1.33 mg., belladonna pow. ext. 0.16 mg./Tab. Bot. 100s.

Use: (2 q. 3 h.) Antipyretic & diaphoretic.

ACOBRYONNA COMP. (Lemonon) Aconite pow. 4.66 mg., belladonna pow. ext. 0.24 mg., byronia pow. 1.33 mg., mercury iodide 1/100 gr., Sod. salicylate 1 gr./Tab. Bot. 100s.

Use: (1-3 q. 4 h.) Follicular tonsillitis.

ACOGESIC (Strasenburgh) Phenyltoloxamine dihydrogen citrate 25 mg., acetyl-p-aminophenol 300 mg., salicylamide 200 mg., dl-Amphetamine phosphate 2 mg., metropine 0.5 mg./Cap. Bot. 100s

Use: (1-2 q. 4 h.) Analgesic

ACOLAC BACTERIAL VACCINE. (U. S. Std. Products) Corynebacterium acnes 1000 million, Staphylococcus aureus 500 million, Staphylococcus albus 500 million/cc. Vial 10 cc., 30 cc.

Use: Acute or chronic infections caused by acne.

ACON. N.N.R. (Endo) Water soluble vitamin A. Cap. (25,000 & 50,000 units/Cap.) Bot. 100s, 1000s. Soln. (25,000 units/cc.) Dropper Bot. 30 cc.

Use: Vitamin A therapy.

ACONITE POWDER W/Belladonna pow. ext. & byronia pow. See: Acobryonna, Tab. (Lemonon).

W/Atropine sulf., aminophylline, veratum V. See: Aminite, Tab. (Mallard).

ACR-ALLANTOMIDE. (National Drug) Aminocyclidine HCl 1:500, sulfanilamide 10%, allantoin 2%, in a water-miscible base. Tube 1 oz., 1 lb.

Use: Topically, bacteriostatic.

ACRIPLAVINE. N.F., A.D.R., N.N.R. 1952. (Euflavine, Gonacrine, Neutreflavine.) Acriflavine neutral; a mixture of 2,6-diamino-10-methylacridinium chloride & 2,8-diaminoacridine.

Use: Antiseptic.

ACRIPLAVINE, BRILLIANT-GREEN, METHYLROSANILINE CHLORIDE MIXTURE, N.N.R. 1953. See: Dymixal, Pow. (McNeil).

ACRIFLAVINE HYDROCHLORIDE, N.F., A.D.R., N.N.R. 1950. (Various Mfr.) Acid acriflavine, acid trypaflavine, flavine, trypaflavine

National Aniline—Pow., Bot. (1, 5, 10, 25 & 50 Gm.). Tab. (1½ gr.) Bot. 50s & 100s. Use: Bacteriostatic.

ACRISAN EAR DROPS. (Recsei) Aminoacridine caprylate 0.25% & benzyl alcohol 1% in propylene glycol. Ear drops, ½ oz. Use: Anti-infective.

ACRISAN THROAT SPRAY. (Recsei) Aminoacridine ascorbate 0.2% w/benzyl alcohol 0.5%. Bot. 4 oz.

Use: Anti-infective

A.C.T. (Tilden) Pot. bromide 11 mg., ammonium bromide 11 mg., Sod. bromide 11 mg., aspirin 0.33 Gm., caffeine 33 mg./Tab. Bot. 100s, 1000s.

Use: (1-2 as needed) Analgesic w/sedative.

ACTAMER. (Monsanto Chem.) See: Bithionol, U.S.P.

ACTH, N.N.R. (Adrenocorticotrophic hormone. Adrenocorticotropic(h)in). See: Corticotropin, U.S.P.

ACTH. (National Drug) Colloidal solution of ACTH. Injectosol (20 U.S.P. units/cc.) 2 cc. & 10 cc. I.M. or I.V. drip.

ACTH GEL, PURIFIED. (National Drug) Corticotropin (40 u./cc.) Vial. 1 cc., 5 cc.; (80 u./cc.) Vial 5 cc.

ACTH (PITUITARY ADRENOCORTICO-TROPIC HORMONE). (Parke, Davis) Vial 25 units, 40 units (one unit is equivalent to 1 mg. international standard).

ACTHAR, N.N.R. (Armour) Corticotropin, U.S.P. Vial. 10, 25, 40 mg. (Lyophilized)

ACTHAR GEL. (Armour & National Drug) Corticotropin, U.S.P. Disp. Cart. (20 mg./cc.) 1 cc., Vial 5 cc.; Disp. Cart. (40 mg./cc.) 1 cc., Vial 5 cc.

ACTHAR GEL HIGHLY PURIFIED. (Armour) H. P. acthar gel, a new, highly purified, repository ACTH preparation that may be given subcutaneously or intramuscularly. 20, 40 & 80 u. per cc., 5 cc. multiple dose vials. Supplied in a sterile 1 cc. B-d cartridge with B-2 disposable cartridge syringe, in potencies of 40 Armour u. per cc. See: Corticotropin, U.S.P.

Use: Stimulant of adrenal cortex, anti-inflammatory, antiallergic & useful in collagen diseases.

ACTHAR SOLUTION. (Armour) Corticotropin, 10 u./cc. Amp. 1 cc., Vial 10 cc.

Use: I.V., I.M., S.C. corticotropin therapy.

ACTIVATED CARBON-SILICON. (Inorganic Bioelements Inc.) Activated carbon 50%, activ. silicon 27.5%, oxides of aluminum

6%, mag. 3.5%, iron 1.5%. Pow. Bot. 3 oz. Use: Irritations of G.I. tract.

ACTIVATED 7-DEHYDROCHOLESTEROL, U.S.P., A.D.R. See: Vitamin D-3. (Various Mfr.)

ACTIVATED ERGOSTEROL. See: Calciferol, U.S.P.

ACTIVIN. (Bischoff) (Not listed) Purified casein with sodium iodohydroxyquinoline sulfonate in an injectable solution. Amp. 1 cc., Bot. 6s, 12s, 100s. Vials 10 cc., 25 cc. Use: I.M. Nonspecific protein therapy.

ACTO-CORTIN GEL. (Myers-Carter) Adrenal cortex ext. 40 & 80 mcg./cc. Vials 5 cc. Also w/Vit. B-12 1000 mcg./cc. Vial 5 cc.

ACTOPHEN. (Biochemical) Vit. D 50,000 u., bile salts 32 mg./Cap. Bot. 50s, 100s, 500s. Use: Choleretic with vitamin D supplement.

ACTROPE. (United Lab.) Corticotropin, U.S.P. Vials 25, 40 & 100 mg./cc.

Use: Adrenocorticotropic therapy.

ACTYLATE TABLETS. (Kinney) Ammonium salicylate 1½ gr., Pot. salicylate 1½ gr., strontium salicylate 1½ gr., p-aminobenzoic acid 4 gr., Vit. C ½ gr./Tab. Bot. 100s. Use: (2 q. 3 hr.) Rheumatic disorders.

ACTYLATE PLUS E. (Kinney) Cortisone acetate 5 mg., amm. salicylate 1½ gr., pot. salicylate 1½ gr., strontium salicylate 1½ gr., pot. p-aminobenzoate 5 gr., vit. C ½ gr./tab. Bot. 50s.

Use: (1-2 t.i.d.) Treatment of rheumatoid arthritis.

ACLYANID TABLETS. (Sandoz) Acetyl-digitoxin. Tab. (0.1 mg.) pink, 100s; (0.2 mg.), white, 100s, 1000s

Use: Digitoxin therapy.

ACLYPYRIN. See: Aspirin. (Various Mfr.)

A & D CAPSULES. (Rowell) Vit. A 5000 u., Vit. D 500 u./Cap. Bot. 100s, 1000s.

Use: (1 cap. t.i.d.) Vitamins A & D therapy.

A-D DROPS. (Walker) Vit. A 2000 u., Vit. D 300 u./drop. Bot. w/dropper, 15 cc., 60 cc.

Use: (m.p.) Vitamin A & D deficiencies.

A & D VITAMIN CONCENTRATE W/VIosterol. (Horton & Converse) Vit. A 5000 u., Vit. D 1700 u./Tab. Bot. 100s, 250s, 500s, 1000s.

Use: Nutritional supplement.

ADALIN, N.N.R. 1947. (Winthrop Labs.) (Carbromal) Bromodiethylacetylcarbamide.

Tab. (5 gr.) Bot. 25s, 500s.

Use: (1-2 m.p.) Sedative & hypnotic.

ADAMIN. (Gold Leaf) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 2 mg./Cap. Bot. 100s, 1000s.

Use: (1 daily) Dietary supplement.

ADANON HYDROCHLORIDE, N.N.R. (Winthrop Labs.) Methadone HCl, 6-dimethyl-

amino-4,4-diphenyl-3-heptanone hydrochloride.

Amp.: (5 mg./cc.) 2 cc. Box 10s, 25s, 100s.

Elixir: (5 mg./5 cc.) Bot. 1 pt.

Syr.: (1.65 mg./5 cc.) Bot. 1 pt.

Tab.: (2.5, 5.0, 7.5, 10 mg.) Bot. 100s, 500s.

Vial: (5 & 10 mg./cc.) 20 cc. Box 1s.

Use: Narcotic, analgesic.

ADA-TABS. (G. F. Harvey) Dihydroxy aluminum aminoacetate 500 mg./tab. Bot. 100s, 1000s.

Use: Antacid.

ADC DROPS. (Parke, Davis) Vit. A 5000 u., D 1000 u., C 50 mg./0.6 cc. Bot. w/dropper, 15 cc., 50 cc.

Use: Dietary supplement.

ADCAPS. (Blue Line) Vit. A 5000 u., Vit. D 500 u./Cap. Bot. 100s, 1000s.

Use: (1 daily) Vitamin A & D.

ACETS. (Walker) Vit. A 5000 u., D 1000 u., C 50 mg./VitaspHERE. Bot. 50s, 250s

Use: (1 daily) Dietary supplement.

ADEF. (Endocrine) Vit. A 20,000 u., D 4000 u., E 10 u., F (Vitagen concentrate) 300 mg./1 cc. Bot. w/dropper 30 cc., 4 oz.

Use: (1 or 2 cc. daily) Fat & water metabolism disorders.

ADEF-C. (Endocrine) Vit. A 5000 u., D 1000 u., E 5 u., F (Vitagen concentrate) 100 mg., C 50 mg./Cap. Bot. 100s.

Use: (1 to 2 daily) Mucous membrane & skin allergies.

ADENINE SULFATE. (Schwarz) Pow. Bot. 5 Gm.

Use: Granulocytopenia.

ADENO-B-12. (Kirk) Adenosine 25 mg., vit. B-12 60 mcg./cc. Vial. 10 cc.

Use: Treatment of tendinitis, bursitis, neuritis.

ADENOLIN W/B-12 GEL. (Lincoln) Adenosine-5-monophosphate 25 mg., vit. B-12 150 mcg., & niacin 10 mg. in gelatin/cc. Vial 15 cc.

Use: Neuritis, bursitis & related conditions

ADENOPLEX. (U. S. Standard Products) Adenosine 5-monophosphate 25 mg., Vit. B-12 50 mcg., niacin 25 mg., B-1 10 mg./cc. Vial 10 cc.

Use: I.M., treatment of collagen diseases

ADENOSINE W/Vit. B-12. See: Adeno-B-12, Vial. (Kirk)

ADENOSINE COMP. GEL. (Vitamix) Adenosine-5-monophosphate sod. 25 mg. & sod. nicotinate 25 mg. & B-1 50 mg. w/gelatine/cc. Vial 10 cc.

Use: I.M., bursitis, neuritis & others

ADENOSINE DIPHOSPHATE. (Schwarz) (ADP, Adenyldiphosphoric acid, Adenosine-diphosphoric acid.) Pow. Bot. 2 Gm.

ADENOSINEDIPHOSPHORIC ACID. See: Adenosine Diphosphate. (Schwarz).

ADENOSINE-5-MONOPHOSPHATE.

See: My-B-Den, Amp. (Bischoff)

W/Heart Muscle ext. See: Cardimone, Amp. (Endo)

W/Nicotinic acid, Vit. B-12.

See: Vastran, Amp. (Wampole Labs.) Adenolin-B-12 Gel (Lincoln)

W/Vit. B-12.

See: Adenovite, Vial. (Metropolitan Labs.) Cobaden, Amp. (Rand)

Cyanosine, Gel. (Vitamix)

Harvaden, Vial (G. F. Harvey)

Mycasine, Vial (Myers-Carter)

Thybaden, Vial. (Bell Pharmacal)

W/Vit. B-12, niacin, B-1

See: Adenoplex, Vial (U.S. Standard Products)

Adenosine Comp. Gel (Vitamix)

B-Denose, Vial (B. F. Ascher)

W/Vit. B-12, nicotinic acid, procaine HCl, and benzyl alcohol. See: Vibruden, Vial. (Philadelphia Ampoule Labs.)

W/Vit. B-12, sod. nicotinate, procaine. See: Bastran, Amp. (Wampole)

ADENOSINE-5-MONOPHOSPHATE AS THE SODIUM SALT. See: Cardiomone, Amp. (Endo).

ADENOSINE TRIPHOSPHATE. (Various Mfr.) (ATP, Adenyltriphosphoric acid, Adenylpyrophosphoric acid.) Pow. Bot. 1, 5, 100 Gm.

ADENOVIRUS VACCINE. (Parke, Davis) Contains adenoviruses serotypes 3, 4, and 7 or adenoidal-pharyngeal-conjunctival viruses. Vial

Use: Immunization (1 cc. I.M.) against colds

ADENOVITE. (Metropolitan Labs.) Adenosine 5-monophosphate 25 mg., Vit. B-12 60 mcg./cc. Vial 10 cc., Box 10s.

Use: Treatment of bursitis, pruritus, dermatitis, ulcerations accompanying varicose veins.

ADENYLDIPHOSPHORIC ACID. See: Adenosine Diphosphate. (Schwarz).

ADENYL PYROPHOSPHORIC ACID. See: Adenosine Triphosphate.

ADENYLTRIPHOSPHORIC ACID. See: Adenosine Triphosphate.

ADEPSINE OIL. See: Petrolatum Liquid. (Various Mfr.).

ADESTRIN OINTMENT. (Breon) Fish liver oils rich in vitamins A & D. Oint. Tubes 1 oz., 1/4 lb. Jar 1 lb.

Use: Emollient in the treatment of wounds.

ADETATE. (Merck Sharp & Dohme) d-Amphetamine sulfate 30 mg., vit. B-1 30 mg., sod. glycerophosphate 240 mg., pot. glyc-