


**LECTURE NOTES ON
General Surgery**

HAROLD ELLIS

and

ROY YORKE CALNE

SIXTH EDITION



Introduction

The ideal medical student at the end of his clinical course will have written his own textbook – a digest of the lectures and tutorials he has assiduously attended and of the textbooks he has meticulously read. Unfortunately few men are perfect, and most approach the qualifying examinations depressed by the thought of the thousands of pages of excellent and exhaustive textbooks wherein lies the wisdom required of them by the examiners.

We believe that there is a serious need in these days of widening knowledge and expanding syllabus for a book which will set out briefly the important facts in general surgery which are classified, analysed and as far as possible rationalized for the revision student. These lecture notes represent our own final year teaching; they are in no way a substitute for the standard textbooks but are our attempts to draw together in some sort of logical way the fundamentals of general surgery.

Because this book is written at student level, principles of treatment only are presented, not details of surgical technique.

These notes cover general surgery; ophthalmology and ENT are already dealt with by lecture notes published by our colleagues, Mr Trevor Roper and Mr Miles Foxen, at Westminster Hospital.

The need for a sixth edition has enabled us to insert a new chapter on post-operative fluid and electrolyte balance, to carry out a detailed revision of the whole text and to add to, and revise, the illustrations.

HAROLD ELLIS
ROY YORKE CALNE

Westminster Medical School, London
Addenbrooke's Hospital, Cambridge
1982

Acknowledgments

We are grateful to our colleagues – registrars, housemen and dressers at Westminster and Addenbrooke's – who have read and criticized this text during its production and to many readers and reviewers for their constructive criticisms.

We have received particular help in this edition from Mr Peter Blenkinsopp (Oral Cavity), Mr John Hollingdale (Spine and Peripheral Nerves), Mr Russell Lock (Urology) and Mr John Shaw (Fluid Balance and Thyroid).

Finally, we should like to acknowledge the great and continued help given by Mr Per Saugman and his staff at Blackwell Scientific Publications.

H.E.
R.Y.C.

Contents

Introduction	v
Acknowledgments	vi
1 Acute Infections	1
2 Specific Infections	4
3 Shock	10
4 Burns	13
5 Post-operative Complications	19
6 Post-operative Fluid and Electrolyte Balance	31
7 Tumours	35
8 Chest and Lungs	40
9 The Heart and Great Vessels	54
10 The Peripheral Arteries	70
11 The Veins of the Legs	85
12 The Lymph Nodes and Lymphatics	92
13 The Brain and Meninges	95
14 Head Injuries	106
15 The Spinal Cord	120
16 Peripheral Nerve Injuries	132
17 The Mouth and Tongue	140
18 The Gums and Jaws	153
19 The Salivary Glands	157
20 The Oesophagus	164
21 The Diaphragm	174
22 The Stomach and Duodenum	180

23	Mechanical Intestinal Obstruction	202
24	Specific and Special Forms of Obstruction	208
25	The Small Intestine	217
26	Acute Appendicitis	222
27	The Colon	228
28	The Rectum and Anal Canal	242
29	Peritonitis	257
30	Paralytic Ileus	266
31	Hernia	269
32	The Liver	281
33	The Gall Bladder and Bile Ducts	301
34	The Pancreas	312
35	The Spleen	321
36	The Kidney and Ureter	324
37	The Bladder	345
38	The Prostate	353
39	The Male Urethra	364
40	The Penis	368
41	The Testis	372
42	The Neck	383
43	The Thyroid	387
44	The Parathyroids	402
45	The Adrenal Medulla	405
46	The Adrenal Cortex	407
47	The Thymus	411
48	The Breast	413
49	The Skin and its Adnexae	430
50	Transplantation Surgery	450
	Index	456

CHAPTER 1

Acute Infections

There is an important general principle in treating acute infection anywhere in the body: antibiotics are invaluable when the infection is spreading through the tissues (e.g. cellulitis, peritonitis, pneumonia), but drainage is essential when abscess formation has occurred.

CELLULITIS

Cellulitis is a spreading inflammation of connective tissues. It is generally subcutaneous, but the term may also be applied to pelvic, perinephric, pharyngeal and other connective tissue infections. The common causative agent is the β haemolytic streptococcus.

The invasiveness of this organism is due to the production of hyaluronidase and streptokinase, which respectively dissolve the intercellular matrix and the fibrin inflammatory barrier.

Characteristically the skin is dark red with local oedema and heat. There may be vesicles and, in severe cases, cutaneous gangrene. Cellulitis is often accompanied by lymphangitis and lymphadenitis, and there may be an associated septicaemia.

Treatment

Immobilization, elevation and antibiotics. If a local abscess forms, this must be drained.

ERYSIPELAS

Erysipelas is a diffuse streptococcal infection of the skin and its underlying lymphatics. It is a notifiable disease. The streptococci enter through a minute breach in the skin, although occasionally a wound or burn may be so affected. It particularly occurs on the face and neck. There is intense local pain, the skin is red and raised and there is profound toxæmia.

Treatment

Antibiotic therapy.

ABSCESS

An abscess is a localized collection of pus, usually, but not invariably, produced by pyogenic organisms; occasionally a sterile abscess results from the injection of irritants into soft tissues (for example, thiopentone).

An abscess commences as a hard, red, painful swelling which then softens and becomes fluctuant. If not drained, it may discharge spontaneously onto the surface or into an adjacent viscus or body cavity. There are the associated features of bacterial infection; a swinging fever, malaise, anorexia and sweating with a polymorph leucocytosis.

Treatment

An established abscess in any situation requires drainage. Chemotherapeutic agents cannot diffuse in sufficient quantity to sterilize an abscess completely. Pus left undrained continues to act as a source of toxæmia and becomes surrounded by dense fibrous tissue.

The technique of abscess drainage depends on the site. The classical method, which is applicable to a superficial abscess, is to wait until there is fluctuation and to insert the tip of a scalpel blade at this point. The track is widened by means of sinus forceps, which can be inserted without fear of damaging adjacent structures. If there is room, the surgeon's finger can be used to explore the abscess cavity and break down undrained loculi. Drainage is then maintained until the abscess cavity heals from below upwards, otherwise the superficial layers can close over, with recurrence of the abscess. The cavity is therefore kept open by means of a gauze wick, a corrugated drain or a tube; the drain is gradually withdrawn until complete healing is achieved.

BOILS

A boil (furuncle) is an abscess, usually due to the pyogenic staphylococcus, which involves a hair follicle and its associated glands. It is therefore not found on the hairless palm or sole, but is usually encountered where the skin is hairy, injured by friction, or dirty and macerated by sweat;

thus it occurs particularly on the neck, axilla and the perianal region. Occasionally a furuncle may be the primary source of a staphylococcal septicaemia and be responsible for osteomyelitis, perinephric abscess or empyema, particularly in debilitated patients. On the face it may be complicated by a cavernous sinus thrombosis, via the facial veins.

Differential diagnosis

Multiple infected foci in the axillae or groins due to infection of the eccrine sweat glands of these regions (*hydradenitis suppurativa*) are usually misdiagnosed as boils. They do not respond to antibiotic therapy and can only be treated effectively by excision of the affected skin area; if this is extensive, the defect may require grafting.

Treatment

When pus is visible the boil should be incised. Recurrent crops of boils should be treated by improving the general hygiene of the patient, and by the use of ultra-violet light and hexachlorophene baths, but systemic chemotherapy is not indicated.

CARBUNCLES

A carbuncle is an area of subcutaneous necrosis which discharges onto the surface through multiple sinuses. It is usually staphylococcal in origin. The subcutaneous tissues become honeycombed by small abscesses separated by fibrous strands. The condition is often associated with general debility and particularly with diabetes. The urine should always be tested for sugar in this or any other septic condition.

Treatment

Surgery is rarely indicated initially. Chemotherapy is given and the carbuncle merely protected with sterile dressings. Occasionally a large sloughing area eventually requires excision and a skin graft. Diabetes, if present, should be controlled.

CHAPTER 2

Specific Infections

TETANUS

Pathology

Tetanus is caused by the *Clostridium tetani*, an anaerobic, flagellated, exotoxin-secreting and gram-positive bacillus, which forms a characteristic terminal spore ('drumstick'), and which is a normal inhabitant of soil and faeces. The organism remains at the site of inoculation and produces a powerful exotoxin which acts upon the motor cells in the CNS and which is probably conveyed along the peripheral nerves directly from the affected part.

Tetanus follows the implantations of spores into a deep, devitalized wound where anaerobic conditions occur. Infection is related less to the severity of the wound than to its nature; thus an extensive injury which has received early and adequate wound toilet is far less at risk than a contaminated puncture wound which has been neglected. Occasionally dressings or catgut which have been contaminated with tetanus spores are the source of infection of surgical wounds. In primitive communities, where dung is used to dress the umbilical cord in the newborn, *tetanus neonatorum* may occur.

Clinical features

The incubation time is 24 hours to 24 days. Muscle spasm first develops at the site of inoculation and then involves the facial muscles and the muscles of the neck and spine. As a rule it is the trismus of the facial spasm (producing the typical 'risus sardonicus', or 'lock-jaw' to the layman) which is the first reliable indication of developing tetanus. This may be so severe that it becomes impossible for the patient to open his mouth. The period of spasm is followed, except in mild cases, by violent and extremely painful convulsions which occur within 24-72 hours of the onset of symptoms and may be precipitated by some trivial stimulus, such as a sudden noise. The convulsions, like the muscle spasm, affect the muscles of the neck, face and trunk. Characteristically, the muscles

remain in spasm between the convulsions. The temperature is a little elevated but the pulse is rapid and weak.

In favourable cases the convulsions, if present at all, become less frequent and then cease and the tonic spasm gradually lessens. It may however, be some weeks before muscle tone returns to normal and the risus sardonicus disappears. In fatal cases paroxysms become more severe and frequent; death occurs from asphyxia due to involvement of the respiratory muscles or from exhaustion, inhalation of vomit, or pneumonia.

The prognosis is serious when the incubation period from the time of injury to the onset of spasm is under 5 days and when convulsions occur within 48 hours of the onset of muscle spasm.

Differential diagnosis

1. Tetany – which characteristically affects the limbs, producing carpopedal spasm.
2. Strychnine poisoning – flaccidity occurs between convulsions whereas in tetanus the spasm permits.
3. Meningitis – because of the neck stiffness.
4. Epilepsy.
5. Hysteria.

Treatment

Prophylaxis

Active immunization comprises two initial injections of tetanus toxoid (formalin treated exotoxin) at an interval of 6 weeks. Booster doses are given at intervals of not more than 7 years, or at the time of any injury. Toxoid should be given to any population at risk of injury, for example, service personnel.

The risk of tetanus can be reduced almost to zero if penetrating and contaminated wounds are adequately excised to remove all dead tissue and a course of prophylactic penicillin (or tetracycline in sensitive patients) is given. Patients who have previously received toxoid should be given a booster dose. If toxoid has not been given in the past, human gamma globulin, prepared from fully immunised subjects, should be given if the wound is heavily contaminated.

There is no justification for the use of anti-tetanus serum as a passive immunization agent. There has never been a controlled trial of the value of ATS, severe reactions may occur, particularly if serum therapy has

been given in the past, and skin sensitivity tests to a small subcutaneous dose give no reliable guide to subsequent severe reactions. Tetanus may occur after ATS has been given and, at present, it seems that the risk of mortality from serum is of the same order as that of an unimmunized subject acquiring tetanus after injury.

Curative treatment

1. *Control convulsions.* The patient is nursed in isolation, quiet and darkness and is heavily sedated with phenobarbitone or chlorpromazine. In severe cases curarization with tracheostomy and intermittent positive pressure artificial respiration is required and this may have to be continued for up to 4 weeks. It is terminated when the spasms and rigidity are absent during a trial period without relaxants. These serious cases are best transferred to a special respiratory unit.

2. *Control the local infection.* Excision and drainage of any wound is carried out under a general anaesthetic. Penicillin or tetracycline are administered and these will also act as a prophylactic against pulmonary infection.

3. *Maintain the general condition and electrolyte balance* of the patient by naso-gastric tube feeding.

The value of large doses of ATS (100 000 units intramuscularly and 100 000 units intravenously) is not established and its use carries with it the danger of a severe serum reaction. However, toxoid is given if previous active immunization has been carried out. Human gamma globulin is safe and effective.

GAS GANGRENE

Pathology

Results from infection by *Clostridia welchii*, *septicum* and *sporogenes*; anaerobic, encapsulated, spore forming, gas-producing, gram-positive organisms which produce an exotoxin. This group includes both proteolytic and saccharolytic organisms. The characteristic gas formation in the tissues is produced by the liberation of CO_2 , H_2S and NH_3 by protein destruction. The organisms are found in soil and in faeces.

Typically gas gangrene is an infection of deep penetrating wounds, particularly of war, but sometimes involvement of the abdominal wall or cavity may follow operations upon the alimentary system. Occasionally gas gangrene complicates amputation of an ischaemic lower limb, or follows abortion or puerperal infection.

Clinical features

The incubation period is about 24 hours. Toxaemia is severe with tachycardia, shock and vomiting. The temperature is first somewhat elevated and then becomes sub-normal. The affected tissues are swollen and crepitant due to gas. The skin becomes gangrenous and the infection spreads along the muscle planes, producing at first dark red swollen muscle and then frank gangrene.

Treatment

Prophylaxis

Consists of adequate excision of wounds which removes both the organisms and the dead tissues which are essential for their anaerobic growth. Seriously contused wounds (such as those produced by a gunshot wound) or contaminated wounds are left open and lightly packed with gauze. Delayed primary suture can then safely be performed after 5 or 6 days, by which time the wound is usually healthy and granulating. Penicillin given in all heavily contaminated wounds and to patients with atherosclerosis undergoing amputation of the leg.

Curative

In the established case, all involved tissue must be excised. Implication of all muscle groups in a limb is an indication for amputation. Penicillin and blood transfusion are given. Hyperbaric oxygen therapy, to eliminate the anaerobic environment, is theoretically sound but as it is combined with all the other modalities of treatment, its efficacy cannot be judged. If a hyperbaric chamber is available, it should certainly be employed.

The value of anti-gas gangrene serum both as a prophylactic and curative measure is not established.

ANTHRAX

Caused by the bacillus anthracis which is gram-positive, encapsulated and spore forming. It is a disease of cattle and sheep, which affects men coming into contact with these animals, e.g. leather workers, wool workers and veterinary surgeons. Anthrax is usually a skin infection, but occasionally it is transmitted by inhalation.

The incubation period is 1-2 days with the formation of a 'malignant pustule', which is a central black slough surrounded by vesicles and

oedema. Inhalation of the organisms produces a highly fatal pneumonia ('wool-sorters' disease').

Diagnosis is confirmed by examination of a swab from the pustule or the sputum.

Treatment

A vaccine is available for workers at risk of anthrax. Penicillin in large doses is now the treatment of choice, and since the antibiotic era, the mortality is under 1 per cent.

ACTINOMYCOSIS

Pathology

Actinomycosis is an infection produced by the actinomyces israelii or ray fungus, so called because the mycelial threads may be seen radiating from the main fungal mass in culture. The fungus is micro-aerophilic and exists as a saprophyte in the mouth (especially where there is dental caries) and in the alimentary canal. Infection may occur via a breach in the mucous membrane, for example following dental extraction, and produces a dense fibrous tissue reaction within which pockets of pus develop. The pus contains typical 'sulphur granules', which are yellow specks of mycelium. The infection spreads along the fascial planes and occasionally by the blood-stream, but not via the lymphatics.

Clinical features

Actinomycosis can be classified into three main groups: cervico-facial, abdominal and pulmonary.

Cervico-facial

This form occurs typically after dental extraction or tonsillitis. Although actinomyces do grow on grasses and decayed vegetable matter, these varieties are not pathogenic in man and the infection does not occur, as was once taught, by chewing contaminated straw. Nor is the disease transmitted from cattle or horses to man. Swelling occurs over the angle of the jaw and the adjacent tissues become greatly indurated. The skin develops a typical bluish discoloration, then sinuses appear, which discharge thin pus. Pain may or may not be a feature, but there is usually marked trismus. Spread may occur by direct infiltration to the orbit, base of skull, jaw, or mediastinum.

Abdominal actinomycosis

Usually located in the ileo-caecal region and follows upon an attack of perforative appendicitis, a perforated peptic ulcer or an abrasion of the alimentary mucosa by some foreign body. A hard fibrous mass, honey-combed with abscess cavities, develops in the right iliac fossa and multiple sinuses may appear on the abdominal wall. Spread may occur via the portal vein producing a portal pyaemia, the liver being riddled with abscess cavities.

Pulmonary actinomycosis

May follow inhalation of fungus from the infected mouth. Spread occurs through the lung to the pleura and eventually the chest wall. Pulmonary disease may also occur secondary to spread from the neck via the mediastinum, or from the abdominal cavity through the diaphragm.

Treatment

Comprises a 12-week course of daily injections of penicillin. Obvious collections of pus should be drained. The actinomyces should be tested for sensitivity and occasionally other antibiotics, e.g. tetracycline, may be required.

CHAPTER 3

Shock

Shock is the term used to describe a clinical state comprising pallor, sweating, coldness and peripheral cyanosis. The pulse is usually rapid and the blood pressure low. In severe cases there may be dyspnoea, thirst, nausea or vomiting. The patient may be confused and restless or be semi-conscious.

Aetiology

Shock is produced by a wide variety of circumstances, the common factor being a *reduction in the effective circulating blood volume*. This clinical picture may be seen in:

1. *Severe haemorrhage* – an actual reduction of blood volume.
2. *Extensive fluid loss* as a result of exudation of plasma from burns, or loss of extra-cellular fluid in severe vomiting or diarrhoea.
3. *The vasovagal syndrome*, produced by severe pain or emotional disturbance. The mechanism of this is reflex vasodilation in muscle together with vagal cardiac slowing. This syndrome can be recognized because the shock picture is accompanied by *slowing* of the heart and responds to the simple measure of lying the patient flat with elevation of the legs.
4. *Severe toxæmia*, as in peritonitis, septicaemia (particularly gram-negative organisms) or pancreatitis. Here there is a combination of fluid loss into the extravascular space, pain and the effect of chemical or bacterial toxins on the heart; once again the shock picture is produced by circulatory failure.
5. *Heart failure* from myocardial infarction or pulmonary embolus.
6. *Sympathetic interruption* which reduces the effective blood volume by widespread vasodilatation; for example, the spinal shock following transection of the spinal cord, or after a high spinal anaesthetic.

The physiological basis of haemorrhagic shock

Severe haemorrhage produces the following chain of events: reduction in blood volume – diminution in the venous return to the heart – fall in cardiac output (Starling's law: the output depends on the degree of

stretch of the heart muscle in diastole) – fall in blood pressure – this is counteracted by the carotid sinus and aortic arch reflexes, which increase the heart rate, and by sympathetic vasoconstriction of the splanchnic bed and of the peripheral cutaneous vessels.

This mechanism maintains essential coronary, cerebral and lung blood flow. The blood pressure remains at first relatively normal, but continued haemorrhage eventually reaches a stage which can no longer be compensated and the blood pressure then falls.

The clinical features of shock which have already been described are thus easily explicable on this physiological basis. The intense peripheral vasoconstriction produces the cold, pale skin. The rapid pulse and low blood pressure are typical features of the impaired cardiac output. The sweating results from sympathetic overactivity. The cerebral disturbances follow inadequate perfusion of the brain.

A continued low blood pressure produces a series of irreversible changes so that the patient may die in spite of later blood replacement. The oxygen lack affects all the vital organs; there may be tubular necrosis of the kidney resulting in renal failure, the adrenals may lose their normal reaction to stress, the heart may fail due to inadequate coronary perfusion and there may be damage to the cardiac and vasomotor centres in the medulla. In the tissues themselves anoxia produces capillary paralysis and dilatation so that a copious fluid loss occurs into the interstitial spaces.

With intensive treatment, life may be preserved despite continuing disease. In such instances, the condition of 'shock lung' may develop. There is progressive impairment of pulmonary gaseous perfusion which may result from multiple blood transfusions causing platelet and fibrin pulmonary microemboli, toxicity from prolonged oxygen therapy and opportunist lung infection. Eventually the build up of 'sick cells' and tissue necrosis results in death – *irreversible shock*.

Treatment

This depends on diagnosis. A vasovagal attack, or faint, rapidly responds to lowering the head and elevating the legs; if the patient does not improve after this, it is suggestive that some complicating factor such as internal haemorrhage coexists with the vasovagal syndrome. It is important to know that considerable loss of blood into the tissues occurs with major fractures of the limbs even if these are not compound.

Where haemorrhage is the cause of the shock the following steps are taken:

1. Further haemorrhage is arrested; this may require direct pressure to

a wound or surgical exploration where continued bleeding is the result of a peptic ulcer haemorrhage, ruptured spleen or ruptured ectopic pregnancy.

2. Blood transfusion to replace the blood loss.
3. Relief of pain by means of an injection of morphia where pain is a marked feature.
4. Elevation of the foot of the bed is a quick and effective temporary means of raising the blood pressure and is a useful emergency measure.

Excessive warmth should be avoided; this produces vasodilatation of the skin vessels thereby diverting available blood from the vital tissues.

Oxygen is seldom required since the blood is usually fully oxygenated unless there is an associated chest injury or respiratory depression due to a head injury.

Shock from other causes may require appropriate fluid replacement; plasma or plasma substitute in the case of burns, or saline in severe vomiting or diarrhoea.

In long continued stress there may be adrenocortical depletion and intravenous hydrocortisone should be administered. In toxæmic patients, for example those with peritonitis, antibiotics are given in heavy dosage.

Regular and frequent monitoring of the central venous pressure is an aid in the correct replacement of fluids.

The management and monitoring of the critically-ill patient

The severely-shocked patient should be admitted to an intensive care ward where his continuous supervision by specially-trained nursing staff is available. As well as careful clinical surveillance, the following need to be monitored:

Rectal temperature, pulse, respiration rate and blood pressure.

Central venous pressure.

Urine output (via catheter).

ECG.

Serum electrolytes, haemoglobin and white blood count.

Arterial P_{O_2} , P_{CO_2} , pH.

The frequency of these measurements depends on the patient's condition and response to treatment. It is particularly important to remember that, in this environment of recording machinery and scientific nursing, the patient remains a human being, who deserves to be treated with dignity and tenderness. If he is conscious he may well be terrified, in pain and acutely aware of all that is going on around him.