

The Autonomic Nervous System

An introduction to basic and clinical concepts

Third revised and enlarged edition

OTTO APPENZELLER

Foreword to the third revised edition

It has been my privilege to pen a few orienting remarks to each edition of Professor Appenzeller's opus on *The Autonomic Nervous System*. And it has permitted me the chance to follow the prodigious advances in this field of neurology during more than a decade. No doubt when looked at in historical perspective at some future time the scholarly efforts of Professor Appenzeller will be recognized for having stimulated and maintained interest in this field.

Since the appearance of the second edition of this book the most exciting developments in studies of the nervous system have been the discovery of the role of many neuropeptide transmitters and hormones. The synthesis, control mechanisms, function, and pathological changes in these complex molecules are coming to the forefront of neurology. In a sense these new insights have in large measure been derived from studies of the central and peripheral autonomic nervous system. Actually, the first specific markers for choline acetyltransferase, the enzyme that synthesizes acetylcholine in the presynaptic sympathetic terminals, and tyrosine hydroxylase, the rate-limiting enzyme in catecholamine biosynthesis, were worked out in relation to the autonomic nervous system. These have been applied to the study of the development of the autonomic nervous system as well as many of its diseases.

The "neuropeptide era" of brain and nerve anatomy and physiology holds interesting prospects for students of neurology, and in this edition Professor Appenzeller undertakes the formidable task of presenting all these new developments in neurobiology in lucid and concise form. But surely he is equal to the challenge, if past attempts are any measure of his capacity.

Raymond D. Adams, M.D.
Bullard Professor of Neuropathology
Emeritus
Harvard Medical School
Cambridge, MA
U.S.A.

Foreword (second revised edition)

A great part of man's nervous system is concerned with activities of which he is essentially unconscious and over which he has little or no voluntary control. Claude Bernard opined that perhaps a divine providence considered them too important to entrust to a capricious will. "We are thrust into this world by smooth muscle, which is under the control of the autonomic nervous system. From moment to moment we are dependent for our conscious existence on the moderate contraction of the blood vessels, routinely kept in this state by autonomic impulses. Most of the complicated processes of digestion, from the initial outpouring of saliva to the final riddance of waste, require the participation of autonomic nerves. Any vigorous exercise in which we may engage depends upon cooperation of autonomic government of appropriate effectors; thus throughout eons of past time the physical struggle for existence has been made possible by that government . . . which preserves the stable states of the fluid matrix that are required for ready response to every call to action."

Once the brilliant studies of Gaskell and Langley had brought to light the importance of the involuntary or vegetative nervous system to the efficient functioning of the human organism, physiologists the world over, led by Walter B. Cannon whose ideas are quoted above, set themselves to elucidate the principles of its action in such fundamental activities as the maintenance of "le milieu interne" necessary for health, the continuation of the race and the natural reflex mechanisms of controlling glands and smooth muscles. But until comparatively recent times, relatively neglected was the manner in which the autonomic nervous system participates in medical and neurological diseases, or itself may be selectively deranged by pathologic processes.

In the past decade or two, clinical investigators have begun to rectify this deficiency. A number of singular advances in the neurology of the autonomic nervous system have been made. A form of pure autonomic polyneuropathy has been identified and in several other types of polyneuropathy the autonomic and pain fiber bear the brunt of the disease. A series of central nervous system diseases have been discovered in which spinal and hypothalamic centers degenerate. And for the first time the pathology of the sympathetic nervous system has

been surveyed in most of the important medical and neurological diseases. Neuropathologic experiments have been done on the autonomic nervous system in animals in the hope of clarifying human diseases.

Professor Appenzeller has presented the medical profession with a valuable monograph devoted to these and other pertinent topics. The warm reception given the first edition and the surge of new information that he has garnered from the recent literature have encouraged him to update the subject and favor us with a second edition. His efforts should reward the interested reader whether of neurological or general medical bent.

Raymond D. Adams, M.D.

Foreword (first edition)

It seems paradoxical that progress in understanding of diseases of different parts of man's nervous system is almost in inverse proportion to the complexity of the structures involved. More explicitly, we appear to be much more familiar with diseases of the complex cerebral and cerebellar cortices or the basal ganglia than we do about those of the peripheral nervous system, which is infinitely simpler and more uniform in structure. And, of the many diseases of the peripheral nervous system those of the autonomic parts are certainly the most obscure. It is not that medical scientists have neglected the autonomic nervous system, for its anatomy and physiology are the subjects of a voluminous literature, but rather that its clinical disorders and their pathology have not been studied systematically by appropriate histological techniques. The few articles that have appeared have consisted of little more than blind surveys of random samples of sympathetic ganglia and nerves and, as expected, were notably uninformative.

Professor Appenzeller reminds us that in recent years this unfortunate state of affairs is being repaired to some extent. Gradually and almost unobtrusively new diseases that exert their predominant effects on the sympathetic and parasympathetic nervous system have been discovered. When he places them alongside one another, they obviously have come to represent a fairly respectable body of clinical medicine and neuropathology. Moreover, it appears that scattered studies of the autonomic nervous system, in some of the standard diseases of man's brain and spinal cord, are beginning to yield unexpected results. In Parkinson's disease, for example, den Hartog Jager has found Lewy type of inclusion bodies in the sympathetic ganglion cells in a large proportion of cases, which may explain their well-known tendency to orthostatic hypotension and fainting; and other examples are to be found.

The chief merit of this work is that it brings together in the confines of one small monograph the salient facts about this new group of autonomic nervous diseases. But it does more for the neurologically oriented reader. Being himself knowledgeable about the physiological techniques whereby this part of the nervous system can be evaluated in living man the author is able to review in a useful fashion some of the current clinical neurophysiological methods for the study of this part of the nervous system. And, finally in

the interest of presenting a well-rounded view of his subject, he concludes with an extensive discussion of the role of the central integrative centers of the autonomic nervous system in some of the more complex biological activities of man.

This small monograph should prove to be a welcome addition to a growing number of systematic studies of the peripheral nervous system.

Raymond D. Adams, M.D.

Introduction to the third revised edition

This new edition needs no apology. Interest in the autonomic nervous system has continued since the appearance of the Second Edition in 1976 and enthusiastic delving into basic concepts related to autonomic nervous system function and their clinical application has led to giant strides in understanding of the complexity and the importance of the autonomic nervous system in homeostasis. Insight into new central and peripheral neurotransmitters closely associated with autonomic activity has revolutionized our concepts of pain perception, of hunger, thirst, and of the adaptation of the human body to exercise, an activity now enthusiastically embraced by millions of previously physically inactive citizens. This new knowledge has, fortunately, been easily superimposed upon the foundations of basic and clinical concepts of autonomic nervous function which have stood the test of time and continue to occupy their rightful place in this new edition. Further impetus to the revision of the Second Edition was given by the influence of basic aspects of autonomic activity on other branches of clinical medicine apart from neurology and by the birth of new disciplines. A host of specialists working in the areas of pain, headache, and sports medicine where autonomic nervous system function is of great importance need up-to-date access to material scattered in widely different publications. For them, the Third Edition will provide an introduction to autonomic nervous system activity and to the concepts of its role and importance in adaptation and homeostasis.

The newly recognized central neurotransmitters have opened the possibility of novel therapeutic approaches, and mention of some tentative beginnings in these areas has been made.

The usual medical curriculum teaches the autonomic nervous system in fragments, the basic sciences touching upon it in physiology, pharmacology, and anatomy; the clinical sciences in neurology and cardiovascular and respiratory "blocks", but it is during the teaching of neurology that an integrated view of autonomic nervous system activity is best provided. Unfortunately, most medical schools' curricula do not allow sufficient time to cover this important area. Because of this, the interested student and physician will find the Third

Edition a useful source and an overview to basic and clinical concepts of autonomic nervous system function.

Otto Appenzeller
Albuquerque, New Mexico
July 1981

Introduction (second revised edition)

The rather modest size of the first edition published in 1970 has grown to the present volume. This increase in size requires some explanation. Over the past few years, clinical and pharmacologic research has led to new insights into the pathogenesis and therapeutic possibilities of disorders affecting the autonomic nervous system. A particularly noticeable advance was made in the understanding of neurochemical processes and clinical manifestations associated with deficiencies of some central and peripheral neurotransmitter mechanisms. Impetus to the publication of the second edition was given by the widespread appreciation of autonomic dysfunction in disorders of many systems and the need to attend to autonomic clinical deficits in the management of patients with a variety of diseases.

In the past, the assessment of autonomic deficits was largely based on anecdotal accounts and patients with predominant autonomic dysfunction were often reported as medical exotica. This early stage of the advancement of knowledge is now fortunately past and objective tests of autonomic function can now be applied to help pinpoint the site of lesions. Therapeutic possibilities have become a reality and the wide dissemination of knowledge to the lay public about autonomic dysfunction has made it necessary for practicing physicians to become thoroughly acquainted with this part of nervous activity in health and disease. These considerations and the wide scatter of recent advances in this area in disparate publications has led to the collection of up-to-date relevant information in the second edition.

Though the plan of the first edition has been preserved, new chapters have been added including some on biofeedback, monoamines and autonomic dysfunction in hypertension. Because of widespread autonomic dysfunction in patients with spinal lesions, a chapter is devoted to the clinical significance of these disorders. Once again, I am most grateful to ASP Biological and Medical Press for their expert help and unflagging support in bringing this project to fruition.

Otto Appenzeller

Introduction (first edition)

This book does not intend to give a complete survey of the autonomic nervous system, rather it is the aim to present to the student and practicing physician, useful information which might help in the understanding of autonomic function and the application of this information to relevant clinical settings.

In the last decade, tremendous advances have been made with the use of sophisticated methods of investigation. Clinicians, however, have developed a fairly rigid view of autonomic function, apparently uninfluenced by experimental results. It is the purpose of this book to select experimental findings of clinical significance and put them into perspective by comparing them with clinical data on autonomic dysfunction.

Anatomically, the term 'system', as distinct from an organ, is used to designate tissue which might be in different topographic areas of the body, but which is functionally related. In the autonomic nervous system, two functional subdivisions can be recognized. The ergotropic system which is concerned with preparedness of the body and so-called 'animal functions', and the trophotropic system which is mainly responsible for the fluctuating activity of organs. The ergotropic system corresponds peripherally with the sympathetic nervous system, whereas the trophotropic system is predominantly parasympathetic. In the central nervous system this anatomical distinction becomes blurred, but the structures concerned with sleep may be part of the trophotropic system.

The autonomic nervous system is concerned with the regulation of three main functions; the somatic or animal function, by which responses to changes in environment are made predominantly by appropriate active measures; the regulation of organ function, which is concerned with rest and activity of organs; and endocrine function, which influences the entire organism. The endocrine system is closely integrated with nervous activity through the hypothalamus and pituitary centrally, and also peripherally through the release of neurohumors in response to appropriate stimuli. This system is dealt with extensively in neuroendocrine texts, and will only briefly be considered in this book.

While in the past physiological experimentation was closely tied to morphology, more

recently pharmacology and neurochemistry have been used to investigate the function of the autonomic nervous system. Nowadays electronmicroscopy and histochemistry once again allow a close association between morphology and physiology advancing knowledge at the cellular level. For this reason, in this book anatomical and physiological considerations precede clinical discussions.

Otto Appenzeller

Acknowledgements

The Third Edition owes its appearance to selfless contributions and continued interest of many faculty members of the University of New Mexico School of Medicine, of the resident staff, particularly Edward Collins, D.O., and of interested students. Their number makes it impossible to acknowledge them, but one individual whose encouragement and interest was particularly helpful must be mentioned. Yen Tsai, M.D., Professor of Neurology at the Third People's Hospital in Shanghai and visiting World Health Organization Fellow in the Department of Neurology, has given impetus to updating and improving this edition in areas in which the Second Edition was deficient, and to him I extend special thanks.

The understanding support of my wife and family has continued throughout the revision and my gratitude to them must be recorded. Katherine Miller has devoted many hours and meticulous care towards the production of this volume, and she deserves all the credit for its timely appearance.

Many illustrations have been obtained from colleagues and other sources. Reference to their generosity and to the original publications appears in the legends. The full source of previously published material is acknowledged in the references at the end of the book. I am indebted to Elsevier Biomedical Press for their interest, support, and encouragement in bringing this new edition to the reader before it is outdated.

Acknowledgements (second revised edition)

This revised and enlarged edition would not have been possible without generous contributions and helpful discussions with many students and residents in Neurology. My indebtedness is particularly great to Mr. Gary Ogin. Thanks are due to Mrs. Connie Sokolowski for typing the manuscript. Again, I am most grateful for the continued support of my wife and family. Lastly, the publishers of the first edition have generously allowed the use of material previously published.

Some illustrations have been obtained from other sources. In all cases reference is made to the original publication in the legend. The full source is recorded in the reference list at the end of the book. Permission for reproducing this material is gratefully acknowledged.

Acknowledgements

(first edition)

This book would not have been written without the understanding support of my wife, to whom it is dedicated. I am also grateful to her for reading the manuscript. My thanks are due to Mrs. Grace Wilson, Mrs. Polly Gauthier and Mrs. Vi Farmer for typing the manuscript. Many students and residents in neurology have contributed by helpful discussions, but my indebtedness is particularly great to Drs. T.K. von Storch and G.B. Marcus, and to Mr. D. Scott. Finally, the editors and publishers of the Handbook of Clinical Neurology have generously allowed the use of material previously published.

Contents

<i>Foreword to the third revised edition</i> by Raymond D. Adams, M.D.	v
<i>Foreword (second revised edition)</i> by Raymond D. Adams, M.D.	vii
<i>Foreword (first edition)</i> by Raymond D. Adams, M.D.	ix
<i>Introduction to the third revised edition</i>	xi
<i>Introduction (second revised edition)</i>	xiii
<i>Introduction (first edition)</i>	xv
<i>Acknowledgements</i>	xvii
<i>Acknowledgements (second revised edition)</i>	xix
<i>Acknowledgements (first edition)</i>	xxi
<i>Chapter 1</i> Anatomy and histology	1
<i>Chapter 2</i> Temperature regulation	31
<i>Chapter 3</i> The neurogenic control of the circulation	109
<i>Chapter 4</i> The autonomic nervous system and hypertension	201
<i>Chapter 5</i> Catecholamines and neurologic disease	209
<i>Chapter 6</i> Operant conditioning and biofeedback	217
<i>Chapter 7</i> Trophic disorders	223
<i>Chapter 8</i> Sleep and wakefulness	237
<i>Chapter 9</i> Respiration and disorders of ventilation	251
<i>Chapter 10</i> Disturbances of water and electrolyte balance and disorders of innervation of the gastrointestinal tract	267
<i>Chapter 11</i> Circadian rhythms	301
<i>Chapter 12</i> The autonomic nervous system and the pineal gland	305

Chapter 13	Pain syndromes	309
Chapter 14	The normal pupil and some pupillary abnormalities	319
Chapter 15	Neurogenic control and disorders of micturition	339
Chapter 16	The reflex control of copulatory behavior and neurogenic disorders of sexual function	381
Chapter 17	Autonomic function, its pathophysiology and clinical assessment in paraplegia and tetraplegia	397
References		411
Subject index		475

CHAPTER 1

Anatomy and histology

Galen (quoted in *Opus de Usu Partium Corporis Humani*, Paris, 1528) seems to have been the first to refer to a part of the autonomic nervous system when he described a nerve trunk lying along rib heads and recognized its connecting fibers with the spinal cord. He thought that this was a branch of the vagus and believed that through it the viscera received sensitivity from the brain and power to move from the spinal cord. Galen also observed at least three enlargements along the course of the sympathetic chain and described the ganglion at the entrance of the chain into the abdomen, which might have been the semilunar ganglion of the celiac plexus. He suggested, a widely accepted belief thereafter, that there was 'sympathy' or 'consent' between body parts and regarded the peripheral nerves as tubes through which animal spirits were distributed to bring about this 'sympathy'.

Galen's belief that the vagi and sympathetic trunks were a single unit, both functionally and anatomically, was shared by subsequent anatomists until Estienne (1545) recognized the sympathetic trunks as a distinct anatomical structure. Willis (1664) gave the name of 'intercostal nerve' to the sympathetic trunk and believed that the cerebellum was responsible for involuntary movements which he distinguished from voluntary motion.

Although Willis recognized the innervation of the heart by the vagus nerve, the functional significance of this was not discovered until Lower's (1669) description of the effects of vagus section on heart rate.

The suggestion that involuntary movements are initiated by local stimulation due to nerve irritation is attributed to Whytt (1751) and he used this as an explanation for the reaction of the pupil to light. Whytt's importance to neurophysiology is further enhanced by his suggestion that all 'sympathy' or 'consent' must be referred to the central nervous system since it occurs between body parts whose nerves make no connection with each other so that the transmission of 'sympathy' cannot involve the flux of matter and must therefore occur in an area where all nerves have their origin (Whytt, 1765).

Du Petit (1727) pointed out that the sympathetic trunk was not directly connected with the brain and was a separate structure from the vagus, but it was Winslow (1732) who first designated the paravertebral chain as 'the great sympathetic nerve'.

In 1764 Johnstone said that the movements of the heart and intestine are involuntary

because the sympathetic ganglia blocked the actions of the will and prevented them from reaching these structures and that this blockade also accounted for the relative insensitivity of the viscera.

Studies by Bichat in 1802 led to the concept of 'animal life' and 'organic life'. He pointed to the continuous action which was the hallmark of organic life distinct from the intermittent activity of animal life, a concept which is still widely expressed by the terms 'visceral' and 'somatic' respectively. Bichat was aware of the different appearances of gray and white rami communicantes but did not recognize their significance.

The term 'vegetative nervous system' was used by Reil (1857). He thought that the rami communicantes served as connectors between the animal and vegetative nervous systems. An early description of nerve cell bodies in sympathetic ganglia appeared in Ehrenberg's (1833) writing, together with comments on the microscopic structure of nerve fibers. In the 19th century the ciliary, sphenopalatine, otic and submandibular ganglia were thought of as part of the autonomic nervous system but their functional significance was not appreciated.

Meissner's (1857) mention of the subarachnoid plexus and the description of the myenteric plexus by Auerbach (1864) in the second half of the 19th century concluded the anatomic studies which paved the way for the physiologic work on vasomotor function by Bernard (1878). His studies led to the concept of sympathetic vasoconstrictor action but not until some time later was he able to demonstrate vasodilator nerves in arteries supplying the submandibular gland after stimulation of the chorda tympani. Bernard thought that the sympathetic reflexes were mediated by the spinal cord and that on stimulation of some areas of the brain impulses were discharged through the sympathetic fibers.

Gaskell (1886) gave a detailed description of the anatomy of the rami communicantes and recognized that the efferent fibers within these nerves arise in the spinal cord and that corresponding fibers can be found in some cranial nerves. He also recognized the connection with the central nervous system of the peripherally located ganglion masses through medullated fibers and divided these fibers into bulbar, thoracolumbar and sacral groups. Langley and Dickinson (1889) used the action of nicotine on ganglia to study the relation of the nerve fibers to the peripheral ganglion cells and proposed the name 'autonomic nervous system'. When this term was coined the different distribution and functional effects of the thoracolumbar and craniosacral outflows were known and Langley separated the former from the rest of the autonomic nervous system. After the discovery of substances which either produced actions similar to those obtained by stimulation of the thoracolumbar or of the craniosacral outflows he coined the term 'parasympathetic' for the latter (Langley, 1901).

ANATOMY OF THE AUTONOMIC NERVOUS SYSTEM

Body functions, which can proceed independently of volitional activity, are regulated at least in part by reflex mechanisms served by afferents, efferents and central integrating structures which are included in the autonomic or vegetative nervous system. Although the activity of this system is essentially autonomous, it is not entirely free from voluntary control. The vegetative nervous system is made up of all neurones which lie outside the central nervous system and is concerned with visceral innervation. The only exceptions are those that are part of the afferent system of the cerebrospinal nerves and are located in the posterior root ganglia or some sensory cranial nerve ganglia. The neurones in the brain, brainstem and spinal cord,

through which these autonomic neurones are functionally connected, are also included in the vegetative nervous system.

There are two divisions of this system, the sympathetic and parasympathetic, each usually made up of preganglionic and postganglionic neurones. The cell bodies of the preganglionic neurones lie in the brain or spinal cord and those of the postganglionic neurones are in the thoracic and upper lumbar cord and this part of the vegetative nervous system is also called the thoracolumbar division (Fig. 1). Preganglionic neurones of the parasympathetic nervous system are in the brainstem and sacral cord and this is also termed the craniosacral division (Figs. 2 and 3). The viscera are mostly innervated by both sympathetic and parasympathetic fibers. Many structures supplied by the vegetative nervous system have, however, a single innervation only, such as some blood vessels and sweat glands.

The preganglionic fibers of the sympathetic nervous system arise in the intermediolateral and intermediomedial cell columns of the spinal cord and join the ventral roots of T1 to L2.




Fig. 1. Diagram to show the pre- and postganglionic fibers of the autonomic innervation of the thoracic and abdominal viscera. Dotted lines: postganglionic fibers of the thoracolumbar division. Short solid lines on viscera: postganglionic fibers of craniosacral division. Drawn by M. Norviel.