

Connective Tissues

Biochemistry and Pathophysiology

Edited by R. Fricke and F. Hartmann

Editorial Board

E. Buddecke · R. Fricke · F. Hartmann · H. Muir · K. Kühn



Connective Tissues

Biochemistry and Pathophysiology

Edited by R. Fricke and F. Hartmann

Editorial Board

E. Buddecke · R. Fricke · F. Hartmann · H. Muir · K. Kühn

With 121 Figures



Springer-Verlag Berlin Heidelberg New York 1974

Professor Dr. Reinhard Fricke, Medizinische Poliklinik,
Hannover-Kleefeld

Professor Dr. Fritz Hartmann, Medizinische Poliklinik,
Hannover-Kleefeld

Illustration on the cover designed by Leonhard Fricke by modification of an electron microscopic picture (Wellauer, P., Wyler, T., Buddecke, E.: Electron microscopic and physicochemical studies on bovine nasal cartilage proteoglycan. Hoppe-Seyler's Z. Physiol. Chem. 353, 1043-1052 (1972)) of a proteoglycan molecule by Dr. P. Wellauer, Institute of Zoology, University of Bern, and by incorporating a figure of R. I. G. Morrison, this book, page 151.

ISBN 3-540-06673-X Springer-Verlag Berlin · Heidelberg · New York
ISBN 0-387-06673-X Springer-Verlag New York · Heidelberg · Berlin

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically those of translation, reprinting, re-use of illustrations, broadcasting, reproduction by photocopying machine or similar means, and storage in data banks.

Under § 54 of the German Copyright Law, where copies are made for other than private use, a fee is payable to the publisher, the amount of the fee to be determined by agreement with the publisher.

The use of general descriptive names, trade names, trade marks etc. in this publication, even if the former are not especially identified, is not to be taken as a sign that such names, as understood by the Trade Marks and Merchandise Marks Act, may accordingly be used freely by anyone.

© by Springer-Verlag Berlin · Heidelberg 1974. Library of Congress Catalog Card Number 74-414. Printed in Germany.
Offsetprinting: Julius Beltz, Hemsbach/Bergstr.

Bookbinding: Brühlsche Universitätsdruckerei, Gießen

Preface

Connective tissues are essential for the physical functioning of the animals's body. The condition of the various connective tissues is governed by biochemical factors, anabolism and catabolism, that are controlled by specific enzymes. Any change outside the normal range of metabolism, for instance induced by immunological reactions, may induce a pathological disturbance. The result can be acute or chronic inflammation, or loss of normal function, expressed in loosening, dilatation, breaking, wear, stiffness, shrinking, scars, stenosis, and cirrhosis or any other kind of fibrosis.

A first step toward improving our understanding of the feedback mechanism that maintains the biological status and texture of a given connective tissue is to combine what is known about synthesis and enzymatic degradation of the components of fibers and ground substance. Common pathological phenomena like chronic inflammation of immune reactions can be either the result of the cause of disturbances in the sensitive balance of connective tissue metabolism. Nowadays connective tissues are less and less regarded as bradytrophic tissue but rather as a stimulating and many-sided problem of research.

Before we can understand the pathogenesis of the connective tissue diseases that result in the destructive processes mentioned above, basic research will be necessary. This research will be furthered by a constant exchange of information and the results of observations.

To promote this exchange of information between scientists, symposia on connective tissue research are organized at regular intervals. However, a symposium is not the place to present surveys of all the themes discussed. Therefore, at the symposium on connective tissue research held in Hannover, Germany, some time ago we seized the opportunity to invite specialists to write review articles on the field of research on which they presented papers at the symposium. A synopsis of the present state of knowledge in the field is a necessity for the promotion of research. We are therefore glad to be able to present review articles and reports of new data on selected topics in connective tissue.

Spring 1974

R. Fricke and F. Hartmann

Contributors

- Adam, M., Research Institute for Rheumatic Diseases, Prague/
Czechoslovakia
- Anderson, A.J., Lower Flat, 9 Welbeck Avenue, Hove, Sussex/Great
Britain
- Barnes, M.J., Dunn Nutritional Laboratory, University of Cambridge
and Medical Research Council, Milton Road, Cambridge CB4 1XY/Great
Britain
- Bates, C.J., Dunn Nutritional Laboratory, University of Cambridge,
Milton Road, Cambridge CB4 1XJ/Great Britain
- Baxter, Elsmaree, Department of Medicine, Royal Melbourne Hospital,
University of Melbourne, P.O. Victoria 3050/Australia
- Bazin S., Service de Pathologie Expérimentale, Institut Pasteur,
92380 Garches/France
- Bollwahn, W., Tierärztliche Hochschule Hannover, Klinik für kleine
Klauentiere, Bischofsholer Damm 15, 3000 Hannover/Federal Republic
of Germany
- Bornstein, P., Departments of Biochemistry and Medicine, University
of Washington, School of Medicine, Seattle, WA 98105/USA
- Buddecke, E., Physiologisch-Chemisches Institut der Universität,
Waldeyerstr. 15, 4400 Münster/Federal Republic of Germany
- Codington, J.F., Massachusetts General Hospital, Laboratory for Carbo-
hydrate Research, Boston, MA 02114/USA
- Dabew, D., Zentrallaboratorium, Städtisches Krankenhaus Holweide,
Neufelderstr. 32, 5000 Köln 80/Federal Republic of Germany
- Delaunay, A., Institut Pasteur, Service de Pathologie Expérimentale,
Annexe de Garches 92 (Hauts-De-Seine)/France
- Deyl, Z., Institute of Physiology, Czechoslovak Academy of Science,
Prague/Czechoslovakia
- Eberhard, A., Klinisch-Chemisches Zentrallaboratorium an der Medizi-
nischen Fakultät der RWTH Aachen, Goethestr. 27/29, 5100 Aachen/
Federal Republic of Germany
- Erl, D., Pathologisches Institut der Universität Hamburg, Martini-
str. 52, 2000 Hamburg 20/Federal Republic of Germany
- Evanson, J.M., Department of Medicine, University Hospital of South
Manchester, Nell Lane, Manchester M20 8LR/Great Britain
- Figura, K.v., Physiologisch-Chemisches Institut der Universität,
Waldeyerstr. 15, 4400 Münster/Federal Republic of Germany
- Fischer-Szafarz, B., Fondation Curie, Institut du Radium, 15, Rue
G.-Clémenceau, Bâtiment 111-91405 Orsay/France
- Fricke, R., Medizinische Klinik, Medizinische Hochschule, Karl-Wiechert-
Allee 9, 3000 Hannover-Kleefeld/Federal Republic of Germany
- Gerlach, U., Medizinische Universitäts-Klinik, Westring 3, 4400 Mün-
ster/Federal Republic of Germany
- Gotte, L., Istituto di Istologia-Embriologia, Università di Padova,
Via Falloppio 16, 35100 Padova/Italy
- Gottschalk, A. †, Max-Planck-Institut für Virusforschung, Spemann-
str. 35, 7400 Tübingen/Federal Republic of Germany
- Grade, J., Pathologisches Institut, Universitäts-Krankenhaus Eppen-
dorf, Martinistr. 52, 2000 Hamburg 20/Federal Republic of Germany

- Grasedyck, K., I. Medizinische Klinik, Universitäts-Krankenhaus Eppendorf, Martinistr. 52, 2000 Hamburg 20/Federal Republic of Germany
- Greiling, H., Klinisch-Chemisches Zentrallaboratorium der Medizinischen Fakultät an der RWTH Aachen, Goethestr. 27/29, 5100 Aachen/Federal Republic of Germany
- Gries, G., Perfallstr. 1, 8000 München 80/Federal Republic of Germany
- Hall, D.A., The Department of Medicine, University of Leeds, The General Infirmary, Leeds LS1 3EX/Great Britain
- Happey, F., Postgraduate School of Studies in Textiles, University of Bradford, Bradford, Yorkshire BD7 1DP/Great Britain
- Hardingham, T.E., The Mathilda and Terence Kennedy Institute of Rheumatology, Hammersmith, London W6 7DW/Great Britain
- Hartmann, F., Medizinische Klinik, Medizinische Hochschule, Karl-Wiechert-Allee 9, 3000 Hannover-Kleefeld/Federal Republic of Germany
- Hauss, W.H., Medizinische Universitätsklinik und Poliklinik, Westring 3, 4400 Münster/Federal Republic of Germany
- Hörmann, H., Max-Planck-Institut für Biochemie, 8033 Martinsried bei München/Federal Republic of Germany
- Jeanloz, R.W., Laboratory for Carbohydrate Research, Massachusetts General Hospital, Boston MA 02114/USA
- Kennedy, J.F., Department of Chemistry, University of Birmingham, Edgbaston, Birmingham B15 2TT/Great Britain
- Kisters, R., Klinisch-Chemisches Zentrallaboratorium der Medizinischen Fakultät an der RWTH Aachen, Goethestr. 27/29, 5100 Aachen/Federal Republic of Germany
- Kivirikko, K.I., Department of Medical Chemistry, University of Oulu, Kajaanintie 52 A, 90220 Oulu 22/Finland
- Kleine, T.O., Universitäts-Nervenklinik, Ortenbergstr. 48, 3550 Marburg/Federal Republic of Germany
- Kölln, H., Pathologisches Institut, Universitäts-Krankenhaus Eppendorf, Martinistr. 52, 2000 Hamburg 20/Federal Republic of Germany
- Kresse, H., Physiologisch-Chemisches Institut der Universität, Waldeyerstr. 15, 4400 Münster/Federal Republic of Germany
- Leibovich, S.J., School of Medicine, University of Washington, Seattle WA/USA
- Levene, C.I., Dunn Nutritional Laboratory, University of Cambridge, Milton Road, Cambridge CB4 1XJ/Great Britain
- Lindner, J., Pathologisches Institut, Universitäts-Krankenhaus Eppendorf, Martinistr. 52, 2000 Hamburg 20/Federal Republic of Germany
- Lindsley, H., Department of Medical Zoology, Walter Reed Army Institute of Research, Walter Reed Army Medical Center, Washington, DC 20012/USA
- Lindy, S., Department of Medical Chemistry, University of Helsinki, Siltavuorenpenger 10, 00170 Helsinki 17/Finland
- Mannik, M., Department of Medicine, University of Washington, Seattle, WA 98195/USA
- Merkel, Mathilde, Institut für Pharmakologie der Ruhr-Universität, Im Lottental, 4630 Bochum/Federal Republic of Germany
- Messow, C., Pathologisches Institut, Tierärztliche Hochschule Hannover, Bischofsholer Damm 15, 3000 Hannover/Federal Republic of Germany
- Moczar, E., Laboratoire de Biochimie du Tissu Conjonctif, Université de Paris-Val de Marne, Faculté de Médecine, 6 Rue du Général Sarrail, 94-Créteil/France
- Moczar, M., Laboratoire de Biochimie du Tissu Conjonctif, Université de Paris-Val de Marne, Faculté de Médecine, 6 Rue du Général Sarrail, 94-Créteil/France
- Morrison, R.I.G., Strangeways Research Laboratory, Wort's Causeway, Cambridge CB1 4RN/Great Britain
- Muir, Helen, The Mathilda and Terence Kennedy Institute of Rheumatology, Hammersmith, London W6 7DW/Great Britain

- Naylor, A., Bradford Royal Infirmary, Bradford, Yorkshire BD7 1DP/
Great Britain
- Németh-Csóka, M., Zentral Laboratorium Fővárosi, Tétényi-uti-Kórház,
tétényi-u 14 - 16, Budapest XI/Hungary
- Osborn, J.M., Postgraduate School of Studies in Textiles, University
of Bradford, Bradford, Yorkshire BD7 1DP/Great Britain
- Palframan, J., Postgraduate School of Studies in Textiles, University
of Bradford, Bradford, Yorkshire BD7 1DP/Great Britain
- Pearson, C.H., Department of Oral Biology, Faculty of Dentistry,
University of Alberta, Edmonton, Alberta/Canada
- Prinz, G., Pathologisches Institut, Universitäts-Krankenhaus Eppen-
dorf, Martinistr. 52, 2000 Hamburg 20/Federal Republic of Germany
- Prokop, O., Institut für Gerichtliche Medizin der Humboldt-Universität,
Hannoversche Str. 6, X104 Berlin/German Democratic Republic
- Reed, F.B., Department of Biochemistry, The Royal Dental Hospital,
University of London, School of Dental Surgery, 32 Leicester Square,
London WC2H 7LJ/Great Britain
- Render, R.M., Postgraduate School of Studies in Textiles, University
of Bradford, Bradford, Yorkshire BD7 1DP/Great Britain
- Robert, B. †, Laboratoire de Biochimie du Conjonctif, Faculté de
Médecine, Université de Paris-Val de Marne, 6 Rue de Gal-Sarrail,
94 Crèteil/France
- Robert, L., Laboratoire de Biochimie du Conjonctif, Faculté de Médecine,
Université de Paris-Val de Marne, 6 Rue du Gal-Sarrail, 94 Crèteil/
France
- Rodén, L., University of Alabama in Birmingham, University Station,
OBSB, Birmingham, AL 35294/USA
- Rosmus, J. †, Research Institute of Food Industry, Czech Agricultural
Academy, Prague/Czechoslovakia
- Rothe, A., Abteilung für Experimentelle Innere Medizin der Medizini-
schen Universitäts-Klinik, Kerpenerstr. 15, 5000 Köln 41/Federal
Republic of Germany
- Schauer, R., Institut für Physiologische Chemie, Ruhr-Universität
Bochum, Buscheystraße, 4630 Bochum-Querenburg/Federal Republic of
Germany
- Schoop, H.J., Zentrallabor der Krankenanstalten, Bogenstraße,
2850 Bremerhaven/Federal Republic of Germany
- Schwartz, Nancy B., Department of Pediatrics, University of Chicago,
5825 Maryland Avenue, Chicago, IL 60637/USA
- Segeth, G., Medizinische Klinik der Universität, Westring, 4400 Münster/
Federal Republic of Germany
- Stockhorst, Eva H., Sixth Street 554, Ann Arbor, MI 48103/USA
- Struck, H., II. Chirurgischer Lehrstuhl, Biochemische Abteilung an
der Chirurgischen Universitäts-Klinik, Ostmerheimerstr. 200,
5000 Köln-Merheim/Federal Republic of Germany
- Stuhlsatz, H.W., Klinisch-Chemisches Zentrallaboratorium der Medizi-
nischen Fakultät an der RWTH Aachen, Goethestr. 27/29, 5100 Aachen/
Federal Republic of Germany
- Szafarz, D., Fondation Curie, Institut du Radium, 15 Rue G.-Clemenceau,
Bâtiment, 111-91405 Orsay/France
- Turner, R.L., Postgraduate School of Studies in Medical and Surgical
Sciences, University of Bradford, Bradford, Yorkshire BD7 1DP/
Great Britain
- Turto, H., Department of Medical Chemistry, University of Helsinki,
Siltavuorenpenger 10, 00170 Helsinki/Finnland
- Uhlenbruck, G., Abteilung für Experimentelle Innere Medizin der Medi-
zinischen Universitäts-Klinik, Kerpenerstr. 15, 5000 Köln 41/
Federal Republic of Germany
- Uitto, J., Department of Biochemistry, Rutgers Medical School, College
of Medicine and Dentistry of New Jersey, Piscataway, NJ 08854/USA
- Vainio, K., Department of Surgery, Rheumatism Foundation Hospital,
Heinola/Finnland

Weiss, Jacqueline B., Department of Medical Biochemistry, University of Manchester, York Place, Manchester M13 9PT/Great Britain

Wollmer, A., Abteilung Physiologische Chemie der Medizinischen Fakultät an der RWTH Aachen, Melatener Str. 211/213, 5100 Aachen/Federal Republic of Germany

Wulff, U., Kinderklinik der Medizinischen Akademie, Ratzeburger Allee 160, 2400 Lübeck/Federal Republic of Germany

Table of Contents

I. Structure

E. Buddecke, H. Kresse, G. Segeth, and K. v. Figura Metabolic Heterogeneity of Acidic Glycosaminoglycans in Bovine Arterial Tissue.....	3
T.E. Hardingham, E. Baxter, and H. Muir Chemical and Molecular Heterogeneity of Cartilage Proteoglycans	7
T.O. Kleine Proteochondroitin Sulfates, Chemical and Metabolic Heterogeneity, Immunology and Biosynthesis.....	16
H.W. Stuhlsatz, R. Kisters, A. Wollmer, and H. Greiling Isolation of Corneal Keratan-Chondroproteoglycans and Their Com- position.....	37
M. Moczar and E. Moczar Glycopeptide Pattern of the Polymeric Macromolecular Complex of the Stroma Layers and of the Endothelial Basement Membrane (Descemet's Membrane) of the Embryonic and Postnatal Calf Cornea	50
J.F. Codington and R.W. Jeanloz The Chemistry of Glycoproteins at the Surfaces of Tumor Cells..	55
M. Németh-Csóka The Relationship between Chemical Structure and Fibrogenic Character of Glycosaminoglycans.....	61
H. Hörmann Periodate Sensitivity of Cross-Links of Mature Collagen.....	64
F. Happey, A. Naylor, J. Palftaman, C.H. Pearson, R.M. Render, and R.L. Turner Variations in the Diameter of Collagen Fibrils, Bound Hexose and Associated Glycoproteins in the Intervertebral Disc.....	67

VI

II. Synthesis

L. Rodén and N.B. Schwartz	
Biosynthesis of Connective Tissue Proteoglycans.....	73
C.J. Bates and C.I. Levene	
Synthesis of Sulfated Glycosaminoglycans by Cultured Cells, with Particular Reference to the Mouse Fibroblast Line 3T6.....	85
R. Schauer	
The Subcellular Site of the Biosynthesis of N-Glycolyl- and N-Acetyl-O-Acetylneuraminic Acid Components of Porcine, Bovine and Equine Submaxillary Gland Glycoproteins.....	89
H.J. Schoop, E.H. Stockhorst, and M. Merkel	
N-Acetyl-Hydroxylation in the Biosynthesis of Glycoproteins....	102
K.I. Kivirikko	
Biosynthesis of Collagen.....	107
K. Grasedyck, U. Wulff, D. Erl, and J. Lindner	
Studies on Collagen Synthesis Applying Labelled Proline.....	122

III. Degradation

E. Buddecke and H. Kresse	
Mammalian Enzymes Degrading Glycosaminoglycans.....	131
J.F. Kennedy	
The Production and Use of Water-Insoluble Derivatives of En- zymes.....	146
R.I.G. Morrison	
The Enzymatic Degradation of Proteoglycans.....	150
C.H. Pearson, F. Happey, A. Naylor, J.M. Osborn, and R.L. Turner	
Lysosomal Enzymes and Proteoglycan Degradation in the Human Intervertebral Disc.....	158
H. Greiling, R. Kisters, H. Stuhlſatz, and A. Eberhard	
Studies on Lysosomal Proteoglycan Degradation.....	166
J.M. Evanson	
Action of Mammalian Collagenases.....	173
G. Gries	
The Mechanism of Collagen Decomposition in Mammals.....	177
J. Lindner, K. Grasedyck, G. Prinz, J. Grade, and H. Kölln	
Studies on Collagen Peptidases.....	180

A.J. Anderson	
Degradation of Insoluble Collagen and other Proteins by Rat Leucocytes.....	191
D. Dabew	
Enzymatic Degradation of the "Collagen-Like-Protein" in Blood Serum.....	193
L. Gotte	
Sequential Degradation and Phagocytosis of Heterologous Elastin	196
IV. Immunobiology	
G. Uhlenbruck, A. Rothe, and O. Prokop	
Immunobiology of Glycoproteins.....	201
A. Gottschalk	
Immunological Properties of Ovine Submaxillary Glycoproteins...	221
P. Bornstein, H. Lindsley, and M. Mannik	
The Distribution of Antigenic Determinants in Collagen as Determined by a Radioimmunoassay.....	225
H. Struck	
Antigenicity of Soluble Collagen Fractions of Different Species	232
V. Pathophysiology	
F. Hartmann	
Questions of Medicine to Molecular Biology of Connective Tissues.....	237
L. Robert and B. Robert	
Structural Glycoproteins of Connective Tissue: Their Role in Morphogenesis and Immunopathology.....	240
R. Fricke, W. Bollwahn, and C. Messow	
Experimental Chronic Arthritis in Pigs after a Single Intra-articular Injection of Heterologous Proteoglycans.....	257
B. Fiszer-Szafarz and D. Szafarz	
Hyaluronidase Activity in Lysosomes and Extracellular Fluid of Tumor Cells.....	261
U. Gerlach and W.H. Hauss	
Clinical and Experimental Investigations on the Metabolism of Collagen, Ground Substance, and Mesenchymal Cells.....	262

VIII

A. Delaunay and S. Bazin	
Collagen Maturation in Granulation Tissue.....	273
J.B. Weiss and S.J. Leibovich	
A Negative Role for Collagenase in Observed Differences between Human Normal and Rheumatoid Polymeric Collagens.....	278
M. Adam, Z. Deyl, and J. Rosmus	
The Possible Role of Cross-Linking of Collagen in Rheumatoid Arthritis.....	281
J. Uitto, S. Lindy, H. Turto, and K. Vainio	
Studies on Collagen Biosynthesis in Rheumatoid Synovial Tissue	288
F.B. Reed and D.A. Hall	
Changes in Skin Collagen in Osteoporosis.....	290
M.J. Barnes	
Collagen and Elastin Metabolism in Scurvy: Studies on the Nature of the Collagen Lesion.....	300
Subject Index.....	305

I. Structure

1. 1. 1. 1. 1.

Metabolic Heterogeneity of Acidic Glycosaminoglycans in Bovine Arterial Tissue

E. Buddecke, H. Kresse, G. Segeth, and K. v. Figura

Mammalian arterial tissue is known to contain hyaluronate (HA), chondroitin 4(6)-sulfate (CS) dermatan sulfate (DS) heparan sulfate (HS) (Kaplan and Meyer, 1960; Antonopoulos et al., 1965; Dunstone, 1967; Nakamura et al., 1968; Buddecke and Kresse, 1960; Kresse and Buddecke, 1970) and their protein complexes respectively (Jacobs and Muir, 1963; Buddecke et al., 1963; Fransson and Havsmark, 1970) in a total amount of 1 - 2% of the dry weight of tissue.

Studies on age dependent and pathological changes in the distribution pattern of acidic glycosaminoglycans (GAG) in human arterial tissue gave inconsistent results (Kaplan and Meyer, 1960; Manley et al., 1969; Dalferes et al., 1971). However, the distribution pattern will be of particular interest, as an increased incorporation of ^{35}S -sulfate into the sulfated GAG has been described to be one of the first metabolic alterations in spontaneous and experimental arteriosclerosis (Hauss et al., 1968; Sanwald et al., 1971).

Since difficulties in distinguishing between normal human arterial tissue and barely detectable arteriosclerotic lesions prevented a systematic investigation of the metabolic feature of GAG, we studied the metabolism of the different types of GAG and proteoglycans in bovine aorta as a model for the normal.

1. Metabolic Heterogeneity of Aortic GAG

When segments of calf aorta were incubated under appropriate conditions in vitro in the presence of $\text{U-}^{14}\text{C}$ -glucose or ^{35}S -sulfate over a period of 12 h, significant differences in the rate of ^{35}S - and ^{14}C -incorporation into the GAG are observed. The relative rates of biosynthesis of the individual GAG were calculated on the basis of relative proportions of HA : CS : DS : HS and their specific radioactivities. The highest rate of synthesis was found for HA and

Table. Relative rates of biosynthesis of arterial tissue GAG following in vitro incubation in the presence of $\text{U-}^{14}\text{C}$ -glucose and ^{35}S -sulfate resp. (Kresse and Buddecke, 1970)

Type of GAG	mg/100 mg total GAG	Relative rate of synthesis following incorporation of		Relative turnover rate following incorporation of	
		^{14}C	^{35}S	^{14}C	^{35}S
HA	30	58		1.9	
CS	49	21	41	0.4	0.3
DS	9.5	10	29	1.0	1.0
HS	11.5	11	30	0.9	0.9

decreasing amounts for the sulfated GAG in the order CS, HS, and DS. The Table summarizes the results (Kresse and Buddecke, 1970). HA and HS were found to be synthesized from two differently labelled glucosamine precursor-pools (Figura et al, 1973).

2. Age Dependent Changes in Specific Radioactivity of GAG

A study of age dependent changes of GAG metabolism revealed the following results: in bovine aortae the total GAG content increases from 1.19% to 1.70% over the first 3 years and decreases subsequently to 1.2% up to the 10th year, all values referring to the dry weight of tissue. No major changes in the distribution pattern of GAG are observed after this period. The specific radioactivity for all sulfated GAG decreases exponentially during the first 2 years but remains nearly constant for the following years, identical results being obtained for ¹⁴C- and ³⁵S-GAG. Surprisingly, however, it was found that the specific ¹⁴C-radioactivity of the HA increases twofold during the first 3 years reaching a plateau for the following years (Buddecke and Segeth, in press). The figure shows the age profile of ¹⁴C-labelled HA and DS, the latter being representative for all sulfated GAG.




Fig. Age dependent changes in specific radioactivity of ¹⁴C-hyaluronic acid(▽) and ¹⁴C-dermatansulfate (O) after in vitro incubation in the presence of U-¹⁴C-glucose


The reason for the higher rate of biosynthesis of HA and its further age dependent increase is still obscure, but a selective regulation of the biosynthesis of the individual GAG must be concluded from these results. An independent control of biosynthesis of the different GAG in arterial tissue is also reflected by the fact that a selective increase in the rate of synthesis of DS and HS of arterial tissue during genetic and experimental hypertonus in the rat has been observed (Kresse et al., 1971b).

3. Metabolic Heterogeneity of Bovine Aorta CS-DS-Proteoglycans

The observed differences between CS and DS in the rates of biosynthesis are especially striking with regard to the well established finding that in the aorta a large proportion of DS contains CS as well as DS disaccharide units in a copolymeric structure (Fransson and Roden, 1967; Fransson, 1968; Fransson and Havsmark, 1970).

Studying the chemical and metabolic features of bovine arterial tissue proteoglycans, we isolated and purified a CS-DS-proteoglycan containing CS as well as DS, both attached to the same protein core. This CS-DS-proteoglycan behaved as a single compound by several criteria of homogeneity. On incubation in vitro in the presence of ^{35}S -sulfate the CS-DS-proteoglycan was obtained in a ^{35}S -sulfate labelled form. From light scattering measurements a molecular weight of 2.0×10^6 and a random coil shape were calculated, chemical analysis indicated a protein content of 20.5% and a polysaccharide content of 79.5%, the latter consisting of 73.0% CS and 27.0% DS.

An exhaustive degradation of the ^{35}S -SD-DS-proteoglycan by testes hyaluronidase and pronase produces ^{35}S -DS and a series of ^{35}S -saccharide fragments ranging from disaccharides to decasaccharides. The tetrasaccharide fraction was found to contain iduronic acid (IdUA) besides glucuronic acid (GlcUA) indicating a copolymeric structure of CS and DS units. The occurrence of IdUA in the tetrasaccharide fraction (as referred to the total uronic acid content) is conclusive for the presence of tetrasaccharide molecules having the structure



From these data and from the results of alkali degradation the macromolecular structure of the CS-DS-proteoglycan may be summarized as follows: about 80 polysaccharide chains at a molecular weight between 15,000 and 20,000 are attached to a protein core by alkali sensitive O-seryl (threonyl) bonds giving a random coiled shaped macromolecule with a molecular weight of 2.0×10^6 . It appears that (a) the CS and DS polymer chains are attached to a single protein core and (b) that CS and DS chains occur partly as copolymers.

Determinations of the specific radioactivity and the iduronic acid content of the DS fraction and the DS and CS units containing saccharide fragments obtained from testes hyaluronidase degradation showed that the specific radioactivity of the sulfate ester group of the DS units was about three times higher than those of the CS units (Kresse et al., 1971).

Further studies on ^{14}C -GlcN-labelled CS-DS-preparations revealed a varying degree of hybridization between CS and DS. A highly labelled ^{14}C -DS fraction hybridized with 19 % of a likewise highly labelled CS and a lower labelled CS fraction containing 20 % of a lower labelled DS as copolymer constituent were isolated by subfractionation of the aortic tissue GAG. These results are conclusive for the assumption that two topographically different sites exist for the biosynthesis of DS and CS, one delivering a highly labelled DS(CS)-hybrid, the other a lower labelled CS(DS)-hybrid (v. Figura et al., unpublished).