

AMERICAN DRUG INDEX 1960

CHARLES O. WILSON, Ph.D.

*Dean and Professor of Pharmaceutical Chemistry, School of Pharmacy,
Oregon State College*

AND

TONY EVERETT JONES, Ph.D.

*Associate Professor of Pharmaceutical Chemistry, School of Pharmacy,
University of Colorado; Director of
Pharmaceutical Research, Carbisulphoil Company*

Copyright © 1960, by J. B. Lippincott Company

Copyright 1956, 1958, 1959, by J. B. Lippincott Company

This book is fully protected by copyright and, with the exception of brief extracts for review, no part of it may be reproduced in any form without the written permission of the publishers

**Distributed in Great Britain by
Pitman Medical Publishing Co., Limited, London**

Library of Congress Catalog Card Number 55-6286

Printed in the United States of America

PREFACE

The *American Drug Index* has been prepared for the identification and correlation of the many pharmaceuticals available to the medical and allied professions. The need for this index has become acute as the number of drugs and drug products in all their tremendous variety has multiplied.

The organization of the index is fundamentally alphabetical with extensive cross-indexing. Names listed are generic; brand (trade-mark, proprietary or specialty), chemical, U.S.P., N.F., N.N.D. and A.D.R. Synonyms that are in general use also are included. All names used for a pharmaceutical occur in alphabetical order with the pertinent data given under the brand name and the name by which it is made available. Data included are generic name, chemical name, manufacturer, pharmaceutical forms, size, dosage and use. Generic names occur in alphabetical order and here the recognition of the drug is indicated by U.S.P. (United States Pharmacopeia), N.F. (National Formulary), N.N.D. (New and Nonofficial Drugs) and A.D.R. (Accepted Dental Remedies). The information is in accord with the latest editions of these books: U.S.P. XV, N.F. X, N.N.D. 1959, A.D.R. 1959.

Because of the multiplicity of specialty names (brand names) used for the same therapeutic agent or the same combination of therapeutic agents, it was apparent that some correlation could be done. As an example of this, please turn to Aluminum Hydroxide Gel. Here under the generic name are given the brand names. Following are combinations of aluminum hydroxide gel organized in a manner to point out relationships among the many products. Reference then is made to the specialty name or names having the indicated composition. Under the specialty name are given manufacturer, composition, available forms, sizes, dosage and use.

The *American Drug Index* should be most useful to physicians, pharmacists, dentists, nurses, and to sales personnel, students and teachers in the fields incorporating pharmaceuticals. The cross-indexing feature permits the finding of drugs or drug combinations when only one major ingredient is known. For example, a combination of aluminum hydroxide gel and magnesium trisilicate is available. This combination can be found by looking under the name of either of the two ingredients, and in each case the specialty names are given.

CHARLES O. WILSON
TONY E. JONES

A.A.S. COMPOUND. (Tilden) Aspirin 5 gr., nux vomica ext. 1/10 gr., hyoscyamus ext. ¼ gr., gelsemium tr. 3 min., cascara sagrada ext. 1 gr./Tab. Bot. 100s, 1000s.

Use: Analgesic.

ABAC. (Massengill) Aspirin 5 gr., caffeine ¼ gr., Pot. bromide ¼ gr., Sod. bromide ¼ gr./Tab. or Cap. Bot. 100s, 1000s. Tab. Bot. 500s.

Use: (1-2 q. 3 h.) Analgesic sedative.

ABAC FOR CHILDREN. (Massengill) Aspirin 1 gr., caffeine 1/10 gr., Pot. bromide 1/20 gr., Sod. bromide 1/20 gr., aromatics q.s./Tab. Bot. 500s, 1000s, 5000s.

Use: (1 q. 4 h.) Analgesic.

ABACTAL (Massengill) Aspirin 5 gr., caffeine ¼ gr., Pot. bromide ¼ gr., Sod. bromide ¼ gr., phenobarbital ½ gr./Tab. Bot. 100s, 1000s.

Use: (1-2 t.i.d.) Analgesic sedative.

ABASIN. (Winthrop Labs.) Acetylcarbromal, Tab. (4 gr.) Bot. 100s.

Use: (1-2 t.i.d.) Sedative.

ABOCCILLIN 800 M., A.D.R. (Abbott) Procaine penicillin G 600,000 u., buffered Pot. penicillin G 200,000 u. for aq. inj. Vial, 1 cc., Box 25s, 5 cc., 1s, 25s.

Use: I.M., bacterial infections.

ABOCCILLIN-DC, A.D.R. (Abbott) Procaine penicillin G in aq. susp. 600,000 u. 1 cc. Abobject disposable syr. w/needle, 1s, 25s.

Use: I.M., penicillin therapy.

ABBO-VAC. See: A.C.D. Solution. (Abbott).

ABDEC DROPS. (Parke, Davis) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 0.4 mg., B-6 1 mg., Sod. pantothenate 2 mg., nicotinamide 5 mg., Vit. C 50 mg./0.6 cc. Bot. w/dropper 15 cc., 50 cc.

Use: Vitamin deficiencies.

ABDEC DROPS MODIFIED. (Parke, Davis) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 1.2 mg., B-6 1 mg., sod. pantothenate 5 mg., nicotinamide 10 mg., B-12 2 mcg., C 50 mg./0.6 cc. Bot. 15 cc., 50 cc.

Use: Vitamin deficiencies.

ABDEC KAPSEALS. (Parke, Davis) Vit. A 10,000 u., D 1000 u., B-1 5 mg., B-2 3 mg., B-6 1.5 mg., B-12 2 mcg., Sod. pantothenate 5 mg., nicotinamide 25 mg., Vit. C 75 mg., mixed tocopherols 5 mg./Cap. Bot. 50s, 100s, 250s, 1000s.

Use: (1 daily) Multivitamin therapy.

A-B-D-G. (Blue Line) Vit. A 5000 u., D 1000 u., B-1 1 mg., B-2 2 mg./Cap. Bot. 100s.

Use: (1 daily) Dietary supplement.

A-B-D-G VITAMIN. (Horton & Converse) Vit. A 5000 u., D 1000 u., B-1 333 u., riboflavin 2000 mcg./Tab. Bot. 100s, 250s, 500s, 1000s.

Use: (1 daily) Multivitamin therapy.

A-B-D-G VITAMINS. (Upjohn) Vit. A 2500 u., D 250 u., B-1 1.5 mg. B-2 1 mg., nicotinamide 10 mg./Cap. Bot. 50s, 100s, 250s, 1000s.

Use: (1 daily) Multivitamin therapy.

ABDOL W/Minerals. (Parke, Davis) Vit. C 50 mg., B-2 2.5 mg., B-12 1 mcg., B-6 0.5 mg., B-1 2.5 mg., nicotinamide 20 mg., Vit. A 5000 u., D 1000 u., Cal. pantothenate 2.5 mg., folic acid 0.1 mg., iodine 0.15 mg., Mang. 1 mg., cobalt 0.1 mg., Pot. 5.0 mg., molybdenum 0.2 mg., iron 15 mg., copper 1 mg., zinc 0.5 mg., Mag. 1 mg., Cal. 44.0 mg., Phos. 34 mg./Cap. Bot. 100s, 250s.

Use: (1 daily) Vitamin-mineral deficiency states.

ABDOL W/MINERALS for Children. (Parke, Davis) Vit. C 40 mg., B-2 2 mg., B-12 1.5 mcg., B-6 0.5 mg., B-1 1 mg., nicotinamide 12 mg., A 3000 u., D 1000 u., cal. pantothenate 2.5 mg., folic acid 0.1 mg., w/iron, copper, zinc, magnesium, calcium, phosphorus, iodine, manganese, cobalt, and pot./Cap. Bot. 100s.

Use: Vitamin-mineral supplement.

ABDOL W/Vit. C Cap. (Parke, Davis) Vit. A 5000 u., D 1000 u., B-1 2.5 mg., B-2 2.5 mg., B-6 0.5 mg., B-12 1 mcg., nicotinamide 20 mg., Vit. C 50 mg., pantothenic acid (as Cal. salt) 5 mg./Cap. Bot. 25s, 100s, 250s, 500s.

Use: (1 daily) Vitamin deficiency.

A-B-M-C OINTMENT. See: Lembrose, Oint. (Wyeth).

ABMINTHIC. (Pfizer) Dithiazanine iodide. 3 3'-Dithiylthiadiazocyanine iodide. 200 mg./Tab. Bot. 100s.

Use: Anthelmintic.

ABRODIL. See Methiodal Sodium. (Winthrop Labs.).

A-B-S & C. (Horton & Converse) Aloin 1/5 gr., strychnine 1/120 gr., belladonna lvs ext. ½ gr., cascara sagrada ½ gr./Tab. Bot. 100s, 1000s.

Use: (1 h.s.) Laxative.

ABSORBABLE CELLULOSE COTTON OR GAUZE, N.N.D. 1959. See: Oxidized Cellulose. (Various Mfr.).

ABSORBABLE DUSTING POWDER, U.S.P., N.N.D. 1959. See: Bio-Sorb, Pow. (Ethicon)

ABSORBABLE GELATIN FILM, N.N.D. 1959. Sterile nonantigenic, absorbable, water-insoluble, gelatin film. See: Gelfilm. (Upjohn)

ABSORBABLE GELATIN SPONGE, U.S.P., N.N.D. 1959, A.D.R. Gelatin sponge. See Gelfoam, Preps. (Upjohn).

ABSORBABLE SURGICAL DRESSINGS. See: Hemo-Pak. (Johnson & Johnson).

ABSORBABLE SURGICAL SUTURE, U.S.P.,

ABSORBENT BASE

2

A.D.R. (Various Mfr.) Surgical Gut, Surgical Catgut, Catgut suture.

ABSORBENT BASE. (Upsler-Smith) Oint. Base: Cetyl alcohol, stearyl alcohol, lauryl sulfate, methyl p-hydroxybenzoate & hydrocarbons. Jar 1.5, 6, 25 lb.
Use: Ointment base.

ABSORBENT OINTMENT BASE.

See: Plastibase Hydrophilic (Squibb).
Polysorb (Fougera).

ABTEN. (Paul Maney) Rauwolfia serpentina alkaloids ext. Tab. (1 mg. & 2 mg.) Bot. 100s, 1000s.

Use: (2 daily) Hypotensive and tranquillizing agent.

ABUPHENINE CITRATE. (Chemo Puro) Diethylaminoethyl phenylethylacetate citrate. Pow. for mfg.

Use: Antispasmodic.

ABUPHENINE HYDROCHLORIDE. (Chemo Puro) Diethylaminoethyl phenylethylacetate hydrochloride. Pow. for Mfg.

Use: Antispasmodic.

A-C TROCHES. (Abbott) Anesthesin $\frac{1}{4}$ gr., calcidin & licorice. Bot. 24s, 100s, 1000s.

Use: Throat & mouth infections.

ACACIA GUM, U.S.P., A.D.R. (Penick) Acacia senegal, gum arabic. Mucilage U.S.P.; pow., $\frac{1}{4}$ lb., 1 lb., tears, $\frac{1}{4}$ -1 lb., syr. U.S.P.; whole, $\frac{1}{4}$ -1 lb.

Use: Demulcent, emulsifier.

W/Mag. oxide & veg. mucinoid. See: Acarosil, Pow. (Sherman).

W/Veg. mucinoid. See: Acarojel, Pow. (Sherman).

A.C.A. COMPOUND. (Rowell) Aspirin $3\frac{1}{2}$ gr., phenacetin $2\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./Cap. or Tab. Bot. 100s, 1000s. Also with codeine phos. $\frac{1}{8}$, $\frac{1}{4}$, $\frac{1}{2}$ and 1 gr. See: A.P.C.

Use: (1-2 q. 3 h.) Analgesic & antipyretic.

ACAP. (Barre) Vit. A 50,000 u./Cap. Bot. 100s.

Use: Vitamin A deficiencies.

ACAPNOID. (Chemico) Belladonna pow. $\frac{1}{10}$ gr., phenobarbital $\frac{1}{3}$ gr., stramonium pow. 1 gr., ephedrine sulfate $\frac{1}{4}$ gr./Tab. Bot. 100s, 500s, 1000s.

Use: Sedative & antispasmodic.

ACARALUM. (Sherman) Carbohydrate uronides (veg. mucinoid) 50%, acacia 20%, aluminum hydroxide gel 25%, Mag. oxide 5%. Pow. Can 8 oz.

Use: (2 dr. p.r.n.) Gastric antacid.

ACAROJEL. (Sherman) Carbohydrate uronides (veg. mucinoid) 50%, acacia 50%. Pow. Can 8 oz.

Use: Laxative.

ACAROSIL. (Sherman) Carbohydrate uro-

nides (veg. mucinoid) 50%, acacia 25%, Mag. oxide. 25%. Pow. Can 8 oz.

Use: (1 dr. p.r.n.) Gastric antacid.

ACCESSORONE. (Upjohn) Vit. A 22,600 u., D 267 u., red bone marrow ext. 0.62 cc., iron & ammonium citrate 0.259 Gm., malt ext., yeast ext., citrum conc. q.s./fl. oz. Bot. 10 fl. oz., 24 fl. oz.

Use: Dietary supplement.

ACCO. (Assoc. Concentrates) Protein, 32%, carbohydrate 54%, ash 5.75%, fat 0.75%, crude fiber 1.61%, Vit. B-1 2 mg., B-2 3 mg., nicotinic acid 15 mg. Jar 15 oz.

Use: Nutritional supplement.

A-C-D SOLUTION, U.S.P. Sod. citrate, citric acid & dextrose in sterile pyrogen-free solution.

Abbott—(Abbo-Vac) 150, 250, 500 cc. (non-Vac) 250 & 500 ml. (Pliapab) w/Blood collection set 500 cc.

Baxter—600 cc. bot. with 70, 120 or 300 cc. of sol.; 1000 cc. with 500 cc. of sol.

Cutter—500 cc. bot. with 75 or 120 cc. of sol.; 650 cc. bot. with 80 or 130 cc. of sol.

Use: Anticoagulant for preparation of plasma or whole blood.

ACE-CAF-EDINE, A.D.R. 1955. (Paul Maney) Aspirin $3\frac{1}{2}$ gr., acetophenetidin $2\frac{1}{2}$ gr., caffeine $\frac{1}{2}$ gr./cap. or tab. Cap., Bot. 100s, 1000s; Tab., No. 1, Bot. 100s, 1000s; also w/phenobarbital, Bot. 1000s. See: A.P.C.

Use: (1-2 q. 3 h.) Analgesic & antipyretic.

ACECOLINE. (Anglo-French) Acetylcholine chloride 20, 50, 100 or 200 mg. w/saligenin 40 mg./cc. in propylene glycol. Amp., 1 cc. Box 6s.

Use: S.C., I.M.; parasympathetic agent.

ACE-JEL. (Ortho) Acid vaginal jelly buffered to pH of 4.0, containing ricinoleic acid, boric acid, oxyquinoline sulfate & propyl ester parahydroxybenzoic acid in water-dispersible base. Tube with or without applicator.

ACENOCOUMARIN. See: Sintrom, Tab. (Geigy)

ACENOCOUMAROL, N.N.D. 1959. See: Sintrom, Tab. (Geigy)

ACEPROMAZINE. 10(3-dimethylaminopropyl)-3-acetylphenothiazine maleate. See: Atravet (Ayerst). Plegical (Ayerst)

Use: Tranquillizer.

ACETABAR. (Columbus) Phenobarbital $\frac{1}{4}$ gr., acetophenetidin $2\frac{1}{2}$ gr., acetylsalicylic acid $3\frac{1}{2}$ grs., caffeine $\frac{1}{4}$ gr./Tab. or wafer. Bots. 100s, 1000s.

Use: (1-2 q. 4 h.) Analgesic-sedative.

ACETALAX. (G. F. Harvey) Diacetylhydroxyphenolisatin. Cap. 5 mg., Bot. 100s, 1000s.

Use: (1-2 h.s.) Laxative.

- p-ACETAMIDO BENZOIC ACID SALT** of 2-dimethylaminoethanol. See: Deaner, Tab. (Riker)
- 3-ACETAMIDO-2,4,6-TRIIODOBENZOIC ACID.** See: Acetizoic acid, U.S.P. (Mallinckrodt).
- 2-ACETAMIDO-5-NITROTHIAZOLE.** See: Tritheon (Ortho)
- ACETAMINOPHEN.** See: N-Acetyl-p-aminophenol.
- ACETAMINOPHENOL.** See: N-acetyl-p-aminophenol.
- ACETANILID,** N.F., A.D.R.; Tablets, N.F. (Various Mfr.). (Acetylaminobenzene, acetylaniline, antifebrin) N-Phenylacetamide Cry. Bot. 1 oz., $\frac{1}{4}$ lb., 1 lb.
Use: Antipyretic, analgesic.
Powder Compound, N.F. (Various Mfr.).
W/Acetophenetidin, salol & caffeine. See: Ferbrisol, liq. (Tilden).
- W/Anesthasin, ephedrine alkaloid, bismuth subnitrate, petrolatum, menthol, eucalyptol. See: Nasalene, No. 1, Liq. (Abbott).
- W/Quinine sulfate, podophyllin, aloin, acornite root, capsicum, atropine sulfate, and camphor. See: C.C.G., Tab. (Noyes).
- ACETARSOL.** See: Acetarsons, Tabs. (Abbott)
- ACETARSONE,** N.F., N.N.R. 1950; Tablets, N.F. (Abbott) 3-Acetamido-4-hydroxyphenylarsonic acid (Acetarzol, Acetphenarsine, Amarsan, Dynarsan, Ehrlich 594, Limarsol, Orarsan, Osarsal, Paroxyl, Stavarsol.) Tab. (0.25 Gm.). Bot. 100s, 500s.
Use: Amebiasis & trichomoniasis.
- W/Hydrolyzed carbohydrates. See: Devegan, Tabs. (Winthrop Labs.).
- W/Polymyxin B, Neomycin. See: Neosporin vag. Supp. (Burroughs Wellcome).
- ACETARSONE SALT OF ARECOLINE.** See: Drocabil, N.F. (Winthrop Labs.)
- ACETAZOLAMIDE.** N.N.D. 1959. See: Diamox, Tabs. (Lederle).
- ACET-DIA-MER-SULFONAMIDE,** N.F., N.N.R. 1955. Sulfacetamide, Sulfadiazine & Sulfamerazine.
See: Buffonamide, Susp., N.N.R. (Tutag)
Cetazine, Tab., N.N.R. (Bowman Bros.).
Dorsulfas, Susp. & Tabs., N.N.R. (Smith-Dorsey).
Incorposul, Susp. & Tabs., N.N.R. (Blue Line).
Pansulfa, Susp., Tabs. (Merrell).
Sulfadiazine, sulfamerazine & sulfacetamide, Susp., N.F.; Tab. N.F.
Tilamide, Liq. (Tilden).
Tricombisul, Susp. & Tabs., N.N.R. (Schering).
Trixyf, Susp. & Tabs. (Central).
- W/Gantresin & Pot. penicillin G. See: Trionacil, Tab. (Flint, Eaton)
- W/Hyoscyamus Ext. See: Suromate, Tab. (Patch)
- W/Sulfamethazine.
See: Delomets, Tab. (Durst).
Deltamide, Tab. or Susp. (Armour)
Katrasul, Tabs., Susp. (Chicago).
Quad-Sulfa, Tab. (Zimmer).
- W/Sulfamethazine, Pot. penicillin G. See: Deltamide w/Penicillin, Tab. or Supp. (Armour)
- W/Urea, neocalamine, talc, glycerin, camphor, menthol. See: Neofonamide, Liq. (McNeil)
- ACETEST.** (Ames). Sod. nitroprusside, disodium phosphate, aminoacetic acid. Tab. Bot. 100s, 250s.
Use: Test for acetone or diacetic acid in urine.
- ACETICYL.** See: Acetylsalicylic Acid.
- ACETIDINE,** A.D.R. (Merck Sharp & Dohme) Aspirin 6 pt., phenacetin 4 pt., caffeine 1 pt. Caps., pink, 5 gr., Bot. 100s, 1000s; white, 5 gr., Bot. 12s, 36s, 100s, 1000s; Tab., pink or white, 5 gr., Box, Bot. 100s, 1000s. See: A.P.C.
Use: (1-2 q. 3 h.) Analgesic, antipyretic.
- W/Codeine phosphate $\frac{1}{4}$ gr., A.D.R. Tab. 100s, 500s.
Use: (1-2 q. 3 h.) Analgesic, antipyretic.
- W/Codeine phosphate $\frac{1}{2}$ gr., A.D.R. Tab. 100s, 500s.
Use: (1-2 q. 3 h.) Analgesic, antipyretic.
- ACETILUM ACIDULATUM.** See: Acetylsalicylic Acid. (Various Mfr.).
- ACETOBRO, PEDIATAB No. 1.** (Columbus Pharm.) Bromisovalum $\frac{1}{2}$ gr., ethyl salicylate carbonate 1 gr., acetophenetidin $\frac{3}{4}$ gr., caffeine $\frac{1}{16}$ gr./tab. Bot. 100s, 1000s.
Use: (1-2 q. 3 hr.) Sedative.
- ACETO-CORT.** (Mallard) Hydrocortisone 1.5 mg., aspirin 5 gr., aluminum hydroxide g:1 100 mg./Tab. Bot. 100s, 1000s.
Use: Treatment of mild to moderate rheumatoid arthritis.
- ACETOL.** See: Acetylsalicylic Acid. (Various Mfr.).
- ACETOLACTIN.** (G. F. Harvey) Acetphenol-satin 5 mg./Cap. Bot. 100s, 1000s
Use: Laxative or purgative
- ACETOMEROCTOL,** N.N.D. 1959. See: Merbak, Prep. (Schieffelin)
- ACETOMIN.** (Central) Salicylamide 60 mg., acetophenetidin 30 mg., atropine sulfate 0.03 mg., camphor monobromated 8 mg., phenolphthalein 6 mg./Palatab. Bot. 100s, 1000s
Use: Analgesic, antipyretic
- ACETONE or DIACETIC ACID TEST.** See: Acetest, Tab. (Ames).
- ACETONYL.** (Upjohn) Granular efferv. salt.

- Aspirin as sod. salt $7\frac{1}{2}$ gr. w/Pot. & Sod. citrates & tartrates, calcium lactate, Sod. phosphate & magnesium sulfate/2 oz. Bot. 4 oz.
- Use: (1 tps.) Analgesic & alkaliizer.
- 3-(α -ACETONYLBENZYL)-4-HYDROXY-COUMARIN.** See: Warfarin sod. N.N.R.
- 3-(α -ACETONYL-4-NITRO-BENZYL)-4-HYDROXYCOUMARIN.** See: Sintrom, Tab. (Geigy)
- ACETOPHEN.** (Smith-Dorsey) Phenobarbital $\frac{1}{4}$ gr., aspirin 3 gr., acetophenetidin 2 gr., hyoscyamus ext. $\frac{1}{8}$ gr./Cap. Bot. 100s, 1000s.
- Use: (1 i.d.) Sedative—analgesic.
- ACETOPHENETIDIN,** U.S.P., N.N.R. 1941, A.D.R.; Tablets, U.S.P. (Various Mfr.) (Phenacetin, p. Ethoxyacetanilide.) Cry. or Pow. 1 oz., $\frac{1}{4}$ lb., 1 lb., 5 lb.
- Use: Antipyretic, analgesic.
- W/Aspirin, caffeine. See: A.P.C.
- Acetophenetidin Combinations on pp. 5 to 9.**
- ACETOPHENETIDIN PHENYL SALICYLATE TABLETS,** N.F.
- ACETOSAL.** See: Acetylsalicylic Acid. (Various Mfr.).
- ACETOSALIC ACID.** See: Acetylsalicylic Acid. (Various Mfr.).
- ACETOSALIN.** See: Acetylsalicylic Acid. (Various Mfr.).
- ACETOSULFONE SODIUM.** 4,4'-diaminodiphenylsulfone-2-N-acetylsulfonamide. See: Promacetin, Tab. (Parke, Davis)
- ACETO-VER.** (Mallard) Aspirin 5 gr., veratrum viride Pow. 1 gr., rutin 20 mg., Vit. C 30 mg./Tab. Bot. 100s, 500s, 1000s.
- Use: Capillary diseases.
- ACETPHEN-ACETYL COMPOUND.** (First Texas) Aspirin 3 gr., acetophenetidin 2 gr., caffeine $\frac{1}{2}$ gr., gelsemium tr. 1 min./Cap. Bot. 100s, 500s, 1000s. See A.P.C. w/gelsemium
- Use: (2-3 cap. daily) Analgesic & antipyretic.
- ACETPHENARSINE.** See: Acetarsona, Tab. (Abbott).
- ACETPHENOLISATIN.** (Chemo Puro) Pow. Pkg. for mfg.
- See: Diacetylhydroxyphenylisatin, Prep. (Various Mfr.).
- Use: Laxative.
- ACETPYROGALL,** N.N.R. 1951. Pyrogallol triacetate.
- ACETRIZOATE SODIUM,** N.N.D. 1959. Injection, U.S.P. 3-Acetylamino-2,4,6-triiodobenzoate sodium.
- See: Thixokon (Mallinckrodt)
- Urekon Sodium, Amp. (Mallinckrodt)
- W/Polyvinylpyrrolidone. See: Salpixon, Vial (Ortho).
- ACETRIZOIC ACID,** U.S.P., N.N.R. 1951. 3-Acetylamino-2,4,6-triiodobenzoic acid. See: Acetrizoate, Sod. (Mallinckrodt).
- ACET-THEOCIN SODIUM.** See: Theophylline Sod. Acetate. (Various Mfr.).
- ACETYCOL.** (Warner-Chilcott) Colchicine salicylated 0.25 mg., p-aminobenzoic acid 162 mg., aspirin 325 mg., thiamine 5 mg., ascorbic acid 20 mg., niacin 15 mg./Tab. Bot. 100s, 500s, 1000s.
- Use: (1 i.d.) Analgesic in arthritis, gout & neuritis.
- ACETYL ADALIN.** See: Acetylcarbromal. (Various Mfr.).
- p-ACETYLAMINO BENZALDEHYDE THIOSEMICARBAZONE.** (Amithiozone, Antib, Berculon A, Benzothiazon, Conteben, Myuzone, Neustab, Tebethion, Thiomicid, Thioparamizone, Thiacetazone)
- See: Panrone, Tab. (Panray).
- Teebazone, Tab. (Consol. Midland)
- Tibione, Tab. (Schenley).
- Use: With streptomycin in the treatment of tuberculosis.
- ACETYLAMINO BENZENE.** See: Acetanilid. (Various Mfr.).
- 2-ACETYLAMINO-4-(N-(p-CHLOROPHENYL)-ACETAMIDO)-s.TRIAZINE.** (Searle) Under study.
- Use: Oral diuretic.
- 2-ACETYLAMINO-5-NITROTHIAZOLE.** See: Tritheton, Tab. (Ortho)
- N-ACETYL-p-AMINOPHENOL.** Acetaminophen.
- See: Amdil, Tab. (Breon)
- Anelix, Elix. (Kirkman)
- Apamide, Tab. (Ames)
- Elixodyne, Liq. (Lemmon)
- Febrolin, Elix. (Buffington)
- Fendon, Tab. (A.P.C.)
- Lyteca Syrup. (Lloyd, D. & W.)
- Metalid, Tab. (Columbus)
- Nacetyl (Allison)
- Nebes, Tab. (Norwich Pharmacal)
- Pedituss (Sutliff & Case)
- Tempra, 5 gr. (Mead)
- Tussapap (Sutliff & Case)
- Tylenol, Elix. (McNeil)
- BUFFERED TABLETS.**
- See Apamide-Ves, Tab. (Ames)
- W/Acetophenetidin, phenylpropanolamine HCl, phenyltoloxamine dihydrogen citrate. See: Sinutab, Tab. (Warner-Chilcott)
- W/Acetylcarbromal. See: Apromal, Tab. (Ames).
- W/A.P.C. See: Doloro Compound, Tab. (Buffington)
- W/Aspirin & caffeine.
- See: Acetyl duo, Tab. (Chicago Pharm.).
- Trigesic, Tab. (Squibb).

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
16 mg. 4 gr.	Acetanilid 10 mg., caffeine 2 gr., salol 4 gr. Acetanilid 10 gr., NaBr. 20 gr., caffeine 2 gr.	Tilden Lilly	Ferbrisol, Tab. Acet. Brom. w/Phenacetin Cpd. Elix. Sinufab, Tab.
150 mg.	N-Acetyl-p-aminophenol 150 mg., phenylpropanolamine HCl 25 mg., phenyltoloxamine dithydrogen citrate 22 mg.	Warner-Chilcott	Achrocidin, Tab.
120 mg.	Achromycin HCl 125 mg., caffeine 30 mg., salicylamide 150 mg., chlorothen citrate 25 mg.	Lederle	Achrocidin, Tab.
120 mg.	Achromycin HCl 125 mg., salicylamide 150 mg., Vit. C 25 mg., pyrilamine maleate 15 mg.	Lederle	Achrocidin, Syr./5 cc.
2 gr.	Aluminum aspirin 3½ gr., caffeine ¾ gr., isobutylalyl barbituric acid ¾ gr.	L. L. Drug Co.	Alatal, Tab.
3 gr.	Alurate 1 gr.		
1 gr.	Amidopyrine 3½ gr., caffeine ½ gr., hyoscyamus ext. ¼ gr.	Roche Lilly	Allonal, Tab. Amidophen, Pulv.
1 gr.	Aminopyrine 3½ gr., caffeine ½ gr., hyoscyamus ext. ¼ gr.	Massengill	Andofive, Cap.
1½ gr.	Amidopyrine 2 gr., caffeine ¾ gr., phenobarbital ½ gr.	Chicago	Nalgised, Tab.
1½ gr.	Amidopyrine 3 gr., caffeine ¾ gr., phenobarbital 1/6 gr., Mag. oxide	Harco	Compirin, Tab.
2½ gr.	Amobarbital ½ gr., d-amphetamine sulf. 5 mg., aspirin 2½ gr.		
2½ gr.	Amobarbital ½ gr., d-amphetamine 5 mg., aspirin 2½ gr.	Smith, Kline & French	Daprisal, Tab.
2½ gr.	d-Amphetamine phosphate 1/24 gr., aspirin 2½ gr.	Smith, Kline & French	Daprisal, Tab.
2 gr.	rac-amphetamine 1/26 gr., aspirin 2½ gr.	Neisler	Threazie, Tab.
2 gr.	Amphetamine sulfate 2.5 mg., aspirin 4 gr.	Harvey	Harvodyne # 4, Tab.
2½ gr.	Amphetamine sulfate 2.5 mg., aspirin 2½ gr.	Smith, Kline & French	Edrisal, Tab.
2½ gr.	d-Amphetamine sulfate 2.5 mg., aspirin 3½ gr.	Pitman-Moore	Alcefain, Tab.
2½ gr.	d-Amphetamine sulfate 2.5 mg., pentobarbital sod. 3½ gr.	Arnar-Stone	Polygesic, Tab.
1½ gr.	Aspirin 3 gr., opium pow. 1/20 gr., ipecac pow. 1/20 gr., atropine sulf. 1/500 gr., camphor ½ gr.	Blue Line	Daphenco, Cap.
194 mg.	Aspirin 162 mg., phenobarbital 16.2 mg., hyoscyamine sulf. 0.031 mg., prophenpyridamine maleate 12.5 mg., phenylephrine HCl 10 mg.	Robins	Pi-naphen Plus, Tab.
2.5 gr.	Aspirin 3 gr., promethazine HCl 6.25 mg., mephentermine sulf. 7.5 mg.	Ives-Cameron	Synalgos, Cap.
2 gr.	Aspirin 4 gr., racephedrine HCl 1/12 gr., pyrilamine maleate 1/5 gr.	Uimer	Hiscaphen, Tab.

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
2 gr.	Sorbital 1½ gr., aspirin 3½ gr.	Warner	Peralga, Tab.
120 mg.	Belladonna alkaloids 0.0625 Gm., aspirin 180 mg., phenobarbital 16 mg.	Rorer	Paadon, Cap.
½ gr.	Belladonna tr. 3.8 min., aspirin 3½ gr., caffeine ½ gr.	Vanpelt & Brown	Bellaspro, Tab.
2 gr.	Belladonna ext. 1/5 gr., Cal. iodide ½ gr., Sod. salicylate 2 gr., camphor ½ gr., capsicum 1/10 gr.	Superior Pharmacial	Cepheb, Tab.
2½ gr.	Benzedrine sulfate 2.5 mg., aspirin 2½ gr.	Smith, Kline & French	Edrisal, Tab.
	Codeine sulfate ½ gr.	Smith, Kline & French	w/Codeine
¾ gr.	Codeine sulfate ¼ gr., ½ gr.	Smith, Kline & French	w/Codeine
	Bromisovalum ½ gr., ethyl salicylate carbonate 1 gr., caffeine 1/16 gr.	Columbus Pharmacial	Acetobro, Pediatob No. 1
½ gr.	Caffeine 1/10 gr., sod. bicarbonate 1/5 gr., amm. carbonate 1/32 gr., licorice ext.	Jenkins Labs.	Anodyne Jr., Tabs.
1 gr.	Colchicine salicylate 1/700 gr., aspirin 5 gr., caffeine ½ gr.	First Texas	For-Dyne, Cap. & Tab.
150 mg.	Chlorcyclizine HCl 15 mg., pseudoephedrine HCl 20 mg., aspirin 200 mg., caffeine 30 mg.	Burroughs Wellcome	Emprazil, Tab.
320 mg.	Chlorothen citrate 25 mg., caffeine 32 mg.	Whittier	Caubren Comp., Tab.
320 mg.	Chlorothenylpyramine 25 mg., caffeine 32 mg.	Breon	Fenoxydyne Comp., Tab.
16 mg.	Chlor-Trimeton maleate .75 mg., aspirin 80 mg.	Schering	Coricidin Pediatric Mediflets
130 mg.	Chlor-Trimeton maleate 4 mg., salicylamide 190 mg., caffeine 30 mg., ascorbic acid 50 mg., methamphetamine HCl 1.25 mg.	Schering	Coricidin Forte, Cap.
2 gr.	d-Desoxyephedrine HCl 1/24 gr., aspirin 4 gr., butabarb ¼ gr.	Central	Asphedex, Tab.
200 mg.	di-Desoxyephedrine HCl 1 mg., Sod. gentisate 200 mg.	Crookes	Gentexin, Tab.
2 gr.	Diallylbarbital ¾ gr., aspirin 3½ gr.	Ulmer	Alprine, Tab.
160 mg.	Dihydrohydroxycodone HCl 4.5 mg., dihydrohydroxycodoneone terephthalate 0.38 mg., hexobarbital 100 mg., homatropine terephthalate 0.38 mg., aspirin 224 mg., caffeine 32 mg.	Endo	Percobarb, Cap.
1 gr.	Dover's pow. 3/16 gr., aspirin 3 gr., caffeine ¼ gr.	Smith, Miller & Patch	Acetyl-Caf-Phen, Cap.
1½ gr.	Dover's pow. ¼ gr., aspirin 3½ gr., caffeine ½ gr., camph. ½ gr.	Harvey	Harvodyne # 3, Tab.
1½ gr.	Dover's pow. ¼ gr., aspirin 2 gr., caffeine ¼ gr., camphor monobromide ¼ gr.	Warren-Teed	Oveen, Tab.
1½ gr.	Dover's pow. ¼ gr., aspirin 2 gr., caffeine ¼ gr., atropine sulfate 1/500 gr., camphor ¼ gr.	Warren-Teed	Oyofen, Cap. & Tab.
1½ gr.	Dover's pow. ¼ gr., aspirin 2 gr., atropine sulfate 1/500 gr., camphor monobromide ¼ gr.	Lemmon	Aspidyne Comp., Cap. & Tab.

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
1½ gr.	Dover's pow. ½ gr., aspirin 2 gr., caffeine ½ gr., atropine sulfate 1/500 gr., camphor ¼ gr., Sod. sulfate 7½ gr.	Hance	Dovium, Cap.
1½ gr.	Ephedrine HCl ½ gr.	Abbott	Dysco, Tab.
3 gr.	Ephedrine HCl ¾ gr., phenobarbital ¼ gr.	Mallard	Phenodrin, Cap.
160 mg.	Ephedrine ethylenediamine 10.6 mg., theobromine sod. acetate 25 mg., caffeine 50 mg.	Field Labs.	Kradon, Tab.
130 mg.	Ergotamine tartrate 1 mg., belladonna alkaloids 0.1 mg., caffeine 100 mg.	Organon	Wigraine, Tab.
2 gr.	Gelsemium 1/10 gr., aspirin 3 gr., caffeine ½ gr.	Smith-Dorsey	Phen-Acetyl, Cap.
2 gr.	Gelsemium 3/10 gr., aspirin 4 gr., caffeine ¼ gr.	Smith-Dorsey	Saliphen, Tab.
2 gr.	Gelsemium ½ gr., aspirin 4 gr., caffeine ½ gr., strychnine sulfate 1/240 gr., camphor monobromide ½ gr.	Elder	Naretic, Tab.
2½ gr.	Gelsemium 1/5 gr., aspirin 2½ gr., colchicine salicylate, camphor	First Texas	Formasal & Camphor Comp., Tab.
130 mg.	Gelsemium ext. 2.2 mg., aspirin 130 mg., caffeine 16 mg., hyoscyamus 11 mg.	Drug Products	Henasphen, Cap.
2½ gr.	Gelsemium tr. 2 min., aspirin 3½ gr., colchicine salicylate ¼ gr.	Strasenburgh	Semapirin, Tab.
2 gr.	Gelsemium tr. 3 min., aspirin 4 gr., colchicine salicylate ¼ gr.	Central	Asphac-G, Tab.
2 gr.	Gelsemium tr. 2 min., aspirin 3 gr., colchicine salicylate ¼ gr., methyl salicylate	Noyes	Aidant, Tab.
2½ gr.	Gelsemium tr. 1 min., aspirin 3½ gr., colchicine salicylate ¼ gr., diallylbarbital ¼ gr.	Buffington	Doradol, Tab.
120 mg.	Hesperidin 100 mg., vit. C 50 mg., phenylephrine HCl 5 mg., prophenpyridamine maleate 6.25 mg., methapyrilene HCl 8.33 mg., pyrilamine maleate 8.33 mg., salicylamide 200 mg., caffeine 30 mg.	Boyle	Boyle Citra, Cap.
1 gr.	Hyoscyamus ext. ¼ gr., amidopyrine 3½ gr., caffeine ½ gr.	Lilly	Amidophen, Pulv.
2 gr.	Methopryliline fumarate 8.3 mg., phenyltoloxamine dihydrogen citrate 8.3 mg., pyrilamine maleate 8.3 mg., aspirin 3 gr., caffeine ½ gr.	Walker Corp.	Trist-A-Lin, Tab.
100 mg.	Pamabrom 50 mg.	Whittier	M-Minus-5, Tab.
80 mg.	Penicillin V Pot. 100,000 u., Pyronil 7.5 mg., histadyl 12.5 mg., clopane 6.25 mg., aspirin 114 mg., caffeine 16 mg.	Lilly	V-Kor, Tab.
130 mg.	Penicillin V 100,000 u., salicylamide 194 mg., promethazine HCl 6.25 mg., mephentermine sulf. 3 mg.	Wyeth	Pen-Vee-Cidin, Cap.

Acetophenetidin Combinations

[8]

Acetophenetidin	Others	Manufacturer	Brand
2 1/2 gr.	Pentobarbital 1/4 gr., aspirin 4 gr.	Smith-Dorsey	Angesal, Tab.
2 1/2 gr.	Pentobarbital sod. 1/4 gr., dextroamphetamine sulfate 2.5 mg., aspirin 3 1/2 gr.	U.S. Standard Prods.	Domitral, Tab.
2 gr.	Phenobarbital 1/4 gr., aspirin 4 gr.	Warren-Teed	Axotal, Tab.
2 gr.	Phenobarbital 1/4 gr., aspirin 4 gr.	Harvey	Harvadal, Tab.
2 gr.	Phenobarbital 1/4 gr., aspirin 4 gr., atropine sulfate 1/500 gr.	Warren-Teed	Axopin, Tab.
120 mg.	Phenobarbital 15 mg., aspirin 180 mg., belladonna alkaloids 0.0625 Gm.	Rorer	Paadon, Cap. & Tab.
1 1/2 gr.	Phenobarbital 1/2 gr., amidopyrine 2 gr., caffeine 3/4 gr.	Chicago	Nalgised, Tab.
1 1/2 gr.	Phenobarbital 1/6 gr., amidopyrine 3 gr., caffeine 3/4 gr., Mag. oxide	Harco	Compirin, Tab.
3 gr.	Phenobarbital 1/4 gr., ephedrine HCl 3/4 gr.	Mallard	Phenadrin, Cap.
120 mg.	Phenylephrine HCl 5 mg., pyrilamine maleate 12 mg., salicylamide 250 mg., dried aluminum hydroxide gel 120 mg., caffeine 15 mg.	Central	Cenagesic, Tab.
150 mg.	Phenyltoloxamine dihydrogen citrate 44 mg., salicylamide 150 mg., caffeine 32 mg.	Organon	Medache, Tab.
150 mg.	Phenyltoloxamine dihydrogen citrate 44 mg., salicylamide 150 mg., caffeine 32 mg.	Philadelphia Ampoule	Achease, Tab.
100 mg.	Pot. penicillin G 250,000 u., salicylamide 100 mg., phenyltoloxamine dihydrogen citrate 25 mg.	Smith, Miller & Patch	Elpafec, Tab.
100 mg.	Pyribromide 50 mg.	Whitier	M-Minus-4, Tab.
2 1/2 gr.	Pyranisamine maleate 25 mg., aspirin 2 1/2 gr., gelsemium 1/5 gr., colchicine salicylate camphor	First Texas	Formasal & Campher Comp. w/Antihistamine
2 1/2 gr.	Pyrilamine maleate 25 mg., aspirin 3 1/2 gr., caffeine 1/2 gr.	Horton & Converse	Histapyran Comp., Tab.
2 1/2 gr.	Pyrilamine maleate 25 mg., aspirin 3 1/2 gr., caffeine 1/2 gr., quinine sulfate 1/2 gr.	Horton & Converse	Hista-Pacq., Tab.
2 1/2 gr.	Pyrilamine maleate 25 mg., aspirin 3 1/2 gr., caffeine 1/2 gr.	Hance	Hista-Pac, Tab. & Cap.
1 1/2 gr.	Pyrilamine maleate 25 mg., aspirin 3 1/2 gr., caffeine 1/2 gr.	Rexall	Monhislin-A.P.C., Tab.
2 1/2 gr.	Pyrilamine maleate 10 mg., aspirin 3 1/2 gr., PABA 1/2 gr.	Gold Leaf	Parisamine, Tab.
2 1/2 gr.	Pyrilamine maleate 5/12 gr., aspirin 3 gr., caffeine 1 gr., Vit. C 2 1/2 gr.	Brewer	Pyrahist Comp. w/Vit. C Cap.
2 1/2 gr.	Pyrilamine maleate 25 mg., aspirin 3 1/2 gr., caffeine 1/2 gr., hyoscyamine HBr 0.4 mg., hyoscine HBr .008 mg.	Meyer	Almegestic, Tab.

Acetophenetidin Combinations

Acetophenetidin	Others	Manufacturer	Brand
2 gr.	Pyritamine maleate 25 mg., salicylamide 3 gr., caffeine ½ gr., atropine sulf. 1/600 gr., quinine sulf. ½ gr., ascorbic acid 25 mg.	Taylor	Histaquin, Cap.
2½ gr.	Salicylamide 3½ gr., caffeine ½ gr.		
2.5 gr.	Salicylamide 3.5 gr., caffeine 0.5 gr., ascorbic acid 50 mg., pyritamine maleate 25 mg., hesperidin 25 mg.	Rowell Bryant	Salprin, Tab. Biocol, Cap.
2½ gr.	Salicylamide 3½ gr., dl-desoxyephedrine HCl 1/30 gr.	Wampole	Tryads, Tab.
2 gr.	Salicylamide 1 gr., dl-desoxyephedrine HCl 5 mg., Mag. gluconate 3 gr.	Mallard	Sedonate, Tab.
120 mg.	Salicylamide 230 mg., propenpyridamine maleate 10 mg., caffeine 30 mg., ascorbic acid 20 mg., hesperidin purified 20 mg.	Roerig	Caryban, Cap.
120 mg.	Salicylamide 250 mg., racephedrine HCl 5 mg., pyritamine maleate 12 mg., ascorbic acid 50 mg., Dover's powder 24 mg.	Central	Centamal, Cap.
65 mg.	Salicylate sod. 320 mg., salicylic ethyl ester carbonate 32 mg., calchicum root pow. ext. 1.6 mg.	Endo	Tetrogen, Tab
¼ gr.	Salicylic acid ½ gr., sod. bicarbonate 3 gr., mag. salicylate 2 gr., strontium salicylate 2 gr., methyl salicylate, pancreatin, diastase	Jenkins Labs.	Foursalco, Tab.
2½ gr.	Salysal 2½ gr., caffeine ½ gr.	Rare-Galen	Salyphec, Tab.
2 gr.	Sandotal ¾ gr., aspirin 3 gr., caffeine ¾ gr.	Sandoz	Florinal, Tab.
200 mg.	Sod. genitrate 200 mg., dl-desoxyephedrine HCl 1 mg.	Crookes	Genutexin, Tab.
2½ gr.	Sod. salicylate, PABA, Sod. bicarb.	Anglo-French	Encynex, Tab.
120 mg.	Tao 125 mg., salicylamide 150 mg., buclizine 15 mg., caffeine 30 mg.	Roerig	TAO-AC, Cap.
120 mg.	Tetracycline HCl 125 mg., caffeine 30 mg., salicylamide 150 mg., buclizine HCl 15 mg.	Pfizer	Tetracyclin, Tab.
125 mg.	Tetracycline phosphate 125 mg., aspirin 150 mg., caffeine 30 mg., phenyltoloxamine HBr 25 mg.	Bristol	Tetrex APC w/Bristamin, Cap.
2½ gr.	Thenylpyramine HCl 25 mg., aspirin 3¾ gr., d-amphetamine sulfate 1 mg.	Pitman-Moore	Cohistine, Tab.
2½ gr.	Vit. C 25 mg., Cal. glutamate 2 gr., aspirin 5 gr., caffeine 1 gr.	Testagar	Nuprin Comp., Tab.

- W/Aspirin & caffeine with codeine. See: Trigesic w/Codeine, Tab. (Squibb).
- W/Aspirin, phenylephrine HCl, noscapine, caffeine, pyrilamine maleate, methapyrilene fumarate. See: Tussapap, Tab. (Sutliff & Case)
- W/Aspirin, trimethyl xanthine, chlorpheniramine maleate, phenylephrine HCl. See: Thorexin Cold Capsules. (Gillette Labs.)
- W/Butabarbital, d-amphetamine sulfate, acetylsalicylic acid, salicylamide. See: Stuart Prolar, Tab. (Stuart)
- W/Butabarbital sod.
See: Butapap (Physicians Products) Nacetyl, Tab. (Allison)
- W/Butabarbital Sod. & methamphetamine HCl. See: Algoson, Tab. (McNeil).
- W/Chlorphenyridamine maleate, phenylephrine HCl, phenylpropranolamine HCl, ephedrine sulfate. See: Pyrets, Tab. (Bernhoff)
- W/Demerol. See: Demerol APAP (Breon)
- W/Dihydrocodeinone bitartrate, meperridine HCl. See: Demerol Compound, Tab. (Breon)
- W/Dihydrocodeinone bitartrate, phenylephrine HCl, pyrilamine maleate, pot. guaiacolsulfonate, sod. citrate, citric acid, chloroform. See: Codigesic, Syr. (Central)
- W/Flexin. See: Flexilon, Tab. (McNeil)
- W/Flexin, hydrocortisone. See: Flexilon-HC, Tab. (McNeil)
- W/d-Methorphan HBr, chlorpheniramine, phenylephrine. See: Romilar-CF, Syr. (Hoffmann-La Roche)
- W/Methyliso-octenylamide mucate, salicylamide. See: Migrade Tab. (Crystal)
- W/Methyliso-octenylamine mucate, salicylamide, and caffeine. See: Nygram, Cap. (Dumas-Wilson)
- W/Neo-synephrine HCl, thenyldiamine HCl, caffeine. See: Neo-Synephrine Compound Cold Tablets, (Winthrop Labs.)
- W/Phenindamine tartrate, sulfadimethoxine, caffeine. See: Madricidin, Cap. (Roche)
- W/Phenylephrine HCl, chlorpheniramine maleate, noscapine, salicylamide, Vit. C. See: Noscaps, Cap. (Table Rock)
- W/I-Phenylephrine HCl, isothipendyl HCl, dimethoxanate HCl, ammonium chloride, sodium citrate, chloroform. See: Cothera Compound, Liq. (Ayerst)
- W/Phenylephrine, d-methorphan HBr, pyrilamine maleate, potassium guaiacolsulfonate, alcohol. See: Thormal, Liq. (Central)
- W/Phenylpropranolamine HCl, phenyltoloxamine dihydrogen citrate, pot. quaiacolsulfonate. See: Conex Syrup (Lloyd, Dabney & Westerfield)
- W/Phenyltoloxamine dihydrogen citrate & phenylpropranolamine HCl. See: Conex Plus, Tab. (Lloyd, Dabney & Westerfield)
- W/Phenyltoloxamine dihydrogen citrate, salicylamide, di-amphetamine phosphate, metropine. See: Acogesic, Cap. (Strassenburgh)
- W/Phenyltoloxamine dihydrogen citrate, salicylamide, Vit. C, citrus bioflavonoids, caffeine, racemic amphetamine sulfate. See: Euphenex, Tab. (Lloyd, Dabney & Westerfield)
- W/Salicylamide. See: Peditagesic (Columbus)
- W/Salicylamide, acetylsalicylic acid, butabarbital, d-amphetamine sulfate. See: Stuart Prolar, Tab. (Stuart)
- W/Salicylamide, acetylsalicylic acid, d-amphetamine sulfate, butabarbital, hyoscyamine sulfate, atropine sulfate, hyoscyne hydrobromide. See: Stuart Prolar-B, Tab. (Stuart)
- W/Salicylamide, caffeine, phenylephrine HCl, tristamine. See: Contramal-CP, Caps. (Physicians Products)
- W/Salicylamide, chlorpheniramine maleate, phenylephrine HCl, phenylpropranolamine HCl, caffeine, ascorbic acid. See: Tristamine Forte, Cap. (Barre)
- W/Salicylamide, codeine phosph., caffeine. See: Cylamide, Tab. (Chicago Pharmacol Co.) Also with phenobarb.
- W/Salicylamide, glyceryl guaiacolate, pyrilamine maleate, phenylephrine HCl. See: Guaiagesic, Liq. (Columbus)
- W/Salicylamide, phenobarbital, caffeine. See: Cylamide w/Phenobarbital, Tab. (Chicago Pharmacol Co.)
- W/Sod. salicylate, phenylephrine, glyceryl guaiacolate. See: Pedituss, Syr. (Sutliff & Case)
- W/Thephorin, madribon, caffeine. See: Madricidin, Cap. (Roche)
- 3-ACETYLAMINO-2,4,6-TRIIODOBENZOATE SODIUM.** See: Acetrisoate Sod. (Mallinckrodt).
- 3-ACETYLAMINO-2,4,6-TRIIODOBENZOIC ACID.** See: Acetrisoic Acid, U.S.P. (Mallinckrodt).
- ACETYLANILINE.** See: Acetanilid. (Various Mfr.).
- ACETYL-BETA-METHYLCHOLINE CHLORIDE,** U.S.P., N.N.R. 1947. See: Mecholy Chloride, Amp. (Merck Sharp & Dohme)
- ACETYL-BROMO-DIETHYLACETYL-CARBAMIDE.** See: Acetylcarbromal. (Various Mfr.).
- ACETYL-CAF-PHEN.** (Smith, Miller & Patch) Aspirin 3 gr., acetophenetidin 1 gr., caffeine 1/8 gr., Dover's Pow. 3/16 gr./Cap. Bot. 100s, 1000s.
Use: (1 t.i.d.) Analgesic.

- ACETYL-CARB.** (Sherman) Aspirin & Sod. bicarb. Tab. 500s, 1000s.
Use: (1-2 q. 3 h.) Analgesic.
- ACETYLCARBOMAL.** (Chemo Puro, Meyer) Acetyl adalin, acetyl-bromo-diethylacetyl-carbamide. Pow. for mfg.
See: Abasin. (Winthrop Labs.)
Sedamyl. (Schenley).
Use: Sedative.
- W/N-acetyl-p-aminophenol. See: Apromal, Tab. (Ames)
- W/Apamide. See: Apromal, Tab. (Ames).
- W/Bromisovalum.
- See: Daysed, Tab. (Tailby-Nason).
Sedayson, Tab. (Tailby-Nason).
- W/d-Desoxyephedrine HCl.
See: Euphased-5, Tab. (Schenley)
- W/Hyoscyamine sulfate, atropine sulfate & hyoscine HBr. See: Disperdon, Tab. (Standard).
- W/Mephenesin & reserpine. See: Amril, Tab. (Leeming)
- ACETYLCOLINE CHLORIDE.** (Various Mfr.).
Leader—Amp. (0.1 Gm., 0.2 Gm.) 1 cc., 12s.
Merck—Amp. (0.1 Gm.). Vial 10 Gm.
Use: S.C., I.M., I.V. Parasympathomimetic agent & vasodilator. Paroxysmal tachycardia.
- W/Saligenin. See: Acecoline, Amp. (Anglo-French).
- ACETYLCOLINE-LIKE THERAPEUTIC AGENTS.** See: Cholinergic agents.
- ACETYLDIGITOXIN,** N.N.D. 1959. See: Acyanid, Tab. (Sandoz).
- N¹-(ACETYL-3,4-DIMETHYL-5-ISOXAZOYL) SULFANILAMIDE.** See: Gantrisin Acetyl, Prep. (Roche Labs.)
- ACETYLDUO.** (Chicago Pharm.) Acetyl-p-aminophenol 2.5 gr., aspirin 3.5 gr., caffeine 0.5 gr./Tab. Bot. 100s, 1000s.
Use: (1 q. 3-4 h.) Analgesic.
- ACETYLIC COMPOUND.** (Central) Salicylamide 30 mg., acetophenetidin 30 mg., atropine sulfate 0.05 mg., aconite root 1.5 mg., camphor monobromated 8.0 mg., phenolphthalein 6.0 mg./Tab. Bot. 100s, 1000s
Use: Analgesic, antipyretic
- 5-ACETYLMIMINO-4-METHYL- Δ^2 -1,3,4-THIADIAZOLINE-2-SULFONAMIDE.** See: Methazolamide Neptazane (Lederle)
- ACETYLIN.** See: Acetylsalicylic Acid. (Various Mfr.).
- ACETYLMETHIONINE.** (Chemo Puro) Methionine 78%. Pow. for mfg.
Use: Nutrient.
- ACETYLMETHIONINE CALCIUM.** (Chemo Puro) Methionine 71%, calcium 9.5%. Pow. for mfg.
Use: Nutrient.
- ACETYL MIDICEL.** See: Midicel Acetyl Susp. (Parke, Davis)
- ACETYLPHENYLISATIN.** See: Diacetylhydroxyphenylistan.
- ACETYLRESORCINOL.** See: Resorcinol Monoacetate. (Various Mfr.).
- ACETYLSAL.** See: Acetylsalicylic Acid.
- ACETYLSALICYLIC ACID,** U.S.P., A.D.R., N.N.R. 1947; Capsules, N.F.; Tablets, U.S.P. (Various Mfr.) Acidum acetylsalicylicum, 2-acetoxybenzoic acid, Acetilum Acidulatum, Acetophen, Acetol, Acetosal, Acetosalic Acid, Acetosalin, Acetylcy, Acetylin, Acetylsal, Acylpyrin, Aspra, Helicon, Rhodine, Salacelin, Salcetogen, Saletin.
See: Aspirin Preps. (Various Mfr.).
Empirin (Burroughs Wellcome).
Salts.
See: Aluminum Acetylsalicylate. A.S.A., Tab. (Lilly)
Calcium Acetylsalicylate.
Use: Analgesic & antipyretic.
- W/Acetophenetidin & caffeine. See: A.P.C. preparations. (Various Mfr.).
- W/Mixtures. See: Aspirin preparations. (Various Mfr.).
- ACETYLSALICYLIC ACID, ACETOPHENETIDIN AND CAFFEINE,** N.F. See: A.P.C., Prep. (Various Mfr.).
- ACETYLSALICYLIC ACID ENTERIC COATED.** See: Asteric, Tab. (Brewer).
- ACETYLISTROBELINE.** See: Asmatone, Cap. (National Research).
- N-ACETYL SULFAMETHOXYPYRIDAZINE,** N.N.D. 1959. 3-[N-Acetylsulfanilamide]-6-methoxy-pyridazine. See: Kynex Acetyl Pediatric Suspension (Lederle)
- N¹-ACETYLSULFANILAMIDE.** (Albucid; p-Aminobenzenesulfonacetamide; Sulfacet; Sulfacetamide, incorrectly called A.P.C.; N-Sulfanilylacetamide.) See: Sulamyd. (Schering).
Use: Sulfonamide therapy.
- ACETYL SULFISOXAZOLE,** N.N.D. 1959. N₁-acetyl-3,4-dimethyl-5-sulfanilamidoisoxazole. See: Gantrisin Acetyl. (Hoffmann-La Roche)
- ACETYLTANNIC ACID,** N.N.R. 1941.
- ACHEASE.** (Philadelphia Ampoule) Phenyltoloxamine dihydrogen citrate 44 mg., acetophenetidin 150 mg., salicylamide 150 mg., caffeine 32 mg./Tab. Bot. 100s, 1000s.
Use: Tension headaches.
- ACHLORHYDRIA DETERMINATION.** See: Diagnex.
- ACHLORHYDRIA THERAPY.**
See: Acidol. (Various Mfr.)
Acidol-Pepsin, Cap. (Winthrop Labs.)
Acid-V, Cap. (Merck Sharp & Dohme)
Aclor, Cap. (Cole Chem. Co.)

Chlor-Ethamine, Tab. (Pitman-Moore)
Glutamic Acid HCl

ACHROCIDIN. (Lederle) **Tab.** Achromycin HCl 125 mg., phenacetin 120 mg., caffeine 30 mg., salicylamide 150 mg., chlorothen citrate 25 mg./tab. Bot. 24s.

Syr.: Achromycin HCl 125 mg., phenacetin 120 mg., salicylamide 150 mg., Vit. C 25 mg., pyrilamine maleate 15 mg., methylparaben 4 mg., propylparaben 1 mg./5 cc. **Syr. Bot.** 4 fl. oz.

Use: Antibiotic, antihistamine, analgesic therapy.

ACHROMYCIN, U.S.P. N.N.D. 1959. (Lederle) Tetracycline HCl.

Cap.: (50 mg. & 100 mg.) Bot. 25s & 100s; (250 mg.) Bot. 16s & 100s.

Ear Solution: (50 mg. vial and a 10 cc. vial of 5% benzocaine in propylene glycol).

Intravenous: (100, 250 & 500 mg.) Vial.

Intramuscular: (100 & 250 mg.) Vial.

Oral Susp.: (250 mg./5 cc.) Bot. 1.5 Gm., dry pow. 1 oz.

Oint.: (3%) Tube $\frac{1}{2}$ & 1 oz.

Ophthalm. Oil Susp.: 1%, Bot. w/dropper 4 cc.

Ophthalm. Oint.: (1%) Tube $\frac{1}{8}$ oz.

Ophthalm. Sterilized: Vial (25 mg.) w/dropper.

Pediatric Drops, Liquid: (100 mg./cc.) Bot. w/dropper 10 cc.

Pharyngets: (15 mg.) Pkg. 10s.

Spersoids: Choc. flavor (50 mg./tsp.) Bot. 12 & 25 doses.

Surgical Powder: Vial (200 mg./Gm.) 5 Gm.

Syr.: (125 mg./5 cc.) Bot. 2 & 16 oz.

Tab.: (50 & 100 mg.) Bot. 25s & 100s; (250 mg.) Bot. 50s, 250s & 1000s.

Tab. Soluble: (50 mg.) Tube 40s, 100s.

Troches: (15 mg.) Pkg. 25s, 250s.

W/Hydrocortisone Oint. (Lederle). Tetracycline HCl 3%, Hydrocortisone 2%. Tube 5 Gm.

Use: Pyogenic infections of the skin.

W/Hydrocortisone (Lederle) Ophthalm. Oint. Tetracycline HCl 1%, Hydrocortisone 1.5%. Tube $\frac{1}{8}$ oz.

Use: Eye infections and skin infections

W/Phenacetin, caffeine, salicylamide, chlorothen citrate. See: Achrocidin, Syr., Tab. (Lederle)

ACHROMYCIN NASAL SUSPENSION.

(Lederle) Tetracycline HCl 3.75 mg., hydrocortisone acetate 2 mg., phenylephrine HCl 0.125% /cc. Bot. squeeze 15 cc.

Use: Upper respiratory infections.

ACHROMYCIN SF CAPSULES. (Lederle)

Achromycin 250 mg., Vit. C 75 mg., B-1 2.5 mg., B-2 2.5 mg., niacinamide 25 mg.,

Vit. B-6 0.5 mg., Cal. pantothenate 5 mg., Vit. B-12 1 mcg., folic acid 0.375 mg., Vit. K 0.5 mg./Cap. Bot. 16s, 100s.

Use: Antibiotic with stress formula vitamins.

ACHROMYCIN SYRUP. (Lederle) Tetracycline HCl 125 mg., methylparaben 0.08%, propylparaben 0.02%, alcohol 1%/5 cc. Bot. 2 oz., 1 pt.

Use: Antibiotic therapy.

ACHROMYCIN-V. (Lederle) Achromycin 250 mg. buffered w/citric acid & Sod. citrate/Cap. Bot. 16s, 100s. Pink Caps.: Achromycin 100 mg. w/citric acid & Sod. citrate/Cap. Bot. 25s, 100s. **Syr.:** Achromycin 125 mg. buffered w/citric acid & Sod. citrate/5 cc. Bot. 2 oz. 1 pt

Pediatric Drops: Achromycin 100 mg. w/citric acid & Sod. citrate/cc. Dropper Bot. 10 cc.

Use: Treatment of infections caused by both gram-negative and gram-positive bacteria

ACHROSTATIN-V. (Lederle) Tetracycline citric acid & Sod. citrate buffered 250 mg., nystatin 250M u./Cap. Bot. 16s, 100s. Oral Susp. tetracycline citric acid & Sod. citrate buf. 125 mg. & nystatin 125M u./5 cc. Bot. 60 cc.

Use: Broad-spectrum antibiotic w/antifungal therapy

ACIBAN. (Lederle) Calcium caseinate 0.4 Gm., calcium carb. 0.1 Gm./Tab. Bot. 50s, 250s, 1000s.

Use: (1-2 t.i.d.) Antacid.

ACICHEK. (Valentine) Dihydroxy aluminum aminoacetate 250 mg., Sod. carboxymethylcellulose 150 mg., glycine 30 mg./Tab. Bot. 100s, 1000s.

Use: Gastric hyperacidity in peptic ulcer.

ACID ACRIFLAVINE. See: Acriflavine Hydrochloride. (Various Mfr.).

ACID CARBOLIC, NUTGALL, AND ATROPINE COMPOUND. (Upjohn) Phenol 6 gr., nutgall (equiv.) 36 gr., atropine $\frac{1}{4}$ gr., benzocaine 10%₁₀ gr./oz. Oint. Tube 1 oz. Jar 1 lb.

Use: Rectally, local anesthetic, analgesic & antipruritic.

ACID CITRATE DEXTROSE SOLUTION. See: A.C.D. Sol.

ACID CLOAK CREME. (Dome) pH 5.0 Creme. Tube 1 oz., Jar 4 oz., 1 lb.

Use: Restore skin to normal pH.

ACI-DERMA. (Pincer) An aqueous emulsion of glyceryl monostearate, cetyl alcohol, sodium lauryl sulfate, propyl and benzyl p-hydroxy benzoates and propylene glycol, buffered to pH 2-3. Bot. 4 oz.

Use: Restore the acid mantle of the skin

ACID HISTAMINE PHOSPHATE. See: Histamine Phosphate. (Various Mfr.).

ACIDIRON. (Walker Labs.) Ferrous sulfate

- exsic. 3 gr., hydrochloric acid diluted 125 mg./tab. Bot. 100s.
Use: Anemias.
- ACID MANTLE CREME.** (Dome) Aluminum acetate in specially prep. water-soluble hydrophilic cream at pH 4.2 Tube 1 oz. Jar 4 oz., 1 lb.
Use: Skin prophylactic agent in restoring normal skin acidity.
- ACID MANTLE LOTION.** (Dome) Aluminum acetate lotion. pH 4.5. Bot. plastic, 1 oz. Bot. 4 oz. squeeze, 1 pt., ½ and 1 gal.
Use: Dermatitis.
- ACIDOL.** See: Betaine Hydrochloride. (Various Mfr.).
- ACIDOLATE.** (White) (Hypoallergenic liquid skin cleanser.) Sulfated vegetable oils, liquid petrolatum & water. Bot. 8 oz., & gal.
Use: Cleanses skin surfaces without irritation.
- ACIDOL-PEPSIN.** (Winthrop Labs.) Betaine HCl 4 gr., pepsin 3.5 gr./Cap. Bot. 50s.
Use: (1-2 t.i.d.) Achlorhydria
- ACIDORIDE.** (Abbott) d-Glutamic acid HCl. Cap. (5 gr.), Bot. 100s, 500s, 1000s.
Use: Achlorhydria.
- ACIDOTHYIN.** (Flint-Eaton) Glutamic acid HCl. (Each cap. is equivalent to 10 min. of Dilute Hydrochloric Acid U.S.P.) Bot. 100s, 500s, 1000s.
Use: (1-2) Achlorhydria.
- ACID TRYPAPFLAVINE.** See: Acriflavine Hydrochloride. (Various Mfr.).
- ACIDULIN.** (Lilly) Glutamic acid HCl. (Equivalent to 0.62 cc. [10-min.] of diluted hydrochloric acid.) Pulvules. Bot. 100s, 500s, 1000s.
Use: (1-2) Achlorhydria.
- ACIDUSOL.** (Texas) Oxyquinoline sulfate, lactic acid, zinc sulfocarbonate, citric acid, dextrose, buffered solution. Bot. 8 oz.
Use: Vaginal cleanser.
- ACIGLUMIN.** See: Glutamic Acid Hydrochloride. (Various Mfr.).
- ACI-JEL.** (Ortho) Acetic acid, ricinoleic acid, oxyquinoline sulfate, and glycerine in a vegetable gum base. Jelly. Tube with applicator, 3 oz.
Use: Vaginal antiseptic.
- ACINTRAZOLE.** See: Tritheton (Ortho)
- ACITAMIN.** (Massengill) Glutamic acid HCl 0.324 Gm., thiamine HCl 1 mg., riboflavin 1 mg., nicotinamide 5 mg./cap. Bot. 100s, 500s, 1000s.
Use: (1-2) Achlorhydria.
- A-C-K.** (G. F. Harvey) Aspirin 5 gr., menadione 1/200 gr., ascorbic acid ½ gr./Tab. Bot. 100s, 1000s
Use: Salicylate therapy.
- A-C-K BUFFERED.** (G. F. Harvey) Aspirin 333 mg., ascorbic acid 33.3 mg., menadione 0.33 mg., calcium carbonate 60 mg./tab. Bot. 100s, 1000s
Use: (1-2 q. 3 h.) Analgesic.
- ACLETIN.** (Associated Concentrates) L-cithin, soya 8 gr. with vitamins/Cap. Bot. 100s, 250s.
Use: (1 to 2 cap. daily) Fat metabolism.
- ACLOR.** (Cole) Glutamic acid HCl. Cap. (5 gr.) Bot. 100s
Use: (1-2) Achlorhydria.
- ACNAVEEN.** (Aveeno Corp.) Colloidal oatmeal, kaolin, sulfur 3%, and hexachlorophen 0.5%. Lot., Pow., Pkg. 4 oz.
Use: Topically, cleansing agent in the treatment of acne
- ACNE CORT DOME.** (Dome) Hydrocortisone ¼%, sulfur 4%, 2,2'-thiobis (4,6-dichlorophenol) 1%, resorcinol monoacetate 3%. Cream, Tube 1 oz. Lotion, Squeeze Bot. 1 oz.
Use: Treatment of acne.
- ACNEDERM.** (Lannett) 5% Dispersible sulfur, 3% zinc sulfate. Bot. 2 oz.
Use: Skin conditions
- ACNE-DOME.** (Dome) Regular: sulfur 4% (bithional), 2,2'-thiobis (4,6-dichlorophenol) 1%, resorcinol monoacetate 3%; Double strength: sulfur 8%, 2,2'-thiobis (4,6-dichlorophenol) 1%, resorcinol monoacetate 3%. Creme 1 oz.
Use: Treatment of acne.
- ACNE MIXED "U.B.A."** (Lilly) Udenaturec antigens of bacillus acne & staphylococci standardized to 0.05 mg. nitrogen/cc. Vial 20 cc.
Use: Pustular acne.
- ACNE SEROBACTERINE MIXED VACCINE.** (Merck Sharp & Dohme) Acne bacillus 800 million, Staphylococcus aureus 2000 million, Staphylococcus albus 2000 million/cc. Vial 5 cc.
Use: Certain types of acne.
- ACNE VACCINES.** (Sherman) Acne bacillus 100 million, Colon bacillus 300 million Staphylococcus albus 1000 million, Staphylococcus aureus 1000 million/cc. Vial 12. cc., 20 cc.
Use: S.C. with increasing dosage, acute & chronic acne infections.
- ACNOMEL.** (Smith, Kline & French) Resorcinol 2%, sulfur 8%, hexachlorophene 0.25% w/bentonite, propylene glycol, alcohol & titanium dioxide. Cream. Tube 1½ oz., also available in cakes.
Use: Acne therapy.
- ACOBRYONNA.** (Lemmon) Aconite pow. 1.33 mg., byronia pow. 1.33 mg., belladonna pow. ext. 0.16 mg./Tab. Bot. 100s.
Use: (2 q. 3 h.) Antipyretic & diaphoretic.

- ACOBRYONNA COMP.** (Lemmon) Aconite pow. 4.66 mg., belladonna pow. ext. 0.24 mg., byronia pow. 1.33 mg., mercury iodide 1/100 gr., Sod. salicylate 1 gr./Tab. Bot. 100s.
Use: (1-3 q. 4 h.) Follicular tonsillitis.
- ACOGESIC** (Strassenburgh) Phenyltoloxamine dihydrogen citrate 25 mg., acetyl-p-aminophenol 300 mg., salicylamide 200 mg., dl-Amphetamine phosphate 2 mg., metoprine 0.5 mg./Cap. Bot. 100s
Use: (1-2 q. 4 h.) Analgesic
- ACOLAC BACTERIAL VACCINE.** (U. S. Std. Products) *Corynebacterium acnes* 1000 million, *Staphylococcus aureus* 500 million, *Staphylococcus albus* 500 million/cc. Vial 10 cc., 30 cc.
Use: Acute or chronic infections caused by acne.
- ACON, N.N.D.** 1959. (Endo) Water soluble vitamin A. Cap. 25,000 & 50,000 units./Cap.) Bot. 100s, 1000s. Soln. (25,000 units./cc.) Dropper Bot. 30 cc.
Use: Vitamin A therapy.
- ACONITE POWDER** W/Belladonna pow. ext., & byronia pow. See: *Acobryonna*, Tab. (Lemmon).
- W/Atropine sulf., aminophylline, veratum V. See: *Aminite*, Tab. (Mallard).
- ACR-ALLANTOMIDE.** (National Drug) Aminoacridine HCl 1:500, sulfanilamide 10%, allantoin 2%, in a water-miscible base. Tube 1 oz., Jar 1 lb.
Use: Topically, bacteriostatic.
- ACRIFLAVINE, N.F., A.D.R., N.N.R.** 1952. (Eufflavine, Gonacrine, Neutroflavine.) *Acriflavine neutral*; a mixture of 2,8-diamino-10-methylacridinium chloride & 2,8-diaminoacridine. (Lilly) Tab. 1½ gr. Bot. 100s, 1000s
Use: External antiseptic.
- ACRIFLAVINE, BRILLIANT-GREEN, METHYLOSOANILINE CHLORIDE MIXTURE, N.N.R.** 1953. See: *Dymixal*, Pow. (McNeil).
- ACRIFLAVINE HYDROCHLORIDE, N.F., A.D.R., N.N.R.** 1950. (Various Mfr.) Hydrochloride form of acriflavine, N.F. Acid *acriflavine*, acid *tryptaflavine*, *flavine*, *tryptaflavine*
National Aniline—Pow., Bot. (1, 5, 10, 25 & 50 Gm.). Tab. (1½ gr.) Bot. 50s & 100s.
Use: External bacteriostatic.
- ACRISAN EAR DROPS.** (Recsei) Aminoacridine propionate 0.35% & Tri-Histin 1% in propylene glycol. Ear drops, ½ oz.
Use: Anti-infective.
- ACRISAN THROAT SPRAY.** (Recsei) Aminoacridine ascorbate 0.2% w/benzyl alcohol 0.5%. Bot. 4 oz.
Use: Anti-infective
- ACRYLIC CATIONIC RESIN.**
See: *Carbo Resin* (Lilly)
Vacid (Fesler)
- A.C.T. (Tilden)** Pot. bromide 11 mg., ammonium bromide 11 mg., Sod. bromide 11 mg., aspirin 0.33 Gm., caffeine 33 mg./Tab. Bot. 100s, 1000s.
Use: (1-2 as needed) Analgesic w/sedative.
- ACTAMER.** (Monsanto Chem.) See: *Bithional*, U.S.P.
- ACTASAL PEDIATRIC DROPS.** (Purdue Frederick.) Choline salicylate 104.5 mg./0.6 ml. Bot. W/calibrated dropper—60 cc.
Use: Antipyretic, anti-inflammatory and analgesic for infants and children.
- ACTASE.** (Ortho) Fibrinolysin (human) 50,000 u./vial with water 10 cc.
Use: (Dissolve in 10 cc. of water, dilute to 250 cc. for I.V.) Intravascular blood clots.
- ACTH, N.N.D.** 1959. (Adrenocorticotrophic hormone. Adrenocorticotrop(h)in). See: *Corticotropin*, U.S.P.
- ACTH.** (National Drug) Colloidal solution of ACTH. *Injectosol* (20 U.S.P. units/cc.) 2 cc. & 10 cc. I.M. or I.V. drip.
- ACTH GEL, PURIFIED, N.N.D.** 1959. (National Drug) Corticotropin (40 u./cc.) Vial 1 cc., 5 cc.; (80 u./cc.) Vial 5 cc.
- ACTH (PITUITARY ADRENOCORTICOTROPIC HORMONE).** (Parke, Davis) Vial 25 units, 40 units (one unit is equivalent to 1 mg. international standard).
- ACTHAR, N.N.D.** 1959. (Armour) Corticotropin, U.S.P. Vial 10, 25, 40 mg. (Lyophilized)
- ACTHAR GEL.** (Armour & National Drug) Corticotropin, U.S.P. Disp. Cart. (20 mg./cc.) 1 cc., Vial 5 cc.; Disp. Cart. (40 mg./cc.) 1 cc., Vial 5 cc.
- ACTHAR GEL HIGHLY PURIFIED.** (Armour) H. P. acthar gel, a new, highly purified, repository ACTH preparation that may be given subcutaneously or intramuscularly. 20, 40 & 80 u. per cc., 5 cc. multiple dose vials. Supplied in a sterile 1 cc. B-d cartridge with B-2 disposable cartridge syringe, in potencies of 40 Armour u. per cc. See: *Corticotropin*, U.S.P.
Use: Stimulant of adrenal cortex, anti-inflammatory, antiallergic & useful in collagen diseases.
- ACTHAR SOLUTION.** (Armour) Corticotropin. 10 u./cc. Amp. 1 cc., Vial 10 cc.
Use: I.V., I.M., S.C., corticotropin therapy.
- ACTIDIL.** (Burroughs Wellcome) *Tripolidine* HCl. Tab. (2.5 mg.). Bot. 100s. Syr. (1.25 mg./5 cc.). Bot. 1 pt.
Use: Antihistamine.
- ACTIFED.** (Burroughs Wellcome) Tabs. Bot. 100s, Syr. Bot. 1 pt.