

11th edition

review of Medical Physiology


WILLIAM F. GANONG, MD

197-10
LA
11th edition

review of Medical Physiology

WILLIAM F. GANONG, MD

Jack and DeLoris Lange Professor of Physiology
Chairman, Department of Physiology
University of California
San Francisco, California



LANGE Medical Publications

Los Altos, California 94022

Preface

This book is designed to provide a concise summary of mammalian and, particularly, of human physiology which medical students and others can supplement with readings in current texts, monographs, and reviews. Pertinent aspects of general and comparative physiology are also included. Summaries of relevant anatomic considerations will be found in each section, but this book is written primarily for those who have some knowledge of anatomy, chemistry, and biochemistry.

Examples from clinical medicine are given where pertinent to illustrate physiologic points. Physicians desiring to use this book as a review will find short discussions of important symptoms produced by disordered function in several sections.

It has not been possible to be complete and concise without also being dogmatic. I believe, however, that the conclusions presented without a detailed discussion of the experimental data on which they are based are those supported by the bulk of the currently available evidence. Much of this evidence can be found in the papers cited in the credit lines of the illustrations. Further discussions of particular subjects and information on subjects not considered in detail in this book can be found in the references listed at the end of each section. Information about serial review publications that provide up-to-date discussions of various physiologic subjects is included in the note on general references in the appendix.

In the interest of brevity and clarity, I have in most instances omitted the names of the many investigators whose work made possible the view of physiology presented here. This is in no way intended to slight their contributions, but including their names and specific references to original papers would greatly increase the length of this book.

I am greatly indebted to many individuals who helped in the preparation of this book. Those whom I wish especially to thank for their help with the eleventh edition include Drs. William Silen, John Baxter, John Diana, and Lelio Orci. I am also indebted to Dr. Karen Sweadner, who gave permission to use the illustration on the front cover; to my wife, who labored long hours typing corrections; and to Annette Lowe, who drew many of the illustrations. I also wish to thank all the students and others who took the time to write to me offering helpful criticisms and suggestions. Such comments are always welcome, and I solicit additional corrections and criticisms, which may be addressed to me at the Department of Physiology, University of California, San Francisco, California 94143, USA. Many associates and friends provided unpublished illustrative materials, and numerous authors and publishers generously granted permission to reproduce illustrations from other books and journals.

The eleventh edition of this book appears on the 20th anniversary of the publication of the first edition, in 1963. I am pleased to be able to announce that during these 20 years, the following translations have been published: Portuguese (third edition), German (fourth edition), Italian (sixth edition), Spanish (eighth edition), Japanese (seventh edition), Polish, Czech, Chinese, Greek, Serbo-Croatian, French, Turkish, and Indonesian. A translation into Bahasa Malaysian is under way. The book has also appeared in various foreign English-language editions and has been recorded in English on tape for use by the blind. The tape recording is available from Recording for the Blind, Inc., 215 E. 58th Street, New York 10022. The German edition is also being recorded on tape.

William F. Ganong

Table of Contents

Preface	xi
---------	----

SECTION I. INTRODUCTION

1. Physiologic Principles	1
---------------------------	---

Functional Morphology of the Cell	1	The Capillary Wall	23
Body Fluid Compartments	13	Sodium & Potassium Distribution & Total Body	
Units for Measuring Concentration of Solutes	15	Osmolality	23
Composition of Body Fluids	16	pH & Buffers	25
Forces Producing Movement of Substances		Intercellular Communication	25
Between Compartments	16	Homeostasis	29
Cell Membrane & Resting Membrane Potentials	20	Aging	29

Section I References: 30

SECTION II. PHYSIOLOGY OF NERVE & MUSCLE CELLS

2. Excitable Tissue: Nerve	31
----------------------------	----

Nerve Cells	31	Nerve Fiber Types & Function	42
Electrical Phenomena in Nerve Cells	34	Nerve Growth Factor	44
Ionic Basis of Excitation & Conduction	39	Glia	44
Properties of Mixed Nerves	41		

3. Excitable Tissue: Muscle	45
-----------------------------	----

Skeletal Muscle	46	Cardiac Muscle (cont'd)	
Morphology	46	Mechanical Properties	55
Electrical Phenomena & Ionic Fluxes	47	Metabolism	58
Contractile Responses	48	Pacemaker Tissue	58
Energy Sources & Metabolism	52	Smooth Muscle	58
Properties of Muscles in the Intact Organism	53	Morphology	58
Cardiac Muscle	55	Visceral Smooth Muscle	59
Morphology	55	Multi-Unit Smooth Muscle	61
Electrical Properties	55		

4. Synaptic & Junctional Transmission	62
---------------------------------------	----

Synaptic Transmission	62	Neuromuscular Transmission	81
Functional Anatomy	62	The Myoneural Junction	81
Electrical Events at Synapses	64	Nerve Endings in Smooth & Cardiac Muscle	82
Inhibition & Facilitation at Synapses	67	Denervation Hypersensitivity	83
Chemical Transmission of Synaptic Activity	71		

5. Initiation of Impulses in Sense Organs	84
-------------------------------------------	----

Sense Organs & Receptors	84	Electrical & Ionic Events in Receptors	86
The Senses	84	"Coding" of Sensory Information	88

Section II References: 90

SECTION III. FUNCTIONS OF THE NERVOUS SYSTEM

6. Reflexes	91
The Reflex Arc 91 Monosynaptic Reflexes: The Stretch Reflex 91	Polysynaptic Reflexes: The Withdrawal Reflex 96 General Properties of Reflexes 98
7. Cutaneous, Deep, & Visceral Sensation	99
Pathways 99 Touch 101 Proprioception 102 Temperature 102 Pain 102	Differences Between Somatic & Visceral Sensory Mechanisms 104 Visceral Pain 105 Referral & Inhibition of Pain 106 Other Sensations 108
8. Vision	109
Anatomic Considerations 109 The Image-Forming Mechanism 113 The Photoreceptor Mechanism: Genesis of Action Potentials 116	Responses in the Visual Pathways & Cortex 120 Other Aspects of Visual Function 122 Color Vision 125 Eye Movements 126
9. Functions of the Ear	128
Anatomic Considerations 128 External & Middle Ear 128 Inner Ear 129 Hearing 132 Auditory Responses 132 Sound Transmission 133 Electrical Phenomena 135 Brain Mechanisms 136	Hearing (cont'd) Deafness 137 Vestibular Function 138 Responses to Rotational & Linear Acceleration 138 Orientation in Space 139 Effects of Labyrinthectomy 139
10. Smell & Taste	140
Smell 140 Receptors & Pathways 140 Physiology of Olfaction 141	Taste 142 Receptor Organs & Pathways 142 Physiology of Taste 143
11. The Reticular Activating System, Sleep, & the Electrical Activity of the Brain	146
The Reticular Formation & the Reticular Activating System 146 The Thalamus & the Cerebral Cortex 146 Evoked Cortical Potentials 147	The Electroencephalogram 148 Physiologic Basis of the EEG & Consciousness 149 Sleep 153 Modulation of Sensory Input 155
12. Control of Posture & Movement	156
Pyramidal System 156 Anatomy 156 Function 158 Extrapyramidal Mechanisms 159 Spinal Integration 160 Medullary Components 162 Midbrain Components 164	Extrapyramidal Mechanisms (cont'd) Cortical Components 164 Basal Ganglia 165 Cerebellum 167 Anatomic & Functional Organization 167 Physiology 170
13. Efferent Pathways to Visceral Effectors	174
Anatomic Organization of Autonomic Outflow 174 Chemical Transmission at Autonomic Junctions 177	Responses of Effector Organs to Autonomic Nerve Impulses 177

14. Central Regulation of Visceral Function 180

- | | |
|----------------------------------------------------|----------------------------------------------|
| Medulla Oblongata 180 | Hypothalamus (cont'd) |
| Hypothalamus 181 | Relation to Cyclic Phenomena 185 |
| Anatomic Considerations 181 | Hunger 185 |
| Hypothalamic Function 183 | Thirst 187 |
| Relation of Hypothalamus to Autonomic Function 183 | Control of Posterior Pituitary Secretion 188 |
| Relation to Sleep 184 | Control of Anterior Pituitary Secretion 192 |
| | Temperature Regulation 195 |

15. Neurophysiologic Basis of Instinctual Behavior & Emotions 200

- | | |
|-----------------------------|--------------------------------------------------------------------------------------|
| Anatomic Considerations 200 | Motivation 205 |
| Limbic Functions 201 | Brain Chemistry, Behavior, & Synaptic Transmission in the Central Nervous System 206 |
| Sexual Behavior 202 | |
| Fear & Rage 204 | |

**16. "Higher Functions of the Nervous System":
Conditioned Reflexes, Learning, & Related Phenomena 211**

- | | |
|--------------|--------------------------------|
| Learning 212 | Functions of the Neocortex 215 |
| Memory 214 | |

Section III References: 219

SECTION IV. ENDOCRINOLOGY & METABOLISM

17. Energy Balance, Metabolism, & Nutrition 221

- | | |
|-----------------------------|----------------------------------|
| Energy Metabolism 221 | Intermediary Metabolism (cont'd) |
| Metabolic Rate 221 | Protein Metabolism 234 |
| Energy Balance 225 | Fat Metabolism 240 |
| Intermediary Metabolism 225 | Nutrition 249 |
| General Considerations 225 | Essential Dietary Components 249 |
| Carbohydrate Metabolism 227 | |

18. The Thyroid Gland 254

- | | |
|------------------------------------------------|---------------------------------------------|
| Anatomic Considerations 254 | Mechanism of Action of Thyroid Hormones 262 |
| Formation & Secretion of Thyroid Hormones 255 | Regulation of Thyroid Secretion 262 |
| Transport & Metabolism of Thyroid Hormones 258 | Clinical Correlates 263 |
| Effects of Thyroid Hormones 260 | |

19. Endocrine Functions of the Pancreas & the Regulation of Carbohydrate Metabolism 268

- | | |
|-------------------------------------------------------------|-----------------------------------------------------|
| Islet Cell Structure 268 | Regulation of Insulin Secretion 279 |
| Structure, Biosynthesis, & Secretion of Insulin 269 | Glucagon 282 |
| Fate of Secreted Insulin 270 | Other Islet Cell Hormones 284 |
| Consequences of Insulin Deficiency & Actions of Insulin 272 | Endocrine Regulation of Carbohydrate Metabolism 285 |
| Insulin Excess 278 | Hypoglycemia & Diabetes Mellitus in Humans 286 |
| Mechanism of Action of Insulin 279 | |

20. The Adrenal Medulla & Adrenal Cortex 289

- | | |
|-------------------------------------------------------------------|------------------------------------------------------------------------------|
| Adrenal Morphology 289 | Adrenal Cortex (cont'd) |
| Adrenal Medulla 290 | Physiologic Effects of Glucocorticoids 300 |
| Structure & Function of Medullary Hormones 290 | Pharmacologic & Pathologic Effects of Glucocorticoids 302 |
| Regulation of Adrenal Medullary Secretion 293 | Regulation of Glucocorticoid Secretion 304 |
| Adrenal Cortex 293 | Effects of Mineralocorticoids 307 |
| Structure & Biosynthesis of Adrenocortical Hormones 293 | Regulation of Aldosterone Secretion 309 |
| Transport, Metabolism, & Excretion of Adrenocortical Hormones 297 | Role of Mineralocorticoids in the Regulation of Salt Balance 312 |
| Effects of Adrenal Androgens & Estrogens 299 | Summary of the Effects of Adrenocortical Hyper- & Hypofunction in Humans 312 |

21. Hormonal Control of Calcium Metabolism & the Physiology of Bone	313
Calcium & Phosphorus Metabolism 313	
Bone Physiology 314	
Vitamin D and the Hydroxychole- calciferols 316	
The Parathyroid Glands 318	
Calcitonin 320	
Effects of Other Hormones on Calcium Metabolism 322	
22. The Pituitary Gland	323
Morphology 324	
Intermediate Lobe Hormones 325	
Growth Hormone 326	
Physiology of Growth 331	
Pituitary Insufficiency 333	
Pituitary Hyperfunction in Humans 335	
23. The Gonads: Development & Function of the Reproductive System	336
Sex Differentiation & Development 336	
Chromosomal Sex 336	
Embryology of the Human Reproductive System 339	
Aberrant Sexual Differentiation 342	
Puberty 343	
Precocious & Delayed Puberty 344	
Menopause 345	
Pituitary Gonadotropins & Prolactin 345	
The Male Reproductive System 347	
Structure 347	
Gametogenesis & Ejaculation 347	
The Male Reproductive System (cont'd)	
Endocrine Function of the Testes 350	
Control of Testicular Function 353	
Abnormalities of Testicular Function 353	
The Female Reproductive System 354	
The Menstrual Cycle 354	
Ovarian Hormones 358	
Control of Ovarian Function 362	
Abnormalities of Ovarian Function 364	
Pregnancy 365	
Lactation 367	
24. Other Organs With Established or Suggested Endocrine Functions	370
The Endocrine Functions of the Kidneys: Renin & Erythropoietin 370	
Pineal 374	

Section IV References: 375

SECTION V. GASTROINTESTINAL FUNCTION

25. Digestion & Absorption	377
Carbohydrates 378	
Proteins & Nucleic Acids 380	
Lipids 381	
Absorption of Water & Electrolytes 383	
Absorption of Vitamins & Minerals 384	
26. Gastrointestinal Secretion & Motility	387
Gastrointestinal Hormones 388	
Mouth & Esophagus 392	
Stomach 394	
Regulation of Gastric Secretion & Motility 396	
Other Functions of the Stomach 398	
Exocrine Portion of the Pancreas 399	
Liver & Biliary System 401	
Small Intestine 406	
Colon 409	

Section V References: 413

SECTION VI. CIRCULATION

27. Circulating Body Fluids	414
The Circulatory System 414	
Blood 414	
Bone Marrow 414	
White Blood Cells 415	
Immune Mechanisms 417	
Platelets 421	
Blood (cont'd)	
Red Blood Cells 422	
Blood Types 426	
Plasma 429	
Hemostasis 430	
Lymph 433	

28. The Heart: Origin of the Heartbeat & the Electrical Activity of the Heart	434
Origin & Spread of Cardiac Excitation 434	Electrocardiographic Findings in Other Cardiac
The Electrocardiogram 436	& Systemic Diseases 447
Cardiac Arrhythmias 441	
29. The Heart as a Pump	451
Mechanical Events of the Cardiac Cycle 451	Cardiac Output 455
30. Dynamics of Blood & Lymph Flow	462
Anatomic Considerations 462	Lymphatic Circulation & Interstitial Fluid
Biophysical Considerations 464	Volume 473
Arterial & Arteriolar Circulation 469	Venous Circulation 475
Capillary Circulation 472	
31. Cardiovascular Regulatory Mechanisms	477
Local Regulatory Mechanisms 477	Systemic Regulatory Mechanisms 478
32. Circulation Through Special Regions	488
Cerebral Circulation 488	Cerebral Circulation (cont'd)
Anatomic Considerations 488	Brain Metabolism & Oxygen Requirements 496
Cerebrospinal Fluid 489	Coronary Circulation 497
The Blood-Brain Barrier 491	Splanchnic Circulation 500
Cerebral Blood Flow 493	Circulation of the Skin 501
Regulation of Cerebral Circulation 495	Placental & Fetal Circulation 503
33. Cardiovascular Homeostasis in Health & Disease	506
Compensations for Gravitational Effects 506	Fainting 514
Exercise 508	Heart Failure 515
Hemorrhage & Hemorrhagic Shock 510	Hypertension 516
Other Forms of Shock 513	

Section VI References: 518

SECTION VII. RESPIRATION

34. Pulmonary Function	520
Properties of Gases 520	Pulmonary Circulation 530
Mechanics of Respiration 521	Other Functions of the Respiratory
Gas Exchange in the Lung 529	System 531
35. Gas Transport Between the Lungs & the Tissues	533
Oxygen Transport 533	Carbon Dioxide Transport 538
Buffers in Blood 536	
36. Regulation of Respiration	540
Neural Control of Breathing 540	Chemical Control of Breathing 542
Regulation of Respiratory Center Activity 542	Nonchemical Influences on Respiration 546
37. Respiratory Adjustments in Health & Disease	549
Effects of Exercise 549	Oxygen Treatment 557
Hypoxia 551	Hypercapnia & Hypocapnia 558
Hypoxic Hypoxia 552	Effects of Increased Barometric Pressure 559
Other Forms of Hypoxia 556	Artificial Respiration 560

Section VII References: 562

SECTION VIII. FORMATION & EXCRETION OF URINE

38. Renal Function 563

Functional Anatomy 563	Acidification of the Urine & Bicarbonate Excretion 580
Renal Circulation 566	Regulation of Na ⁺ & Cl ⁻ Excretion 582
Glomerular Filtration 568	Regulation of K ⁺ Excretion 584
Tubular Function 570	Diuretics 584
Water Excretion 575	Effects of Disordered Renal Function 585

39. Micturition 587

Filling of the Bladder 587	Abnormalities of Micturition 588
Emptying of the Bladder 587	

40. Regulation of Extracellular Fluid Composition & Volume 590

Defense of Tonicity 590	Defense of Specific Ionic Composition 591
Defense of Volume 590	Defense of H ⁺ Concentration 591

Section VIII References: 597

Appendix 599

General References 599	Standard Respiratory Symbols 605
Normal Values & the Statistical Evaluation of Data 599	Equivalents of Metric, United States, & English Measures 605
Appendix References 601	Greek Alphabet 606
Abbreviations & Symbols Commonly Used in Physiology 602	

Index 607

Tables

Atomic Weights	Inside Front Cover
Ranges of Normal Values in Human Whole Blood, Plasma, or Serum	Inside Back Cover

Section I. Introduction

Physiologic Principles | 1

In unicellular organisms, all vital processes occur in a single cell. As the evolution of multicellular organisms has progressed, various cell groups have taken over particular functions. In humans and other vertebrate animals, the specialized cell groups include a gastrointestinal system to digest and absorb food; a respiratory system to take up O_2 and eliminate CO_2 ; a urinary system to remove wastes; a cardiovascular system to distribute food, O_2 , and the products of metabolism; a reproductive system to perpetuate the species; and nervous and endocrine systems to coordinate and integrate the functions of the other systems. This book is concerned with the way these systems

function and the way each contributes to the functions of the body as a whole.

FUNCTIONAL MORPHOLOGY OF THE CELL

Revolutionary advances in the understanding of cell structure and function have been made through the use of electron microscopy, x-ray diffraction, and the other techniques of modern cellular and molecular biology. The specialization of the cells in the various organs is very great, and no cell can be called "typi-




Figure 1-1. Ultrastructure of the common cell organelles and inclusions. The endoplasmic reticulum illustrated here is the granular type, with ribosomes attached to it. Some cells also contain tubes of membrane without ribosomes (agranular endoplasmic reticulum). The pores in the nuclear envelope are closed by a thin, homogeneous membrane. (Modified and reproduced, with permission, from Bloom W, Fawcett WD: *A Textbook of Histology*, 10th ed. Saunders, 1975.)

cal'' of all cells in the body. However, a number of structures (**organelles**) are common to most cells. These structures are shown in Fig 1-1.

Cell Membrane

The membrane that surrounds the cell is a remarkable structure. It is not only semipermeable, allowing some substances to pass through it and excluding others, but its permeability can be varied. It is generally referred to as the **plasma membrane**. The nucleus is also surrounded by a membrane, and the organelles are surrounded by or made up of membrane.

Although the chemical structure of membranes and their properties vary considerably from one location to another, they have certain common features. They are generally about 7.5 nm (75 Angstrom units) thick. They are made up primarily of protein and lipids. The chemistry of proteins and lipids is discussed in Chapter 17. The major lipids are phospholipids such as phosphatidylcholine and phosphatidylethanolamine. The shape of the phospholipid molecule is roughly that of a clothespin (Fig 1-2). The head end of the molecule contains the phosphate portion, is




Figure 1-2. Biologic membrane. **Top:** Model of membrane structure. The phospholipid molecules each have 2 fatty acid chains (wavy lines) attached to a phosphate head (open circle). The proteins, indicated by the long lines, are partly in the α -helical configuration and partly folded, with the charged ends (+ and - signs) on the exterior or interior sides of the membrane. **Bottom:** Three-dimensional view of membrane. Proteins are shown as irregular shaded globules. (Reproduced, with permission, from Singer SJ, Nicolson GL: The fluid mosaic model of the structure of cell membranes. *Science* 1972;175:720. Copyright 1972 by the American Association for the Advancement of Science.)

positively charged, and is quite soluble in water (polar, **hydrophilic**). The tails are quite insoluble (nonpolar, **hydrophobic**). In the membrane, the hydrophilic ends are exposed to the aqueous environment that bathes the exterior of the cells and the aqueous cytoplasm; the hydrophobic ends meet in the water-poor interior of the membrane. However, there is a degree of asymmetry in the distribution of lipid in the membrane; in human red cells, for example, there is more phosphatidylethanolamine and phosphatidylserine in the inner lamella and more lecithin and sphingomyelin in the outer lamella. The significance of this asymmetry is unknown. In **prokaryotes** (cells like bacteria in which there is no nucleus), phospholipids are generally the only membrane lipids, but in **eukaryotes** (cells containing nuclei), cell membranes also contain cholesterol (in animals) or other steroids (in plants). The cholesterol/phospholipid ratio in the membrane is inversely proportionate to the fluidity of the membrane. Variations in the ratio lead to abnormalities of cell function. In healthy animals, the ratio is maintained at a nearly constant level by a variety of regulatory mechanisms.

There are many different proteins embedded in the membrane. They exist as separate globular units and stud the inside and outside of the membrane in a random array (Fig 1-2). Some are located in the inner surface of the membrane; some are located on the outer surface; and some extend through the membrane (**through and through proteins**). In general, the uncharged, hydrophobic portions of the protein molecules are located in the interior of the membrane and the charged, hydrophilic portions are located on the surfaces. Some of the proteins contain lipids (lipoproteins) and some contain carbohydrates (glycoproteins). There are 5 types of proteins in the membrane. In addition to structural proteins, there are proteins that function as pumps, actively transporting ions across the membrane. Other proteins function as passive channels for ions that can be opened or closed by changes in the conformation of the protein. A fourth group of proteins functions as receptors that bind neurotransmitters and hormones, initiating physiologic changes inside the cell. A fifth group functions as enzymes, catalyzing reactions at the surfaces of the membrane.

The protein structure—and particularly the enzyme content—of biologic membranes varies not only from cell to cell but also within the same cell. For example, there are different enzymes embedded in cell membranes than in mitochondrial membranes; in epithelial cells, the enzymes in the cell membrane on the mucosal surface differ from those in the cell membrane on the lateral margins of the cells. The membranes are dynamic structures, and their constituents are being constantly renewed at different rates. In addition, some proteins move laterally in the membrane. For example, receptors move in the membrane and aggregate at sites of endocytosis (see below). There is evidence that the lateral movement of components in the membrane is not random but is controlled

by intracellular mechanisms that probably involve microfilaments and microtubules (see below).

Underlying most cells is a thin, fuzzy layer plus some fibrils that collectively make up the **basement membrane** or, more properly, the **basal lamina**. The material that makes up the basal lamina has been shown to be made up of a collagen derivative plus 2 glycoproteins.

Intercellular Connections

Two types of junctions form between the cells that make up tissues: junctions that fasten the cells to one another and to surrounding tissues, and junctions that permit transfer of ions and other molecules from one cell to another. The junctions that tie cells together and endow tissues with strength and stability are **tight junctions** and **desmosomes** (Fig 1–3). Desmosomes are subdivided into **belt desmosomes**, **spot desmosomes**, and **hemidesmosomes**. The junctions by which molecules are transferred are **gap junctions**.

Tight junctions characteristically surround the apical margins of the cells in epithelia such as the intestinal mucosa, the walls of the renal tubules, and the choroid plexus. They are made up of ridges—half from one cell and half from the other—that adhere so strongly at cell junctions that they almost obliterate the space between the cells. The ridges were thought to be made up of protein, but recent evidence suggests that instead they may be long tubes made up of membrane lipids. In addition to tying the cells together, tight junctions form a barrier to the movement of ions and

other solutes from one side of the epithelium to the other. Some epithelia are more “leaky” to solutes than others, but the tight junctions in all of them play an important role in their physiologic function.

Belt desmosomes and spot desmosomes are characterized by apposed thickenings of the membranes of 2 adjacent cells. Attached to the thickened area in each cell are cytoplasmic fibrils, some running parallel to the membrane and others radiating away from it. Between the 2 membrane thickenings, there is filamentous material in the intercellular space. Belt desmosomes, also known as **intermediate zones**, form bands that link epithelial cells just below and parallel to their tight junctions. The intracellular filaments associated with these junctions contain the protein **actin** (see below and Chapter 3) and are capable of contraction. However, the function of their contractions is unknown. The spot desmosomes are small, buttonlike points of attachment between the membranes of adjacent cells that are not limited to their apical regions. They have been likened to spot welds or rivets that hold tissues together. At spot desmosomes, the intercellular space is actually widened to 30 nm from its normal width of about 25 nm, and the filamentous material between the thickenings of the membranes of the 2 cells forms a dense central line called the **central stratum**. The cytoplasmic fibers associated with spot desmosomes are thicker than those associated with belt desmosomes and are not contractile. **Hemidesmosomes** anchor epithelial cells to the underlying connective tissue.

At gap junctions, the intercellular space narrows from 25 nm to 3 nm, and hexagonal arrays of protein units called **connexons** in the membrane of each cell are lined up with one another. Each connexon is made up of 6 subunits surrounding a channel that, when lined up with the channel in the corresponding connexon in the adjacent cell, permits substances to pass between the cells without entering the extracellular fluid. The diameter of the channel is normally about 2 nm, which permits the passage of sugars, amino acids, and other solutes with molecular weights up to about 800. Gap junctions thus permit the rapid propagation of electrical activity from cell to cell (see Chapter 4) and the exchange of various chemical messengers. However, the diameter of each channel is regulated by intracellular Ca^{2+} , and an increase in Ca^{2+} causes the subunits to slide together, closing the channel. Consequently, an increase in Ca^{2+} in an individual cell uncouples it from its neighbors.

Mitochondria

Although their morphology varies somewhat from cell to cell, each mitochondrion (Figs 1–1, 1–4) is in essence a sausage-shaped structure. It is made up of an outer membrane and an inner membrane that is folded to form shelves (**cristae**). The mitochondria are the power-generating units of the cell and are most plentiful and best developed in parts of cells where energy-requiring processes take place. The chemical reactions occurring in them are discussed in detail in




Figure 1–3. Intercellular junctions in the mucosa of the small intestine. The various types of desmosomes are not shown in detail. (Reproduced, with permission, from Weinstein RS, McNutt SN: Cell junctions. *N Engl J Med* 1972;286:521.)




Figure 1-4. Cutaway drawing of a mitochondrion, showing the inner and outer membranes. The inner membrane is folded, forming shelves (cristae). The coiled structures represent possible arrangements of the mitochondrial DNA. Mitochondrial branching of the type shown here is not present in all cells. (Reproduced, with permission, from Nass MMK: Mitochondrial DNA: Advances, problems and goals. *Science* 1969;165:25. Copyright 1969 by the American Association for the Advancement of Science.)

Chapter 17. The outer membrane of each mitochondrion is studded with the enzymes concerned with biologic oxidations, providing raw materials for the reactions occurring inside the mitochondrion. The interior of the mitochondrion contains the enzymes concerned with the citric acid cycle and the respiratory chain enzymes by which the 2-carbon fragments produced by metabolism are burned to CO_2 and water (see Chapter 17). In this process, electrons are transferred along the respiratory enzyme chain. Associated with the electron transfer is **oxidative phosphorylation**, the synthesis of the high-energy phosphate compound **adenosine triphosphate (ATP)**. This ubiquitous molecule is the principal energy source for energy-requiring actions in animals and plants. The inner membrane of the mitochondrion appears to be made up of repeating units, each of which contains a basepiece, a stalk, and a spherical headpiece. The basepieces contain the enzymes of the electron transfer chain, and the stalks and headpieces contain adenosine triphosphatase and other enzymes concerned with the synthesis and metabolism of ATP.

Mitochondrial DNA is discussed below.

Lysosomes

In the cytoplasm of the cell, there are large, somewhat irregular structures surrounded by membrane that may contain fragments of other cell structures. These organelles are the **lysosomes**. Some of the granules of the granulocytic white blood cells are lysosomes. Each lysosome contains a variety of enzymes (Table 1-1) that would cause the destruction of most cellular components if the enzymes were not separated from the rest of the cell by the membrane of the lysosome.

The lysosomes function as a form of digestive system for the cell. Exogenous substances such as bacteria that become engulfed by the cell end up in membrane-lined vacuoles. A vacuole of this type (**phagocytic vacuole**) may merge with a lysosome,

Table 1-1. Some of the enzymes found in lysosomes and the cell components that are their substrates.

Enzyme	Substrate
Ribonuclease	RNA
Deoxyribonuclease	DNA
Phosphatase	Phosphate esters
Glycosidases	Complex carbohydrates: glycosides and polysaccharides
Arylsulfatases	Sulfate esters
Collagenase	Proteins
Cathepsins	Proteins

permitting the contents of the vacuole and the lysosome to mix within a common membrane. Some of the products of the "digestion" of the engulfed material are absorbed through the walls of the vacuole, and the remnants are dumped from the cell (exocytosis; see below). The lysosomes also engulf worn-out components of the cell in which they are located, forming **autophagic vacuoles**. When a cell dies, lysosomal enzymes cause autolysis of the remnants. In vitamin A intoxication and certain other conditions, lysosomal enzymes are released to the exterior of the cell, with resultant breakdown of intercellular material. There is evidence that in gout, phagocytes ingest uric acid crystals, and that such ingestion triggers the release of lysosomal enzymes that contribute to the inflammatory response in the joints. When one of the lysosomal enzymes is congenitally absent, the lysosomes become engorged with one of the materials they normally degrade. This eventually disrupts the cells that contain the defective lysosomes and leads to one of the **lysosomal storage diseases**. More than 25 such diseases have been described. They are generally rare, but they include such widely known disorders as Tay-Sachs disease.

Microfilaments & Microtubules

Most, if not all, eukaryotic cells contain **microfilaments**, long solid fibers 4-6 nm in diameter, and **microtubules**, long hollow structures with 5-nm walls surrounding a cavity 15 nm in diameter (Fig 1-5).

Microfilaments are made up of **actin**, the protein that by its interaction with myosin brings about contraction of muscle (see Chapter 3). Actin is present in all types of cells so far examined. Myosin is difficult to observe in cells other than muscle and is not arranged with actin in orderly arrays, but it appears to be present nonetheless. Actin microfilaments reach to the tips of the microvilli on the epithelial cells of the intestinal mucosa and bring about their contraction probably via interaction with myosin filaments located at their bases. Microfilaments are also found in association with belt desmosomes (see above), in bundles under the plasma membrane, and scattered in a seemingly random fashion in the cytoplasm.

Microtubules are made up of 2 globular protein





Figure 1-5. *Left:* Electron micrograph of the cytoplasm of a fibroblast, showing microfilaments (MF) and microtubules (MT). (Reproduced, with permission, from Junqueira LC, Carneiro J: *Basic Histology*, 4th ed. Lange, 1983.) *Right:* Distribution of microtubules in fibroblasts. The cells are treated with a fluorescently labeled antibody to tubulin, making microtubules visible as the light-colored structures. (Reproduced, with permission, from Connolly J et al: Immunofluorescent staining of cytoplasmic and spindle microtubules in mouse fibroblasts with antibody to τ protein. *Proc Natl Acad Sci USA* 1977;74:2437.)

subunits, α and β tubulin. The subunits unite to form dimers (Fig 1-6), and the dimers aggregate to form long tubes made up of stacked rings, with each ring usually containing 13 subunits. The tubules also contain other proteins that facilitate their formation. The assembly of microtubules is facilitated by warmth and various other factors, and disassembly is facilitated by cold and other factors. Both processes occur simultaneously in vitro. Assembly is prevented by colchicine and vinblastine.

Microtubules have been called the skeleton of the cell, but because of their constant assembly and disassembly, they are a very dynamic skeleton. They are associated with cell movement, although the actual motion is probably brought about by microfilaments. They appear to provide the tracks along which secretory granules move to the cell membrane. They play key roles in nerve fiber outgrowth, axoplasmic transport (see Chapter 2), maintenance of cell shape, structure and function of cilia, and cell division.

Centrioles & Cilia

In the cytoplasm of most cells there are 2 short cylinders called **centrioles**. The centrioles are located near the nucleus, and they are arranged so that they are at right angles to each other. Microtubules in groups of 3 run longitudinally in the walls of the centriole (Fig 1-1). There are 9 of these triplets spaced at regular intervals around the circumference. **Cilia**, the hairlike motile processes that in multicellular animals extend from various types of epithelial cells, also have an array of 9 tubular structures in their walls, but they have in addition a pair of microtubules in the center, and there are 2 rather than 3 microtubules in each of the




Figure 1-6. Assembly and disassembly of a microtubule by aggregation and disaggregation of dimers made up of α and β tubulin. (From Sloboda RD: The role of microtubules in cell structure and cell division. *Am Sci* 1980;68:290. Reprinted by permission of American Scientist, journal of Sigma Xi, the Scientific Research Society.)




Figure 1-7. Tetranucleotide portion of one strand of deoxyribonucleic acid. The strand is composed of adenine (A), thymine (T), cytosine (C), and guanine (G) deoxynucleotides.




Figure 1-8. Diagram of the Watson and Crick model, slightly modified, of the double helical structure of DNA. On the left, the hydrogen bonding between the bases of the 2 nucleotide chains is shown. A, adenine; T, thymine; C, cytosine; G, guanine; S, deoxyribose; P, phosphate. (Reproduced, with permission, from Harper HA: *Review of Physiological Chemistry*, 15th ed. Lange, 1975.)

9 circumferential structures. The **basal granule**, on the other hand, the structure to which a cilium is anchored, has 9 circumferential triplets, like a centriole.

The centrioles are concerned with the movement of the chromosomes during cell division. They duplicate themselves at the start of mitosis, and the pairs move apart to form the poles of the mitotic spindle. In multinucleate cells, there is a pair of centrioles near each nucleus. The mitotic spindle is made up of microtubules and microfilaments.

Nucleus & Related Structures

A nucleus is present in all animal cells that divide. If a cell is cut in half, the anucleate portion eventually dies without dividing. The nucleus is made up in large part of the **chromosomes**, the structures in the nucleus that carry a complete blueprint for all the heritable species and individual characteristics of the animal. Each chromosome is made up of a giant molecule of **deoxyribonucleic acid (DNA)** that in multicellular organisms is virtually covered with proteins. The molecule has been likened to a string of beads; the beads are balls of histone proteins with DNA wrapped around them, and the string is more DNA with other proteins attached to it. This complex of DNA and proteins is called **chromatin**. During cell division, the pairs of chromosomes become visible, but between cell divisions, only clumps of chromatin can be discerned in the nucleus. Evidence is beginning to accumulate that the structure of chromatin is important in regulating the function of the DNA. The ultimate units of heredity are the **genes** on the chromosomes (see below), and each gene is a portion of the DNA molecule.

During normal cell division by **mitosis**, the chromosomes duplicate themselves and then divide in such a way that each daughter cell receives a full complement (**diploid number**) of chromosomes. During their final maturation, germ cells undergo a division in which half the chromosomes go to each daughter cell (see Chapter 23). This reduction division (**meiosis**) is actually a 2-stage process; but the important consideration is that, as a result of it, mature sperms and ova contain half the normal number (the **haploid number**) of chromosomes. When a sperm and ovum unite, the resultant cell (**zygote**) has a full (diploid) complement of chromosomes, one-half from the female parent and one-half from the male.

The nucleus of most cells contains a **nucleolus** (Fig 1-1), a patchwork of granules rich in **ribonucleic acid (RNA)**. In some cells, the nucleus contains several of these structures. Nucleoli are most prominent and numerous in growing cells. The DNA in the nucleus serves as a template for the synthesis of RNA, which then moves to the cytoplasm, where it regulates the synthesis of proteins by the cell. The nucleolus is probably the site where the RNA found in the ribosomes is synthesized.

The nucleus is surrounded by a **nuclear membrane (envelope)** (Fig 1-1). This membrane is double, and the spaces between the 2 folds are called

perinuclear cisterns. The nuclear membrane is apparently quite permeable, since it permits passage of molecules as large as RNA from the nucleus to the cytoplasm. There are areas of discontinuity in the nuclear membrane, but these "pores" are closed by a thin, homogeneous membrane.

Endoplasmic Reticulum

The endoplasmic reticulum is a complex series of tubules in the cytoplasm of the cell (Fig 1-1). The tubule walls are made up of membrane. In **granular** endoplasmic reticulum, granules called **ribosomes** are attached to the cytoplasmic side of the membrane, whereas in **agranular** endoplasmic reticulum the granules are absent. Free ribosomes are also found in the cytoplasm. The ribosomes are about 15 nm in diameter. Each is made up of a large and a small subunit called, on the basis of their rates of sedimentation in the centrifuge, the 50S and the 30S subunits, respectively. The ribosomes contain about 65% RNA and 35% protein. They are the sites of protein synthesis. The ribosomes attached to the endoplasmic reticulum synthesize proteins such as hormones that are secreted by the cell and proteins that are segregated in lysosomes. The polypeptide chains that form these proteins are extruded into the endoplasmic reticulum. The free ribosomes synthesize cytoplasmic proteins such as hemoglobin (see Chapter 27).

The agranular endoplasmic reticulum is the site of steroid synthesis in steroid-secreting cells and the site of detoxification processes in other cells. As the sarcoplasmic reticulum (see Chapter 3), it plays an important role in skeletal and cardiac muscle.

Golgi Complex

The Golgi complex is a collection of membranous tubules and vesicles. It is usually located near the nucleus, and is particularly prominent in actively secreting gland cells. Hormones and enzymes are stored in protein-secreting cells as membrane-enclosed **secretion granules**, and these granules are produced in the Golgi complex. Thus, the Golgi complex "packages" proteins. The complex is also the site of formation of lysosomes (see below), and it adds certain carbohydrates to proteins to form glycoproteins.

Structure & Function of DNA & RNA

DNA is found in bacteria, in the nuclei of eukaryotic cells, and in mitochondria. It is made up of 2 extremely long nucleotide chains containing adenine (A), guanine (G), thymine (T), and cytosine (C) (Fig 1-7). The chemistry of these purine and pyrimidine bases and of nucleotides is discussed in Chapter 17. The chains are bound together by hydrogen bonding between the bases, adenine bonding to thymine and guanine to cytosine. The resultant double helical structure of the molecule is shown in Fig 1-8. An indication of the complexity of the molecule is the fact that the DNA in the human genome (the total genetic message) is made up of 2.5×10^9 base pairs.

DNA is the component of the chromosomes that

carries the "genetic message," the blueprint for all the heritable characteristics of the cell and its descendants. Each chromosome is in effect a segment of the DNA double helix. The genetic message is coded by the sequence of purine and pyrimidine bases in the nucleotide chains. The text of the message is the order in which the amino acids are lined up in the proteins manufactured by the cell. The message is transferred to the sites of protein synthesis in the cytoplasm by RNA. The proteins formed include all the enzymes, and these in turn control the metabolism of the cell. There is probably one gene for each enzyme or enzyme subunit in the body; indeed, a gene has been defined as the amount of information necessary to specify a single peptide molecule. Mutations occur when the base sequence in the DNA is altered by x-rays, cosmic rays, or other mutagenic agents.

At the time of each somatic cell division (mitosis), the 2 DNA chains separate, each serving as a template for the synthesis of a new complementary chain. DNA polymerase catalyzes this reaction. One of the double helices thus formed goes to one daughter cell and one to the other, so the amount of DNA in each daughter cell is the same as that in the parent cell. In the case of germ cells, reduction division (meiosis) takes place during maturation. The net result is that one DNA chain goes to each mature germ cell; consequently, each mature germ cell contains half the amount of chromosomal material found in somatic cells. Therefore, when a sperm unites with an ovum, the resulting zygote has the full complement of DNA, half of which came from the father and half from the mother. The chromosomal events that occur at the time of fertilization are discussed in detail in Chapter 23.




Figure 1-9. The functions of DNA and RNA in terms of the formation of the individual and the continuity of species. Germ plasm (DNA) passes from generation to generation via the germ cells. From this "germ line" the mortal soma (body) buds off each generation. Formation of the individual depends upon transcription of the genetic message by RNA, thus determining the enzymes formed and consequently controlling the metabolism of the cell. (Reproduced, with permission, from Haggis GH et al: *Introduction to Molecular Biology*. Wiley, 1964.)