

HANDBOOK OF EXPERIMENTAL IMMUNOLOGY
IN FOUR VOLUMES

Volume 1: Immunochemistry

EDITED BY

D. M. WEIR MD, FRCP

CO-EDITORS

L. A. Herzenberg PhD Caroline Blackwell PhD

Leonore A. Herzenberg DSc

FOURTH EDITION

HANDBOOK OF EXPERIMENTAL IMMUNOLOGY
IN FOUR VOLUMES

Volume 1: Immunochemistry

EDITED BY

D. M. WEIR MD, FRCP

*Professor of Microbial Immunology,
University of Edinburgh, Scotland*

CO-EDITORS

L. A. Herzenberg PhD

*Professor of Genetics,
Stanford University, USA*

Caroline Blackwell PhD

*Lecturer, Department of Bacteriology,
University of Edinburgh, Scotland*

Leonore A. Herzenberg DSc

*Senior Research Associate, Department of Genetics,
Stanford University, USA*

FOURTH EDITION

BLACKWELL SCIENTIFIC PUBLICATIONS

OXFORD LONDON EDINBURGH

BOSTON PALO ALTO MELBOURNE

© 1967, 1973, 1978, 1986 by
Blackwell Scientific Publications
Editorial offices:
Osney Mead, Oxford, OX2 0EL
8 John Street, London, WC1N 2ES
23 Ainslie Place, Edinburgh, EH3 6AJ
52 Beacon Street, Boston
Massachusetts 02108, USA
667 Lytton Avenue, Palo Alto
California 94301, USA
107 Barry Street, Carlton
Victoria 3053, Australia

All rights reserved. No part of this
publication may be reproduced, stored in a
retrieval system, or transmitted, in any form
or by any means, electronic, mechanical,
photocopying, recording or otherwise
without the prior permission of the copyright
owner

First published 1967
Second edition 1973
Third edition 1978
Reprinted 1979
Fourth Edition 1986

Printed in Great Britain
at the Alden Press, Oxford

DISTRIBUTORS
USA

Blackwell Scientific Publications Inc
PO Box 50009, Palo Alto
California 94303

Blackwell Mosby Book Distributors
11830 Westline Industrial Drive
St Louis, Missouri 63141

Canada

The C. V. Mosby Company
5240 Finch Avenue East
Scarborough, Ontario

Australia

Blackwell Scientific Publications
(Australia) Pty Ltd
107 Barry Street
Carlton, Victoria 3053

British Library

Cataloguing in Publication Data

Handbook of experimental immunology.—4th ed. 1.
Immunology—Laboratory manuals I. Weir,
D.M. II. Herzenberg, L.A. III. Blackwell,
C. IV. Herzenberg, Leonore A. 599.02'9'028
QR183

ISBN 0-632-01499-7
ISBN 0-632-00975-6 v. 1
ISBN 0-632-01378-8 v. 2
ISBN 0-632-01379-6 v. 3
ISBN 0-632-01381-8 v. 4

Contributors

- C. A. Allen, *Imperial Cancer Research Fund, London, UK*
- G. A. Andres, *Department of Pathology, State University of New York at Buffalo, NY, USA*
- J. D. Ansell, *Department of Zoology, University of Edinburgh, Edinburgh, UK*
- P. K. C. Austwick, *Department of Allergy and Clinical Immunology, Cardiothoracic Institute, London, UK*
- J. Austyn, *Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- T. Bachi, *Institute for Immunology and Virology, University of Zurich, Zurich, Switzerland*
- J. A. Badwey, *Department of Biological Chemistry, Harvard Medical School, Boston, Mass, USA*
- D. Baltimore, *Whitehead Institute for Biomedical Research, Cambridge, Mass, USA*
- M. J. Banda, *Laboratory of Radiobiology and Environmental Health, University of California, San Francisco, Ca, USA*
- A. N. Barclay, *MRC Cellular Immunology Unit, Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- G. R. Barclay, *Blood Transfusion Centre, Royal Infirmary, Edinburgh, UK*
- B. Benacerraf, *Department of Pathology, Harvard Medical School, Boston, Mass, USA*
- P. Beverley, *ICRF Human Tumour Immunology Group, School of Medicine, University College London, London, UK*
- D. E. Bidwell, *Nuffield Laboratories of Comparative Medicine, Zoological Society of London, London, UK*
- C. Caroline Blackwell, *Department of Bacteriology, University of Edinburgh, Edinburgh, UK*
- A. E. Bolton, *Department of Biological Science, Sheffield City Polytechnic, Sheffield, UK*
- J. Braun, *Department of Pathology, University of California, Los Angeles, Ca, USA*
- P. H. Brodeur, *Department of Pathology, Tufts University, Boston, Mass, USA*
- C. J. Burrell, *Division of Virology, Institute of Medical and Veterinary Science, Adelaide, Australia*
- E. C. Butcher, *Department of Pathology, Stanford University Medical School, Stanford, Ca, USA*
- G. C. Butcher, *Institute of Animal Physiology, Agricultural Research Council, Babraham, Cambridge, UK*
- H. Cantor, *Department of Pathology, Harvard Medical School, Boston, Mass, USA*
- P. Chandler, *Transplantation Biology Section, Clinical Research Centre, Harrow, UK*
- M. W. Clark, *Department of Biochemistry and Microbiology, University of Victoria, BC, Canada*
- J. E. Coligan, *National Institute of Allergy and Infectious Diseases, Bethesda, Md, USA*
- Suzanne Cory, *The Walter and Eliza Hall Institute of Medical Research, Royal Melbourne Hospital, Victoria, Australia*
- O. Cromwell, *Department of Allergy and Clinical Immunology, Cardiothoracic Institute, London, UK*
- M. R. Daha, *Department of Nephrology, University Hospital, Leiden, The Netherlands*
- M. M. Davis, *Department of Medical Microbiology, Stanford University Medical School, Stanford, Ca, USA*
- R. Dildrop, *Institute of Genetics, University of Cologne, Cologne, Federal Republic of Germany*
- G. J. Dougherty, *Imperial Cancer Research Fund, London, UK*
- S. Dreborg, *Allergy and Diagnostics Division, Pharmacia AB, Uppsala, Sweden*
- D. W. Dresser, *Division of Immunology, National Institute for Medical Research, Mill Hill, London, UK*
- S. R. Durham, *Department of Allergy and Clinical Immunology, Cardiothoracic Institute, London, UK*
- P. J. Edelson, *Department of Pediatrics, New York Hospital, Cornell Medical Centre, New York, NY, USA*
- A. B. Edmundson, *Department of Biology, University of Utah, Salt Lake City, Utah, USA*
- R. Einärsson, *Biochemistry Division, Pharmacia AB, University of Uppsala, Uppsala, Sweden*
- E. L. Elson, *Department of Biological Chemistry, Washington University Medical Centre, St. Louis, Mo, USA*
- R. A. Elton, *Medical Computing and Statistics Group, University of Edinburgh, Edinburgh, UK*
- Katherine R. Ely, *Department of Biology, University of Utah, Salt Lake City, Utah, USA*
- E. G. Engleman, *Department of Medicine, Stanford University, School of Medicine, Stanford, Ca, USA*
- R. A. B. Ezekowitz, *Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- C. G. Fathman, *Department of Medicine, Stanford University School of Medicine, Stanford, Ca, USA*

- A. S. Fauci, *Laboratory of Immunoregulation, National Institute of Health, Bethesda, Md, USA*
- S. Felder, *Department of Biological Chemistry, Washington University School of Medicine, St. Louis, Mo, USA*
- Lorraine Flaherty, *Centre for Laboratories and Research, New York State Department of Health, Albany, NY, USA*
- The late W. L. Ford, *Department of Immunology, University of Manchester, Manchester, UK*
- J. Forman, *Department of Microbiology, University of Texas Southwestern Medical School, Dallas, Texas, USA*
- Sara Fuchs, *Department of Clinical Immunology, The Weizmann Institute of Science, Rehovot, Israel*
- H. H. Fudenberg, *Department of Basic and Clinical Immunology and Microbiology, Medical University of South Carolina, Charleston, SC, USA*
- W. Gerhard, *Wistar Institute of Anatomy and Biology, Philadelphia, Pa, USA*
- Elizabeth J. Glass, *Animal Breeding Research Organisation, Edinburgh, UK*
- W. Godfrey, *Department of Microbiology, University of California, San Francisco, Ca, USA*
- J. W. Goding, *Department of Pathology and Immunology, Monash Medical School, Victoria, Australia*
- S. Gordon, *Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- D. R. Green, *Department of Immunology, University of Alberta, Edmonton, Canada*
- Margaret C. Green, *The Jackson Laboratory, Bar Harbor, Maine, USA*
- N. M. Gough, *Ludwig Institute for Cancer Research, Royal Melbourne Hospital, Victoria, Australia*
- G. A. Gutman, *Department of Microbiology and Molecular Genetics, University of California, Irvine, Ca, USA*
- J. Guyden, *Department of Microbiology and Immunology, University of California, San Francisco, Ca, USA*
- S. Habu, *Department of Cell Biology, Tokai University, Japan*
- S. Hakomori, *Division of Biochemical Oncology, Fred Hutchinson Cancer Research Center, Seattle, Wash, USA*
- E. Handman, *Laboratory of Immunoparasitology, The Walter and Eliza Hall Institute of Medical Research, Royal Melbourne Hospital, Victoria, Australia*
- R. R. Hardy, *Institute for Molecular and Cellular Biology, Osaka University, Osaka, Japan*
- R. A. Harrison, *Mechanisms in Tumour Immunity Unit, MRC Centre, Cambridge, UK*
- K. Hayakawa, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*
- W. J. Herbert, *Animal Services Unit, University of Dundee, Dundee, UK*
- L. A. Herzenberg, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*
- Leonore A. Herzenberg, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*
- Eadie Heyderman, *Department of Histopathology, St. Thomas's Hospital Medical School, London, UK*
- S. Hirsch, *Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- Nancy M. Hogg, *Imperial Cancer Research Fund, London, UK*
- E. J. Holborow, *Bone and Joint Research Unit, London Hospital Medical College, London, UK*
- L. Hood, *Division of Biology, California Institute of Technology, Pasadena, Ca, USA*
- J. C. Howard, *Institute of Animal Physiology, Agricultural Research Council, Babraham, Cambridge, UK*
- Maureen Howard, *National Institute of Allergy and Infectious Diseases, Bethesda, Md, USA*
- K. C. Hsu, *Department of Microbiology, College of Physicians and Surgeons of New York, NY, USA*
- D. A. Hume, *Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- S. V. Hunt, *MRC Cellular Immunology Unit, Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- W. M. Hunter, *Celltec Ltd, Slough, Berkshire, UK*
- G. M. Iverson, *Dept of Pathology, Howard Hughes Medical Institute, Yale University, New Haven, Conn, USA*
- M. K. Jenkins, *Department of Microbiology-Immunology, Northwestern University, Chicago, Ill, USA*
- G. D. Johnson, *Department of Immunology, University of Birmingham, Birmingham, UK*
- M. J. Johnston, *Department of Genetics, Stanford University Medical School, Stanford, Ca, USA*
- E. A. Kabat, *Department of Microbiology, College of Physicians and Surgeons of Columbia University, New York, NY, USA*
- M. E. Kamarck, *Department of Biology, Yale University, New Haven, Conn, USA*
- R. Kannagi, *Division of Biochemical Oncology, Fred Hutchinson Cancer, Seattle, Wash, USA*
- J. Kappler, *Department of Medicine, National Jewish Hospital and Research Center, Denver, Co, USA*

- M. J. Karnovsky, *Department of Biological Chemistry, Harvard Medical School, Boston, Mass, USA*
- M. L. Karnovsky, *Department of Pathology, Harvard Medical School, Boston, Mass, USA*
- R. A. Kautz, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*
- Paula Kavathas, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*
- A. B. Kay, *Department of Allergy and Clinical Immunology, Cardiothoracic Institute, London, UK*
- B. S. Kim, *Department of Microbiology-Immunology, Northwestern University, Chicago, Ill, USA*
- T. J. Kindt, *National Institute of Allergy and Infectious Disease, Bethesda, Md, USA*
- T. J. Kipps, *Scripps Clinic and Research Foundation, La Jolla, Ca, USA*
- M. Klein, *Department of Immunology, Toronto Western Hospital, Ontario, Canada*
- R. D. Kornberg, *Department of Cell Biology, Stanford University Medical School, Stanford, Ca, USA*
- P. J. Lachmann, *Mechanisms in Tumour Immunity Unit, MRC Centre, Cambridge, UK*
- H. C. Lane, *Laboratory of Immunoregulation, National Institute of Health, Bethesda, Maryland, USA*
- L. L. Lanier, *Becton Dickinson Monoclonal Center Inc, Mountain View, Ca, USA*
- P. C. J. Leijh, *Department of Infectious Diseases, University Hospital, Leiden, The Netherlands*
- Sheelagh Lloyd, *Department of Clinical Veterinary Medicine, University of Cambridge, Cambridge, UK*
- Joan L. Longbottom, *Department of Clinical Immunology, Cardiothoracic Institute, London, UK*
- Eva Lotta Larsen, *Department of Immunobiology, Karolinska Institute, Stockholm, Sweden*
- W. H. R. Lumsden, *19a Merchiston Crescent, Edinburgh, UK*
- W. Lutz, *Medical Computing and Statistics Group, University of Edinburgh, Edinburgh, UK*
- W. H. McBride, *Department of Radiation Oncology, University of California, Los Angeles, USA*
- M. C. McCann, *Blood Transfusion Centre, Royal Infirmary, Edinburgh, UK*
- J. A. McIntyre, *Department of Obstetrics and Gynaecology, Southern Illinois School of Medicine, Springfield, Ill, USA*
- Judith A. MacKay, *Department of Allergy and Clinical Immunology, Cardiothoracic Institute, London, UK*
- Rose A. Mage, *Laboratory of Immunology, National Institute of Allergy and Infectious Diseases, Bethesda, Md, USA*
- O. Mäkelä, *Department of Bacteriology and Immunology, University of Helsinki, Helsinki, Finland*
- Phillipa Marrack, *Department of Medicine, National Jewish Hospital and Research Center, Denver, Co, USA*
- Olivia Martinez, *Department of Microbiology and Immunology, University of California, Berkeley, Ca, USA*
- D. W. Mason, *MRC Cellular Immunology Unit, Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- R. Melvold, *Dept of Microbiology and Immunology, Northwestern University Medical School, Chicago, Ill, USA*
- T. G. Merrett, *RAST Allergy Unit, Benenden Chest Hospital, Cranbrook, Kent, UK*
- H. S. Micklem, *Department of Zoology, University of Edinburgh, Edinburgh, UK*
- R. G. Miller, *Ontario Cancer Institute, University of Toronto, Toronto, Ontario, Canada*
- S. D. Miller, *Department of Microbiology Immunology, The Medical and Dental Schools, Ill, Northwestern University, Chicago, USA*
- C. Milstein, *Laboratory of Molecular Biology, Medical Research Council The Medical School, Cambridge, UK*
- G. F. Mitchell, *Laboratory of Immunoparasitology, The Walter and Eliza Hall Institute of Medical Research, Royal Melbourne Hospital, Victoria, Australia*
- W. A. Moore, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*
- Sherie L. Morrison, *Department of Microbiology, Columbia University College of Physicians and Surgeons, New York, NY, USA*
- D. B. Murphy, *Department of Pathology, Yale University, New Haven, Conn, USA*
- R. M. Nakamura, *Department of Pathology, Scripps Clinic and Research Foundation, La Jolla, Ca, USA*
- Y. Nakao, *Department of Medicine, University of Kobe, Japan*
- C. J. Newby, *Department of Genetics, Stanford University Medical School, Stanford, Ca, USA*
- L. A. Nilsson, *Department of Medical Microbiology, University of Gothenburg, Gothenburg, Sweden*
- G. J. V. Nossal, *Walter and Eliza Hall Institute of Medical Research, Royal Melbourne Hospital, Victoria, Australia*
- V. Oi, *Becton Dickinson Monoclonal Center, Mountain View, Ca, USA*
- K. Okumura, *Department of Immunology, Juntendo University, Tokyo, Japan*

- Ö. Ouchterlony, *Department of Medical Microbiology, University of Gothenburg, Gothenburg, Sweden*
- Z. Ovary, *Department of Pathology, New York University Medical Center, New York, NY, USA*
- W. Page Faulk, *Medi-Search AG, Meiringen, Switzerland*
- P. Parham, *Department of Cell Biology, Stanford University School of Medicine, Stanford, Ca, USA*
- D. R. Parks, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*
- M. Parson, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*
- T. W. Pearson, *Department of Biochemistry and Microbiology, University of Victoria, BC, Canada*
- N. O. Petersen, *Department of Chemistry, University of Western Ontario, London, Ontario, Canada*
- M. Potter, *Laboratory of Genetics, National Cancer Institute, Bethesda, Md, USA*
- I. R. Poxtton, *Department of Bacteriology, University Medical School, Edinburgh, UK*
- A. Radbruch, *Institute of Genetics, University of Cologne, Cologne, Federal Republic of Germany*
- P. Ralph, *Department of Cell Biology, Cetus Corporation, Emeryville, Ca, USA*
- K. Rajewsky, *Institute of Genetics, University of Cologne, Cologne, Federal Republic of Germany*
- U. Ramstedt, *Department of Immunology, Karolinska Institute, Stockholm, Sweden*
- Elizabeth Raveche, *National Institute of Health, Bethesda, Md, USA*
- L. F. Reichardt, *Department of Neurology, University of California, San Francisco, Ca, USA*
- J. Reidler, *Department of Cell Biology, Stanford University School of Medicine, Stanford, Ca, USA*
- R. Riblet, *Department of Immunology, Medical Biology Institute, La Jolla, Ca, USA*
- J. M. Robinson, *Department of Pathology, Harvard Medical School, Boston, Mass., USA*
- F. S. Rosen, *Division of Immunology, Children's Hospital Medical Center, Boston, Mass, USA*
- R. V. Rouse, *Department of Pathology, Stanford University School of Medicine, Stanford, Ca, USA*
- F. A. Ruddle, *Department of Biology, Yale University, New Haven, Conn, USA*
- T. Sasazuki, *Department of Genetics, Medical Institute of Bioregulation, Kyushu University, Fukuoka, Japan*
- M. S. Schanfield, *Genetic Testing Institute, Atlanta, Georgia, USA*
- Beatrice C. Seegal, *Department of Microbiology, College of Physicians and Surgeons of Columbia University, New York, NY, USA*
- M. Sela, *Department of Chemical Immunology, The Weizmann Institute of Science, Rehovot, Israel*
- I. J. T. Seppälä, *Department of Bacteriology and Immunology, University of Helsinki, Helsinki, Finland*
- E. E. Sercarz, *Department of Microbiology, University of California, Los Angeles, Ca, USA*
- Eva Severinson, *Department of Immunobiology, Karolinska Institute, Stockholm, Sweden*
- N. Shastri, *Department of Microbiology, University of California, Los Angeles, Ca, USA*
- J. F. L. Shaw, *Department of Surgery, University of Cambridge, Cambridge, UK*
- S. C. Silverstein, *Laboratory of Cellular Physiology and Immunology, The Rockefeller University, New York, NY, USA*
- R. J. Shaw, *Department of Allergy and Clinical Immunology, Cardiothoracic Institute, London, UK*
- K. Simonovitch, *Department of Immunology, Toronto Western Hospital, Ontario, Canada*
- Elizabeth Simpson, *Transplantation Biology Section, Clinical Research Centre, Harrow, UK*
- L. Smith, *Division of Biology, California Institute of Technology, Pasadena, Ca, USA*
- E. J. L. Soulsby, *Department of Clinical Veterinary Medicine, University of Cambridge, UK*
- D. M. Spalding, *Division of Clinical Immunology and Rheumatology, University of Alabama in Birmingham, Birmingham, Alabama, USA*
- T. A. Springer, *Dana Farber Cancer Institute, Harvard Medical School, Boston, Mass, USA*
- D. R. Stanworth, *Department of Immunology, University of Birmingham, Birmingham, UK*
- Phyllis M. Starkey, *Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- A. D. Steinberg, *National Institute of Health, Bethesda, Md, USA*
- R. M. Steinman, *Laboratory of Cellular Physiology, and Immunology, The Rockefeller University, New York, NY, USA*
- M. Steinmetz, *Basel Institute for Immunology, Basel, Switzerland*
- M. W. Steward, *Department of Medical Microbiology, London School of Hygiene and Tropical Medicine, London, UK*
- C. C. Stewart, *Experimental Pathology Group, Los*

- Alamos National Laboratory, Los Alamos, New Mexico, USA*
- J. Stewart, *Department of Bacteriology, University of Edinburgh, Edinburgh, UK*
- E. J. Stott, *Institute for Research on Animal Diseases, Newbury, Berkshire, UK*
- T. Tada, *Department of Immunology, University of Tokyo, Tokyo, Japan*
- N. Takemura, *Medical Research Institute, Tokyo Medical and Dental University, Tokyo, Japan*
- R. Takemura, *Laboratory of Radiobiology and Environmental Health, University of California, San Francisco, Ca, USA*
- M. Taniguchi, *Department of Immunology, Chiba University, Chiba, Japan*
- Glenys Thomson, *Department of Genetics, University of California, Berkeley, Ca, USA*
- P. W. Tucker, *Department of Microbiology, University of Texas, Southwestern Medical School, Dallas, Texas, USA*
- T. Tokuhisa, *Department of Immunology, Chiba University, Chiba, Japan*
- M. W. Turner, *Department of Immunology, Institute of Child Health, London, UK*
- D. A. J. Tyrrell, *MRC Common Cold Unit, Harvard Hospital, Salisbury, Wiltshire, UK*
- E. R. Unanue, *Department of Pathology, Harvard Medical School, Boston, Mass, USA*
- J. Unkeless, *The Rockefeller University, New York, NY, USA*
- R. J. Urbaniak, *Aberdeen and North East of Scotland Blood Transfusion Service, Aberdeen, UK*
- E. E. Uzgiris, *Department of Cell Biology, Stanford University School of Medicine, Stanford, Ca, USA*
- Karen L. Valentino, *Department of Physiology, University of California School of Medicine, San Francisco, Ca, USA*
- R. van Furth, *Department of Infectious Diseases, University Hospital, Leiden, The Netherlands*
- E. van Loghem, *The Netherlands Red Cross Blood Transfusion Service, Amsterdam, The Netherlands*
- W. C. van Voorhis, *Laboratory of Cellular Physiology, The Rockefeller University, New York, NY, USA*
- Theda L. van Zwet, *Department of Infectious Diseases, University Hospital, Leiden, The Netherlands*
- A. Voller, *Nuffield Laboratories of Comparative Medicine, Zoological Society of London, London, UK*
- C. Waltenbaugh, *Department of Microbiology-Immunology, Northwestern University, Chicago, Ill, USA*
- R. A. Warnke, *Department of Pathology, Stanford University Medical School, Stanford, Ca, USA*
- D. M. Weir, *Department of Bacteriology, University of Edinburgh, Edinburgh, UK*
- Zena Werb, *Laboratory of Radiobiology and Environmental Health, University of California, San Francisco, Ca, USA*
- G. Whalen, *Laboratory of Immunoregulation, National Institute of Health, Md, USA*
- A. G. White, *Department of Surgery, Faculty of Medicine, Kuwait University, Kuwait*
- D. J. G. White, *Department of Surgery, University of Cambridge, Cambridge, UK*
- H. Wigzell, *Department of Immunology, Karolinska Institute, Stockholm, Sweden*
- P. C. Wilkinson, *Department of Bacteriology and Immunology, University of Glasgow, Glasgow, UK*
- A. F. Williams, *MRC Cellular Immunology Unit, Sir William Dunn School of Pathology, University of Oxford, Oxford, UK*
- F. P. Winstanley, *Armed Services University, Washington, DC, USA*
- Janet Winter, *Department of Physiology, University of California, San Francisco, Ca, USA*
- L. Wofsy, *Department of Microbiology and Immunology, University of California, Berkeley, Ca, USA*
- S. D. Wright, *Laboratory of Cellular Physiology and Immunology, The Rockefeller University, New York, NY, USA*
- M. Zauderer, *Department of Microbiology and Oncology, University of Rochester, Rochester, NY, USA*
- D. Zelaschi, *Department of Genetics, Stanford University School of Medicine, Stanford, Ca, USA*

Preface

The pace of progress in immunology has not slackened since the last edition of this handbook. The subject now draws heavily on molecular biology and genetics and this has necessitated the inclusion of an additional volume on **Genetics and Molecular Immunology**. The explosion in the development of hybridoma technology and cell culture, since the last edition, can be seen from the many chapters in each volume that employ monoclonal reagents and cell lines. Some idea of the expansion of the field can be gained from the **Cellular Immunology** volume where contributions on phagocytes and lymphocytes now occupy 30 chapters compared to 12 in the previous edition. A new section on immunoregulation contains 14 chapters and there are now 6 chapters devoted to mammalian cell membrane antigens in the **Immunochemistry** volume.

It is now no longer possible for one editor to keep in touch with the enormous expansion in this field, and I am much indebted to my co-editors Len and Leonore Herzenberg who have joined me in the task of co-opting research workers in the wide range of disciplines now contributing to the field of immu-

nology. I am particularly grateful to my wife Dr Caroline Blackwell for her help with the massive editing task.

Amongst the many new features of this edition is the provision of overviews for many of the sections. I am most grateful to our contributors in the methodology sections for their efforts to achieve a consistent style of presentation of the procedures, and I hope that this will help in the accessibility of the descriptive material. A work of this size inevitably takes a number of years to put together but considerable effort has gone into introducing up to date material into the chapters. This has been achieved by enabling and encouraging contributors to introduce new material and references during the proof stages of their chapters.

I wish to thank Hilary Flénley for her careful and thorough index, and Nigel Palmer and his staff at Blackwell Scientific Publications Edinburgh office without whom production of the new edition would have been impossible. Per Saugman has as always, maintained a benevolent paternal interest in the project.

D.M.W.

Contents

List of contributors xi

Preface xvii

VOLUME 1 IMMUNOCHEMISTRY

Antigens

- 1 Overview: Antigens
M. SELA
- 2 Preparation of synthetic antigens
M. SELA AND SARA FUCHS
- 3 Haptens and carriers
O. MÄKELÄ AND I. J. T. SEPPÄLÄ
- 4 Isolation and identification of bacterial antigens
I. R. POXTON AND C. CAROLINE BLACKWELL
- 5 Preparation of viral antigens
C. J. BURRELL
- 6 Antigens of parasites
T. W. PEARSON AND M. W. CLARK
- 7 Fungal antigens
JOAN L. LONGBOTTOM AND P. K. C. AUSTWICK
- 8 Immunization of experimental animals
D. W. DRESSER
- 9 Carbohydrate antigens in higher animals
S. HAKOMORI AND R. KANNAGI
- 10 The chemistry and standardization of allergens
S. DREBORG, R. EINARSSON, JOAN L. LONGBOTTOM

Immunoglobulins:

Purification and characterization

- 11 Overview: The quest for antibody homogeneity: the *sine qua non* for structural and genetic insights into antibody complementarity
E. A. KABAT
- 12 Immunochemical analysis of human and rabbit immunoglobulins and their subunits
D. R. STANWORTH AND M. W. TURNER

- 13 Purification and characterization of monoclonal antibodies

R. R. HARDY

- 14 Preparation and purification of active fragments from mouse monoclonal antibodies

P. PARHAM

- 15 Determination of the three-dimensional structures of immunoglobulins

A. B. EDMUNDSON AND KATHERYN R. ELY

- 16 Immunoabsorbents

SARA FUCHS AND M. SELA

- 17 Two-dimensional crystals of immunoglobulins
J. REIDLER, E. E. UZGIRIS AND R. D. KORNBERG

- 18 Sources of myeloma proteins, and M-components

M. PÖTTER

Mammalian cell membrane antigens

- 19 Overview: Mammalian cell surface antigens

R. D. KORNBERG

- 20 Membrane and secretory immunoglobulins: structure, biosynthesis and assembly

J. W. GODING

- 21 Class I MHC antigens of mouse and humans

J. E. COLIGAN AND T. J. KINDT

- 22 Glycoprotein antigens of the lymphocyte surface and their purification by antibody affinity chromatography

A. F. WILLIAMS AND A. N. BARCLAY

- 23 Immunofluorescence methods to study cell surface and cytoplasmic proteins, and their dynamics

E. R. UNANUE AND J. BRAUN

- 24 Measurement of lateral diffusion by fluorescence photobleaching recovery

N. O. PETERSEN, S. FELDER AND E. L. ELSON

Antibody interaction with soluble and cellular antigens

- 25 Overview: Introduction to methods used to

- study the affinity and kinetics of antibody-antigen reactions
M. W. STEWARD
 - 26 Radioimmunoassays and related methods
A. E. BOLTON AND W. M. HUNTER
 - 27 Enzyme immunoassays: heterogenous and homogeneous systems
R. M. NAKAMURA, A. VOLLER AND D. E. BIDWELL
 - 28 Preparation and use of fluorochrome conjugates
G. D. JOHNSON AND E. J. HOLBOROW
 - 29 Flow cytometry and fluorescence activated cell sorting (FACS)
D. R. PARKS, L. L. LANIER AND L. A. HERZENBERG
 - 30 Data analysis in flow cytometry
W. A. MOORE AND R. A. KAUTZ
 - 31 Purification and coupling of fluorescent proteins for use in flow cytometry
R. R. HARDY
 - 32 Immunodiffusion and immunoelectrophoresis
Ö. OUCHTERLONY AND L.-Å. NILSSON
 - 33 Passive cutaneous anaphylaxis
Z. OVARY
 - 34 Solid-phase radioimmune assays
C. J. NEWBY, K. HAYAKAWA AND LEONORE A. HERZENBERG
 - 35 Immunological techniques for the identification of antigens and antibodies by electron microscopy
G. A. ANDRES, K. C. HSU AND BEATRICE C. SEGAL
 - 36 Affinity targeting and fusion of vesicles to cells
W. GODFREY, J. GUYDEN AND L. WOFSY
 - 37 Immunotoxins
OLIVIA MARTINEZ AND L. WOFSY
 - 38 Kinetics of antibody reaction and the analysis of cell surface antigens
D. W. MASON AND A. F. WILLIAMS
 - Complement
 - 39 Complement technology
R. A. HARRISON AND P. J. LACHMANN
 - 40 Complement fixation by monoclonal antigen-antibody complexes
R. R. HARDY
- VOLUME 2
CELLULAR IMMUNOLOGY**
- Phagocytes**
- 41 Overview: The function of receptors in phagocytosis
S. D. WRIGHT AND S. C. SILVERSTEIN
 - 42 Overview: The mononuclear phagocyte system
R. VAN FURTH
 - 43 Plasma membrane markers to study differentiation, activation and localization of murine macrophages. Ag F4/80 and the mannosyl, fucosyl receptor
S. GORDON, PHYLLIS M. STARKEY, D. HUME, R. A. B. EZEKOWITZ, S. HIRSCH AND J. AUSTYN
 - 44 Methods for studying the ontogeny of mononuclear phagocytes
C. C. STEWART
 - 45 Macrophage cell lines
P. RALPH
 - 46 *In vitro* determination of phagocytosis and intracellular killing by polymorphonuclear and mononuclear phagocytes
P. C. J. LEIJM, R. VAN FURTH AND THEDA L. VAN ZWET
 - 47 Secreted proteins of resting and activated macrophages
ZENA WERB, M. J. BANDA, R. TAKEMURA AND S. GORDON
 - 48 Macrophage membrane receptors
J. STEWART, ELIZABETH J. GLASS, D. M. WEIR AND M. R. DAHA
 - 49 Dendritic cells
R. M. STERNMAN, W. C. VAN VOORHIS AND D. M. SPALDING

- 50 Reduction and excitation of oxygen by phagocytic leukocytes: biochemical and cytochemical techniques
J. A. BADWEY, J. M. ROBINSON, M. J. KARNOVSKY AND M. L. KARNOVSKY
 - 51 Locomotion and chemotaxis of leukocytes
P. C. WILKINSON
 - 52 Assays for phagocyte ecto-enzymes
P. J. EDELSON
- The lymphoid system**
- 53 Overview: Lymphocytes and their relations
H. S. MICKLEM
 - 54 Physical methods for separation of lymphocyte subpopulations
R. G. MILLER
 - 55 Preparative immunoselection of lymphocyte populations
S. V. HUNT
 - 56 Genetic markers for following cell populations
J. D. ANSELL AND H. S. MICKLEM
 - 57 Following cellular traffic: methods of labelling lymphocytes and other cells to trace their migration *in vivo*
E. C. BUTCHER AND W. L. FORD
 - 58 Human leukocyte subpopulations
P. BEVERLEY
 - 59 Lymphokines
J. KAPPLER AND PHILLIPA MARRACK
 - 60 Natural killer cells
H. WIGZELL AND U. RAMSTEDT
 - 61 Genetics and cell distributions of mouse cell surface alloantigens
LORRAINE FLAHERTY AND J. FORMAN
- Lymphocyte responses**
- 62 Overview: Lymphocyte responses
G. J. V. NOSSAL
 - 63 Lymphocyte responses to polyclonal B and T cell activators
EVA SEVERINSON AND EVA-LOTTA LARSSON
- 64 Assays for immunoglobulin-secreting cells
D. W. DRESSER
 - 65 Limiting dilution analysis of effector cells and their precursors *in vitro*
M. ZAUDERER
 - 66 *In vitro* evaluation of human lymphocyte function
H. C. LANE, GAIL WHALEN AND A. S. FAUCI
 - 67 Assay for *in vivo* adoptive immune response
G. M. IVERSON
 - 68 Analysis of cytotoxic T cell responses
ELIZABETH SIMPSON AND P. CHANDLER
 - 69 T cell lines and hybrids in mouse and man
C. G. FATHMAN AND E. G. ENGLEMAN
 - 70 B cell growth factors
MAUREEN HOWARD
- Immunoregulation**
- 71 Overview: Helper and suppressor T cells
K. OKUMARA AND T. TADA
 - 72 Overview: Ir genes
B. BENACERRAF
 - 73 Overview: Idiotypic regulation
K. RAJEWSKY
 - 74 Overview: Epitope-specific regulation
LEONORE A. HERZENBERG
 - 75 Overview: T cell clones
H. CANTOR
 - 76 Subtractive cDNA hybridization and the T-cell receptor genes
M. M. DAVIS
 - 77 Detection of suppressor cells and suppressor factors for delayed-type hypersensitivity responses
S. D. MILLER AND M. K. JENKINS
 - 78 Detection of suppressor cells and suppressor factors for antibody responses
C. WALTEBAUGH AND B. S. KIM
 - 79 Contrasuppressor T cells: a practical guide to

the identification of contrasuppressive effects in immunoregulatory systems
D. R. GREEN

80 Antigen-specific suppressor molecules produced by T cells
M. TANIGUCHI AND T. TOKUHISA

81 Immunosuppressive agents
D. J. G. WHITE AND J. F. L. SHAW

82 Studies of autoimmune diseases
ELIZABETH S. RAVECHE AND A. D. STEINBERG

83 Primary immunodeficiencies: definitions and diagnosis
F. S. ROSEN

84 Studying immune regulation with protein and peptide antigens
N. SHASTRI AND E. E. SERCARZ

VOLUME 3 GENETICS AND MOLECULAR IMMUNOLOGY

Molecular approaches to immunology

85. Overview: Introductory comments on molecular immunology
D. BALTIMORE

86 Construction and screening of recombinant DNA libraries in bacteriophage lambda
P. W. TUCKER

87 The major histocompatibility complex of the mouse
L. SMITH, M. STEINMETZ AND L. HOOD

88 The murine immunoglobulin heavy chain constant region gene locus
N. M. GOUGH AND SUZANNE CORY

89 The Igh-V genes of the mouse
R. RIBLET AND P. H. BRODEUR

90 Classification of mouse V_H sequences
RENATE DILDROP

91 Transfection for lymphocyte cell surface antigens
PAULA KAVATHAS AND L. A. HERZENBERG

92 Lymphoid cell gene transfer
SHERIE L. MORRISON AND V. T. OI

Immunoglobulin genetics

93 Overview: Allotypes
H. H. FUDENBERG

94 Human immunoglobulin allotypes
M. S. SCHANFIELD AND ERNA VAN LOGHEN

95 New methods for human allotyping
D. ZBLASCHI, M. J. JOHNSTON, C. J. NEWBY AND LEONORE A. HERZENBERG

96 Gm and disease
Y. NAKAO AND T. SASAZUKI

97 Mouse immunoglobulin allotypes
MARILYN PARSON, LEONORE A. HERZENBERG, A. M. STALL AND L. A. HERZENBERG

98 Rat immunoglobulin allotypes
G. A. GUTMAN

99 Rabbit immunoglobulin allotypes
ROSE G. MAGE

Genetics of the major histocompatibility complex

100 Overview: The murine MHC
D. B. MURPHY

101 The MHC of the laboratory rat, *Rattus norvegicus*
G. W. BUTCHER AND J. C. HOWARD

102 Human HLA genetics and disease associations
GLENYS THOMSON

103 Common assays for identifying immune response genes
C. WALLENBAUGH AND S. D. MILLER

Resources

104 The mouse linkage map
MARGARET C. GREEN

105 Sonatic cell genetics and the human gene map
M. E. KAMARCK AND F. H. RUDDLE

- 106 Inbred, congenic, recombinant-inbred and mutant mouse strains
R. W. MELVOLD

**VOLUME 4
APPLICATIONS OF IMMUNOLOGICAL
METHODS IN BIOMEDICAL
SCIENCES**

Monoclonal antibodies

- 107 Overview: Monoclonal antibodies
C. MILSTEIN
- 108 Schemata for the production of monoclonal antibody-producing hybridomas
T. J. KIPPS AND L. A. HERZENBERG
- 109 Hybridoma immunoglobulin isotype switch variant selection with the fluorescence activated cell sorter
T. J. KIPPS AND L. A. HERZENBERG
- 110 Isotype switch variants
A. RADBRUCH
- 111 Growing hybridoma and producing monoclonal antibodies *in vivo*
K. OKUMURA AND S. HABU

**Applications:
Monoclonal Antibodies**

- 112 Neurobiology
JANET WINTER, KAREN L. VALENTINO AND L. F. REICHARDT
- 113 Monoclonal antibodies in the study of parasites and host-parasite relationships
E. HANDMAN AND G. F. MITCHELL
- 114 Application of monoclonal antibodies in bacteriology
C. CAROLINE BLACKWELL AND F. P. WINSTANLEY
- 115 Applications of monoclonal antibodies in virology
W. GERHARDT AND T. BACHI
- 116 Special applications of tissue section immunologic staining in the characterization of monoclonal

al antibodies and in the study of normal and neoplastic tissues
R. V. ROUSE AND R. A. WARNKE

- 117 Monoclonal antibodies directed to carbohydrate antigens
R. KANNAGI AND S. HAKOMORI
- 118 Macrophages
J. C. UNKELESS AND T. A. SPRINGER

**Applications:
Immunological methods**

- 119 Application of immunological techniques in bacteriology
F. P. WINSTANLEY AND C. CAROLINE BLACKWELL
- 120 Application of immunological methods in virology
E. J. STOTT AND D. A. J. TYRRELL
- 121 Applications of immunological methods in mycology
JOAN L. LONGBOTTOM
- 122 Applications of immunological methods in protozoology
E. H. R. LUMSDEN
- 123 Applications of immunological methods in helminthology
W. J. L. SOULSBY AND SHEELAGH LLOYD
- 124 Immunological aspects of human pregnancy
J. A. MACINTYRE AND W. PAGE FAULK
- 125 Applications of immunological techniques to the study of the tumour-host relationship
G. J. DOUGHERTY, C. A. ALLEN AND NANCY M. HOGG

**Additional clinical applications of
immunological methods**

- 126 Tests of immune function
S. J. URBANIAK, M. C. MCCANN, A. G. WHITE, G. R. BARCLAY AND A. B. KAY
- 127 Provocation tests and measurements of media-

tors from mast cells and basophils in asthma and allergic rhinitis

O. CROMWELL, S. R. DURHAM, R. J. SHAW,
JUDITH A. MACKAY AND A. B. KAY

- 128 Methods for detecting immune complexes in biological fluids
M. H. KLEIN AND K. SIMINOVITCH

- 129 Immunohistochemistry in pathology
EADIE HEYDERMAN, I. STRUDLEY AND T. C. RICHARDSON

- 130 Quantification of IgE and IgG4 both as total immunoglobulins and as allerge-specific antibodies
T. G. MERRETT

General methods for immunologic studies

- 131 Statistical aspects of planning and design of immunological experiments
R. A. ELTON AND W. H. MCBRIDE

- 132 Guidelines to statistical analysis
W. LUTZ

- 133 Laboratory animal techniques for immunology
W. J. HERBERT, F. KRISTENSEN, with RUTH M. AITKEN, M. B. ESLAMI, ANNE FERGUSON, KATHLEEN G. GRAY, W. J. PENHALE

Index to Volume 1 xix

Chapter 1

Overview: Antigens

M. SELA

Definitions, 1.1

Molecules as antigens, 1.2

Immunoglobulins, 1.3

Synthetic antigens, 1.4

Complex antigens, 1.5

Definitions

The expression '*antigen*' may have two meanings. On the one hand, we mean by 'antigen' the substance capable of triggering an immune response, but on the other hand, we use the same word for a substance capable of reacting immunospecifically with an antibody, even though sometimes it is not able to trigger a response. To distinguish between these two different notions, we have been using the expressions *immunogenicity* and *antigenic specificity*. By immunogenicity we understand the capacity of a substance to provoke an immune response, whatever its specificity, and such a substance is an *immunogen*. A chemical or physical change in a molecule that will result in an increased immune response will increase its immunogenicity, even though it may or may not have changed its antigenic specificity. It is the *antigenic determinant* (also defined as epitope) which is also responsible for the specificity.

Putting it in other words, immunogenicity is a property of substances that can induce a detectable immune response (humoral, cellular, or, most commonly, both) when introduced into an animal or human being. Such substances are therefore generally called *immunogens* or *antigens*. Ultimately, the ability to mount an immune response depends upon the interplay between the chemistry of the antigen and the physiological state of the host. Thus immunogenicity is operationally dependent on the experimental conditions of the system, including parameters such as the antigen, the mode of immunization, the organism being immunized and its genetic background, as well as the sensitivity of methods used to detect a response.

It is worth stressing that we are using the word *antigen* sometimes for a molecule, sometimes for a virus or a bacterium, and sometimes for an organ or a tissue, but the antibodies—or the sensitized cells—have a *combining site* (also defined as paratope), with all its distinctive features, of a more or less similar size and similar cavity. These combining sites are not complementary to a complete bacterium or a complete

heart, but they are always complementary to a unique antigenic determinant which is of a limited molecular size.

The antigenic determinants may be parts of proteins, nucleic acids, polysaccharides, lipids or glycolipids, or other biological macromolecules. Very often they have unique steric conformations and are part of a native structure. We have thus to distinguish between *sequential determinants* and *conformation-dependent* (or conformational) determinants. The operational definition holds that if antibodies against, e.g. a protein, react well with a tetra-, penta-, or hexapeptide derived from that protein, then the antibody is against a sequential determinant. The antibody, on the other hand, is against a conformation-dependent determinant if it is made against a juxtaposition of atoms in space which results from a unique conformation of the macromolecule, and any peptide derived from such a protein will not be able, after denaturation, to react with the antibody.

It is of interest that for most globular proteins and native nucleic acids, almost all the antigenic determinants are conformation-dependent, whereas for most polysaccharides, fibrillar proteins such as silk fibroin, and single-stranded nucleic acids, the determinants are sequential. The use of homopolymers of amino acids or sugars, and of peptidyl proteins as antigens has established that the determinant is composed of four to six amino acid or sugar residues which contribute unequally to binding with the antibody combining site.

Not all the antigenic determinants express themselves all the time: some are more *immunopotent* than others. Some do not express themselves at all under a certain set of conditions, and we call them *immunosilent*, even though under other conditions they may provoke an efficient immune response. Thus a determinant may be immunosilent within a complete macromolecule, but may be quite immunopotent when a segment of that macromolecule on which it is present is used for immunization. Situations are also known where a determinant is immunosilent but

1.2 Antigens

becomes immunopotent in animals made tolerant to other parts of the immunogenic macromolecule of which the determinant discussed forms a part. We may thus define *immunopotency* as the capacity of a region of an antigen molecule to serve as an antigenic determinant and induce the formation of a specific immune response. For all the above reasons, one should never use the expression immunogenic determinant when discussing a determinant within an immunogen which expresses itself; this expression (*immunogenic determinant*) should be used only to define a portion of an antigen of the size of a specificity determinant which can be cut off by chemical or enzymatic methods, and is still able to provoke an immune response when used for immunization by itself.

The term *hapten*, in its strictest sense, designates any substance, large or small, which does not elicit an immune response by itself but can be shown to react with antibody provoked by immunization with a complete immunogen of which the hapten formed a part. In practice, most haptens investigated are small chemical substances, in most cases of a size definitely smaller than that of a complete antigenic determinant. When attached to a protein, haptens such as dinitrophenyl or penicilloyl might be defined as *immunodominant* parts of an antigenic determinant (an epitope). In studies of determinants of polysaccharide or polypeptide nature it is of interest to establish which is their immunodominant portion.

The common response to an immunogen is either the production of antibodies, which find their way usually into the bloodstream, or exclusively cell-mediated immunity, e.g. delayed hypersensitivity. Still other responses may result in specific *immunological tolerance*, i.e. induction of the inability to form antibodies or cellular reaction towards a specific antigen. Generally an organism distinguishes between 'self' (material which is its own) and 'non-self' (any foreign material). The immune system of an organism reacts against any foreign compound (antigen), and is *tolerant* (unable to react) towards its own body components, which may be good immunogens in other organisms. This *self-tolerance* is acquired during fetal or neonatal development, and the immune system can be made tolerant even to foreign material or tissue introduced during this period. Such material, which can induce immunological tolerance or unresponsiveness, is called a *tolerogen*.

Another phenomenon which should be mentioned here is *antigenic competition*. This may occur between different antigens (intermolecular), or between different specificity determinants on the same antigen (intramolecular), in which case we define it as *competition between antigenic determinants*. The existence of

this phenomenon may account for some determinants being *immunosilent* under certain circumstances. Antigenic competition may be defined as the inhibition of the immune response to one antigen or determinant by the administration of another antigen or determinant.

Molecules as antigens

The two types of natural macromolecules most investigated as antigens are proteins and polysaccharides. These also include glycoproteins, nucleoproteins, lipoproteins, etc. as well as peptidoglycans, glycolipids, and other conjugates. Nucleic acids are also antigenic. Lipids are poor immunogens, but antibodies against them can be obtained, and liposomes play a role here, as they do in enhancing the immunogenicity of various other antigens. Synthetic antigens, especially synthetic polypeptides, have played an important role in the elucidation of the molecular basis of antigenicity and many other immunological phenomena, and they will be discussed here in more detail below, as well as in a separate chapter (Chapter 2). Other synthetic polymers have also been shown to be immunogenic, e.g. polyvinylpyrrolidone.

All proteins are probably immunogenic, although individual proteins differ markedly in their immunogenicities. Similarly, although a foreign protein can induce the synthesis of both circulating and cell-bound antibodies, the relative distribution of these responses varies considerably for different proteins. Denatured proteins are often less immunogenic than the corresponding native proteins. Self-aggregation of a protein is usually associated with a negligible change in its antigenic specificity, but with a considerable increase in immunogenicity. Like other antigens, proteins possess a finite number of antigenic determinants that correspond to discrete portions of the surface structure and that are preferentially located in those regions most exposed to the external environment. The relationship between structure and antigenicity is, however, more complex for globular proteins than for other antigens in that it depends to a very large extent upon the overall conformation of the molecule. The exploration of antigenic regions on the surface of proteins has become easier with the advent of monoclonal antibodies and rapid methods of peptide synthesis. The antigenic sites may be described as surface domains made up from amino acid side-chains which may be distant in sequence but close in space (conformation-dependent determinants). Such domains are probably overlapping and cover most of the protein surface. On the other hand, fibrous proteins possess sequential determinants, whose size may be determined with synthetic peptides to be in the range of three or four, up to five or six amino acid residues.