# TABLES on the THERMOPHYSICAL PROPERTIES of LIQUIDS and GASES

In Normal and Dissociated States

N. B. Vargaftik

+3.81107+ V 297(2)

# TABLES ON THE THERMOPHYSICAL PROPERTIES OF LIQUIDS AND GASES

#### In Normal and Dissociated States

SECOND EDITION

BY

#### N. B. VARGAFTIK


WITH A FOREWORD TO THE ENGLISH EDITION BY

Y. S. Touloukian
PURDUE UNIVERSITY



HEMISPHERE PUBLISHING CORPORATION

Washington - London



Originally published by Nauka Press, Moscow as Spravochnik po Teplofizicheskim Svoitstvam Gazov i Zhidkostey

English translation copyright © 1975 by Hemisphere Publishing Corporation.

All rights reserved. No part of this book may be reproduced in any form, by photostat, microform, retrieval system, or any other means, without the prior written permission of the publisher.

Hemisphere Publishing Corporation 1025 Vermont Aye., N.W., Washington, D.C. 20005

Distributed solely by Halsted Press, a Division of John Wiley & Sons, Inc., New York.

Library of Congress Cataloging in Publication Data

Vargaftik, N B
Tables on the thermophysical properties of liquids and gases.

Translation of Spravochnik po teplofizicheskim svoistvam gazov i zhidkostel.
Bibliography: p.
Includes index.
1. Gases—Thermal properties—Tables, etc.
2. Liquids—Thermal properties—Tables. I. Title.
QC145.4.T5V3713 1975 530.4'2 75-14260
ISBN 0-470-90310-4

Printed in the United States of America

# TABLES ON THE THERMOPHYSICAL PROPERTIES OF LIQUIDS AND GASES

In Normal and Dissociated States

Editors:

#### JAMES P. HARTNETT THOMAS F. IRVINE, JR.

1	Blackshear	<ul> <li>Heat Transfer in Fires: Thermophysics, Social Aspects, Economic Impact*</li> </ul>
H	Afgan and Beer	Heat Transfer in Flames*
Ш	dez #ties and Afgan	<ul> <li>Heat and Mass Transfer in the Biosphere: Part I Transfer Processes in Plant Environment*</li> </ul>
IV	Vargaftik	Tables on the Thermophysical Properties of Liquids and Gases
V	Veziroglu	Remote Sensing: Energy-Related Studies
	Begell	• Glossary of Terms in Heat Transfer: English, Russian, German, French, and Japanese (in preparation)
	Chang	• Control of Flow Separation (in preparation)
	Chi	Heat Pipe Theory and Practice (in preparation)
	Denton	• Future Energy Production* (in preparation)
	Eckert and Goldstein	<ul> <li>Heat Transfer Measurement</li> <li>2nd revised and augmented edition</li> <li>(in preparation)</li> </ul>
	Ginoux	• Two-Phase Flows with Application to Nuclear  Reactor Design Problems† (in preparation)
	Gutfinger .	• Topics in Transport Phenomena: Bioprocesses, Mathematical Treatment, Mechanisms (in preparation)
	Hartnett	<ul> <li>Heat and Mass Transfer Problems in Future Energy Production*         <ul> <li>(in preparation)</li> </ul> </li> </ul>
	Hsu and Graham	<ul> <li>Transport Processes in Boiling and Two-Phase Systems, Including Near-Critical Fluids (in preparation)</li> </ul>

(in preparation)

Theory† (in preparation)

• Advanced Heat Conduction (in preparation)

Pfender

Richards

Sieverding

Yovanovich

DN97/15

• High-Temperature Phenomena in Electric Arcs (in preparation)

Two-Phase Fluid Dynamics in Turbines: Engineering, Instrumentation,

Measurement of Unsteady Fluid Dynamic Phenomena†

<sup>\*</sup>A publication of the International Centre for Heat and Mass Transfer, Belgrade. †A von Karman Institute Book, Brussels.

## FOREWORD TO THE ENGLISH TRANSLATION-SECOND EDITION

I am indeed pleased to write a foreword to this extensive work. This book is a translation of the second, most recent edition of Professor Vargaftik's "Spravochnik po. Teplofizicheskim Svoitstvam Gazov i Zhidkostey," which first appeared in this format in 1963. The primary contribution of the translated version comes from the author's extensive presentation, primarily from the Russian literature, on a wide range of materials of exchnological importance, in the liquid, vapor, gaseous, dissociated, and ionized states. The thermodynamic and transport properties presented, often over a wide range of temperatures and pressures, are basically reproductions from other data sources. In several instances, the labular material is unique, and this reasonably up-to-date compendium would be most welcome by many in the English-speaking world to whom much of the data would otherwise remain unobtainable.

In the absence of any evaluative commentary in the presentation of the data, those who use this work will have to rely at this time on the author's judgment in the selection of the data and his internationally acknowledged reputation for his many valuable and original contributions to the field. I very much hope, however, that this work, together with others, may constitute the basis for future cooperative endeavors for the generation of an internationally agreed-upon set of thermophysical properties tables for gases and liquids such as that currently sponsored by special Task Groups within CODATA/ICSU, IUPAC, and the International Association for the Properties of Steam

In this translation the corrections found on the errata page of the original work have been incorporated as have additional corrections noted since it was published. As an added feature the translated edition presents a thorough index to both the substances and the properties covered in the volume. The reader will find these conveniences most helpful in using this extensive reference work.

Y. S. Touloukian, Director Center for Information and Numerical Data Analysis and Synthes:3 an<sup>3</sup> Distinguished Atkins Professor of Engineerin Purdue University West Lafayette, Indiana

## FOREWORD TO THE SECOND EDITION

Since the publication of the first edition of this Handbook (in 1963) a large amount of material on thermophysical properties of different substances has been published. The following changes and additions have been incorporated in the present edition:

- 1. New, detailed data are given for two types of hydrogen, both in the liquid and in the gaseous states. The present tables contain data up to 6,000°K and 1,000 bar, that is, the data include the region of high temperatures where hydrogen exists in a dissociated state. New tables of transport properties of hydrogen have been provided for both the liquid and the gaseous states at different pressures up to a temperature of 2,000°K.
- 2. The thermophysical properties of nitrogen, oxygen, air and argon are given over a much wider range of parameters. Detailed new data at high temperatures are given for these substances, for both the liquid as well as for the gaseous states, ranging from extremely low to very high pressures (up to 1,000 bar).
- 3. The tables of thermophysical properties of carbon dioxide  $(CO_2)$  have been reworked, expanded and rendered more accurate. This section contains new detailed data for liquid carbon dioxide at high temperatures (up to  $4,000^{\circ}$ K) in the critical region, where it exists in the dissociated state.
- 4. More detailed tables are given for alkali metals and mercury. Thermodynamic properties have been determined on the basis of new, more accurate data on the dissociation energy of diatomic molecules of alkali metals. The thermodynamic properties tables have been expanded to 3,000°K, taking into account both dissociation and ionization. Thermodynamic properties for ionized lithium at high temperatures are given in a separate set of tables. Viscosity and thermal conductivity of alkali metals in the gas phase are reported here for the first time.
- 5. Information on thermophysical properties of monatomic substances has been significantly expanded. In particular, new tables for helium include data for both liquid and gas phases at extremely low and high temperatures (up to 3,000°K) at different pressures. Thermophysical properties of neon, krypton and xenon are given for a wide range of temperatures and pressures.
- 6: Thermophysical properties of water and steam are given here on the basis of new international tables adapted in 1963–1964. For dissociated steam, data are given up to 6,000° K and 1,000 bar taking into account true properties of dissociated steam.

- 7. New data are given on diffusion in binary gas mixtures, on viscosity and thermal conductivity of mixtures and solutions; data at different pressures are included.
- 8. The majority of tables for the remaining substances cite more accurate and newer data over a wider range of parameters.
- 9. The editors have not included in the present edition data on thermophysical properties of compounds of boron, nitrogen oxide, gallium, deuterium and individual hydrocarbons, due to the fact almost no new information for these substances has been published since the appearance of the original edition of this handbook.
- 10. All thermophysical properties are given in SI units, with the exception of the data for ionized lithium, argon and nitrogen.

The author is grateful to the entire faculty of the Department of Physics at Moscow Institute of Aviation for their help in the preparation of this handbook. The author was helped in many aspects in the preparation of the present edition by Prof. P. M. Kesselman, Docents L. D. Volyak, Y. K. Vinogradov and N. D. Kosov and Assistant E. L. Studnikov, to whom he wishes to express his appreciation.

The author is very grateful to Academician A. S. Predvoditelev and Professor L. P. Filippov for valuable comments and to E. I. Gaydul for great help in the preparation of the manuscript

N. B. Vargaftik

## FOREWORD TO THE FIRST EDITION

The requirements of modern science and technology necessitate the knowledge of thermophysical properties of gases and liquids.

This area of science, in addition to experimental work, centers on research in diffusion, thermal conductivity and viscosity of gases, as well as on the derivation of equations of state for real gases. Over the recent years, a large amount of data on thermophysical properties of gases and liquids have been collected; these data are of great practical interest.

The present book attempts to provide a systematic presentation of data obtained over the last 5-10 years. Most reliable data for pure substances, gas mixtures and solutions are presented. The data are essentially based on experimental results. Most often, the data are given for whole-number temperatures and pressures, and in the cases of mixtures for concentrations convenient for practical calculations.

The handbook gives data for thermophysical properties of a number of gases: hydrogen, lithium, nitrogen, argon and steam; data are given at high temperatures and take into account the dissociation of the given gas. Data on thermophysical properties of vapors of lithium, sodium and potassium up to 2,000°K take into account the dimerization of these substances in the gaseous state. The book gives I-S [entropy-enthalpy] charts for these substances in the range of high temperatures.

In the compilation of this handbook, data reported in both monographic and periodical literature has been cited.

The author would like to express his appreciation to the entire faculty of the Department of Physics at the Moscow Institute of Aviation for their help in the preparation of this book.

Special thanks are due to Docent L. D. Volyak who arranged and correlated the data and compiled the tables on thermodynamic properties of potassium in the gas phase, to Assistant Yu. D. Vasilevskaya who worked on diffusion in binary mixtures, to V. V. Rybakov for thermodynamic properties of lithium at high temperatures in the gas phase. Candidate of Technical Sciences L. S. Zaytseva prepared the graphical presentation of thermodynamic properties of air at high temperatures.

The author is deeply grateful to Academician A. S. Predvoditelev and Candidate of Physical-Mathematical Sciences L. P. Filippov for their very valuable comments, and to E. I. Gaydul for great help in the preparation of the manuscript.

#### CONTENTS

21-12
$\mathfrak{A}_{l}$

FOREWORD TO THE ENGLISH TRANSLATION—SECOND EDITION ix
FOREWORD TO THE SECOND EDITION xi
FOREWORD TO THE FIRST EDITION xiii
USEFUL CONVERSION TABLES AND SYMBOLS 1
NOTATION 6
CONVERSION OF UNITS FROM SI TO OTHER SYSTEMS 6
DADET T
PART I PURE SUBSTANCES
1 HYDROGEN AND HYDROGEN COMPOUNDS
2 METALS
3 CARBON COMPOUNDS
19436

4	HYDROCARBONS AND ORGANIC COMPOUNDS
*	Alkanes 211
	Naphthenes 296
	Olefines 306
	Cyclic Aliphatic Compounds 331
	Dienes 333
ì	Alkynes 337
	Aromatic Hydrocarbons 342
	Hydrocarbons of the Diphenylmethane Series (high-temperature
	heat transfer agents [coolants]), 360
	Halogen Substitute Hydrocarbons 362
	Alcohols 404
	Acetone 427
	Carboxylic Acids 429
	Aniline 430
	Nitrobenzene 431
5	NITROGEN AND AMMONIA
	Nitrogen 433
	Ammonia 464
6	OXYGEN
7	SULFUR DIOXIDE
8	HALOGENS
	Fluorine 506
	Chlorine 516
9	MONOATOMIC GASES
	Helium 521
	Neon 536
•	Argon 543
	Krypton 575
	Xenon 580
	PART II
	MIXTURES
10	AIR
- ; -	
11	DIFFUSION IN GASES
12	THERMODIFFUSION IN GASES

101	TEN	ma

	CONTENTS	
13	THERMOPHYSICAL PROPERTIES OF GAS MIXTURES AND SOLUTIONS 666 Viscosity of Binary Gas Mixtures 666 Thermoconductivity of Binary Gas Mixtures 679 Viscosity of Aqueous Solution 683 Thermoconductivity of Aqueous Solution 685	
14	LIQUID FUELS	
15	HIGH-TEMPERATURE HEAT TRANSFER AGENTS	
16	OILS	
REF	FERENCES	
IND	DEX	
	Elements, Compounds, and Mixtures, and Their Properties 719  Properties of Elements, Compounds, and Mixtures 741  Diffusion in Gases 753  Aqueous Solutions and Gas Mixtures and Their Properties 757	

#### USEFUL CONVERSION TABLES AND SYMHOLS

#### **Dimensionless Groups**

Group*	Symbol	Name
$\Delta p/\rho V^2$	Eu	Euler number
$at/r_0^2$	Fo	Fourier number
$(L/d)(k/Vd\rho c_p)$	Gz = (L/d)/RePr	Graetz number
$g\beta(\Delta T) L^3 \rho^2/\mu^2$	Gī	Grashof number
λ/L	Kn	Knudsen number
α/D	Le	Lewis number
V/V sound	Ma	Mach number
hL/k, $hd/k$	Nu	Nusselt number
Vdρc <sub>p</sub> /k	Pe (= RePr)	Peciet number
$c_{p}\mu/k$	Pr	Prandtl number
$g\beta(\Delta T)L^3\rho^2c_p/\mu k$	Ra (= GrPr)	Rayleigh number
ρVD/u, ρVL/μ	Re	Reynolds number
$\mu/\rho D$	Sc	Schmidt number
$h_{p}d/D$	Sh	Sherwood number
$h/c_pG$	St (= Nu/RePr)	Stanton number
$V_{\infty}^2/C_{p}(\Delta T)_{0}$	E	Eckert number
$V^{2}/gL$	Fr	Froude number
$f_r d/V$	St	Strouhal number
$\rho V^2 L/\sigma$	We	Weber number

<sup>\*</sup> $f_r$  = frequency of oscillation  $\sigma = \text{surface tension}$ .

#### Conversion Table

	To convert number of	То	Multiply by	
Length	inch	cm	2.540	
	ft .	m	0.3048	
Area	ft <sup>2</sup>	m²	0.0929	
Volume	ft³	m³	0.02832	
Mass	lbm	kg	0.45359	
	Slugs	kg	14.594	
Force	lbf	Newtons	4.4482	
Density	lbm/ft³	kg/m <sup>3</sup>	16.02	
Work	ft-lbf	mkg	0.1383	
	hp-hr	mkg	273,700	
Heat	Btu	kcal	0.2520	
	Chu	Btu	1.800	
	Btu	Joules	1054.35	
i	Btu	ft-lbf	778.26	
	kw-hr	Btu	3412.75	
Specific heat	Btu/lbm-°F	cal/gC	1.000	
•	Btu/lbm °F	Wsec/kgmC	4184.0	
Pressure	lbf/in <sup>2</sup> , psi	kgf/cm <sup>2</sup>	0.070309	
	psi	atm	0.068046	
	psi	bars .	0.068948	
	psi	dynes/cm <sup>2</sup>	68947.0	
Surface tension	lbf/ft	dynés/cm	6.8519 x 10	

#### Heat Flux, q/A

, 4/						
Multiply number of To obtain	<u>Btu</u> ft <sup>2</sup> -hr	w/cm²	kcał hr-m²	calsec-cm²		
Btu/ft²-hr W/cm² kcal/hr-m² cal/sec-cm²	1 3.154 x 10 <sup>-4</sup> 2.7126 7.536 x 10 <sup>-5</sup>	3,170.75 1 8,500 0.2389	0.36865 1.163 x 10 <sup>-4</sup> 1 36,000	13,277.26 4.1868 2.778 × 10 <sup>-5</sup>		

#### Heat Transfer Coefficient, h

Multip://number of	Btu	₩	cal	keal
	hr-ft <sup>2</sup> .°F	cm²-°C	sec-cm <sup>2</sup> -°C	hr-m²-°C
Btu/hr-ft <sup>2</sup> -°F	1	1761	7376	0.20489
W/cm <sup>2</sup> -°C	5.6785 x 10 <sup>-4</sup>	1	4.186	1.163 × 10 <sup>-4</sup>
cal/sec-cm <sup>2</sup> -°C	1.356 x 10 <sup>-4</sup>	0.2391	1	2.778 × 10 <sup>-5</sup>
kcal/hr-m <sup>2</sup> -°C	4.8826	8600	36000	1

#### Thermal Conductivity, &

Multiply number of	Btu hr-ft-°F	W cm-°C	cal sec-cm-°C	<u>kcal</u> hr-m-°C	Bu in hr-ft <sup>2</sup> ° F	
Btu/hr-ît- r W/cm-°C cal/sec-cm °C cai/it-it- °C Btu in /hr ft <sup>2</sup> °F	1 0.01730 4.134 × 19 <sup>-3</sup> 1.466 12	57.793 1 0.2389 86.01 693.5	241.9 4.186 1 - 360 2903	0.6722 0.01171 2.778 x 10 <sup>-3</sup> 1 8.064	0.08 33 1.44 × 10 <sup>-3</sup> 3.44 × 10 <sup>-4</sup> 0.1240	

#### Viscosity, µ

	•	,,,,			
Multiply number of op.	lbm ft-hr	lbf-sec ft <sup>2</sup>	Centipoise	kgm m-hr	kgf-sec m <sup>2</sup>
lbm/ft-hr lbf-sec/ft <sup>2</sup> Centipoise kgm/m-hr kgf-sec/m <sup>2</sup>	1 0.00000862 0.413 1.49 0.0000421	116,000 1 47,880 172,000 4.882	2.42 0.00002086 1 3.60 0.0001020	0.672 0.00000579 0.278 1 0.0000284	23733 0.2048 9807 35305

100 centipoise = 1 Poise = 1g/sec-cm = 1 dyne sec/cm<sup>2</sup>.

	Kinematic Visc	osity, υ		
Multiply number of dy.	ft² hr	Stokes	m² hr	m² sec
ft²/hr Stokes m²/hr m²/sec	1 0.25806 0.092903 0.00002581	3.875 1 0.3599 10 <sup>-4</sup>	10.764 2.778 1 0.0002778	38,751 10 <sup>4</sup> 3600 1

#### **Conversion Factors**

The following tables of conversion factors are convenient. In order to convert the numerical value of a property expressed in one of the units in the left-hand column of the table to the numerical value of the same property expressed in one of the units in the tow row of the table, multiply the former value by the factor in the block common to both units.

In tables involving energy, cal denotes the thermochemical calorie; IT cal denotes the International Steam Table calorie. The thermochemical calorie (cal) equals 4.184 joule. The International Steam Table calorie (IT cal) equals 4.186 joule. The Btu is the International Steam Table Btu and it equals 1055.04 joule.

#### Conversion Factors for Mass

		lbm	slugs	gm	kg	ton
l lbm	=	· 1	0.03108	453.59	0.45359	0.0005
l slug	=	32.174	1	1.4594 x 10 <sup>4</sup>	14.594	0.016087
l gm	=	2.2046 x 10 <sup>-3</sup>	6.8521 x 10 <sup>-5</sup>	1	10-3	1.1023 x 10 <sup>-6</sup>
	=	2.2046	6.8521 x 10 <sup>-3</sup>	10 <sup>3</sup>	1	1.1023 x 10 <sup>-3</sup>
l ton	=	2000	61.162	9.0718 x 10 <sup>5</sup>	907.18	1

SOURCE: Modified and extended from "Selected Values of Properties of Hydrocarbons," National Bureau of Standards.

Conversion Factors for Density

		ibm/ft <sup>3</sup>	slug/ft <sup>3</sup>	lbm/in. <sup>3</sup>	lbm/gal	gm/cc
lbm/ft3	=	1	0.03108	5.787 × 10 <sup>-4</sup>	0.13368	0.01602
slug/ft <sup>3</sup>	<b>±</b>	32.174	1	0.1862	4.3010	0.51543
lbm/in.3	=	1728	53.706	1	231	27.680
lbm/gal	=	7.4805	0.2325	4.329 x 10 <sup>-3</sup>	ı	0.11983
gm/cc	=	62.428	1.9403	0.03613	8.345	1

SOURCE: Modified and extended from "Selected Values of Properties of Hydrocarbons," National Buresu of Standards.

Conversion Factors for Pressure

				omeont to store to the	•		
	lbf/ftt²	lbf/in.²	atm	in. He	n H.O	mm Ho	
1 lbf/ft² =	-	0.006944	4.726 x 10 <sup>-4</sup>	0.014139	0.19243	0.3591	7 700 10-4
1 lbf/in.2 =	144	1	0.06805	2.036	27.71	\$17.15	1.700 X 10
l atm =	2116.2	14.696	-	29.921	407.18	760	0.00093
l in. Hg =	70.726	0.49116	0.033421	-	17.500		1.01323
			0.0007441	-	13:000	25.40	0.03386
1 m. H <sub>2</sub> O =	5.197	0.036092	0.002456	0.07348		1.8665	0.002488
1 mm Hg =	2.7845	0.019337	1.315 x 10 <sup>-3</sup>	0.03937	0.53577	-	1 333 ~ 10-3
1 mm H <sub>2</sub> O =	2.0886-x 10*	14.504	0.98692	29.530	401.85	750.06	O1 4 CCC
					20:00	30.00	-•

Conversion Factors for Energy

						3			
		abs joule	caj	IT cal	Btu	int. kw-hr	ho-hr	4	liter of
l abs ioule	11		0.239005	0.239005 0.238848	0.047877 . 10-3	7-01 10000			חובו -מוזוו
	Ī			0.500040	01 X /76/4-C'0	. 01 x 15///7 01 x /76/+60	3.72505 x 10 <sup>-7</sup>	0.737561	9.86896 x 10 <sup>-3</sup>
l cal	н	4.18401	1	0.999344	0.999344 3.96572 x 10 <sup>-3</sup>	1.162028 × 10-6	1.162028 × 10-6 1.558566 × 10-6	3.08596	4 12918 \$ 10-2
! IT cal	11	4.18676	1.000657	I	3.96832 × 10 <sup>-3</sup>	1.162791 x 10-6 1 559590 x 10-6	1 559590 - 10-6	2 00700	4 12100 10-2
100		1000	1				01 × 0000001		4.13169 X 1U
nia :	"	1005.045	191.757	251.996		2.93018 x 10 <sup>-4</sup>	2.93018 x 10 <sup>-4</sup> 3.93010 x 10 <sup>-4</sup>	778.16	10.4122A
int kwhr	H	101 km.hr = 2 400 412	373 070	800	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			1	07711.01
The Water		2,000,012	000,000	200,000	3412.76	_	1.341247	2.655.669	35534 3
1 hp-hr	11,	= 2,684,525 641.615	641.615	641.194	2544.46	0.745575		000 000 1	2,000,0
1 60 11	,	1 25 5001	0.000	, 00000		ì		000,000,1	20493.3
25.11	,	1.30000	1.353621 0.324048	0.323836	1.285083 × 10 <sup>-3</sup>	3.76553 x 10 <sup>-7</sup>	5.05051 x 10 <sup>-7</sup>	_	1 338054 ~ 10-2
1 liter-atm	II	101.3278   24.2179	24.2179	24.2020	0.0960412	2 81418 v 10-5 2 77457 10-5	2 77467 10-5	7, 306,	2000
	1					212111	3.1/432 × 10	/4./334	

Conversion Factors for Specific Energy

				The state of the s	cure merigy			
	abs joule/gm	cal/gm	IT cal/gm	Bru/lb	fr.lhf/lhm	int but had	L	
- 1 - 1 - 1 - 4 - 0					1101/1011	DIC. NW-III/BIII	or/su-du	11./sec.
aos jonie/gm =	-	0.2390	0.2388	0.4299	334.53	2.777 × 10-7	1 690 x 10-4	10762
cal/gm =	4.184	1	0.9993	1.7988	1399.75		_	40.00
IT callom =	7817	1,000				2000		.01 × toc.
, car, 6,	7.100	1.000/	-	∞. ∞.	1400.69	1.163 x 10 <sup>-6</sup>	7 074 v 10-4	1 504 104
· Btu/lb =	2 326	0.5550	0.000				2 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	1.200 × 10
	4260	0.3333	0.2220	_	778.16	6.460 × 10 <sup>-7</sup>	3 930 x 10-4	25.027
ft-lbf/lbm =	2.989 x 10 <sup>-3</sup>	7.144 x 10-4 7.139 x 10-4 1.285 x 10-3	7.139 x 10-4	1.285 x 10-3		9 302 3 10-10	7-01	
hor 1012 ( = m2/24 w/d ini	301.0176	27.000			-	0.302 X 10	. 01 × 1c0.c	32.174
111/611	3.010 × 10	800,303	860,000	1.548 × 10°	1.2046 x 109	_	608.4	2 074 1010
hp-hr/lb = =	60105	14145	14136		,,,,			3.070 × 10
1		1414.0	1413.0	7242	1.980 × 10°	0.001644	_	6 370 5 107
11, /scc = =	9.291 x 10 <sup>-5</sup>	$= 9.291 \times 10^{-5}   2.220 \times 10^{-5}   2.219 \times 10^{-5}   3.994 \times 10^{-5}$	2.219 × 10 <sup>-5</sup>	3 994 × 10-5	İ	=		
						7.300 × 10		_

#### Conversion Factors for Specific Energy per Degree

		abs joule/gm-°K	cal/gm-°K	lT cal/gm-°K	Btu/lb-°R	w-sec/kg-°K
abs joule/gm-°I	(=	1	0.2390	0.2388	0.2388	10 <sup>3</sup>
cal/gm-°K	=	4.184	1	0.9993	0.9993	4184
IT cal/gm-°K	=	4.186	1.0007	l	1	4186
Biu/lb-°R	=	4.186	1.0007	1	1	4186
w-sec/kg-°K	=	10-3	$2.390 \times 10^{-4}$	2.388 × 10 <sup>-4</sup>	$2.388 \times 10^{-4}$	1

#### Conversion Factors for Thermal Conductivity

	cal/sec-cm-°C	Btu/hr-ft-°F	Btu/hr-ft2-°F/in.	w/cm-°C
cal/sec-cm-°C =	1	241.9	2903	4.183
Btu/hr-ft-°F =	$4.13 \times 10^{-3}$	1	12	0.0173
Btu/hr-ft2-°F/in.=	3.45 x 10 <sup>-4</sup>	0.0833	1	1.44 x 10 <sup>-3</sup>
$l w/cm^2 \cdot C =$	0.239	57.8	694	1

#### Conversion Factors for Dynamic Viscosity

	poise or g/cm-sec, or dyn-sec/cm <sup>2</sup>	lbm/ft-hr or pdl-hr/ft <sup>2</sup>	lbm/ft-sec or pdl-sec/ft <sup>2</sup>
l poise =	ı	242	0.0672
1 lbm/ft-hr =	4.13 x 10 <sup>-3</sup>	l	2.78 x 10 <sup>-4</sup>
l lbm/ft-sec =	14.87	3600	1

#### Conversion Factors for Kinematic Viscosity

	ft²/hr	stokes	m²/hr	m²/sec
ft²/hr =	l	0.25806	0.092903	2.58 x 10 <sup>-5</sup>
stokes =	3.885	1	0.36	10-4
$m^2/hr =$	10.764	2.778	1	$2.778 \times 10^{-4}$
$m^2/sec =$	38,750	104	3600	1