

HANDBOOK OF CHEMICAL NEUROANATOMY

Edited by A. Björklund and T. Hökfelt

Volume 4:

GABA AND NEUROPEPTIDES IN THE CNS, PART I

Editors:

A. BJÖRKLUND

T. HÖKFELT

HANDBOOK OF CHEMICAL NEUROANATOMY

Edited by A. Björklund and T. Hökfelt

Volume 4:

GABA AND NEUROPEPTIDES IN THE CNS, PART I

Editors:

A. BJÖRKLUND

Department of Histology

University of Lund, Lund, Sweden

T. HÖKFELT

Department of Histology

Karolinska Institute, Stockholm, Sweden



1985

ELSEVIER

Amsterdam – New York – Oxford

© Elsevier Science Publishers B.V., 1985

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without the prior written permission of the publisher, Elsevier Science Publishers B.V./ Biomedical Division, P.O. Box 1126, 1000 BC Amsterdam, The Netherlands.

Special regulations for readers in the USA – This publication has been registered with the Copyright Clearance Center Inc. (CCC), Salem, Massachusetts. Information can be obtained from the CCC about conditions under which photocopies of parts of this publication may be made in the USA. All other copyright questions, including photocopying outside of the USA, should be referred to the publisher.

ISBN 0 444 90353 4

ISBN SERIES 0 444 90340 2

Library of Congress Cataloging-in-Publication Data

Main entry under title:

GABA and neuropeptides in the CNS.

(Handbook of chemical neuroanatomy ; v. 4, etc.)

Includes bibliographies and index.

1. Neuropeptides--Analysis. 2. GABA--Analysis.

3. Central nervous system--Anatomy. I. Björklund, Anders, 1945- II. Hökfelt, Tomas. III. Series.

[DNLM: 1. Central Nervous System. 2. GABA. 3. Nerve Tissue Proteins. W1 HA51J v. 4 etc. / WL 104 G112]

QM451.H24 1983 vol. 4, etc. 616.8 s 85-20596

[QP552.N39] [599'.01'88]

ISBN 0-444-90353-4 (U.S. : pt. 1)

Published by:

Elsevier Science Publishers B.V.

P.O. Box 1126

1000 BC Amsterdam

Sole distributors for the USA and Canada:

Elsevier Science Publishing Co. Inc.

52 Vanderbilt Avenue

New York, NY 10017

Printed in The Netherlands by Casparie -- Amsterdam

List of contributors

G.M. ABRAMS
Department of Neurology
College of Physicians and Surgeons
Columbia University
630 West 168th Street
New York, NY 10032
U.S.A.

J. BARRY
Laboratoire d'Histologie
Faculté de Médecine
Place de Verdun
59045 Lille
France

M.J. BROWNSTEIN
Laboratory of Clinical Science
National Institute of Mental Health
Building 10
National Institutes of Health
Bethesda, MD 20205
U.S.A.

P.C. EMSON
MRC Neurochemical Pharmacology Unit
Medical Research Council Centre
Hills Road
Cambridge, CB2 2QH
U.K.

M. GOEDERT
MRC Neurochemical Pharmacology Unit
Medical Research Council Centre
Hills Road
Cambridge, CB2 2QH
U.K.

G.E. HOFFMAN
Department of Anatomy
University of Rochester
School of Medicine and Dentistry
Rochester, NY 14642
U.S.A.

KANG TSOU
Shanghai Institute of Materia Medica
Chinese Academy of Science
Shanghai 200031
China

H. KHACHATURIAN
Mental Health Research Institute,
The University of Michigan
205 Washtenaw Place
Ann Arbor, MI 48109
U.S.A.

M.E. LEWIS
Mental Health Research Institute
The University of Michigan
205 Washtenaw Place
Ann Arbor, MI 48109
U.S.A.

J.L. MADERDRUT
Department of Anatomy
Bowman Gray School of Medicine
Winston-Salem, NC 27103
U.S.A.

P.W. MANTYH
MRC Neurochemical Pharmacology Unit
Medical Research Council Centre
Hills Road
Cambridge, CB2 2QH
U.K.

I. MERCHENTHALER
Department of Anatomy
University of North Carolina
111 Swing Building
Chapel Hill, NC 27514
U.S.A.

List of contributors

E. MUGNAINI

Laboratory of Neuromorphology
Department of Biobehavioral Sciences
Box 154, University of Connecticut
Storrs, CT 06268
U.S.A.

G. NILAVER

Department of Neurology
College of Physicians and Surgeons
Columbia University
630 West 168th Street
New York, NY 10032
U.S.A.

W.H. OERTEL

Department of Neurology
Technical University of Munich
Moehlstrasse 28
D-8000 München 80
F.R.G.

M. PALKOVITS

Laboratory of Cell Biology
National Institute of Mental Health
Building 36, Rm 3A-17
National Institutes of Health
Bethesda, MD 20205
U.S.A.

P. PETRUSZ

Department of Anatomy
University of North Carolina
111 Swing Building
Chapel Hill, NC 27514
U.S.A.

V. PICKEL

Laboratory of Neurobiology
Department of Neurology
Cornell University Medical College
1300 York Avenue
New York, NY 10021
U.S.A.

M.V. SOFRONIEW

Department of Human Anatomy
South Parks Road
Oxford, OX1 3QX
U.K.

Present address:

Department of Neurosurgery
Meyer 7-109

The Johns Hopkins Hospital
600 North Wolfe Street
Baltimore, MD 21205
U.S.A.

J.J. VANDERHAEGHEN

Laboratory of Neuropathology and
Neuropeptide Research
Faculty of Medicine
Free University of Brussels
Queen Elizabeth Medical Foundation
1 J.J. Crocq Avenue
1020 Brussels
Belgium

S.J. WATSON

Mental Health Research Institute
University of Michigan School of
Medicine
205 Washtenaw Place
Ann Arbor, MI 48109
U.S.A.

S. WRAY

Department of Anatomy
University of Rochester
School of Medicine and Dentistry
Rochester, NY 14642
U.S.A.

E.A. ZIMMERMAN

Department of Neurology
Columbia Presbyterian Hospital
Neurology Institute, Room 216
710 W. 168th Street
New York, NY 10032
U.S.A.

Preface

The first neuroactive peptides to be structurally identified in the central nervous system were vasopressin and oxytocin (Du Vigneaud et al. 1953a,b). At that time, these peptides were viewed as hormones, and the so-called neurosecretory neurons (Bargmann and Scharrer 1951) which produce them were thought to be exceptional in that they secrete their products in an endocrine manner into the bloodstream. The current upsurge of interest in the role of small peptides as transmitter candidates or neuromodulatory compounds dates back to the early 1970's. Of great importance for this development was the discovery by Guillemin, Schally, Vale and their collaborators that the hypothalamic releasing and inhibitory hormones, originally postulated by Harris in 1955 – in terms of their chemistry – belong to the group of small peptides. Thus, thyrotropin releasing hormone (TRH) turned out to be a tripeptide (Bøler et al. 1969; Burgus et al. 1969), the luteinizing hormone releasing hormone (LHRH) a decapeptide (Amoss et al. 1971; Matsuo et al. 1971; Schally et al. 1971), and somatostatin, the growth hormone release inhibiting hormone, a tetradecapeptide (Brazeau et al. 1971).

At about the same time, Leeman and collaborators (Chang et al. 1971) were able to chemically characterize a compound, substance P, which was already discovered in 1931 by Von Euler and Gaddum. Substance P was also found to be a peptide, an undecapeptide. Using radioimmunoassay and immunohistochemistry, it was demonstrated that these peptides occur in neurons not only in the hypothalamus but also in widespread areas of the central nervous system, and in several cases also in peripheral neurons.

These discoveries drew attention to other peptide hormones that had been isolated and structurally characterized much earlier, for example cholecystokinin (CCK), first described by Mutt and Jorpes in 1968. In fact, the demonstration of gastrin/CCK-like immunoreactivity in the brain by Vanderhaeghen and coworkers (1975) showed that a polypeptide which so far had been associated only with endocrine functions in the periphery, could also be present in the central nervous system. These and many other findings were the basis for the growing interest in and formulation of the concept of the brain-gut axis.

During the seventies and eighties, numerous other neuronal peptides have been discovered. Snyder (1980), discussing a possible transmitter role of peptides in the central nervous system, predicted that their number may well be several hundred, and the development since then has supported that prediction. In fact, an important issue has been the realization that families of peptides may exist. One example is the dramatic progress in the field of opioid peptides, starting with the discovery by Hughes, Kosterlitz and collaborators (1975) of two pentapeptides, the enkephalins, with opioid activity in the brain. Thanks to the work of many groups, notably Eipper and Mains and the teams around Chrétien, Goldstein, Herbert and Numa, this research has resulted in less than a decade in a fairly complete knowledge of the precursor molecules not only of these two pentapeptides but also of two other opioid peptide precursors, and the description of a large number of biologically active cleavage products arising from these compounds (see Weber et al. 1983).

Furthermore, evidence has been presented that substance P is not the only tachykinin present in the mammalian nervous system, but that there are several compounds belonging to this peptide family. Thus, substance P and substance K are two peptides coded

Preface

for by the same gene; in addition, there are other tachykinins (Kangawa et al. 1983; Kimura et al. 1983; Nawa et al. 1983; Minamino et al. 1984; Tatemoto et al. 1985).

New powerful techniques have contributed to these advancements and to the explosive development in the peptide field. Thus, in addition to classical purification methods starting with huge amounts of tissue and purification steps monitored by parallel bioassay, there are now more chemically oriented methods, depending on for example, isolation of peptides containing C-terminal amide groups (Tatemoto and Mutt 1980). The DNA recombinant technology is now rapidly being introduced in the field of neurobiology. The discovery of a calcitonin gene related peptide (CGRP) (Amara et al. 1982; Rosenfeld et al. 1983) with powerful actions in the central and peripheral nervous systems represents a major breakthrough demonstrating the potential of this approach.

The *Neuropeptides* volumes of the *Handbook of Chemical Neuroanatomy* deal with the structural organization of neuropeptide-containing neuronal systems in the CNS, as revealed primarily by the immunohistochemical technique. In the second of the two *Neuropeptide* volumes so far planned, we have also included chapters dealing with peptide receptor localization based on the ligand binding techniques. Peptide distributions in peripheral tissues are covered in the *Peripheral Nervous System* volume of This Series.

The present volume also includes a major chapter by Mugnaini and Oertel on GABA neurons in the CNS as revealed by an antiserum raised against the GABA-synthesizing enzyme, glutamic acid decarboxylase (GAD). This chapter was originally planned for Volume 3, but due to the extensive work the authors had to invest in this chapter – reflecting the vast and complex distribution of the GABAergic system throughout all parts of the CNS – we chose to include it in the present volume. This chapter, which should be studied in conjunction with the chapters by Ottersen and Storm-Mathisen and by McGeer and collaborators in *Volume 3*, gives a unique overview of putative GABAergic neurons in the CNS and contains a wealth of previously unpublished material, including the first complete mapping of GAD in the rat CNS.

The term 'neuropeptide' is as yet poorly defined. The compounds included here share localizational features in that: (1) they are neuron-specific (although they occur in endocrine and paracrine types of cells as well); (2) they are characteristic for defined subsets of neurons, i.e. for some but not all neurons; and (3) they are likely to be secretory products of these neuronal subsets. Functionally, the compounds we now include among the neuropeptides may, on the other hand, turn out to be quite heterogeneous. For some of the most studied peptides, such as vasopressin, oxytocin and the hypophysiotrophic peptides, at least part of their biological properties (in particular their neurohormonal actions) are well known. But even so, the role of neuropeptides in neuronal communication is only beginning to be unraveled. This problem is particularly evident for individual members of the 'peptide families' such as the opioid peptide, substance P or the pancreatic polypeptide families mentioned above. It is possible, or perhaps even likely, that only some of these neuropeptides will turn out to possess neurotransmitter- or neuromodulator-type actions, while others may exert e.g. trophic, metabolic or vascular effects.

Regardless of their function, however, the neuropeptides have become important as chemical 'tags' for the study of neuronal circuitries. Thus, the neuropeptides and the neuropeptide families serve to label subsets of neurons that are, or are likely to be, functionally related, and sensitive and specific antibodies raised against the purified or synthetic compounds are today important tools for selective immunohistochemical visualization of such neuronal subsets. The availability of an increasing repertoire of neuropeptide antisera has in fact been of great importance for the rapid development of chemical neuroanatomy over the last decade.

The immunohistochemical visualization of peptide-containing neurons forms the basis of most chapters in the present volume. While this has proved to be a powerful approach, there are definite interpretational problems inherent to the technique which are important to bear in mind, viz. the shortcomings of the immunological recognition mechanisms which underlie the attachment of the antibodies to antigens in the tissue. Thus, most antibodies used recognize only a small part of a peptide molecule, and it is well known that such antigenic determinant parts may be shared by several different (although sometimes related) peptides. Specificity is therefore frequently a difficult problem in the interpretation of immunohistochemical pictures. In some fields, such as the opioid peptide or the pancreatic polypeptide family, this has been gradually sorted out through the combined use of antibodies directed against different parts of a molecule or against different related molecules. However, in other cases, e.g. the gastrin/CCK family of peptides or the tachykinin family of peptides, much work obviously remains in order to establish the diversity and identity of immunohistochemically visualized compounds. This elementary feature of antibody-based staining is the reason why most authors chose to characterize their observed material as (peptide)-like immunoreactivity. This is particularly warranted in microscopic material, where high concentrations of antiserum are used and where the tissue most often is exposed to fixatives (such as formaldehyde or glutaraldehyde) which can modify or significantly alter the chemical structure of the compounds under study.

Peptide research has a strong tradition in Sweden. Substance P, in a broad sense one of the best characterized neuronal peptides, was discovered by Von Euler and Gaddum in 1931. Börje Uvnäs and his collaborators have been working since the sixties on the physiology and pharmacology of gastrointestinal hormones. Viktor Mutt and his colleagues have for decades played a leading role in the work of isolating peptides from the gastrointestinal tract and, much based on their discoveries, an intense peptide research has been going on since the beginning of the seventies at Swedish Universities, especially at the Karolinska Institutet and in Lund.

The two volumes *'Neuropeptides in the CNS'* are dedicated to three of the pioneer Swedish scientists who have been of special importance to us and the development in this research area. They are Rolf Luft, Bengt Pernow and Viktor Mutt. Viktor Mutt, Professor of Biochemistry at Karolinska Institutet, has discovered more than a dozen peptides, mainly by isolation from the gastrointestinal tract. Among these peptides are cholecystokinin, vasoactive intestinal polypeptide (VIP), neuropeptide Y, peptide HI and peptide YY. Through his work and extreme generosity, laboratories not only in Sweden but all over the world have benefited greatly and obtained peptides for their research. Uncountable numbers of papers have been published based on peptides provided by Viktor Mutt and his associates.

Initiation of research in a new area is often the result of the insight and foresight of single individuals. Thus, the explosive development of peptide research in Sweden was much dependent on the initiative of Professor Bengt Pernow and Professor Rolf Luft. In the beginning of the 1970's they were among the first to realize the interesting new opportunities that were opening up in the neuropeptide field. Bengt Pernow, Professor of Clinical Physiology at the Karolinska Hospital, wrote his now classical thesis on substance P (Pernow 1953) in the mid fifties, working in Ulf von Euler's laboratory. When Susan Leeman and her collaborators in 1971 sequenced substance P, Pernow realized the full potential of this discovery and initiated a new dynamic phase of substance P research which was then transmitted to many laboratories in Sweden, including our own. He recently summarized the progress in the substance P field in a review article

Preface

in *Pharmacological Reviews* (Pernow 1983). Rolf Luft was the first Professor of Endocrinology in Sweden, and already in the late forties he carried out work related to hypothalamic control of anterior pituitary hormone secretion. As a leading scientist in the field of diabetes, he became interested in somatostatin and initiated work at the Karolinska Institutet on the distribution and functional role of this peptide not only as a hypothalamic hormone but also, surprisingly, with regard to its involvement in the function of the endocrine pancreas (Luft et al. 1974).

It is an honour and pleasure to dedicate the neuropeptide volumes to these three pioneers and friends, who have been of great importance not only for our own research but for biomedical research in Sweden in general.

Lund and Stockholm in July 1985

ANDERS BJÖRKLUND

TOMAS HÖKFELT

References

- Amara SG, Jonas V, Rosenfeld MG, Ong ES, Evans RM (1982): Alternative RNA-processing in calcitonin gene expression generates mRNAs encoding different polypeptide products. *Nature (London)*, **298**, 240-244.
- Amoss M, Burgus R, Blackwell R, Vale W, Fellows R, Guillemin R (1971): Purification, amino acid composition and N-terminus of the hypothalamic luteinizing hormone-releasing factor (LRF) of bovine origin. *Biochem. Biophys. Res. Commun.*, **44**, 205-210.
- Bargmann W, Scharrer B (1951): The site of origin of the hormones of the posterior pituitary. *Am. Sci.*, **39**, 255-259.
- Bøler J, Enzmann F, Folkers K, Bowers CY, Schally AV (1969): The identity of chemical and hormonal properties of the thyrotropin-releasing hormone and pyroglutamyl-histidyl-proline amide. *Biochem. Biophys. Res. Commun.*, **37**, 705-710.
- Brazeau P, Vale W, Burgus R, Ling N, Butcher M, Rivier R, Guillemin R (1973): Hypothalamic polypeptide that inhibits the secretion of immunoreactive pituitary growth hormone. *Science*, **179**, 77-79.
- Burgus R, Dunn TF, Desiderio D, Guillemin R (1969): Structure moléculaire du facteur hypothalamique hypophysiotrope TRF d'origine ovine: mise en évidence par spectrométrie de masse de la séquence PCA-His-Pro-NH₂. *C.R. Acad. Sci. Paris (Ser. D)*, **269**, 1870-1873.
- Chang MM, Leeman SE, Niall HD (1971): Amino acid sequence of substance P. *Nature New Biol.*, **232**, 86-87.
- Du Vigneaud V, Lawler HC, Popenoe EA (1953a): Enzymatic cleavage of glycylamide from vasopressin and a proposed structure for the pressor-antidiuretic hormone of the posterior pituitary. *J. Am. Chem. Soc.*, **75**, 4880-4881.
- Du Vigneaud V, Ressler C, Swan JM, Roberts CW, Katsoyannis PG, Gordon S (1953b): The synthesis of an octapeptide amide with the hormonal activity of oxytocin. *J. Am. Chem. Soc.*, **75**, 4879-4880.
- Harris GH (1955): *Neural Control of the Pituitary Gland*. Edward Arnold, London.
- Hughes J, Smith TW, Kosterlitz HW, Fothergill LA, Morgan BA, Morris HR (1975): Identification of two related pentapeptides from the brain with potent opiate agonist activity. *Nature (London)*, **258**, 577-579.
- Kangawa K, Minamino N, Fukuda A, Matsuo H (1983): Neuromedin K: a novel mammalian tachykinin identified in porcine spinal cord. *Biochem. Biophys. Res. Commun.*, **114**, 533-540.
- Kimura S, Okada M, Sugita Y, Kanazawa I, Munekata E (1983): Novel neuropeptides, neurokinin α and β , isolated from porcine spinal cord. *Proc. Jap. Acad.*, **59**, 101.
- Luft R, Efendic S, Hökfelt T, Johansson O, Arimura A (1974): Immunohistochemical evidence for the localization of somatostatin-like immunoreactivity in a cell population of the pancreatic islets. *Med. Biol.*, **52**, 428-430.
- Matsuo H, Arimura A, Nair RMG, Schally AV (1971): Synthesis of the porcine LH and FSH releasing hormone by the solid phase method. *Biochem. Biophys. Res. Commun.*, **45**, 822-827.
- Minamino N, Kangawa K, Fukuda A, Matsuo H (1984): A novel mammalian tachykinin identified in porcine spinal cord. *Neuropeptides*, **4**, 157-166.

Preface

- Mutt V, Jorpes JE (1968): Structure of porcine cholecystokinin-pancreozymin. I. Cleavage with thrombin and with trypsin. *Eur. J. Biochem.*, **6**, 156-162.
- Nawa H, Hirose T, Takashima H, Ianyama S, Nakanishi S (1983): Nucleotide sequences of cloned cDNAs for two types of bovine brain substance P precursor. *Nature (London)*, **306**, 32-36.
- Pernow B (1953): Studies on substance P: purification, occurrence and biological actions. *Acta Physiol. Scand.*, **29 (Suppl.)**, 1-105.
- Pernow B (1983): Substance P. *Pharmacol. Rev.*, **35**, 85-141.
- Rosenfeld MG, Mermod J-J, Amara SG, Swanson LW, Sawchenko PE, Rivier J, Vale WW, Evans RM (1983): Production of a novel neuropeptide encoded by the calcitonin gene via tissue-specific RNA processing. *Nature (London)*, **304**, 129-135.
- Schally AV, Arimura A, Baba Y, Nair RMG, Matsuo J, Redding TW, Debeljuk L, White WF (1971): Isolation and properties of the FSH- and LH-releasing hormone. *Biochem. Biophys. Res. Commun.*, **42**, 393-399.
- Snyder SH (1980): Brain peptides as neurotransmitters. *Science*, **209**, 976-983.
- Tatemoto K, Mutt V (1980): Isolation of two novel candidate hormones using a chemical method for finding naturally occurring polypeptides. *Nature (London)*, **285**, 417-418.
- Tatemoto K, Lundberg JM, Jörnvall M, Mutt V (1985): Neuropeptide K: isolation, structure and biological activities of a novel brain tachykinin. *Nature (London)*, submitted for publication.
- Vanderhaeghen JJ, Signeau JC, Gepts W (1975): New peptide in the vertebrate CNS reacting with anti-gastrin antibodies. *Nature (London)*, **257**, 604-605.
- Von Euler US, Gaddum JH (1931): An unidentified depressor substance in certain tissue extracts. *J. Physiol. (London)*, **72**, 74-87.
- Weber E, Evans CJ, Barchas JD (1983): Multiple endogenous ligands for opioid receptors. *TINS*, **6**, 333-336.

Contents

I. DISTRIBUTION OF NEUROPEPTIDES IN THE CENTRAL NERVOUS SYSTEM USING BIOCHEMICAL MICROMETHODS - M. PALKOVITS AND M.J. BROWNSTEIN

1. Introduction	1
2. Comments on the distributional maps	2
2.1. Luteinizing hormone releasing hormone (LH-RH)	2
2.2. Thyrotropin releasing hormone (TRH)	4
2.3. Corticotropin releasing hormone (CRF)	4
2.4. Somatostatin	4
2.5. Oxytocin	4
2.6. Vasopressin	4
2.7. ACTH	4
2.8. Melanocyte-stimulating hormones (MSH's)	5
2.9. β -Endorphin	5
2.10. Enkephalins	5
2.11. Dynorphins	5
2.12. Vasoactive intestinal polypeptide (VIP)	5
2.13. Cholecystokinin (CCK)	6
2.14. Bombesin	6
2.15. Substance P	6
2.16. Neurotensin	6
3. References	70

II. GENERAL MORPHOLOGICAL FEATURES OF PEPTIDERGIC NEURONS - V.M. PICKEL

1. Introduction	72
2. Light microscopy	73
2.1. Regional distribution	73
2.2. Cytology	73
3. Ultrastructural morphology and neuronal interactions	75
3.1. Perikarya and dendrites	75
Hypothalamic magnocellular neurons	75
Peptidergic neurons in striatum and neocortex	75
Dendritic localization	79
3.2. Axon terminals	80
Synaptic interaction with neurons	80
Vesicular localization	83
4. Non-neuronal associations	86
4.1. Vascular	86
4.2. Ventricular	87
5. Conclusion	87
6. References	87

Contents

III. VASOPRESSIN, OXYTOCIN AND THEIR RELATED NEUROPHYSINS – M.V. SOFRONIEW

1. Introduction	93
1.1. Historical perspective	93
1.2. Biosynthesis of vasopressin, oxytocin and their associated neurophysins	94
1.3. Methods of localizing vasopressin, oxytocin or neurophysin	94
Immunohistochemistry	94
Radioimmunoassay	95
2. Distribution of neurons producing vasopressin, oxytocin and neurophysin	95
2.1. Hypothalamic magnocellular neurons	110
Supraoptic nucleus	111
Paraventricular nucleus	111
Accessory magnocellular groups	114
2.2. Parvo- and mediocellular neurons	115
Septal region	115
Bed nucleus (n. interstitialis) of the stria terminalis	115
Suprachiasmatic nucleus	117
Hypothalamic paraventricular nucleus	117
Posterior hypothalamus	117
Medial amygdala	117
Locus ceruleus	117
2.3. Comparative differences between mammalian species	119
2.4. Co-localization of other substances in vasopressin or oxytocin neurons	120
Magnocellular supraoptic and paraventricular nucleus neurons	120
Parvocellular paraventricular nucleus neurons	121
Locus ceruleus	122
3. Distribution of fibers and terminals	122
3.1. Forebrain	122
Neocortex, claustrum and related areas	122
Olfactory bulb, olfactory nuclei and olfactory tubercle	125
Septal region	125
Preoptic area and hypothalamus	125
Thalamus	126
Amygdala	128
Hippocampus and entorhinal cortex	128
Organum vasculosum of the lamina terminalis (OLVT) and subfornical organ (SFO)	128
Other forebrain areas	130
3.2. Brainstem	130
Tegmentum and colliculi	130
Central gray	131
Raphe nuclei	131
Parabrachial region and locus ceruleus	131
Nucleus tractus solitarius and dorsal vagal nucleus	132

Lateral reticular nucleus	132
Substantia gelatinosa trigemini	133
Other brainstem regions	134
3.3. Spinal cord	135
4. Nature of terminals in target areas	136
4.1. Vascular targets	136
4.2. Neural targets	137
5. Experimentally established projections	138
5.1. Vascular projections	138
Posterior pituitary	139
Median eminence	139
Other circumventricular organs (CVO's)	141
5.2. Neural projections	141
Projections to the medulla oblongata and spinal cord	142
Projections arising from parvocellular vasopressin neurons in the suprachiasmatic nucleus	144
Projections arising from parvocellular vasopressin neurons in the bed nucleus of the stria terminalis (BNST)	145
6. Vasopressin, oxytocin and neurophysin in cerebrospinal fluid	145
7. Functional considerations	146
7.1. Electrophysiology of vasopressin and oxytocin neurons	147
7.2. Release of vasopressin and oxytocin from nerve terminals	147
7.3. Authenticity of central vasopressin and oxytocin	147
7.4. Evidence for central vasopressin and oxytocin receptors	148
7.5. Effects of vasopressin and oxytocin at the cellular level	149
7.6. Neuroanatomical potential for interaction with other neuroactive substances	149
7.7. Vasopressin, oxytocin and behavior	149
7.8. Vasopressin, oxytocin and central cardiovascular regulation	150
7.9. Central autonomic and neuroendocrine regulation	150
7.10. Vasopressin, oxytocin and pain	151
7.11. Vasopressin and fever	151
7.12. Vasopressin, suprachiasmatic nucleus and biological rhythms	151
8. Summary and concluding remarks	152
9. Acknowledgments	152
10. References	152

IV. LHRH-CONTAINING SYSTEMS—J. BARRY, G.E. HOFFMAN AND S. WRAY

1. Introduction	166
1.1. LHRH immunocytochemistry	166
2. The atlas of rat LHRH	168
3. LHRH cells	179
3.1. Morphology of reactive LHRH perikarya	179
Mammals	179
Birds	180
Reptiles and amphibians	180

Contents

Fishes	182
Ultrastructure of LHRH cells	182
3.2. Topography of reactive LHRH perikarya	182
Mammalian	182
Birds	184
Reptiles	184
Amphibians	184
Fishes	184
4. LHRH pathways	185
4.1. Hypothalamic reactive LHRH pathways	185
Infundibular tracts	185
Preoptico-terminal tract	189
Preoptico-supraoptic tract	192
Ventricular tract	192
Intrahypothalamic tracts	193
4.2. Extrahypothalamic LHRH projections	193
Olfactory projections	193
Other limbic system projections	195
Centers controlling reproductive behavior	202
Other projections	205
5. Concluding remarks	205
6. Acknowledgments	207
7. References	207
V. β -ENDORPHIN, α -MSH, ACTH, AND RELATED PEPTIDES—	
H. KHACHATURIAN, M.E. LEWIS, KANG TSOU AND S.J. WATSON	
1. Introduction	216
2. Background	217
2.1. Pituitary gland	218
2.2. Central nervous system	221
3. Immunocytochemistry of POMC peptides	224
3.1. Anatomical distribution of neuronal perikarya	224
3.2. Projectional systems	226
Rostral projections	226
Lateral projections	229
Caudal projections	231
3.3. Distribution of fibers and terminals	234
4. The non-POMC α -MSH-like immunoreactive system in brain	242
4.1. Distribution of α -2 perikarya	242
4.2. The α -2 projections	243
5. Physiology and pathophysiology of POMC peptides	244
5.1. Analgesia	244
5.2. Cardiovascular regulation	245
5.3. Respiratory regulation	245
5.4. Neuroendocrine regulation	245
5.5. Thermoregulation	246
5.6. Consummatory behavior	247
5.7. Sexual behavior	248

5.8. Aggressive and defensive behavior	248
5.9. Locomotion and grooming	248
5.10. Mediation of endogenous reward	249
5.11. Learning and memory	249
5.12. Toward an integrative physiology of brain POMC systems	250
6. Acknowledgments	264
7. References	264
VI. DISTRIBUTION OF ENKEPHALIN-CONTAINING NEURONS IN THE CENTRAL NERVOUS SYSTEM – P. PETRUSZ, I. MERCHENTHALER AND J.L. MADERDRUT	
1. Introduction	273
2. Materials and methods	276
2.1. Sources of tissues	276
2.2. Preparation of tissues	276
2.3. Primary antisera	277
2.4. Immunocytochemical methods	277
2.5. Controls and definitions	278
3. Results	279
3.1. General	279
3.2. Telencephalon	280
3.3. Diencephalon	287
Thalamus	287
Hypothalamus	288
Posterior pituitary	295
Subthalamus and metathalamus	295
3.4. Mesencephalon	295
3.5. Metencephalon	296
3.6. Myelencephalon	297
3.7. Spinal cord	299
3.8. Comparison of the distribution of proenkephalin-, prodynorphin- and proopioidmelanocortin-derived peptides	303
4. Discussion	308
4.1. Comparison with earlier immunocytochemical studies	308
4.2. Enkephalin-containing pathways	310
4.3. Co-localization of enkephalins with other putative transmitters	314
4.4. Relationship of enkephalin-containing pathways to the localization of opiate receptors and biosynthetic enzymes	317
4.5. Distribution of enkephalin-containing neurons in submammalian vertebrates	319
5. Conclusions	322
6. Acknowledgments	323
7. References	323
VII. VIP-CONTAINING NEURONS – G.M. ABRAMS, G. NILAVER AND E.A. ZIMMERMAN	
1. Introduction	335

Contents

2. Anatomy	337
2.1. General overview	337
2.2. Radioimmunoassay	338
2.3. Immunocytochemistry	338
3. Localization of cell bodies	344
3.1. Cerebral cortex	344
3.2. Hippocampus	344
3.3. Amygdala	346
3.4. Other telencephalic areas	346
3.5. Hypothalamus	346
3.6. Mesencephalon	346
3.7. Other non-telencephalic areas	346
4. Fibers	349
4.1. Cortex	349
4.2. Hippocampus	349
4.3. Amygdala	349
4.4. Hypothalamus	349
4.5. Striatum	350
4.6. Thalamus	350
4.6. Mesencephalon	350
4.8. Other non-telencephalic areas	350
5. Non-neural innervation	351
6. Conclusions	351
7. Acknowledgments	351
8. References	351

VIII. NEUROTENSIN-CONTAINING NEURONS – P.C. EMSON, M. GOEDERT AND P.W. MANTYH

1. Introduction	355
2. Structure-activity relationships and mechanism of action of neurotensin	357
3. Neurotensin in the central nervous system	357
3.1. Retina	359
3.2. Olfactory system	360
3.3. Septum, nucleus accumbens and basal forebrain	365
3.4. Amygdala, cingulate gyrus, cerebral cortex and hippocampus	372
3.5. Caudate-putamen, substantia nigra and ventral mesencephalon	373
3.6. Hypothalamus and thalamus	382
3.7. Brainstem and spinal cord	387
4. Neurotensin in peripheral tissues	391
4.1. Gastrointestinal tract	392
4.2. Anterior pituitary gland	395
4.3. Adrenal gland	397
4.4. Other tissues	397
5. Discussion	398
6. Acknowledgments	399
7. References	400

IX. NEURONAL CHOLECYSTOKININ – J.J. VANDERHAEGHEN

1. Introduction	406
2. Materials and methods	406
2.1. Production of antisera	407
2.2. Preparation of the tissue extracts	407
2.3. Radioimmunoassay procedure	407
2.4. Specificity of antisera	407
2.5. Immunohistochemical staining procedures	408
Experimental animals	408
Tissue preparation	408
Staining methods	408
Controls	408
3. Distribution of gastrin and cholecystokinin in the central nervous system and hypophysis studied by radioimmunoassay – results and discussion	409
3.1. CCK in the nervous system of vertebrates and invertebrates	409
3.2. CCK in the developing nervous system	409
3.3. CCK distribution in rat brain	410
3.4. G-CCK in posterior hypophysis	411
3.5. G-CCK in adenohypophysis	411
3.6. CCK distribution in human brain	411
4. Distribution of gastrin and cholecystokinin in the central nervous system and hypophysis studied by immunohistochemistry – results and discussion	412
4.1. Retina	412
4.2. Olfactory bulbs, nuclei and tracts	412
Nuclei olfactorii	412
Main olfactory bulbs	412
Lamina medullaris interna bulbi olfactorii	413
4.3. Cerebral cortex	413
Neocortex	413
Cingulate gyri, prepyriform, periamygdaloid and entorhinal cortex	415
Hippocampus	416
4.4. Claustrum	416
4.5. Amygdaloid complex	416
4.6. Basal ganglia	416
Caudate-putamen	416
Nucleus accumbens	416
Nucleus interstitialis stria terminalis	416
Nucleus septi lateralis	417
4.7. Thalamus	417
4.8. Prehypothalamic structures	417
4.9. Hypothalamus	417
Magnocellular neurons	417
Median eminence	417
Periventricular nucleus	418
Dorsomedial and ventromedial nuclei	419
Suprachiasmatic nuclei	419