

SECOND EDITION

STATISTICAL MECHANICS

KERSON HUANG

SECOND EDITION

STATISTICAL MECHANICS

KERSON HUANG

Massachusetts Institute of Technology

JOHN WILEY & SONS

New York • Chichester • Brisbane • Toronto • Singapore

Copyright © 1963, 1987, by John Wiley & Sons, Inc.

All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of this work beyond that permitted by Sections 107 and 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful. Requests for permission or further information should be addressed to the Permissions Department, John Wiley & Sons.

Library of Congress Cataloging-in-Publication Data:

Huang, Kerson, 1928-

Statistical mechanics.

Bibliography: p.

Includes indexes.

1. Statistical mechanics. I. Title.

QC174.8.H83 1987 530.1'3 86-32466

ISBN 0-471-81518-7

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

FROM THE PREFACE TO THE FIRST EDITION

This book is an outgrowth of a year course in statistical mechanics that I have been giving at the Massachusetts Institute of Technology. It is directed mainly to graduate students in physics.

In this book the starting point of statistical mechanics is taken to be certain phenomenological postulates, whose relation to quantum mechanics I try to state as clearly as I can, and whose physical consequences I try to derive as simply and directly as I can.

Before the subject of statistical mechanics proper is presented, a brief but self-contained discussion of thermodynamics and the classical kinetic theory of gases is given. The order of this development is imperative, from a pedagogical point of view, for two reasons. First, thermodynamics has successfully described a large part of macroscopic experience, which is the concern of statistical mechanics. It has done so not on the basis of molecular dynamics but on the basis of a few simple and intuitive postulates stated in everyday terms. If we first familiarize ourselves with thermodynamics, the task of statistical mechanics reduces to the explanation of thermodynamics. Second, the classical kinetic theory of gases is the only known special case in which thermodynamics can be derived nearly from first principles, i.e., molecular dynamics. A study of this special case will help us understand why statistical mechanics works.

A large part of this book is devoted to selected applications of statistical mechanics. The selection is guided by the interest of the topic to physicists, its value as an illustration of calculating techniques, and my personal taste.

To read the first half of the book the reader needs a good knowledge of classical mechanics and some intuitive feeling for thermodynamics and kinetic theory. To read the second half of the book he needs to have a working knowledge of quantum mechanics. The mathematical knowledge required of the reader does not exceed what he should have acquired in his study of classical mechanics and quantum mechanics.

KERSON HUANG

*Cambridge, Massachusetts
February 1963*

PREFACE

This book is intended for use as a textbook for a one-year graduate course in statistical mechanics, and as a reference book for workers in the field.

Significant changes and additions have been made at various places in the text and in the problems to make the book more instructive. The general approach remains the same as that stated in the preface to the first edition.

The most substantive change is the addition of the last three chapters on the Landau-Wilson approach to critical phenomena. I hope that these chapters will provide the uninitiated reader with an introduction to this fascinating and important subject, which has developed since the first edition of this book.

To make room for the additions, I have omitted or abridged some of the original material, notably the sections on many-body theory and "rigorous" statistical mechanics, which by now have become separate fields.

All the material in this book probably cannot be covered in a one-year course (even if that were deemed desirable). It might be helpful, therefore, to list the chapters that form the "hard core" of the book. They are the following: Chapters 3, 4, 6, 7, 8, 11, 12 (excepting 3.5, 7.6, 7.7, 7.8).

KERSON HUANG

Marblehead, Massachusetts

CONTENTS

PART A THERMODYNAMICS AND KINETIC THEORY 1

CHAPTER 1 THE LAWS OF THERMODYNAMICS 3

- 1.1 Preliminaries 3**
- 1.2 The First Law of Thermodynamics 7**
- 1.3 The Second Law of Thermodynamics 9**
- 1.4 Entropy 14**
- 1.5 Some Immediate Consequences of the Second Law 19**
- 1.6 Thermodynamic Potentials 22**
- 1.7 The Third Law of Thermodynamics 25**

CHAPTER 2 SOME APPLICATIONS OF THERMODYNAMICS 31

- 2.1 Thermodynamic Description of Phase Transitions 31**
- 2.2 Surface Effects in Condensation 35**
- 2.3 Van der Waals Equation of State 38**
- 2.4 Osmotic Pressure 43**
- 2.5 The Limit of Thermodynamics 48**

CHAPTER 3 THE PROBLEM OF KINETIC THEORY 52

- 3.1 Formulation of the Problem 52**
- 3.2 Binary Collisions 56**
- 3.3 The Boltzmann Transport Equation 60**
- 3.4 The Gibbsian Ensemble 62**
- 3.5 The BBGKY Hierarchy 65**

CHAPTER 4 THE EQUILIBRIUM STATE OF A DILUTE GAS 73

- 4.1 Boltzmann's H Theorem 73**
- 4.2 The Maxwell-Boltzmann Distribution 75**
- 4.3 The Method of the Most Probable Distribution 79**
- 4.4 Analysis of the H Theorem 85**
- 4.5 The Poincaré Cycle 90**

CHAPTER 5 TRANSPORT PHENOMENA 93

- 5.1 The Mean Free Path 93**
- 5.2 Effusion 95**
- 5.3 The Conservation Laws 96**
- 5.4 The Zero-Order Approximation 100**
- 5.5 The First-Order Approximation 104**

- 5.6 Viscosity 108
- 5.7 Viscous Hydrodynamics 111
- 5.8 The Navier-Stokes Equation 113
- 5.9 Examples in Hydrodynamics 117

PART B STATISTICAL MECHANICS 125

CHAPTER 6 CLASSICAL STATISTICAL MECHANICS 127

- 6.1 The Postulate of Classical Statistical Mechanics 127
- 6.2 Microcanonical Ensemble 130
- 6.3 Derivation of Thermodynamics 135
- 6.4 Equipartition Theorem 136
- 6.5 Classical Ideal Gas 138
- 6.6 Gibbs Paradox 140

CHAPTER 7 CANONICAL ENSEMBLE AND GRAND CANONICAL ENSEMBLE 143

- 7.1 Canonical Ensemble 143
- 7.2 Energy Fluctuations in the Canonical Ensemble 145
- 7.3 Grand Canonical Ensemble 149
- 7.4 Density Fluctuations in the Grand Canonical Ensemble 152
- 7.5 The Chemical Potential 154
- 7.6 Equivalence of the Canonical Ensemble and the Grand Canonical Ensemble 157
- 7.7 Behavior of $W(N)$ 161
- 7.8 The Meaning of the Maxwell Construction 163

CHAPTER 8 QUANTUM STATISTICAL MECHANICS 171

- 8.1 The Postulates of Quantum Statistical Mechanics 171
- 8.2 Density Matrix 174
- 8.3 Ensembles in Quantum Statistical Mechanics 176
- 8.4 The Third Law of Thermodynamics 178
- 8.5 The Ideal Gases: Microcanonical Ensemble 179
- 8.6 The Ideal Gases: Grand Canonical Ensemble 185
- 8.7 Foundations of Statistical Mechanics 189

CHAPTER 9 GENERAL PROPERTIES OF THE PARTITION FUNCTION 193

- 9.1 The Darwin-Fowler Method 193
- 9.2 Classical Limit of the Partition Function 199
- 9.3 Singularities and Phase Transitions 206
- 9.4 The Lee-Yang Circle Theorem 210

CHAPTER 10 APPROXIMATE METHODS 213

- 10.1** Classical Cluster Expansion 213
- 10.2** Quantum Cluster Expansion 220
- 10.3** The Second Virial Coefficient 224
- 10.4** Variational Principles 228
- 10.5** Imperfect Gases at Low Temperatures 230

CHAPTER 11 FERMI SYSTEMS 241

- 11.1** The Equation of State of an Ideal Fermi Gas 241
- 11.2** The Theory of White Dwarf Stars 247
- 11.3** Landau Diamagnetism 253
- 11.4** The De Haas-Van Alphen Effect 260
- 11.5** The Quantized Hall Effect 261
- 11.6** Pauli Paramagnetism 267
- 11.7** Magnetic Properties of an Imperfect Gas 272

CHAPTER 12 BOSE SYSTEMS 278

- 12.1** Photons 278
- 12.2** Phonons in Solids 283
- 12.3** Bose-Einstein Condensation 286
- 12.4** An Imperfect Bose Gas 294
- 12.5** The Superfluid Order Parameter 298

PART C SPECIAL TOPICS IN STATISTICAL MECHANICS 309**CHAPTER 13 SUPERFLUIDS 307**

- 13.1** Liquid Helium 307
- 13.2** Tisza's Two-Fluid Model 311
- 13.3** The Bose-Einstein Condensate 313
- 13.4** Landau's Theory 315
- 13.5** Superfluid Velocity 317
- 13.6** Superfluid Flow 321
- 13.7** The Phonon Wave Function 325
- 13.8** Dilute Bose Gas 329

CHAPTER 14 THE ISING MODEL 341

- 14.1** Definition of the Ising Model 341
- 14.2** Equivalence of the Ising Model to Other Models 344
- 14.3** Spontaneous Magnetization 348
- 14.4** The Bragg-Williams Approximation 352
- 14.5** The Bethe-Peierls Approximation 357
- 14.6** The One-Dimensional Ising Model 361

CHAPTER 15 THE ONSAGER SOLUTION 368**15.1 Formulation of the Two-Dimensional Ising Model 368****15.2 Mathematical Digression 374****15.3 The Solution 378****CHAPTER 16 CRITICAL PHENOMENA 392****16.1 The Order Parameter 392****16.2 The Correlation Function and the Fluctuation-Dissipation Theorem 394****16.3 Critical Exponents 396****16.4 The Scaling Hypothesis 399****16.5 Scale Invariance 403****16.6 Goldstone Excitations 406****16.7 The Importance of Dimensionality 407****CHAPTER 17 THE LANDAU APPROACH 416****17.1 The Landau Free Energy 416****17.2 Mathematical Digression 418****17.3 Derivation in Simple Models 420****17.4 Mean-Field Theory 422****17.5 The Van der Waals Equation of State 426****17.6 The Tricritical Point 428****17.7 The Gaussian Model 434****17.8 The Ginzburg Criterion 437****17.9 Anomalous Dimensions 438****CHAPTER 18 RENORMALIZATION GROUP 441****18.1 Block Spins 441****18.2 The One-Dimensional Ising Model 443****18.3 Renormalization-Group Transformation 446****18.4 Fixed Points and Scaling Fields 449****18.5 Momentum-Space Formulation 452****18.6 The Gaussian Model 455****18.7 The Landau-Wilson Model 458****APPENDIX N-BODY SYSTEM OF IDENTICAL PARTICLES 468****A.1 The Two Kinds of Statistics 468****A.2 N-Body Wave Functions 470****A.3 Method of Quantized Fields 477****A.4 Longitudinal Sum Rules 484****INDEX 487**

A

THERMODYNAMICS AND KINETIC THEORY

THE LAWS OF THERMODYNAMICS

1.1 PRELIMINARIES

Thermodynamics is a phenomenological theory of matter. As such, it draws its concepts directly from experiments. The following is a list of some working concepts which the physicist, through experience, has found it convenient to introduce. We shall be extremely brief, as the reader is assumed to be familiar with these concepts.

- (a) A *thermodynamic system* is any macroscopic system.
- (b) *Thermodynamic parameters* are measurable macroscopic quantities associated with the system, such as the pressure P , the volume V , the temperature T , and the magnetic field H . They are defined experimentally.
- (c) A *thermodynamic state* is specified by a set of values of all the thermodynamic parameters necessary for the description of the system.
- (d) *Thermodynamic equilibrium* prevails when the thermodynamic state of the system does not change with time.
- (e) The *equation of state* is a functional relationship among the thermodynamic parameters for a system in equilibrium. If P , V , and T are the thermodynamic parameters of the system, the equation of state takes the form

$$f(P, V, T) = 0$$

which reduces the number of independent variables of the system from three to two. The function f is assumed to be given as part of the specification of the system. It is customary to represent the state of such a system by a point in the three-dimensional P - V - T space. The equation of state then defines a surface in this space, as shown in Fig. 1.1. Any point lying on this surface represents a state in equilibrium. In




Fig. 1.1 Geometrical representation of the equation of state.

thermodynamics a state automatically means a state in equilibrium unless otherwise specified.

- (f) A *thermodynamic transformation* is a change of state. If the initial state is an equilibrium state, the transformation can be brought about only by changes in the external condition of the system. The transformation is *quasi-static* if the external condition changes so slowly that at any moment the system is approximately in equilibrium. It is *reversible* if the transformation retraces its history in time when the external condition retraces its history in time. A reversible transformation is quasi-static, but the converse is not necessarily true. For example, a gas that freely expands into successive infinitesimal volume elements undergoes a quasi-static transformation but not a reversible one.
- (g) The *P-V diagram* of a system is the projection of the surface of the equation of state onto the *P-V* plane. Every point on the *P-V* diagram therefore represents an equilibrium state. A reversible transformation is a continuous path on the *P-V* diagram. Reversible transformations of specific types give rise to paths with specific names, such as *isotherms*, *adiabatics*, etc. A transformation that is not reversible cannot be so represented.
- (h) The concept of *work* is taken over from mechanics. For example, for a system whose parameters are *P*, *V*, and *T*, the work dW done by a system in an infinitesimal transformation in which the volume increases by dV is given by

$$dW = PdV$$

- (i) *Heat* is what is absorbed by a homogeneous system if its temperature increases while no work is done. If ΔQ is a small amount of the heat absorbed, and ΔT is the small change in temperature accompanying the absorption of heat, the *heat capacity* *C* is defined by

$$\Delta Q = CAT$$

The heat capacity depends on the detailed nature of the system and is given as a part of the specification of the system. It is an experimental fact that, for the same ΔT , ΔQ is different for different ways of heating

up the system. Correspondingly, the heat capacity depends on the manner of heating. Commonly considered heat capacities are C_V and C_P , which respectively correspond to heating at constants V and P . Heat capacities per unit mass or per mole of a substance are called *specific heats*.

- (j) A *heat reservoir*, or simply *reservoir*, is a system so large that the gain or loss of any finite amount of heat does not change its temperature.
- (k) A system is *thermally isolated* if no heat exchange can take place between it and the external world. Thermal isolation may be achieved by surrounding a system with an *adiabatic wall*. Any transformation the system can undergo in thermal isolation is said to take place *adiabatically*.
- (l) A thermodynamic quantity is said to be *extensive* if it is proportional to the amount of substance in the system under consideration and is said to be *intensive* if it is independent of the amount of substance in the system under consideration. It is an important empirical fact that to a good approximation thermodynamic quantities are either extensive or intensive.
- (m) The *ideal gas* is an important idealized thermodynamic system. Experimentally all gases behave in a universal way when they are sufficiently dilute. The ideal gas is an idealization of this limiting behavior. The parameters for an ideal gas are pressure P , volume V , temperature T , and number of molecules N . The equation of state is given by Boyle's law:

$$\frac{PV}{N} = \text{constant} \quad (\text{for constant temperature})$$

The value of this constant depends on the experimental scale of temperature used.

- (n) The equation of state of an ideal gas in fact defines a temperature scale, the *ideal-gas temperature* T :

$$PV = NkT$$

where

$$k = 1.38 \times 10^{-16} \text{ erg/deg}$$

which is called Boltzmann's constant. Its value is determined by the conventional choice of temperature intervals, namely, the Centigrade degree. This scale has a universal character because the ideal gas has a universal character. The origin $T = 0$ is here arbitrarily chosen. Later we see that it actually has an absolute meaning according to the second law of thermodynamics.

To construct the ideal-gas temperature scale we may proceed as follows. Measure PV/Nk of an ideal gas at the temperature at which water boils and at




Fig. 1.2 The ideal-gas temperature scale.

which water freezes. Plot these two points and draw a straight line through them, as shown in Fig. 1.2. The intercept of this line with the abscissa is chosen to be the origin of the scale. The intervals of the temperature scale are so chosen that there are 100 equal divisions between the boiling and the freezing points of water. The resulting scale is the Kelvin scale (K). To use the scale, bring anything whose temperature is to be measured into thermal contact with an ideal gas (e.g., helium gas at sufficiently low density), measure PV/Nk of the ideal gas, and read off the temperature from Fig. 1.2. An equivalent form of the equation of state of an ideal gas is

$$PV = nRT$$

where n is the number of moles of the gas and R is the gas constant:

$$\begin{aligned} R &= 8.315 \text{ joule/deg} \\ &= 1.986 \text{ cal/deg} \\ &= 0.0821 \text{ liter-atm/deg} \end{aligned}$$

Its value follows from the value of Boltzmann's constant and Avogadro's number:

$$\text{Avogadro's number} = 6.205 \times 10^{23} \text{ atoms/mol}$$

Most of these concepts are properly understood only in molecular terms. Here we have to be satisfied with empirical definitions.

In the following we introduce thermodynamic laws, which may be regarded as mathematical axioms defining a mathematical model. It is possible to deduce rigorous consequences of these axioms, but it is important to remember that this model does not rigorously correspond to the physical world, for it ignores the atomic structure of matter, and will thus inevitably fail in the atomic domain. In the macroscopic domain, however, thermodynamics is both powerful and useful. It enables one to draw rather precise and far-reaching conclusions from a few seemingly commonplace observations. This power comes from the implicit assumption that the equation of state is a regular function, for which the thermodynamic laws, which appear to be simple and naive at first sight, are in fact enormously restrictive.

1.2 THE FIRST LAW OF THERMODYNAMICS

In an *arbitrary* thermodynamic transformation let ΔQ denote the net amount of heat absorbed by the system and ΔW the net amount of work done by the system. The first law of thermodynamics states that the quantity ΔU , defined by

$$\Delta U = \Delta Q - \Delta W \quad (1.1)$$

is the same for all transformations leading from a given initial state to a given final state.

This immediately defines a state function U , called the internal energy. Its value for any state may be found as follows. Choose an arbitrary fixed state as reference. Then the internal energy of any state is $\Delta Q - \Delta W$ in *any* transformation which leads from the reference state to the state in question. It is defined only up to an arbitrary additive constant. Empirically U is an extensive quantity. This follows from the saturation property of molecular forces, namely, that the energy of a substance is doubled if its mass is doubled.

The experimental foundation of the first law is Joule's demonstration of the equivalence between heat and mechanical energy—the feasibility of converting mechanical work completely into heat. The inclusion of heat as a form of energy leads naturally to the inclusion of heat in the statement of the conservation of energy. The first law is precisely such a statement.

In an infinitesimal transformation, the first law reduces to the statement that the differential

$$dU = dQ - dW \quad (1.2)$$

is exact. That is, there exists a function U whose differential is dU ; or, the integral $\int dU$ is independent of the path of the integration and depends only on the limits of integration. This property is obviously not shared by dQ or dW .

Given a differential of the form $df = g(A, B) dA + h(A, B) dB$, the condition that df be exact is $\partial g / \partial B = \partial h / \partial A$. Let us explore some of the consequences of the exactness of dU . Consider a system whose parameters are P, V, T . Any pair of these three parameters may be chosen to be the independent variables that completely specify the state of the system. The other parameter is then determined by the equation of state. We may, for example, consider $U = U(P, V)$. Then*

$$dU = \left(\frac{\partial U}{\partial P} \right)_V dP + \left(\frac{\partial U}{\partial V} \right)_P dV \quad (1.3)$$

The requirement that dU be exact immediately leads to the result

$$\frac{\partial}{\partial V} \left[\left(\frac{\partial U}{\partial P} \right)_V \right]_P = \frac{\partial}{\partial P} \left[\left(\frac{\partial U}{\partial V} \right)_P \right]_V \quad (1.4)$$

The following equations, expressing the heat absorbed by a system during an infinitesimal reversible transformation (in which $dW = P dV$), are easily ob-

*The symbol $(\partial U / \partial P)_V$ denotes the partial derivative of U with respect to P , with V held constant.

tained by successively choosing as independent variables the pairs (P, V) , (P, T) , and (V, T) :

$$dQ = \left(\frac{\partial U}{\partial P} \right)_V dP + \left[\left(\frac{\partial U}{\partial V} \right)_P + P \right] dV \quad (1.5)$$

$$dQ = \left[\left(\frac{\partial U}{\partial T} \right)_P + P \left(\frac{\partial V}{\partial T} \right)_P \right] dT + \left[\left(\frac{\partial U}{\partial P} \right)_T + P \left(\frac{\partial V}{\partial P} \right)_T \right] dP \quad (1.6)$$

$$dQ = \left(\frac{\partial U}{\partial T} \right)_V dT + \left[\left(\frac{\partial U}{\partial V} \right)_T + P \right] dV \quad (1.7)$$

Called dQ equations, these are of little practical use in their present form, because the partial derivatives that appear are usually unknown and inaccessible to direct measurement. They will be transformed to more useful forms when we come to the second law of thermodynamics.

It can be immediately deduced from the dQ equations that

$$C_V \equiv \left(\frac{\Delta Q}{\Delta T} \right)_V = \left(\frac{\partial U}{\partial T} \right)_V \quad (1.8)$$

$$C_P \equiv \left(\frac{\Delta Q}{\Delta T} \right)_P = \left(\frac{\partial H}{\partial T} \right)_P \quad (1.9)$$

where $H = U + PV$ is called the enthalpy of the system.

We consider the following examples of the application of the first law.

(a) *Analysis of Joule's Free-Expansion Experiment.* The experiment in question concerns the free expansion of an ideal gas into a vacuum. The initial and final situations are illustrated in Fig. 1.3.

Experimental Finding. $T_1 = T_2$.

Deductions. $\Delta W = 0$, since the gas performs no work on its external surrounding. $\Delta Q = 0$, since $\Delta T = 0$. Therefore, $\Delta U = 0$ by the first law.




Fig. 1.3 Joule's free-expansion experiment.