

---

**IMMUNOCHEMICAL TECHNIQUES  
FOR THE IDENTIFICATION AND  
ESTIMATION OF MACROMOLECULES**

---

2nd Fully Revised Edition

J. Clausen

---

# IMMUNOCHEMICAL TECHNIQUES FOR THE IDENTIFICATION AND ESTIMATION OF MACROMOLECULES

---

2nd Fully Revised Edition

J. Clausen

*The Neurochemical Institute  
58 Rådmandsgade  
Copenhagen  
and  
Laboratory for  
Biochemistry and  
Toxicology  
Roskilde University  
4000 Roskilde  
Denmark*



1981

ELSEVIER/NORTH-HOLLAND BIOMEDICAL PRESS  
AMSTERDAM • NEW YORK • OXFORD

© Elsevier/North-Holland Biomedical Press, 1981

*All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright owner*

Library of Congress Catalog Card Number: 68-54514

ISBN for the series: 0 7204 4200 1

Vol. 1, part 3, paperback: 0-444-80244-4

Vol. 1, part 3, hardbound: 0-444-80245-2

1st edition: 1969

2nd edition: 1981

*Published by:*

Elsevier/North-Holland Biomedical Press

1 Molenwerf

1014 AG Amsterdam

The Netherlands

*Sole distributors for the U.S.A. and Canada:*

Elsevier/North-Holland Inc.

52 Vanderbilt Avenue

New York, N.Y. 10017

---

**This book is the revised edition of Volume 1, Part III, of the series 'Laboratory Techniques in Biochemistry and Molecular Biology'.**

**Volume I of the series contains the following parts:**

**Part I** Electrophoresis of proteins in polyacrylamide and starch gels  
by A.H. Gordon – 2nd Edition 1978

**Part II** An introduction to gel chromatography  
by L. Fischer – 2nd Edition 1980 (Gel filtration chromatography)

**Part III** Immunochemical techniques for the identification and estimation of macromolecules  
by J. Clausen – 2nd Edition 1981

---

*Printed in The Netherlands*

## Preface

The aim of this manual is to present practical advice on application of the rapidly expanding mass of data on immunochemistry, and more specifically to examine the possibilities of employing immunological microtechniques in biochemical and histochemical analyses. The terminology in immunochemistry and protein chemistry has been bewildering (Grabar 1963); however, in 1964 WHO and in 1975 The International Union of Immunological Sciences recommended rules on the nomenclature of immunoglobulins (cf. p. 9). The World Health Organization terminology has been used throughout the manual, although this terminology has not, however, been generally accepted in all laboratories working in the field of immunochemistry (WHO 1964).

It is not the object of this manual to present an entire and comprehensive work in the field of immunochemistry, competing with the classical work of Kabat and Mayer (1961), but rather to describe the practical application of immunochemical analysis for laboratory use and especially to stress for the reader the possible errors and misinterpretations.

It is a pleasure to participate in creating this series of 'Laboratory Manuals' in collaboration with Drs. T.S. and E. Work.

The rapid expansion in the field of immunochemistry has necessitated the publication of a new edition of the present manual. Several chapters i.e. those dealing with lectins, HLA-antigens and with the application of crossed immunoelectrophoresis for characterization of membrane proteins, have been extensively revised and extended. Furthermore, new chapters dealing with the complement system, enzyme immuno-assay systems, immuno-fluorescence methods and immunoadsorbent techniques have been added and the recent developments in the methodology

already presented in the first edition have been included.

Cellular immunity has been described as a basis for the use of lymphocytic systems for the tracing of antigens giving rise to delayed hypersensitivity reactions.

Gratitude is expressed to Doctors T.S. and E. Work for their critical evaluation.

JØRGEN CLAUSEN

Copenhagen 1981

# Contents

<i>Preface</i> . . . . .	xii
<i>Chapter 1. Introduction</i> . . . . .	1
1.1. The antigen-antibody reaction . . . . .	1
1.1.1. Physicochemical aspects of the antigen-antibody reaction . . . . .	1
1.2. Antigen-antibody reaction in gels . . . . .	4
1.3. Antibodies; the immunoglobulins . . . . .	8
1.4. Antigens . . . . .	18
1.4.1. Polysaccharide antigens . . . . .	18
1.4.2. Glycopeptide antigens . . . . .	19
1.4.3. Glycoprotein and histocompatibility antigens . . . . .	21
1.4.4. Lipopolysaccharide and lipoprotein antigens . . . . .	22
1.4.4.1. Bacterial cell-wall components . . . . .	23
1.4.5. Protein antigens . . . . .	28
1.4.6. Virus antigens . . . . .	29
1.5. Immune defence reactions . . . . .	29
1.6. Lectins . . . . .	40
1.7. The complement system . . . . .	41
Guide to further reading . . . . .	48
<i>Chapter 2. Immunodiffusion in gels</i> . . . . .	49
2.1. Diffusion in one dimension . . . . .	49
2.1.1. Single (simple) diffusion in one dimension (Oudin's technique) . . . . .	49
2.1.2. Single radial diffusion (Petrie plate technique) . . . . .	50
2.2. Double diffusion . . . . .	52
2.2.1. Double diffusion in one dimension . . . . .	52
2.2.2. Double diffusion in two dimensions . . . . .	53
2.3. Determination of diffusion constants . . . . .	54

2.4. Immunoelectrophoresis . . . . .	59
2.4.1. General principles . . . . .	59
2.4.2. Types of supporting media . . . . .	60
2.4.2.1. Agar (agarose and agaro-pectin) . . . . .	60
2.4.2.2. Cellulose acetate. . . . .	61
2.4.2.3. Starch gel. . . . .	62
2.4.2.4. Polyacrylamide gel . . . . .	62
2.4.2.5. Fibrin-agar . . . . .	63
2.4.3. Electrophoresis in antibody-containing media (Laurell technique) . . . . .	63
2.4.3.1. General Aspects . . . . .	63
2.4.3.2. Crossed and rocket immunoelectrophoresis and related techniques . . . . .	65
<i>Chapter 3. Antisera</i> . . . . .	72
3.1. Preparation and choice of antigen . . . . .	73
3.1.1. General principles . . . . .	73
3.1.2. Whole organ antigens . . . . .	74
3.1.3. Haptens. . . . .	75
3.2. Preparation of antiserum . . . . .	77
3.3. Use of adjuvants . . . . .	79
3.4. Absorption of antisera . . . . .	79
3.5. Monoclonal antibodies (the hybridoma technology) . . . . .	80
<i>Chapter 4. Techniques of immunodiffusion</i> . . . . .	82
4.1. Choice of method and its limitations . . . . .	82
4.2. Choice of antiserum . . . . .	83
4.3. Choice of amount of antigen . . . . .	83
4.4. Choice of hole patterns in double diffusion . . . . .	84
4.5. Quantitative methods . . . . .	86
4.5.1. Single diffusion techniques. . . . .	86
4.5.2. (Semi-) quantitative estimation of antigens by double diffusion. . . . .	87
<i>Chapter 5. Techniques of immunoelectrophoresis</i> . . . . .	89
5.1. Choice of buffers . . . . .	89
5.2. Choice of method . . . . .	90
5.2.1. General considerations . . . . .	90
5.2.2. Choice of type of pattern. . . . .	92
5.2.2.1. Comparison of antigenic composition of two or more samples . . . . .	92
5.2.2.2. Comparison of composition of antibodies in different antisera . . . . .	93
5.2.2.3. Comparison of one or two antigen solutions with two antisera . . . . .	94
5.3. Antigens . . . . .	94
5.4. Reference markers. . . . .	96

5.5. Steps in electrophoresis and diffusion. . . . .	97
5.5.1. The electrophoretic stage. . . . .	97
5.5.2. Diffusion stage. . . . .	98
5.6. Quantitative methods . . . . .	99
5.6.1. Electrophoresis prior to immunoprecipitation. . . . .	99
5.6.2. Electrophoresis in a medium containing antibodies . . . . .	99
5.6.3. Crossed immuno-electrophoresis. . . . .	100
5.6.3.1. Estimation of antigens . . . . .	100
5.6.3.2. Identification of soluble protein antigens . . . . .	101
5.6.3.3. Identification and characterization of membrane proteins . . . . .	102
5.6.3.3.1. Analysis of (partially) intact membrane structures . . . . .	102
5.6.3.3.2. Solubilization of membrane proteins for immunochemical analysis . . . . .	104
5.6.3.3.3. Conditions for crossed immunoelectrophoretic characterization of membrane constituents . . . . .	106
<i>Chapter 6. Visualization and interpretation of precipitates in gels. . . . .</i>	<i>120</i>
6.1. Direct identification and interpretation of immunoprecipitates . . . . .	120
6.1.1. Identification by absorption of antisera . . . . .	120
6.1.2. Morphology of precipitates. . . . .	121
6.1.2.1. Double diffusion . . . . .	121
6.1.2.2. Immunoelectrophoresis. . . . .	123
6.2. Staining gels for proteins, lipids and carbohydrates . . . . .	127
6.3. Enzymic reactions in gels. . . . .	129
6.3.1. Hydrolases (EC 3). . . . .	130
6.3.1.1. Carboxylic esterases (EC 3.1.1). . . . .	130
6.3.1.2. Phosphatases (EC 3.1.3). . . . .	133
6.3.1.3. Proteolytic enzymes (EC 3.4). . . . .	133
6.3.1.4. Amylases (EC 3.2.1.1 and EC 3.2.1.2) and other glycoside hydrolases (EC 3.2.1.1). . . . .	134
6.3.2. Oxidoreductases (EC 1). . . . .	134
6.3.2.1. Oxidases (EC 1.4). . . . .	134
6.3.2.2. Peroxidases (EC 1.9 and EC 1.11). . . . .	135
6.3.2.3. Dehydrogenases with NAD <sup>+</sup> and NADP <sup>+</sup> as acceptor (EC 1.1.1). . . . .	135
6.4. Fluorescent techniques . . . . .	136
6.5. Isotopic methods . . . . .	136
<i>Chapter 7. Immunoabsorbent techniques. . . . .</i>	<i>139</i>
7.1. General considerations . . . . .	139
7.2. Formation of immunoabsorbents . . . . .	139
7.2.1. Protein copolymers . . . . .	139



7.2.2. Matrix-bound antigens and antibodies . . . . .	140
7.3. Evaluation of quality of immunoadsorbents . . . . .	143
7.4. Immunoadsorption by Staphylococcus A-antigen, with binding to Fc fragments. . . . .	144
<i>Chapter 8. Radio-immunochemical techniques . . . . .</i>	<i>146</i>
8.1. Radio-immunoassay . . . . .	147
8.1.1. Difficulties . . . . .	151
8.2. Double antibody precipitation in radioimmunoassay . . . . .	152
8.3. Radioimmunoassay of membrane-bound immunoglobulin . . . . .	153
8.4. Solid-phase radio-immunochemical techniques. . . . .	153
<i>Chapter 9. Enzyme- and metallo-immunoassays . . . . .</i>	<i>156</i>
9.1. Enzyme-immunoassay (EIA) techniques . . . . .	156
9.1.1. Histological localization . . . . .	157
9.1.2. Quantitative assays: General considerations . . . . .	157
9.1.2.1. Coupling of enzymes to proteins . . . . .	159
9.1.2.2. Coupling enzymes to haptens . . . . .	160
9.1.2.3. Purification and properties of conjugates. . . . .	161
9.1.2.4. Types of EIA assay . . . . .	161
9.1.2.4.1. Competitive assay systems . . . . .	162
9.1.2.4.2. Non-competitive assay systems (sandwich technique) . . . . .	163
9.1.2.4.3. Coenzyme labelling . . . . .	165
9.1.2.5. Choice of method of labelling . . . . .	166
9.1.2.6. The influence of the immune reaction on the enzymic activity. . . . .	166
9.1.2.7. Separation techniques . . . . .	167
9.1.2.8. The enzymic assay in EIA . . . . .	167
9.1.2.9. Artifacts in EIA . . . . .	168
9.1.2.10. Example of EIA: The sandwich procedure . . . . .	168
9.2. Metallo-immunoassay (MIA) . . . . .	170
9.2.1. Coupling with albumin and immunization . . . . .	171
9.2.2. Metal labelling . . . . .	172
9.2.3. Immunochemical procedure and chemical assay of MIA . . . . .	172
9.2.4. Precision of MIA. . . . .	172
<i>Chapter 10. Difficulties and artifacts arising in gel diffusion and in immunoelectrophoresis. . . . .</i>	<i>173</i>
10.1. The antibody-antigen ratio. . . . .	173
10.2. Position of immunoprecipitates . . . . .	173
10.3. Variation in agar . . . . .	174

10.4. Effects of molecular size on electrophoretic mobility . . . . .	175
10.5. Antigens . . . . .	176
10.6. Antisera . . . . .	177
10.7. Various artifacts . . . . .	178
10.7.1. Interaction of antigens with non-antibody substances . . . . .	178
10.7.2. Artifacts in double immunodiffusion techniques . . . . .	179
10.7.3. Abnormal binding of antigens to antibodies . . . . .	180
<i>Chapter 11. Immunochemical reactions in free buffer systems . . . . .</i>	<i>182</i>
11.1. Qualitative and structural analysis of antigens . . . . .	182
11.1.1. Analytical procedures . . . . .	182
11.1.2. Preparative procedures . . . . .	183
11.2. Quantitative analysis . . . . .	184
11.2.1. Direct precipitation . . . . .	184
11.2.2. Salting out procedures . . . . .	185
11.2.3. The quantitative precipitin reaction combined with tracer technique . . . . .	186
11.2.4. Laser scattering immunoassay . . . . .	186
11.2.5. Free-radical immunoassay techniques . . . . .	187
11.2.6. Phage-immunoassays . . . . .	188
11.2.7. Electro-immunoassay . . . . .	188
<i>Chapter 12. Immunofluorescence techniques . . . . .</i>	<i>189</i>
12.1. General principles . . . . .	189
12.2. Fluorescent markers . . . . .	189
12.3. Antibodies for immunofluorescence assays . . . . .	190
12.4. Immunofluorescence methods . . . . .	192
12.5. Equipment for immunofluorescence microscopy . . . . .	193
12.6. Artifacts (autofluorescence) . . . . .	196
12.7. Application of immunofluorescence methods . . . . .	196
12.8. Quantitative immunofluorescence . . . . .	198
<i>Chapter 13. Complement fixation assays . . . . .</i>	<i>201</i>
<i>Chapter 14. Application of lymphocytes for tracing antigens . . . . .</i>	<i>203</i>
14.1. General considerations . . . . .	204
14.2. Methods for assay of lymphocyte activation . . . . .	204
14.3. HL-antigens . . . . .	206
<i>Chapter 15. Miscellaneous topics in immunochemistry . . . . .</i>	<i>212</i>
15.1. Methods for elucidation of the conformational structure of proteins (inhibition assay of refolding) . . . . .	212

15.2. Determination of virus and tumor antigens . . . . .	214
15.2.1. Immunoprecipitation . . . . .	215
15.2.2. Cytotoxicity test . . . . .	216
15.2.3. Labelled antibody binding assay . . . . .	216
<i>Chapter 16. Appendices</i> . . . . .	217
1 Preparation of Freund's adjuvant . . . . .	217
2 Preparation of antisera and purification of antibodies . . . . .	218
3 Concentration of protein solutions (antigens and antibodies) . . . . .	222
4 Specific absorption of antisera . . . . .	225
5 Purification of agar . . . . .	225
6 Preparation of agarose . . . . .	226
7 Preparation of agar gel for immunodiffusion or immunoelectrophoresis . . . . .	227
8 Simple linear immunodiffusion (Oudin technique) . . . . .	228
I Single radial immunodiffusion in agar gel . . . . .	229
II Single radial immunodiffusion on cellulose acetate . . . . .	230
10 Preparation of agar slides for immunoelectrophoresis and double immunodiffusion . . . . .	231
11 Double immunodiffusion (Ouchterlony technique) . . . . .	233
12 Techniques for immunoelectrophoresis . . . . .	234
13 Immunoelectrophoresis in an antibody-containing medium for quantitative estimation of antigens (Laurell technique) . . . . .	235
14 Crossed immunoelectrophoresis . . . . .	237
15 Crossed immunoelectrophoresis in a system combining thin-layer electrofocussing in polyacrylamide gel with immunoelectrophoresis in an agarose gel . . . . .	240
16 Fibrin-agar micro electrophoresis . . . . .	242
17 Buffers . . . . .	243
18 Reference markers for immunoelectrophoresis . . . . .	245
19 Washing, drying, photographing and storing of completed slides or plates . . . . .	246
20 Protein stains . . . . .	247
21 Lipid stains . . . . .	248
22 Double stains for protein and lipid . . . . .	249
23 Polysaccharide stains . . . . .	250
24 Staining for oxidase activity (including ceruloplasmin) . . . . .	251
25 Staining for peroxidase activity . . . . .	252
26 Staining of lactate dehydrogenase isoenzymes and other NAD- or NADP-dependent dehydrogenases . . . . .	253
27 Staining for esterase activity . . . . .	253
28 Staining for phosphatase activity . . . . .	254
29 Staining for amylase activity and glycosidase activities . . . . .	255
30 Labelling of antibodies with fluorescein isocyanate . . . . .	256

31	Radio-immunoassay of protein or peptide hormones . . . . .	257
32	The double antibody technique for determination of insulin and other protein hormones . . . . .	259
33	Lactoperoxidase-catalyzed iodination of proteins . . . . .	264
34	'Sandwich' solid-phase radioimmunoassay, using plastic tubes or discs, for the estimation of human immunoglobulins. . . . .	265
35	Radioimmunoabsorbent assay, using antigen-conjugated paper discs to estimate allergic antibodies . . . . .	268
36	Radioimmunoabsorbent assay, using antiserum-coated plastic tubes to estimate carcinoembryonic antigen . . . . .	270
37	Immunoperoxidase histochemical technique . . . . .	271
38	Preparation and use of cross-linked protein immunoabsorbents . . . . .	274
	I Ethylchloroformate or glutaraldehyde as cross-linking agents . . . . .	274
	II Isolation of antigens or antibodies . . . . .	277
	III Properties of the various protein immunoabsorbents . . . . .	278
39	Hapten (steroid and other hormones) conjugation to macromolecules. . . . .	281
40	Enzyme-linked sandwich immunoassay of macromolecular antigens. . . . .	284
	I Antibody-coupled glass rod immunoabsorbent. . . . .	284
	II Antibody-coupled cellulose immunoabsorbent. . . . .	289
41	Use of an antigen-antibody precipitate as an intermediate in the purification of human $\alpha$ -foetoprotein. . . . .	294
42	Preparation and use of various immunoabsorbents for purification of antigens and antibodies . . . . .	295
	I Myoglobin-linked cellulose derivatives and other polymers . . . . .	295
	II Anti $\alpha$ -foetoprotein Sepharose immunoabsorbent . . . . .	298
43	Staphylococcal protein A-antibody adsorbent in the isolation of cell membrane antigens . . . . .	301
44	Immunochemical characterization of membrane constituents . . . . .	305
45	Immunofixation . . . . .	311
46	Quantitative micro-complement fixation and its use in the study of antigenic structure by specific antigen-antibody inhibition. . . . .	312
47	Isolation of lymphocytes . . . . .	318
48	Separation of B- and T-lymphocytes by column chromatography . . . . .	319
	I Immunoabsorbent columns . . . . .	320
	II Lectin-affinity chromatography . . . . .	323
49	Isolation of T- and B-lymphocytes by erythrocyte adsorption . . . . .	324
50	Assay of lymphocyte B- and T-cells by rosette formation . . . . .	326
51	Lymphocyte reactivity assay by [ $^3\text{H}$ ] thymidine incorporation. . . . .	328
52	Lymphocyte reactivity assay by [ $^3\text{H}$ ] myoinositol incorporation. . . . .	329
53	Assay of virus and tumour antigens by $^{51}\text{Cr}$ ium-release. . . . .	330
54	Assay of virus and tumour antigens by immunoprecipitation . . . . .	334
	<i>References</i> . . . . .	338
	<i>Subject index</i> . . . . .	362

## Introduction

### *1.1. The antigen—antibody reaction*

Since the initial discovery of infectious agents and of the acquired defense reactions towards these agents (Pasteur 1876; Von Behring and Kitasato 1890; Koch 1891; Metchnikoff 1892), followed by the demonstration of the presence of specific serum factors neutralizing the infectious agents (Ehrlich and Morgenroth 1900), immunochemistry has developed into a well-defined physicochemical discipline. This subject was initiated by the discovery of precipitating antibodies (Kraus, 1897) and greatly advanced by the discovery of the blood group antigens (Landsteiner and van Der Scheer, 1928). The discovery of the precipitin reaction arose from the demonstration of the precipitate formed when cell-free filtrates of typhoid cultures were mixed with the corresponding antiserum. This technique was made quantitative by Heidelberger and Kendall (1935), thus permitting detection of less than 0.1  $\mu\text{g}$  antigen in solution.

During recent years, precipitation reactions in which antigens and antibodies diffuse through and react in semi-solid matrices (e.g. agar gel) have become essential tools in biochemical analysis. These techniques generally fall into three distinct categories, simple diffusion, double diffusion and immunoelectrophoresis. Before describing them in detail, some understanding is required of the physicochemical factors involved.

#### *1.1.1. Physicochemical aspects of the antigen—antibody reaction*

As early as 1902 Arrhenius and Madsen described how the antigen—antibody reaction may be treated on the basis of the law of mass action:

the combination of antibody (Ab) and antigen (Ag) was considered as a reversible bimolecular reaction:





where  $k_1$  and  $k_2$  are the rate-constants of the forward and backward reactions. According to the law of mass action, the following relationship holds:

$$\frac{(\text{AbAg})}{(\text{Ab}) \times (\text{Ag})} = \frac{k_1}{k_2} = K \quad (1.2)$$

where (Ab), (Ag) and (AbAg) represent the concentration of reactants and products and  $K$  is the equilibrium or association constant. The determination of the value  $K$  for any particular antigen-antibody reaction at different temperatures is necessary for determination of thermodynamic constants. The thermodynamic data indicate the nature of the bonds established between the two reactants (see p. 4).

The quantitative nature of the antigen-antibody reaction was first exploited by Heidelberger and Kendall (1929, 1935) who found that it is possible to isolate and estimate the immunoprecipitate formed by adding increasing amounts of an antigen to a fixed amount of antibody. The amount of the precipitated complex increases with rising quantities of antigen up to a certain point. This region of increasing precipitation is referred to as the interval, or zone, of antibody excess. Above this point there is no further increase in precipitated complex as long as some degree of equivalence exists between the two reactants (zone of equivalence). With increasing quantities of added antigen, a so-called zone of inhibition of complex formation is reached (antigen excess): here, there is a progressive decrease in amount of antigen-antibody complex precipitated (fig. 1.1).

These results led to the postulate that the formation of immunoprecipitates is based on a series of bimolecular reactions following the law of mass action:



Using light-scattering and fluorescence-quenching measurements, Ten-gerdy and Small (1966) showed that the primary Ab-Ag complex is established within 10 sec, while equilibrium-dialysis has demonstrated that the final polymerized antigen-antibody complex is formed during the first half hour (Hughes-Jones et al. 1963). Heidelberger's basic concept was further modified by Eisen and Keston (1949) who demonstrated the existence of polyvalent binding sites on the antigen. The antibody-antigen reaction may be considered similar to the binding of drugs to e.g. serum albumin (Klotz 1946; Clausen 1966b).

The antibody-antigen reaction shows temperature dependence of




Fig. 1.1. Precipitation curves of three protein antigens. The three curves illustrate the immunoprecipitation in a free buffer solution of human albumin (Alb), IgG globulin (IgG glob) and human chorionic gonadotropin (HCG) with corresponding specific antibodies. *Abcissa*:  $\mu\text{g}$  of protein present per 500  $\mu\text{l}$  antigen solution (in 0.9% NaCl). *Ordinate*: mg protein precipitated by corresponding specific antisera per 100 ml solution.

the rate of association of the two reactants between about 0 to 40°C. Various investigators have found  $Q_{10}$  values varying from about 2 (Burnet et al. 1937) to about 1.4. Values intermediate between these extremes may be the most correct. Jerne and Skovsted (1953) demonstrated that the reaction was maximal at a low salt concentration. Finally, the antigen-antibody reaction is dependent on pH, showing an optimum at 7.0 (Kleinschmidt and Boyer 1952). The immunoprecipitate is split below pH 4.5 and above pH 10.0.

From a physicochemical point of view the antigen-antibody reaction is characterized by small changes in the thermodynamic constants. Thus, the change in Gibbs' free energy  $\Delta G^0$  seems of the order of -9 kcal/mol. The change in entropy  $\Delta S^0$  is positive and the change in enthalpy  $\Delta H^0$  is about -9 kcal/mol (Wurmser and Filitti-Wurmser 1957; Karush 1958; see also the survey by Schultze 1962). These values are of the same order as the changes in the thermodynamic constants for binding of drugs to serum albumin (Klotz 1946; Clausen 1966b).

The small changes in the thermodynamic constants and the positive entropy change may be partly related to displacement of water from the antibody site and to conformational changes of the reactants after establishment of the antigen-antibody complex. These changes may be related to formation of hydrophobic bonds (Scheraga 1963).

The physicochemical event during the antigen-antibody reaction depends on the type (horse or rabbit) of antibody used. The antigen-antibody complex formed is usually characterized by being soluble in excess of antigen and also, in the case of horse type antibody only, in excess of antibody (Kleinschmidt and Boyer 1952).

A more detailed description of the antigen-antibody reaction based on physicochemical studies is not the object of this book; the interested reader is referred to Goldberg (1952) and Singer and Campbell (1952).

## *1.2. Antigen-antibody reaction in gels*

Early in this century Bechhold (1905) discovered that immunochemical reactions could also be performed in gels. The 'gel diffusion' method was further developed by Arrhenius (1907) who showed that it could fractionate antigen mixtures such as tetanus and diphtheria toxins in complex with their corresponding antibodies. Qualitative methods for



identification of antigens and antibodies were introduced by Nicolle et al. (1920). Both Petrie (1932) and Sia and Chung (1932) further developed the gel method as a means for identification of bacterial antigens. Later, Maegraith (1933) and Hanks (1935) suggested gel diffusion as a microanalytical tool for demonstration of the antigenic composition of unknown samples.

In 1946 Oudin elaborated the physicochemical and mathematical basis for employing the diffusion in gels of a single antigen and/or antibody for quantitative purposes. The basic work of Oudin (1946; review, 1952) was extended and modified by Ouchterlony (1949) and Elek (1948). In 1953 Grabar and Williams described 'immunoelectrophoresis' which combined the electrophoretic separation of proteins in agar gel with a subsequent antigen-antibody reaction performed by immunodiffusion in the same agar gel. Although the basic principle of this method had been independently developed by Poulik (1952) who combined filter paper electrophoresis with agar gel immunodiffusion, the recent expansion in analytical immunochemistry was initiated by the work of Grabar and Williams. On the basis of their method, Scheidegger (1955) developed his micro-immunoelectrophoretic method which made it possible to separate and identify a large number of antigens in a volume of about 1.5  $\mu\text{l}$ .

Electrophoresis combined with the antigen-antibody reaction was also the basis for the development in recent years of radio-immunochemical techniques (Berson and Yalow 1961; Berson et al. 1964). These techniques have provided the impetus for development of ultra-micro-methods for estimation of proteins and peptide hormones in biological fluids (see Chard (1978)).

Gel diffusion reactions are dependent on the high water content (often as high as 99%) of the gelatinous supporting media. This permits the antigens and antibodies to migrate through the gel by diffusion (or be accelerated by an electric field) almost independently of the supporting medium, provided that the diameter of the reactants is less than the width of the microstructure of the gels (Wieme 1957, 1959). Under these conditions the free diffusion of both antigens and antibodies conforms to laws of gaseous diffusion and thus follows Fick's law:

$$dx/dt = Dd^2c/dt^2, \quad (1.4)$$

*Subject index p.363*