

*American*  
**DRUG INDEX**  
*1978*

NORMAN F. BILLUPS, Ph.D., R.Ph.

*American*  
**DRUG INDEX**  
*1978*

**NORMAN F. BILLUPS, Ph.D., R.Ph.**

Dean and Professor of Pharmacy  
College of Pharmacy  
The University of Toledo

Associate Editor

**SHIRLEY M. BILLUPS, R.N.**

**J. B. LIPPINCOTT COMPANY**  
Philadelphia · Toronto

**Copyright © 1978, by J. B. Lippincott Company**

Copyright © 1956, 1958, 1959, 1960, 1961, 1962, 1963, 1964,  
1965, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1975, 1976,  
1977 by J. B. Lippincott Company

This book is fully protected by copyright and, with the exception  
of brief excerpts for review, no part of it may be reproduced  
in any form by print, photoprint, microfilm or any other means,  
without the written permission of the publishers.

**Second Printing**

**ISBN 0-397-50401-2**

**Library of Congress Catalog Card Number 55-6286**

**Printed in the United States of America**

# Preface

The *American Drug Index* has been prepared for the identification, explanation and correlation of the many pharmaceuticals available to the medical, pharmaceutical and allied health professions. The need for this index has become acute as the variety and number of drugs and drug products has continued to multiply. For this reason the *American Drug Index* should be most useful to physicians, pharmacists, dentists, nurses, sales personnel, students and to teachers in the fields incorporating pharmaceuticals.

The organization of the *ADI* falls into the following seven major sections: Monographs of Drug Products, Pharmaceutical Manufacturer and/or Drug Distributor Addresses, Common Abbreviations Used in Medical Orders, Common Systems of Weight and Measure, Approximate Practical Equivalents, Glossary, and Pharmaceutical Company Labeler Code Index Numbers. The Glossary section has been added for the first time this year to make the *American Drug Index* an even more complete and valuable reference tool.

**MONOGRAPHS:** The organization of the monograph section of the *ADI* is fundamentally alphabetical with extensive cross-indexing. Names listed are generic (also called non-proprietary, public name or common name); brand (also called trade-mark, proprietary or specialty); and chemical. Synonyms that are in general use also are included. All names used for a pharmaceutical occur in alphabetical order with the pertinent data given under the brand name by which it is made available.

The monograph for a typical brand name product consists of the manufacturer, generic and/or chemical names, composition and strength, pharmaceutical forms available, package size, dosage and use.

Generic names occur in alphabetical order, followed by the corresponding recognition of the drug to the U.S.P. (United States Pharmacopeia), N.F. (National Formulary), and USAN (United States Adopted Names). Each of these officially recognized generic names is preceded by a bullet at the beginning of each entry. The information is in accord with the U.S.P. XIX, N.F. XIV, and the USAN-1976.

British Approved Names, indicated by (B.A.N.), are included in this edition with the permission of the Controller of Her Britannic Majesty's Stationery Office.

Because of the multiplicity of specialty names (brand names) used for the same therapeutic agent or the same combination of therapeutic agents, it was apparent that some correlation could be done. As an example of this, please turn to tetracycline HCl. Here under the generic name are listed the various brand names. Following are combinations of tetracycline HCl organized in a manner to point out relationships among the many products. Reference then is made to the specialty name or names having the indicated composition. Under the

specialty name are given manufacturer, composition, available forms, sizes, dosage and use.

The multiplicity of generic names for the same therapeutic agent has complicated the nomenclature of these agents. Examples of multiple generic names for the same chemical substance are (1) parabromdylamine, brompheniramine; (2) acetaminophen, p-hydroxyacetanilid, N-acetyl-p-aminophenol; (3) d-isoeephedrine, d-pseudoephedrine; (4) methapyrilene HCl, thenylpyramine HCl; and (5) pyrillamine, pyranisamine, pyranilamine, pyraminyl and anisopyradamine.

The cross-indexing feature of the *ADI* permits the finding of drugs or drug combinations when only one major ingredient is known. For example, a combination of aluminum hydroxide gel and magnesium trisilicate is available. This combination can be found by looking under the name of either of the two ingredients, and in each case the specialty names are given.

**ABBREVIATIONS:** A listing of Common Abbreviations Used in Medical Orders is included as an aid to interpreting medical orders. The Latin or Greek word and abbreviation are both given with the meaning.

**WEIGHT AND MEASURE:** Tables containing the Common Systems of Weight and Measure are included to aid the practitioner in calculating dosages in the metric, apothecary and avoirdupois systems.

**CONVERSION FACTORS:** The listing of Approximate Practical Equivalents is added as an aid in calculating and converting dosages among the metric, apothecary and avoirdupois systems.

**GLOSSARY:** A listing of commonly used terms, with brief definitions, is included as an aid to interpreting the use given for drug monographs listed in the *ADI*.

**LABELER CODE INDEX:** The Pharmaceutical Labeler Code Index is presented to aid the practitioner in the identification of drug products. In this section the codes are listed in numerical order followed by the name of the manufacturer.

**MANUFACTURER ADDRESSES:** The name, address and zip code of virtually every American Pharmaceutical Manufacturer and/or Drug Distributor are listed in alphabetical order in this section. Additionally, a pharmaceutical labeler code number appears prior to the address for each company as a further aid in the identification of drug products.

A special word of appreciation and acknowledgment is given to my wife, Shirley, who served again this year as my Associate Editor—and helped to compile much of the information that appears in the monograph section of this volume. Special thanks are also extended to the manufacturers who supplied product information and made helpful comments and suggestions for this index. We found it especially rewarding to work with the Medical Publications Division of J. B. Lippincott Company, and we are especially grateful to J. Stuart Freeman,

Senior Editor of Medical Books, for his competent and meticulous attention.

Correspondence or communication with reference to a drug or drug products listed in the *American Drug Index* should be directed to: Norman F. Billups, Ph.D.; Dean, College of Pharmacy; The University of Toledo; Toledo, Ohio 43606.

Norman F. Billups

# Contents

|                                                                   |         |
|-------------------------------------------------------------------|---------|
| 1. Monographs .....                                               | 1-697   |
| 2. Common Abbreviations Used<br>In Medical Orders .....           | 698-699 |
| 3. Common Systems of Weight<br>and Measure .....                  | 700     |
| 4. Approximate Practical<br>Equivalents .....                     | 701     |
| 5. Glossary .....                                                 | 702     |
| 6. Pharmaceutical Company Labeler<br>Code Index .....             | 707-712 |
| 7. Pharmaceutical Manufacturers<br>and/or Drug Distributors ..... | 713-728 |

[•] Denotes Official Name: Generic name or chemical name approved by  
F.D.A., or recognized by the USAN, U.S.P. or N.F.

## A

**A=1 LAXATIVE.** (Western Research) Aloin  $\frac{1}{4}$  gr., pow. ext. stramonium  $\frac{1}{4}$  gr., pow. ext. cascara sagrada  $\frac{1}{2}$  gr., pow. ext. rhubarb  $\frac{1}{2}$  gr., ginger  $\frac{1}{2}$  gr./Tab. Bot. 1000s. Paks 28.

Use: Laxative.

**A-200 PYRINATE.** (Norcliff Thayer) Pyrethrins, piperonyl butoxide, deodorized kerosene. **Liquid**, Bot. 2 oz., 4 oz. **Gel**, Tube 1 oz.

Use: Topically, pediculosis.

**A.A. ANTACID TABLETS.** (Western Research)

Balanced combination of calcium carbonate ppt. 583.2 mg., magnesium carbonate light, 194.4 mg., magnesium trisilicate and aluminum hydroxide dried gel 64.8 mg./Chewable Tab. Strip-pak 28s, 1000s.

Use: Antacid.

**A.A.A. OINTMENT.** (Jenkins) Ammoniated mercury 2%, acid salicylic 1.25%, acid boric 1%, zinc oxide 15%. Jar 1 oz., 1 lb.

Use: Impetigo, nonspecific eczema & minor skin irritations.

**A-A PLUS.** (Galen Labs.) Al hydroxide gel 400 mg., Mg oxide 175 mg., Na lauryl sulfate 25 mg., Me cellulose 100 mg., hyoscyamine sulfate 0.1037 mg., atropine sulfate 0.0194 mg., hyoscyne HBr 0.0065 mg./Tab. Bot. 100s, 1000s. Susp. 1 pt.

**AARANE.** (Syntex) Cromolyn sodium 20 mg./Cap. Bot. 60s, Inhaler 1s.

Use: Bronchial asthma inhibitor.

**ABBOCIN.** Oxytetracycline, B.A.N.

**ABBOKINASE.** (Abbott) Urokinase 250,000 I.U./10 ml. Lyophilized powder.

Use: Plasminogen activator.

**ABCDG VITAMIN CAPSULES.** (Lannett) Vit. A 5000 u., D 400 u., B-1 1 mg., B-2 2 mg., C 30 mg./Cap. Bot. 1000s.

Use: Vitamin supplement.

**ABDEC BABY VITAMIN DROPS.** (Parke, Davis) Vit. A 5000 u., D 400 u., B-1 1 mg., B-2 1.2 mg., B-6 1 mg., sod. pantothenate 5 mg., niacinamide 10 mg., C 50 mg./0.6 cc. Bot.  $\frac{1}{2}$  oz.,  $\frac{1}{4}$  oz. W/dropper.

Use: Vitamin deficiencies.

**ABDEC WITH FLUORIDE BABY VITAMIN DROPS.**

(Parke, Davis) Vit. A 5000 u., D 400 u., C 50 mg., B-1 1 mg., B-2 1.65 mg., B-6 1 mg., nicotinamide 10 mg., sodium pantothenate 5.5 mg., sodium fluoride 1.1 mg./0.6 ml. Bot. 50 ml.

Use: Vitamin therapy with fluoride.

**ABDEC KAPSEALS.** (Parke, Davis) Vit. A 10,000 u., D 400 u., B-1 5 mg., B-2 3 mg., B-6 1.5 mg., B-12 2 mcg., dl-panthanol 10 mg., niacinamide 25 mg., Vit. C 75 mg., E 5 I.U./Cap. Bot. 100s, 250s., unit dose pkg. 100s.

Use: Multivitamin therapy.

**ABDOL W/MINERALS.** (Parke, Davis) Vit. C 50 mg., B-2 2.5 mg., B-12 1 mcg., B-6 0.5 mg., B-1 2.5 mg., niacinamide 20 mg., Vit. A 5000 u., D 400 u., Ca pantothenate 2.5 mg., folic acid 0.1 mg., 10.15 mg., Mn 1 mg., K 5.0 mg., Fe 15 mg., Cu 1 mg., Zn 0.5 mg., Mg 1 mg., Ca 44 mg., P 34 mg./Cap. Bot. 100s, 250s.

Use: Vitamin-mineral deficiency states.

**ABDOL W/VIT. C.** (Parke, Davis) Vit. A 5000 u., D 400 u., B-1 2.5 mg., B-2 2.5 mg., B-6 0.5 mg., B-12 1 mcg., nicotinamide 20 mg., Vit. C 50 mg., pantothenic acid (as Cal. salt) 5 mg./Cap. Bot. 100s.

Use: Vitamin deficiency.

**ABEREL.** (McNeil) Tretinoin (all-trans retinoic acid) 0.05%, w/polyethylene glycol 400, butylated hydroxytoluene, alcohol 55%. Saturated swabs. Box 30s.

Use: Acne.

**ABSORBABLE CELLULOSE COTTON OR GAUZE.**

See: Oxidized Cellulose (Various Mfr.)

• **ABSORBABLE DUSTING POWDER,** U.S.P. XIX.

**ABSORBABLE GELATIN FILM.** Sterile nonantigenic, absorbable, water-insoluble, gelatin film.

See: Gelfilm (Upjohn)

• **ABSORBABLE GELATIN SPONGE,** U.S.P. XIX.

Gelatin sponge.

Use: Surgical aid.

See: Gelfoam, Preps. (Upjohn)

• **ABSORBABLE SURGICAL SUTURE,** U.S.P. XIX. (Various Mfr.) Surgical Gut, Surgical Catgut, Catgut suture.

Use: Surgical aid.

• **ABSORBENT GAUZE,** U.S.P. XIX.

Use: Surgical aid.

**ABSORBINE ARTHRITIC PAIN LOTION.** (Young)

Bot. 2 oz., 4 oz.

**ABSORBINE FOOT POWDER.** (Young) Plastic bot. 2 oz. w/shaker top.

Use: Athlete's foot.

**ABSORBINE, JR.** (Young) Wormwood, thymol, YG-7 (chloroxylenol), menthol, acetone. Liq. Bot. 1 oz., 2 oz., 4 oz., 12 oz. w/applicator.

Use: Muscle soreness, stiffness.

**ABSORBINE MEDICATED SOAP.** (Young) Bar  $3\frac{1}{2}$  oz.

**ABUSCREEN.** (Roche Diagnostics) An immunological and radiochemical assay for morphine and morphine glucuronide in nanogram levels. Utilizes  $^3\text{H}$  labeled morphine assayed with liquid scintillation counters using conventional beta counting procedures and  $^{125}\text{I}$  labeled morphine requiring gamma scintillation equipment. 100 tests.

**A.C.A. CAPS.** (Scrip) Aspirin  $3\frac{1}{2}$  gr., caffeine  $\frac{1}{2}$  gr., acetophenetidin  $2\frac{1}{2}$  gr./Cap. Bot. 100s.

**A.C.A. No. 2.** (Scrip) Aspirin 4 gr., caffeine  $\frac{1}{4}$  gr., acetophenetidin 2 gr./Tab. Bot. 100s.

**A.C.A. No. 5.** (Scrip) Phenacetin 2 gr., salicylamide 2 gr., atropine sulfate  $1/600$  gr., tr. gelsemium 3 min., camphor monobromate  $\frac{1}{2}$  gr./Tab. Bot. 100s.

**ACABEL.** Bevonium Methylsulphate, B.A.N.

**A-CAINE.** (A.V.P.) Benzocaine 2%, cod liver oil, zinc oxide, balsam peru, bismuth subgallate, ephedrine sulfate and lanolin in a special base. Tube w/applicator,  $1\frac{1}{2}$  oz.

Use: Temporary relief of hemorrhoids, minor rectal irritation and as an aid in rectal examination.

**A CAPSULES.** (Stanlabs) Natural Vit. A 5,000 u., 10,000 u., 25,000 u., 50,000 u./Cap. Bot. 100s. Synthetic or s. uble, 10,000 u., 25,000 u., 50,000 u./Cap. Bot. 1: Js.

- **ACACIA**, U.S.P. XIX. (Penick) *Acacia senegal*, gum arabic. Mucilage N.F. XIV; pow., ¼ lb., 1 lb., tears, ¼-1 lb., syr. N.F. XIII. whole, ¼-1 lb.  
Use: Demulcent, emulsifier, suspending agent.
- A.C.A. NASAL DROPS**. (Emerson) Phenylephrine HCl in a saline solution. Dropper bot. 1 oz.  
Use: Relief of nasal blockage in infants and children.
- A-CAP**. (A.V.P.) Vit. A 50,000 u./Cap. Bot. 100s.
- A-CAPS**. (Drug Industries) Vit. A 50,000 u./Cap. Bot. 100s, 500s.  
Use: Vitamin A therapy.
- ACCELERASE**. (Organon) Pancrelipase 165 mg., bile salts 65 mg., cellulase 2 mg., calcium carbonate 20 mg./Cap. Bot. 60s.  
Use: Functional digestive disorders.
- ACCELERASE-PB**. (Organon) Same as Accelerase, with belladonna alkaloids (87.5% hyoscyamine and 12.5% atropine sulf.) 0.2 mg., phenobarbital 16 mg./Cap. Bot. 60s.  
Use: Digestive enzyme.
- ACCOPAIN**. (Vitarine) Codeine phosphate ¼ gr., buffered aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr./Cap. Bot. 24s.  
Use: Analgesic.
- A.C.D. CAPSULES**. (Philips Roxane) Pwd. opium 1.62 mg., phenacetin 97.5 mg., aspirin 97.5 mg., caffeine 4 mg., camphor monobromated 16.2 mg., atropine sulfate 0.065 mg./Cap. Bot. 1000s.
- A-C-D SOLUTION**, U.S.P. Sod. citrate, citric acid & dextrose in sterile pyrogen-free solution.  
Baxter-600 cc. bot. with 70, 120 or 300 cc. of sol.; 1000 cc. with 500 cc. of sol.  
Cutter-500 cc. bot. with 75 or 120 cc. of sol.; 650 cc. bot. with 80 or 130 cc. of sol.  
Diamond-(Abbo-Vac) 250 cc., 500 cc.  
Use: Anticoagulant for preparation of plasma or whole blood.
- A-C-D SOLUTION MODIFIED**. (Squibb) Acid citrate dextrose anticoagulant solution modified.  
Use: Diagnostic agent for cancer.
- ACDRILE**. Methyl Cysteine, B.A.N.
- **ACEBUTOLOL**. B.A.N. (±)-1-(2-Acetyl-4-butyramidophenoxy)-3-isopropylaminopropan-2-ol. (Ives) (±)-N-[3-Acetyl-4-[2-hydroxy-3-[(1-methylethyl)amino]propoxy]-phenyl] butamide.  
Use: Beta adrenergic receptor blocking agent.
- **ACECLIDINE**. (Kingshill) 3-Quinuclidinol acetate (ester). Glauco-stat. Under study.  
Use: Parasympathomimetic.
- **ACEDAPSONE**. 4',4'-Sulfonylbis (acetanilide).  
Use: Antimalarial; antibacterial (leprostatic).
- ACEDOVAL**. (Vale) Dover's powder ¼ gr., phenacetin 2½ gr., caffeine anhydrous ¼ gr., camphor ¼ gr./Tab. Bot. 100s, 1000s, 5000s.
- ACEDYNE**. (Vale) Phenacetin ½ gr., aspirin ½ gr., caffeine anhydrous ½ gr., atropine sulfate ½<sub>1000</sub> gr., magnesium trisilicate 1 gr./Tab. Bot. 100s, 1000s, 5000s.
- ACEFYLLINE PIPERAZINE**. Acepifylline, B.A.N.
- ACELAX**. (Marcen) Danthron 100 mg., prune concentrate 64 mg., *Chondrus drispus* 64 mg./Tab. Bot. 100s, 1000s.  
Use: Laxative.
- ACENOCOUMARIN**.  
See: Sintrom, Tab. (Geigy)
- **ACENOCOUMAROL**. N.F. XIV, Tab. N.F. XIV. N.D. 1967. 3-( $\alpha$ -Acetonyl-4-nitrobenzyl)-4-hydroxy-coumarin. Nicoumalone, B.A.N.  
See: Sintrom, Tab. (Geigy)
- ACEPIFYLLINE**. B.A.N. Piperazine theophyllin-7-ylacetate. Acefylline Piperazine (I.N.N.). Etophylate.  
Use: Spasmolytic.
- **ACEPROMAZINE**. 10(3-Dimethylaminopropyl)-3-acetylphenothiazine maleate. Plegicil. (Ayerst) 1-(10-(3-(Dimethylamino)propyl)-10H-phenothiazin-2-yl)ethanon e(2)-2-butenedioate.  
Use: Tranquillizer.
- ACEPROMAZINE**. B.A.N. 2-Acetyl-10-(3-dimethylaminopropyl)phenothiazine. Notsensil maleate.  
Use: Tranquillizer.
- ACEROLA C**. (Kirkman) Vit. C 100 mg./Wafer. Bot. 100s.
- ACEROLA-C**. (Barth's) Vit. C 300 mg./Wafer. Bot. 30s, 90s, 180s, 360s.
- ACEROLA-PLEX**. (Barth's) Vit. C 100 mg., bioflavonoids 50 mg./Tab. Bot. 100s, 500s.
- ACETA**. (Century) Acetaminophen. Elix. (120 mg./5 cc) Bot. 4 oz., pint, gal. Tabs. (5 gr.) Bot. 100s, 1000s, 3000s, 5000s. Supp. 300 mg., 600 mg. 12 $\frac{1}{2}$   
Use: Analgesic.
- ACETA W/CODEINE**. (Century) Acetaminophen 325 mg., codeine phosphate ½ gr./Tab. Bot. 100s.  
Use: Analgesic.
- ACETABAR**. (Philips Roxane) Phenobarbital ¼ gr., acetophenetidin 2½ gr., acetylsalicylic acid 3½ gr., caffeine ¼ gr./Tab. Bot. 100s, 1000s.  
Use: Analgesic-sedative.
- ACETAGESIC**. (Kay) Acetaminophen 120 mg./5 cc. Elix. Bot. pts., gals.  
Use: Analgesic and antipyretic, in pain and fever syndrome.
- ACETAMIDE**. N-[5-(Aminosulfonyl)-3-methyl-1,3,4-thiadiazol-2(3H)-ylidene]-. Methazolamide, U.S.P. XIX.
- ACETAMIDE, 2,2-DICHLORO-N-[2-HYDROXY-1-(HYDROXYMETHYL)-2-(4-NITROPHENYL)ETHYL]-**. Chloramphenicol, U.S.P. XIX.
- ACETAMIDE, 2-(DIETHYLAMINO)-N-(2,6-DIMETHYL-PHENYL)-**. Lidocaine, U.S.P. XIX.
- ACETAMIDE, N-[5-(AMINOSULFONYL)-1,3,4-THIADIAZOL-2-YL]-**. Acetazolamide, U.S.P. XIX.
- ACETAMIDE, N-(4-HYDROXYPHENYL)-**. Acetaminophen, U.S.P. XIX.
- ACETAMIDE, N-(5,6,7,9-TETRAHYDRO-1,2,3,10-TETRAMETHOXY-9-OXOBENZO[a]-HEPTALEN-7-YL)-**. Colchicine, U.S.P. XIX.
- p-ACETAMIDOBENZOIC ACID SALT** of 2-di-methylaminoethanol.  
See: Deaner, Tab. (Riker).
- 5-ACETAMIDO-2,4,6-TRIIODO-N-METHYLISOPHTHALAMIC ACID**. Iothalamic Acid, U.S.P. XIX.
- 5-ACETAMIDO-2,4,6-TRIIODO-N-METHYLISOPHTHALAMIC ACID, SODIUM SALT**. Iothalamate Sodium Injection, U.S.P. XIX.

• **ACETAMINOPHEN**, U.S.P. XIX. Elix., Tabs: U.S.P. XIX. N-Acetyl-p-amino-phenol. Acetamide, N-(4-hydroxyphenyl)-. 4'-Hydroxyacetanilide.

Use: Analgesic, antipyretic.

See: Aceta, Tab. Elix., Supp. (Century)

Acetagesic (Kay)

Actamin, Tab. (Buffington)

Aminodyne, Elix. (Bowman)

Anapap, Tab. (Fellows)

Anelix, Elix. (Kirkman)

Anuphen, Supp. (Comatic)

Apadon, Tab. (Elder)

Apamide, Tab. (Dome)

Apap, Cap., Tab. (Various Mfr.)

Bayer Acetaminophen Non-Aspirin Pain Reliever, Tab. (Glenbrook)

Bromo-Seltzer, Pow. (Warner-Lambert)

Capital, Tab. (Carrick)

Cap-Apap, Liq. (Central)

Chlor-A-Tyl. (Jenkins)

Dapa, Tab. (Ferndale)

Datril, Elixir, Tab. (Bristol-Myers)

Datril 500, Tab. (Bristol-Myers)

Dimindol, Tab., Elix. (Blue Line)

Dolanex, Elix. (Lannett)

Dularin, Syr. (Donner)

Elixodyne, Liq. (Lemmon)

Febriol, Elix., Tab. (Amid)

Febrogesic, Tab. (First Texas)

Fendon, Tab. (APC)

G-1 (Palmedico)

Janupap, Tab. (Tutag)

Lestemp, Elix. (Reid-Provident)

Liquiprin, Susp. (Norcliff-Thayer)

Lyteca, Syr., Tab. (Westerfield)

Med-Apap, Elix. (Med. Chem.)

Medigesic (Medics)

Menaigesia, Tab. (Otis Clapp)

Nebs, Tab., Liq. (Eaton)

Neopap, Supp. (Webcon)

Nilprin 7½, Tab. (AVP)

Parten, Tab. (Parmed)

Phendex, Tab. (Mallard)

Pinin, Tab. (Scrip)

Proval, Cap., Elix., Drop., Tab. (Reid-Provident)

Pyrpap, Tab., Susp. (Savage)

SK-Apap, Tab., Elix. (SmithKline)

St. Joseph, Prods. (Plough)

Sub-Due, Tab. (Wisconsin)

Taper, Elix., Tab. (Parke, Davis)

Temetan, Elix., Tab. (Nevin)

Tempra, Drop, Syr., Tab. (Mead Johnson)

Tenlap, Elix. (Dow)

Tussapap, Tab. (Sutliff & Case)

Tylenol, Elix., Tab., Chew. Tab., Cap. (McNeil)

Tylenol Extra-Strength, Tab., Cap. (McNeil)

Valadol, Tab., Liq. (Squibb)

Valorin Tab. (Otis Clapp)

**ACETAMINOPHEN W/COMBINATIONS**

Aceta w/Codeine, Tab. (Century)

Akes-N-Pain, Cap. (Edw. J. Moore)

Al-Ay Modified, Tab. (Bowman)

Algecol, Tab. (Winston)

Allylgescic, Tab. (Elder)

Allylgescic W/Ergotamine, Cap. (Elder)

Allylvon, Cap. (Elder)

Alumadrine, Tab. (Fleming)

Amaphen, Cap. (Trimen)

Aminodyne, Elix. (Bowman)

Anaphen, Cap. (Mallard)

Anodynos Forte, Tab. (Buffington)

Apap w/Codeine, Tab. (Central)

Apapalone, Tab. (Amfre-Grant)

Arthol Tabs. (Towne)

Arthralgen, Tab. (Robins)

Bancaps (Westerfield)

Bancaps-C (Westerfield)

Banesin-Forte, Tab. (Westerfield)

Bexahist, Cap. (Bexar)

Biophens-S, Tab. (La Crosse)

Bowman Cold Tab. (Bowman)

B-Pap., Liq. (Wren)

Bromo Quinine Cold Tabs (Bristol-Myers)

Burgescic, Tab. (Burlington)

Butgetic, Tab. (McNeil)

Calm Aids, Cap. (Daniels, Robert)

Capital w/Codeine, Tab., Susp. (Carrick)

Cerose Comp., Cap. (Ives)

Chexit, Tab. (Dorsey)

Christodyne-DHC, Tab. (Canfield)

Coastaldyne, Tab. (Coastal)

Coastalgescic, Tab. (Coastal)

Codalan, Tab. (Lannett)

Codap (Tutag)

Codrene W/Codeine, Syr. (Daniels, Robert)

Codrene-D S.F., Syr. (Daniels, Robert)

Codimal, Tab. (Central)

Coldene, Syr. (Pharmcraft)

Colrex Compound, Elix. (Rowell)

Conar-A, Tab., Susp. (Beecham Labs)

Conex, Preps. (Westerfield)

Contac Jr., Liq. (Menley & James)

Coryban-D, Syr. (Roerig)

Corzans, Tab. (Moore Kirk)

Co-Tylenol, Tab. (McNeil)

Dapase, Tab. (Ferndale)

Darvocet-N, Tab. (Lilly)

Darvocet-N 100, Tab. (Lilly)

Dengesic, Tab. (Scott-Alison)

Desa-Hist AT, Tab. (Pharmics)

Dialog, Prods. (Ciba)

Dolene AP-65, Tab. (Lederle)

Dolopar, Liq. (O'Neal)

Drinacet Cap. (Philips Roxane)

Drinophen, Cap. (Lannett)

D-Sinus, Cap. (Laser)

Dularin-TH, Tab. (Dooner)

Duo-Gescic, Tab. (Trout)

Duoprin, Tab. (Dunhall)

Duo-3X, Tab. (Med. Spec.)

Duramid, Tab. (O'Neal)

Dynosal, Tab. (Mallard)

Enpayne, Tab. (Burgin-Arden)

Enz-Cold, Cap. (Edw. J. Moore)

- Esemgesic, Tab. (Spencer-Mead)  
 Esgic, Tab. (Gilbert)  
 Excedrin, Tab. (Bristol-Myers)  
 Femicin, Tab. (Norcliff Thayer)  
 Fendol, Tab. (Buffington)  
 G-2, Tab. (Palmedico)  
 G-3, Cap. (Palmedico)  
 Gaysal, Tab. (Geriatric)  
 G-Lixir (Palmedico)  
 Guaifamine, Cap. (Sutliff & Case)  
 Hasacode, Tab. (Arnar-Stone)  
 Her-Caps, Cap. (Edw. J. Moore)  
 Histalets, Tab. (Edw. J. Moore)  
 Histogesic, Tab. (Century)  
 Histosal #2, Tab. (Ferdale)  
 Hi-Temp, Tab. (Saron)  
 Hycomine Comp., Tab. (Endo)  
 Intensin, Cap. (McNeil)  
 Isomel, Cap. (Tutag)  
 Kiddies Sialco, Tab. (Foy)  
 Kleer, Syr. (Scrip)  
 Koryza Tab. (Fellows-Testagar)  
 Liquix-C, Liq., Tab. (Elder)  
 Maxigesic, Cap. (W.F. Merchant)  
 Medigesic, Tab. (Medics)  
 Mense, Tab. (Daniels, Robert)  
 Midran, Tab. (Columbia)  
 Midrin, Cap. (Carnrick)  
 Minotal, Tab. (Carnrick)  
 Myocalm, Tab. (Parmed)  
 Myolate, Tab. (Panray)  
 Naldetuss, Syr. (Bristol)  
 N-C-P, Tab. (Wendt-Bristol)  
 N-D Gestic, Tab. (Hyrex-Key)  
 Neo-Pyranistan, Tab. (Standex Labs.)  
 Neo-Vadrin Sinus Headache Formula, Tab. (First Texas)  
 Nokane, Tab. (Wren)  
 Novahistine Singlet, Tab. (Dow)  
 Nylorac PB, Tab. (Beard)  
 Opacedrin, Cap. (Elder)  
 Ornex, Cap. (SmithKline)  
 Panitol H.M.B., Tab. (Wesley)  
 Panritis, Tab. (Pan Amer.)  
 Parafon Forte, Tab. (McNeil)  
 Partuss-A, Tab. (Parmed)  
 Partuss T.D., Tab. (Parmed)  
 Pavadon Elix. (Coastal)  
 Pedituss, Liq. (Sutliff & Case)  
 Pedic, Elix., Waters (Vale)  
 Percogesic, Tab. (Endo)  
 Phenahist, Cap., Elix. (Amid)  
 Phrenilin, Tab. (Carnrick)  
 Presalin, Tab. (Mallard)  
 Prodolor, Tab. (Amfre-Grant)  
 Proval No. 3, Tab. (Reid-Provident)  
 Pyradyne Comp., Tab. (Lemmon)  
 Pyrihist Cold Cap. (Daniels, Robert)  
 Renpap, Tab. (Wren)  
 Rentuss, Cap. Syr. (Wren)  
 Repan, Tab. (Everett)  
 Rhinex, Tab. (Lemmon)  
 Rhinidrin, Tab. (Central)  
 Rhinogesic, Tab. (Vale)  
 Rhinspec, Tab. (Lemmon)  
 S.A.C. Sinus, Tabs. (Towne)  
 Salatin w/Codeine, Tab. (Ferdale)  
 Saleto, Tab. (Mallard)  
 Saleto-D, Tab. (Mallard)  
 Salimeph, Preps. (Kremers Urban)  
 Salphenyl, Cap., Liq. (Mallard)  
 Sanspen, Tab. (Spencer-Mead)  
 Santussin, Prep. (Sandia)  
 Scotgesic, Cap., Elix. (Scott/Cord)  
 Scotuss Pediatric Cough Syr. (Scott Cord)  
 Sedacane, Cap. (Edw. J. Moore)  
 Sedalgesic, Tab. (Table Rock)  
 Sedapap, Elix. (Mayrand)  
 Sedragesic, Tab. (Lannett)  
 Sialco, Tab. (Foy)  
 Sinacet, Tab. (Meyers)  
 Sinarest, Tab. (Pharmacraft)  
 Sinoze, Tab. (Bowman)  
 Sinubid, Tab. (Warner Chilcott)  
 Sinudan, Tab. (Daniels, Robert)  
 Sinulin, Tab. (Carnrick)  
 Sinumal, Tab., Liq. (Cook)  
 Sinus Tab., Tab. (Zenith)  
 Sinutab Preps. (Warner-Chilcott)  
 Sinuwes, Tab. (Western Research)  
 SK-APAP W/codeine, Tab. (SmithKline)  
 SK-65 APAP, Tab. (SmithKline)  
 Solstice Decongestant Tablets (Chattem Labs.)  
 Spantuss, Liq. (Arco)  
 Spendrisin, Tab. (Spencer-Mead)  
 Stopain, Tab. (Columbia)  
 Sunril, Cap. (Emko)  
 Super-Anahist Tablets (Warner-Lambert)  
 Symptomax, Liq. (Reid-Provident)  
 T-Caps, Cap. (Burlington)  
 Tega-Code, Cap. (Ortega)  
 Tegapap, Liq., Tab. (Ortega)  
 Teragen, Tab. (O'Neal)  
 T-Gestic, Cap. (Tennessee)  
 Tnamincin, Tab. (Dorsey)  
 Triaprin, Cap. (Dunhall)  
 Trind, Syr. (Mead Johnson)  
 Trind-DM, Syr. (Mead Johnson)  
 Tussagesic, Tab., Susp. (Dorsey)  
 Two-Dyne, Tab. (Hyrex-Key)  
 Tylenol Preps. (McNeil)  
 Tylox, Cap. (McNeil)  
 Valihist, Cap. (Clapp)  
 Vannor, Cap. (North Amer. Pharm.)  
 Windolor, Tab. (Winston)  
 Wygesic, Tab. (Wyeth)  
 Zenex Comp., Cap. (Zenith)
- ACETAMINOPHENOL**  
 See: Acetaminophen.
- ACETANILID.** (Various Mfr.) (Acetylamino benzene, acetylaniline, antifebrin) N-phenylacetamide cry.  
 Use: Antipyretic, analgesic.
- ACETARSOL**  
 See: Acetarsone, Tabs.

- **ACETARSONE**, (City Chemical) N.F. XI., Tablets, N.F. XI. 3-Acetamido-4-hydroxy-phenylarsonic acid (Acetarsol, Acetphenarsine, Amarsan, Dynarsan, Ehrlich 594, Limarsol, Orarsan, Osarsal, Osvarsan, Paroxyl, Stovarsol.)

# **ACETARSONE SALT OF ARECOLINE.**

See: Drocarbil, N.F.

- **ACETASEM**, (Philips Roxane) Phenacetin 162.5 mg., aspirin 227.5 mg., caffeine 8.1 mg., gelsemium pwd. extract 4.62 mg./Tab. Bot. 1000s.

Use: Analgesic.

- **ACETAZIDE**, Acetazolamide, B.A.N.

- **ACETAZOLAMIDE**, U.S.P. XIX. N-(5-Sulfamoyl-1,3,4-thiadiazol-2-yl) acetamide. Acetamide, N-[5-(amino-sulfonyl)-1,3,4-thiadiazol-2-yl]-. Tab., U.S.P. XIX.

Use: Carbonic anhydrase inhibitor.

See: Diamox, Tab., Sequels (Lederle)

Hydrazol, Tab. (Softcon)

- **ACETAZOLAMIDE**, B.A.N. 5-Acetamido-1,3,4-thiadiazole-2-sulfonamide.

Use: Treatment of glaucoma.

See: Acetazide

Diamox

- **ACETAZOLAMIDE SODIUM**, U.S.P. XVIII. Sterile, U.S.P. XIX.

Use: Carbonic anhydrase inhibitor.

See: Diamox, Inj. (Lederle)

- **ACET-DIA-MER-SULFONAMIDE**, N.F. Sulfacetamide, sulfadiazine & sulfamerazine.

See: Buffonamide, Susp., Tab. (Tutag)

Bur-Sul, Tabs. & Susp. (Sheryl)

Cetazine, Tab. (Bowman)

Coco-Diazine, Susp. (Lilly)

Sulfadiazine, sulfamerazine & sulfacetamide,

Susp., N.F.; Tab. N.F.

- **ACETEST**, (Ames) Sod. nitroprusside, disodium phosphate, aminoacetic acid, lactose. Tab. Bot. 100s, 250s.

Use: Test for acetone or diacetic acid in urine.

- **ACETIC ACID**, U.S.P. XIX, Glacial, U.S.P. XIX. Diluted, N.F. XIV.

Use: Pharmaceutical aid (acidifying agent).

- **ACETIC ACID IRRIGATION**, U.S.P. XIX. 0.25% solution.

Use: Irrigating solution.

- **ACETIC ACID, POTASSIUM SALT**, Potassium Acetate, U.S.P. XIX.

# **ACETICYL**

See: Acetylsalicylic Acid.

# **ACETILUM ACIDULATUM.**

See: Acetylsalicylic Acid. (Various Mfr.)

- **ACETOHEXAMIDE**, U.S.P. XIX, Tab. U.S.P. XIX. N-(p-acetylphenylsulfonyl)-N'-cyclo-hexylurea. 1-((p-acetylphenyl)sulfonyl)-3-cyclohexylurea. Benzenesulfonamide, 4-acetyl-N-[(cyclohexylamino)carbonyl].

Use: Blood sugar lowering compound, antidiabetic.

See: Dymelor, Tab. (Lilly)

- **ACETOJEN, JR.** (Jenkins) Acetylsalicylic acid 1½ gr., phenacetin 1 gr., hyoscyamus ext. 0.0125 gr., (total alkaloids 0.0002 gr.) gelsemium ext. 0.0125 gr./Tab. Bot. 1000s.

- **ACETOLAX**, (Mills) Acetphenylisatin 5 mg., Vit. B-1 1 mg., sod. carboxymethyl cellulose 500 mg./Tab. Bulk Pkg. 1000s.

Use: Laxative.

# **ACETOL.**

See: Acetylsalicylic Acid. (Various Mfr.)

- **ACETOMEROCOTOL**, Acetato [2-hydroxy-5-(1,1,3,3-tetramethylbutyl) phenyl] mercury. Under study.

Use: Topical antiseptic.

- **ACETONE**, N.F. XIX.

# **ACETONE or DIACETIC ACID TEST.**

See: Acetest, Tab. (Ames)

- **ACETONYL**, (Upjohn) Granular efferv. salt. Aspirin as sod. salt 7½ gr. w/pot. & sod. citrates & tartrates having an alkalinity equiv. to 33½ gr. Na bicarbonate, calcium lactate, sod. phosphate & magnesium sulfate/60 gr. Bot. 4 oz.

Use: Analgesic & alkalinizer.

- **3-(alpha-ACETONYLBENZYL)-4-HYDROXYCOUMARIN SODIUM SALT**, Warfarin Sodium, U.S.P. XIX.

- **ACETOPHEN CAPSULE**, (Jenkins) Aspirin 3½ gr., phenacetin 2½ gr., caffeine ½ gr./Cap. Bot. 1000s.

- **ACETOPHEN C.T. GREEN**, (Jenkins) Aspirin 3½ gr., phenacetin 2½ gr., caffeine ¼ gr./Tab. Bot. 1000s.

- **ACETOPHENAZINE MALEATE**, N.F. XIV. Tab. N.F. XIV. 2-Acetyl-10- [3-[4-(β-hydroxyethyl)-piperazinyl] propyl] phenothiazine dimaleate. 10-[3-[4-(2-Hydroxyethyl)-1-piperazinyl]-propyl]-phenothiazin-2-yl methyl ketone maleate (1:2).

Use: Tranquilizer.

See: Tindal Maleate, Tab. (Schering)

- **ACETOPHENETIDIN**, U.S.P. XVI. Tablets. U.S.P. XVI. (Various Mfr.) Phenacetin, Ethoxyacetanilide.

Use: Antipyretic, analgesic.

- **ACETORPHINE**, B.A.N. O<sup>3</sup>-Acetyl-7,8-dihydro-7α(1R)-hydroxy-1-methylbutyl]-O<sup>6</sup>-methyl-6-, 14-endoethenomorphic.

Use: Narcotic analgesic.

# **ACETOSAL.**

See: Acetylsalicylic Acid. (Various Mfr.)

# **ACETOSALIC ACID.**

See: Acetylsalicylic Acid. (Various Mfr.)

# **ACETOSALIN.**

See: Acetylsalicylic Acid. (Various Mfr.)

- **ACETOSULFONE**, 4,4'-Diaminodiphenylsulfone-2-N-acetylsulfonamide sodium. (N<sup>1</sup>-acetyl-6-sulfanilyl-metanilamido) sodium.

- **ACETOTOT**, (Freeport) Aspirin 2.5 gr, chlorpheniramine maleate 1.25 mg., phenylephrine HCl 2.5 mg., buffered with aluminum & magnesium hydroxide/Tab. Bot. 1000s.

Use: Potent pediatric antihistamine & decongestant with buffered analgesic.

- **ACETOXYPHENYLMERCURY**, N.F. XIII.

See: Phenylmercuric acetate.

# **ACETPHENARSINE.**

See: Acetarsone, Tab. (City Chemical)

# **ACETPHENOLISATIN.**

See: Oxyphenisatin Acetate Prep.

W/D-Calcium pantothenate, sodium sulfosuccinate.

See: Peri-Pantyl, Tab. (McGregor)

W/Dioctyl sodium sulfosuccinate, sodium carboxy-

methylcellulose.

See: Scrip-Lax, Tab. (Scrip)

- **ACETRIZOATE SODIUM.** Injection, U.S.P. XVI. 3-Acetamido-2,4,6-triiodobenzoate sodium.

W/Polyvinylpyrrolidone.

See: Salpix, Vial (Ortho)

- **ACETRIZOIC ACID,** U.S.P. XV. 3-Acetylmino-2,4,6-triiodobenzoic acid.

See: Acetrizoate, Sod.

**ACETRYPTIN HYDROCHLORIDE.** (Warner-Chilcott) 5-Acetyltryptamine hydrochloride.

Use: Antihypertensive agent.

**ACET-THEOCIN SODIUM.**

See: Theophylline Sod. acetate (Various Mfr.)

**ACETYCOL** (O'Neal) Colchicine salicylated 0.25 mg., aspirin 325.0 mg., p-aminobenzoic acid 162.0 mg., Vit. C 20.0 mg., B-1 5.0 mg., niacin 15.0 mg./Tab. Bot. 100s.

Use: Analgesic.

**ACETYL ADALIN.**

See: Acetylcarbromal. (Various Mfr.)

**p-ACETYLAMINO BENZALDEHYDE THIOSEMICARBAZONE.** (Amithiozone, Antib, Berculon A, Benzothiozon, Conteben, Myuizone, Neustab, Tebethion, Thiomicid, Thioparamizone, Thiacetazone).

Use: With streptomycin in the treatment of tuberculosis.

**ACETYLAMINO BENZENE.**

See: Acetanilid. (Various Mfr.)

**N-ACETYL-p-AMINOPHENOL.** Acetaminophen, U.S.P. XIX.

**ACETYLANILINE.**

See: Acetanilid. (Various Mfr.)

**ACETYL-BETA-METHYLCHOLINE CHLORIDE,** U.S.P.

See: Mecholyl Chloride, Amp. (Baker, J.T.)

**ACETYL-BROMO-DIETHYLACETYL-CARBAMIDE.**

See: Acetylcarbromal. (Various Mfr.)

**ACETYLCARBROMAL** (Meyer) Acetyl adalin, acetylbromodiethylacetyl-carbamide. Pow. for mfg.

Use: Sedative.

See: Carbased, Tab. (Mallard)

Paxarel, Tab. (Circle)

Sedamyl. (Riker)

W/Bromisovalum, acetaminophen.

See: Isomel, Cap. (Tutag)

Medigesic, Tab. (Medics)

Vannor, Cap. (North Amer. Pharm.)

W/Bromisovalum, scopolamine aminoxide HBr.

See: Tranquinal, Tab. (Barnes-Hind)

**ACETYLCHOLINE CHLORIDE.**

Use: S.C., I.M., I.V. Parasympathomimetic agent & vasodilator. Paroxysmal tachycardia.

See: Miochol Ophthalmic (Smith, Miller & Patch)

**ACETYLCHOLINE-LIKE THERAPEUTIC AGENTS.**

See: Cholinergic agents.

- **ACETYLCYSTEINE,** N.F. XIV, Sol. N.F. XIV N-Acetyl-L-cysteine.

Use: Mucolytic agent.

See: Mucomyst, Sol. (Mead Johnson)

**ACETYLCYSTEINE.** B.A.N. N-Acetyl-L-cysteine. Air-bron

Use: Liquefaction of bronchial secretion.

- **ACETYLDIGITOXIN,** N.F. XIV, Tab, N.F. XIV.  $\alpha$ -Digitoxin monoacetate. Acetyltridigitoxose. The natural-occurring cardiac glycoside defined as the alpha acetyl ester of digitoxin.

Use: Digitoxin therapy. Cardiotonic.

See: Acylanid, Tab. (Sandoz)

**ACETYLDIHYDROCODEINONE.** Thebacon, B.A.N.

**ACETYLIN.**

See: Acetylsalicylic Acid. (Various Mfr.)

**ACETYLMETHADOL,** I.N.N. Methadyl acetate, B.A.N.

**ACETYLPHENYLISATIN.**

See: Oxyphenisatin Acetate.

**1-((p-ACETYLPHENYL)SULFONYL)-3-CYCLO-HEXYLUREA.** Acetohexamide, U.S.P. XIX.

**ACETYLRESORCINOL.**

See: Resorcinol Monoacetate. (Various Mfr.)

**ACETYLSAL.**

See: Acetylsalicylic Acid

**ACETYLSALICYLATE ALUMINUM.**

See: Aluminum Aspirin

- **ACETYLSALICYLIC ACID,** U.S.P. XVII. Capsules, N.F.; Tablets, U.S.P. XVII. (Various Mfr.) Aspirin, U.S.P. XVII. Acidum acetylsalicylicum, 2-acetoxybenzoic acid, Acetillum Acidulatum, Acetophen, Acetol, Acetosol, Acetosalic Acid, Acetosalin, Aceticyl, Acetylsal, Acylpyrin, Aspro, Helicon, Rhodine, Salacetin, Salcetogen, Saletin.

See: Aspirin Preps. (Various Mfr.)

**ACETYLSALICYLIC ACID, ACETOPHENETIDIN AND CAFFEINE,** N.F.

See: A.P.C., Prep. (Various Mfr.)

**ACETYL SULFAMETHOXYPYRIDAZINE.** N.D. 1967.

3-(N-Acetylsulfanilamido)-6-methoxypyridazine.

- **4'-(ACETYLSULFAMOYL)PHTHALANILIC ACID,** N.F. XIII.

See: Phthalylsulfacetamide

**N'-ACETYLSULFANILAMIDE.** (Albucid; p-Aminobenzenesulfonacetamide; Sulfacet; Sulfacetamide, N-Sulfanilylacetamide.)

Use: Sulfonamide therapy.

- **ACETYL SULFISOXAZOLE,** N.F. XIII, Oral Susp. N.F. XIII, N'-Acetyl-N'-(3,4-dimethyl-5-isoxazolyl)-sulfanilamide: N-(3,4-dimethyl-5-isoxazolyl)-N-sulfanilylacetamide.

See: Gantrisin Acetyl, Syr., Susp. (Roche)

Lipo-Gantrisin Acetyl, Liq. (Roche)

**ACETYLTANNIC ACID.** Tannic acid acetate.

Use: Antiperistaltic.

**5-ACETYLTRYPHTAMINE HYDROCHLORIDE.** (Warner Chilcott) Potent, rapid acting oral antihypertensive agent; pending release.

**ACHLORHYDRIA DETERMINATION.**

See: Diagnex Blue, Prep. (Squibb)

**ACHLORHYDRIA THERAPY.**

See: Acidol. (Various Mfr.)

Acidol-Pepsin, Cap. (Winthrop)

Glutamic Acid HCl (Various Mfr.)

Muripsin, Tab. (Norgine)

Normacid, Tab. (Stuart)

**ACHOL.** (Enzyme Process) Vit. A 4000 u., ketocholeonic acids 62 mg./Tab. Bot. 100s, 250s. Use: Vitamin A supplement and therapy.

**ACHOLEST.** (Fougera) Sebatic acid bromthynol blue.  
Kit w/30 tests.

Use: Plasma cholinesterase test kit.

**ACHROMYCIN.** (Lederle) Tetracycline HCl.

**Intramuscular:** (100 mg., 250 mg., buffered w/ascorbic acid) Vial. Box 1s, 100s.

**Intravenous:** (250 mg., 500 mg., buffered w/ascorbic acid) Vial.

**Oint.** (3%) Tube ½ & 1 oz.

**Oint. (Ophth.)** 1% Tubes ½ oz.

**Ophth. Susp.** 1%, Bot. w/dropper 4 cc.

**ACHROMYCIN-V.** (Lederle) Tetracycline HCl 100 mg./

**Cap.** Bot. 100s; 250 mg. Bot. 16s, 100s, 1000s;

Unit of use 50 × 20s, 50 × 28s, 50 × 40s; 500

mg. Bot. 100s, 1000s; Unit of use 50 × 20s; Unit dose 10 × 10s.

**Syr.** 125 mg./5 cc. w/methylparaben 0.12%,

propylparaben 0.03%. Bot. 2 oz., 1 pt.

Use: Treatment of infections caused by both gram-negative and gram-positive bacteria.

**ACHROSTATIN-V.** (Lederle) Tetracycline HCl 250 mg., nystatin 250M u./Cap. Bot. 16s, 100s.

Use: Broad-spectrum antibiotic w/antifungal therapy.

**ACICHEK.** (Valentine) Dihydroxy aluminum aminoacetate 250 mg., Sod. carboxymethylcellulose 150 mg., glycine 30 mg./Tab. Bot. 100s, 1000s.

Use: Gastric hyperacidity in peptic ulcer.

**ACID ACRIFLAVINE.**

See: Acriflavine Hydrochloride (Various Mfr.)

**ACID CITRATE DEXTROSE SOLUTION.**

See: A.C.D. Sol. (Various Mfr.)

**ACID CITRATE DEXTROSE ANTICOAGULANT SOLUTION MODIFIED.**

See: A-C-D Solution Modified (Squibb)

**ACI-DERMA.** (Princer) An aqueous emulsion of glyceryl monostearate, cetyl alcohol, sodium lauryl sulfate,

propyl and benzyl p-hydroxy benzoates and propylene glycol, buffered to pH 2-3. Bot. 4 oz., 1 pt.

Use: Restore the acid mantle of the skin.

**ACID HISTAMINE PHOSPHATE.**

See: Histamine Phosphate. (Various Mfr.)

**ACID MANTLE CREME.** (Dorsey) Aluminum acetate in specially prep. water-soluble hydrophilic cream at pH 4.2. Tube 1 oz., Jar 4 oz., 1 lb.

Use: Skin prophylactic agent in restoring normal skin acidity.

**ACID MANTLE LOTION.** (Dorsey) Aluminum acetate lotion. pH 4.5 Bot. 4 oz., 1 pt.

Use: Dermatitis.

**ACIDOL.**

See: Betaine Hydrochloride (Various Mfr.)

**ACIDOL-PEPSIN.** (Winthrop) Betaine HCl 4 gr., pepsin 3.5 gr./Cap. Bot. 50s.

Use: Achlorhydria.

**ACIDOPHILUS W/PECTIN.** (Barth's) Lactobacillus acidophilus w/natural citrus pectin 100 mg./Cap. Bot. 100s.

Use: Aid to digestion.

**ACID TRYPAFLAVINE.**

See: Acriflavine Hydrochloride. (Various Mfr.)

**ACIDULIN.** (Lilly) Glutamic acid HCl 340 mg./Pulvule.

Bot. 100s, 1000s.

Use: Achlorhydria.

**ACIGLUMIN.**

See: Glutamic Acid Hydrochloride. (Various Mfr.)

**ACI-JEL.** (Ortho) Acetic acid 0.92%, ricinoleic acid 0.70%, oxyquinoline sulfate 0.025%, boric acid 3.0%, and glycerine 5% w/Tragacanth, acacia, propylparaben, pot. bitartrate, stannous chloride, egg albumen, pot. bitartrate. Jelly, Tube with applicator, 85 Gm.

Use: Vaginal antiseptic.

**ACILLIN.** (ICN Pharm.) Ampicillin trihydrate equiv. to ampicillin 250 mg., 500 mg./Cap. Bot. 100s, 1000s.

Use: Penicillin therapy.

**ACINITRAZOLE.** B.A.N. 2-Acetamido-5-nitrothiazole, Aminotroazole (I.N.N.)

Use: Treatment of trichomoniasis.

See: Trichorad

Tritheon

**ACLETIN.** (Associated Concentrates) Lecithin, soya 8 gr. with vitamins/Cap. Bot. 100s, 250s.

Use: Fat metabolism.

**ACLOBROM.** Propantheline Bromide, B.A.N.

**A.C.N.** (Person & Covey) Vitamin A palmitate 25M u., ascorbic acid 250 mg., niacinamide 25 mg., hesperidin 50 mg./Tab. Bot. 100s, 1000s.

Use: Acne vulgaris.

**ACNAVEEN.** (Cooper) Aveeno colloidal oatmeal 8%, sulfur 2%, salicylic acid 2%, in a soap-free cream base. Plastic tube 4 oz. Bar 3.5 oz.

Use: Antipruritic cleanser and shampoo.

**ACNA-VITE.** (Cenci) Vit. A 10,000 u., C 250 mg., hesperidin 50 mg., niacinamide 25 mg./Cap. Bot. 75s.

Use: Acne vulgaris with Vit. A & C deficiency.

**ACNE-AID CREAM.** (Stiefel) Sulfur 2.5%, resorcinol 1.25%, parachlorometaxylene 0.375% w. Micel (brand of microporous cellulose). Tube, 1.8 oz.

Use: Oily acne.

**ACNE-AID DETERGENT SOAP.** (Stiefel) Natural soap with nonirritating detergent. Bar 4.1 oz.

Use: Cleanse oily skin.

**ACNE-AID LOTION.** (Stiefel) Sulfur 10%, alcohol 10% in a washable base. Clear or Tinted Bot. plastic 1 oz. or 2 oz.

Use: Treatment of acne.

**ACNE CREAM.** (De Pree) Sulfur 8%, resorcinol 2%, allantoin, zinc oxide, alcohol 9.25%. Tube 1½ oz.

Use: Dermatologic.

**ACNEDERM.** (Lannett) **Lotion:** Dispersable sulfur 5%, zinc sulfate 1%, zinc oxide 10%, alcohol 21%. Bot. 2 oz. **Soap:** Zinc oxide 2%, zinc sulfate 1%, colloidal sulfur 5%. Cake 3.2 oz.

Use: Skin conditions

**ACNE-DOME CREAM & LOTION.** (Dome) Regular: sulfur 4%, resorcinol monoacetate 3%, in a vehicle buffered to the pH range of normal skin.

**Cream:** Tube 1 oz. **Lotion:** Bot. 2 oz.

Use: Treatment of acne.

**ACNE-DOME MEDICATED CLEANSER.** (Dome) Sulfur 2%, salicylic acid 2% in a colloidal soybean complex. Medicated w/sponge 4 oz.

Use: Medicated skin cleanser.

**ACNESARB LOTION.** (C & M Pharmacal) Salicylic acid 3%, boric acid 2%, hyamine 10X 0.08%, isopropyl alcohol. Bot. 4 oz., pt., gal.  
Use: Anti-acne lotion.

**ACNETONE-ACNE LOTION.** (Whorton) Lot. 4 oz., 16 oz.

**ACNO.** (Cummins) Micronized sulfur 3%, isopropanol 25%, salicylic acid 2%. Bot. 8 oz. F-Lot. & M-Lot., Bot. 2 oz.

Use: Acne & related skin conditions.

**ACNOMEAD.** (Spencer-Mead) Colloidal sulfur 2%, salicylic acid 1% w/soapless cleansers, wetting agents, and polyethylene granules. Tube 1½ oz., jar 1 lb.

Use: Acne, scrub cream.

**ACNOMEL.** (SmithKline) Resorcinol 2%, sulfur 8%, w/bentonite, propylene glycol, alcohol & titanium dioxide.

**Cream.** Tube 1 oz. **Cake.** Sulfur 4%, resorcinol 1%. 1-oz. containers.

Use: Acne therapy.

**ACNOPHILL.** (Torch) Sulfur 5%, Pot. and zinc sulfides and polysulfides 5%, zinc oxide 10% in hydrophilic vehicle. Plain or tinted. Jar 1 lb. and 2 oz.

Use: Acne treatment.

**ACNOTEX.** (C & M Pharmacal) Sulfur 8%, salicylic acid 2.25%, hyamine 10X 0.08%, propylene glycol, powder base, acetone, isopropyl alcohol. Bot. 2 oz., pt., gal.

Use: Anti-acne lotion.

**ACON.** (Endo) Water-miscible vitamin A palmitate (50,000 i.u./Cap.) Bot. 100s.

Use: Vitamin A therapy.

• **ACONITINE.** U.S.P. X.

**ACOTUS.** (Whorton) Phenylephrine HCl 5 mg., guaiacol glyceryl ether 100 mg., menthol 1 mg., alcohol, by vol. 10%/5 cc. Bot. 4 oz., 12 oz., gal.  
Use: Colds, coughs.

• **ACRIFLAVINE.** N.F. X. (Euflavine, Gonacrine, Neutroflavine.) Acriflavine neutral; a mixture of 2, 8-diamino-10-methylacridinium chloride & 2,8-diaminoacridine. (Lilly) Tab. 1½ gr. Bot. 100s.  
Use: External antiseptic.

• **ACRIFLAVINE HYDROCHLORIDE.** N.F.X. (Various Mfr.) Hydrochloride form of acriflavine, N.F. Acid acriflavine, acid tryptaflavine, flavine, tryptaflavine. National Aniline—Pow., Bot. (1, 5, 10, 25, & 50 Gm.). Tab. (1½ gr.) Bot. 50s & 100s.  
Use: External bacteriostatic.

• **ACRISORCIN.** N.F. XIV., Cream, N.F. XIV. 9-Amino-acridinium 4-hexylresorcinolate.  
See: Akrinol Cream (Schering)

Use: Topical, tinea versicolor. Local antifungal.

• **ACRONINE.** (Lilly) 3, 12-Dihydro-6-methoxy-3,3,12-trimethyl-7H-pyrano[2,3-c]acridin-7-one. Under study.

Use: Antineoplastic.

**ACTAGEN.** (Generix) Triprolidine HCl 2.5 mg., pseudoephedrine HCl 60 mg. Tab. Bot. 100s, 1000s.

Use: Symptomatic treatment of allergic rhinitis.

**ACTAMER.** (Monsanto Chem.)

See: Bithionol.

**ACTAMIN.** (Buffington) Acetaminophen 325 mg./Tab. Dispens-A-Kit: 500s, MiniPak 200s.

Use: Pain reliever.

**ACTH-ACTEST GEL.** (Fellows) Repository corticotropin 40 or 80 u./cc. Vial 5 cc.

**ACTH.** (Adrenocorticotrophic hormone).

Adrenocorticotrop(h)in).

See: Corticotropin, U.S.P.

Mallard—(40 u./ml., 5 ml.)

O'Neal, Jones & Feldman—(40 u. or 80 u./ml., 5 ml.)

Parke, Davis—25 units/Vial; 40 units/Vial.

Pharmex—(40 u. or 80 u./ml., 5 ml.)

**ACTH GEL, PURIFIED.** Arcum—40 or 80 units/ml., vial 5 ml. Conal—40 or 80 units/ml., vial 5 ml. Hart Labs.—40 units/ml., vial 5 ml. Bowman—

Adrenocorticotrophic hormone 40 u., aqueous gelatin 16%, phenol 0.5%/cc. Vial 5 cc.

Use: Repository corticotropin.

Arcum—(40 u. or 80 u./ml., 5 ml.)

Bell—(40 u. or 80 u./ml., 5 ml.)

Bowman—(40 u./ml., 5 ml.)

Hyrex-Key—(40 u. or 80 u./ml., 5 ml.)

Jenkins—(40 u. or 80 u./ml., 5 ml.)

North Amer. Pharm.—40 units/ml., 80 units/ml.,

Vial 5 ml.

Rocky Mtn. Pharmacal—40 units/ml., 80 units/ml.

Vial 5 ml.

Wesley Pharm.—40 units/ml., vial 5 ml.; 80 units/ml., vial 5 ml.

Wyeth—(40 u. or 80 u./ml. or Tubex)

• **ACTHAR.** (Armour) Corticotropin Injection U.S.P. XIX. B.A.N. Vial 25, 40 u. (Lyophilized w/gelatin). Pkg. 10s.

**ACTHAR GEL HIGHLY PURIFIED.** (Armour) H.P. acthar gel, a new, highly purified, repository ACTH preparation that may be given subcutaneously or intramuscularly. 40 & 80 u. per cc., 1 cc., 5 cc. multiple dose vials, w/gelatin, phenol, & cysteine.  
Use: Stimulant of adrenal cortex, anti-inflammatory, antiallergic & useful in collagen diseases.

See: Corticotropin, U.S.P.

**ACTICORT LOTION 100.** (Cummins) Hydrocortisone alcohol 1% in a specially formulated vehicle. Bot. 2 oz.

Use: Anti-inflammatory & antipruritic.

**ACTIDL.** (Burroughs Wellcome) Triprolidine HCl Tab. (2.5 mg.) Bot. 100s. **Syr.** (1.25 mg./5 cc.) Bot. 1 pt.  
Use: Antihistamine.

**ACTI-DIONE.** (Upjohn) Cycloheximide. [3-(2-(3,5-Dimethyl-2-oxocyclohexyl)-2-hydroxyethyl)-glutarimide] 85-100% w/w.

Not for medicinal use. For professional use only.

Bot. 4 Gm.

**ACTIFED.** (Burroughs Wellcome) Triprolidine HCl. 2.5 mg., pseudoephedrine HCl. 60 mg./Tab. Bot. 30s, 100s, 1000s. Triprolidine HCl 1.25 mg., pseudoephedrine HCl 30 mg./5 cc. **Syr.** Bot. 4 oz., 1 pt., gal.

Unit dose pack 100s.

Use: Common cold, allergic rhinitis.

**ACTIFED-C EXPECTORANT.** (Burroughs Wellcome)

Codeine phosphate 10 mg., triprolidine HCl 2 mg., pseudoephedrine HCl 30 mg., guaifenesin 100 mg./5 cc. Bot. pt.

Use: Treatment of the common cold.

**ACTINOMYCIN.** B.A.N. Antimicrobial substances with antitumor activity produced by *Streptomyces* antibioticus and *Streptomyces chrysomallus*. Sanamycin is Actinomycin C.

**ACTINOMYCIN C.** Name previously used for Cactinomycin.

• **ACTINOMYCIN D.** Dactinomycin, U.S.P. XIX. See: Cosmegen (Merck Sharp & Dohme)

**ACTINOSPECTOCIN.** Name previously used for Spectinomycin.

**ACTIVATED ATTAPULGITE.**

W/Aluminum hydroxide, magnesium carbonate coprecipitate, compressed gel.

See: Hykasil, Cream (Philips Roxane)

W/Polysorbate 80, colloidal sulfur, salicylic acid, propylene glycol.

See: Sebasorb Lotion (Summers Labs.)

• **ACTIVATED 7-DEHYDROCHOLESTEROL,** U.S.P. XVI.

See: Vitamin D-3 (Various Mfr.)

**ACTIVATED ERGOSTEROL**

See: Calciferol, U.S.P.

**ACTIVOL.** (Carbisulphoil) Liquid soap. Bot. 1 gal, 55 gal. drum.

Use: Surgical Scrub-ups.

**ACTOL EXPECTORANT.** (Beecham Labs) Noscopine 30 mg., guaifenesin 200 mg./5 ml. or Tab. Syr.

W/alcohol 12.5%. Bot. pt. Tab. Bot. 100s.

Use: Treatment of symptoms associated with common cold.

**ACTOMOL.** Mebanazine, B.A.N.

**ACTOQUINOL SODIUM.** 8-Ethoxy-5-quinoline-sulfonic acid sodium salt.

Use: Ultraviolet screen.

**ACTOS 50.** Corticotrophin, B.A.N.

**ACTOSPAR.** (Sandoz) Sparteine sulfate.

**ACTRIOL.** Epioestriol, B.A.N.

**ACTYLATE.** (Kinney) Ammonium salicylate 80 mg., pot. salicylate 80 mg., strontium salicylate 80 mg., pot. para-aminobenzoate 0.32 Gm., ascorbic acid 20 mg./Tab. Bot. 100s.

Use: Rheumatoid arthritis.

**ACUDERM.** (Nortech) Zinc oxide, starch, talc, castor oil, petrolatum. Oint. Tube 1 oz.

**ACUSONE OINTMENT.** (Nortech) Hydrocortisone 0.5%, zinc oxide, starch, talc, castor oil, petrolatum. Oint.: Tube ½ oz., 1 oz.

**ACUTRAN.** (Mallinckrodt) A.nphecloral.

**ACUTUSS EXPECTORANT W/CODEINE.** (Philips Roxane) Codeine phosphate 10 mg., phenylpropanolamine HCl 5 mg., phenylephrine HCl 5 mg., chlorpheniramine maleate 2 mg., guaifenesin 100 mg., alcohol 5%/Tsp. (5 cc.) Bot. 8 oz., gal.

Use: For acute cough.

**ACYLANID.** (Sandoz) Acetyldigitoxin. Tab. 0.1 mg. (pink) 100s; 0.2 mg., (white) 100s.

Use: Digitoxin therapy.

**ACYLPYRIN.**

See: Aspirin

**A AND D CREAM.** (Schering) Emollients, isopropyl myristate & squalene. Tube 2 oz.

Use: Dry skin, sunburn, irritations.

**A AND D HEMORRHOIDAL SUPPOSITORIES WITH HC.** (Schering) Hydrocortisone acetate 10 mg., Vit. A 1500 u., Vit. D-3 200 u., Peruvian balsam, bismuth subgallate, zinc oxide/Supp. Box 12s.

Use: Relief of internal hemorrhoids.

**A AND D OINTMENT.** (Kenyon) Vits. A & D in lanolin-petrolatum base. Jar 1 lb.

**A AND D OINTMENT.** (Schering) Petrolatum-lanolin. Tube 1½, 4 oz.; Jar 1 lb.

**A & D TABLETS.** (Barth's) Vit. A 10,000 u., D 400 u./Tab. Bot. 100s, 500s.

**A AND D VITAMIN CAPSULES.** (Lannett) Vit. A 5000 u., D 400 u./Cap. Bot. 500s, 1000s.

Use: Vit. supplement.

**A AND D VITAMIN CREAM.** (DePree) Bot. 1½ oz., 4 oz.

**A AND D VITAMIN OINTMENT.** (Lannett) In a petrolatum-lanolin base. Jar 1, 5 lb.

**A AND D VITAMIN OINTMENT.** (Ulmer) Jar 1 lb.

**A-D DRESSING.** (La Crosse) Vit. A 6000 u., D 600 u., allantoin 2%. Tubes 1 oz., Jars 1 lb., 5 lb.

Use: Multi-purpose dressing.

**A-D DRESSING.** (La Crosse) Vit. A 6000 u., D 600 u., allantoin 2%/oz. Tube 1 oz. Jar 1 lb., 5 lb.

Use: Antiseptic-antibacterial ointment.

**A-D-A.** (LaCrosse) Amobarbital 60 mg., dextroamphetamine sulfate 15 mg./Cap. Bot. 100s, 1000s.

Use: Obesity.

**ADABEE & ADABEE with MINERALS.** (Robins) Vit. A 10,000 u., C 250 mg., B-1 15 mg., B-2 10 mg., B-6 5 mg., niacinamide 50 mg./Tab. Bot. 100s, 500s.

**Adabee with Minerals.** In addition to above formula, iron 15 mg., calcium 103 mg., phosphorus 80 mg./Tab. Bot. 100s, 500s.

Use: Vitamin supplement.

**ADALAN LANATABS.** (Lannett) Amobarbital 50 mg., homatropine methylbromide 7.5 mg., methamphetamine HCl 10 mg./T.D. Tab. Bot. 100s, 1000s.

**ADAMANTANAMINE HCl.** (Endo) Amantadine HCl. Anti-flu capsule. This drug is thought to protect cells against entry of the flu virus without actually destroying the virus. Also used in the treatment of Parkinson's disease.

See: Symmetrel, Cap., Syr. (Endo)

**ADAMYCIN TABS.** (Lederle) Erythromycin (as stearate) 250 mg., or 500 mg./Tab. Bot. 100s.

**ADANON.** Methadone, B.A.N.

**ADAPETTES.** (Burton, Parsons) Cleans and wets contact lens. Pkg. 15 cc.

**ADAPIN.** (Pennwalt) Doxepin HCl 10 mg., 25 mg. 50 mg./Cap. Bot. 100s, 1000s.

Use: Antianxiety, antidepressant.

**ADAPT.** (Burton, Parsons) Povidone, EDTA 0.1%, thimerosal 0.004%. Wetting sol. Bot. 15 cc.

Use: Contact lens wetting sol.

**ADATUSS.** (W. F. Merchant) Hydrocodone bitartrate 5 mg., glyceryl guaiacolate 100 mg., alcohol 5%/5 ml. Syr. Bot. pt., gal.  
Use: Cough relief.

**A.D.C. DROPS.** (Parke, Davis) Vit. A 5000 u., D 400 u., C 50 mg., B-6 1 mg./0.6 cc. Dropper Bot. ½ oz., 1½ oz.  
Use: Vitamin deficiencies in infants and children.

**A.D.C. DROPS.** (Consol. Midland) Vit. A 3000 u., D 400 u., C 60 mg./0.6 cc. Bot. 60 cc.

**ADC DROPS WITH FLUORIDE.** (Parke, Davis) Vit. A 5000 u., D 400 u., C 50 mg., B-6 1 mg., sodium fluoride 1.1 mg. Dropper. Bot. 50 ml.  
Use: Vitamin therapy with fluoride.

**AD-CEBRIN with FLUORIDE DROPS.** (Lilly) Vitamin A 4,000 u., D 400 u., C 60 mg., fluoride 0.5 mg./0.6 cc. Dropper Bot. 30 ml.  
Use: Dietary supplement & prevention of dental caries.

**ADCCOTYL.** Triamcinolone, B.A.N.

**ADEECON.** (Consol. Midland) Vit. A 5000 u., D 1000 u./Cap. Bot. 1000s.  
Use: Supplement.

**ADEFLO B DROPS.** (Upjohn) Vitamin A 2000 I.U., D 400 I.U., B-1 1 mg., B-2 1 mg., C 50 mg., niacinamide 10 mg., B-6 1 mg., d-pantothenyl alcohol 3 mg., fluoride (from sodium fluoride) 0.5 mg./0.6 cc. Calibrated drop. Bot. 50 cc.  
Use: Aid in the prevention of dental caries w/vitamin supplement.

**ADEFLO CHEWABLE.** (Upjohn) Fluoride 1 mg. & 0.5 mg., Vitamin A 4000 I.U., D 400 I.U., C 75 mg., thiamine monitrate (B-1) 2 mg., B-2 2 mg., niacinamide 18 mg., B-6 1 mg., calcium pantothenate 5 mg., B-12 2 mcg./0.5 or 1 mg. Chewable Tab. Bot. 100s, 500s.  
Use: Prevention of dental caries.

**ADEFLO DROPS.** (Upjohn) Vit. A 2000 I.U., D 400 I.U., C 50 mg., B-6 1 mg., fluoride (as sodium fluoride) 0.5 mg./0.6 cc. 30 or 50 cc. Bot. w/dropper.  
Use: Prevention of dental caries.

**ADEFLO M TABLETS.** (Upjohn) Vit. A 6000 I.U., D 400 I.U., B-1 1.5 mg., B-2 2.5 mg., C 100 mg., niacinamide 20 mg., Vit. B-6 10 mg., calcium pantothenate 10 mg., Vit. B-12 2 mcg., folic acid 0.1 mg., fluoride (as sodium fluoride) 1 mg., calcium (as calcium carbonate) 250 mg., iron 30 mg./Tab. Bot. 100s, 500s.  
Use: Dietary supplement.

**ADENIC.** (Kenyon) Adenosine-5-monophosphate 25 mg., B-12 75 mcg., niacin 20 mg./cc. Bot. 10 cc.  
Use: Anti-inflammatory.

**ADENIC FORTE GEL.** (Myers-Carter) Adenosine-5-monophosphate 25 mg., B-12 300 mcg., niacin 10 mg., methionine 75 mg./cc. Vial 10 cc.  
Use: Anti-inflammatory.

**ADENINE.** (Various Mfr.) 6-Aminopurine.

**ADENO-B.** (Blaine) Adenosine 50 mg., Vit. B-12 120 mcg./cc. Vial 10 cc.

**ADENO TWELVE GEL INJECTION.** (Fellows) Adenosine-5-monophosphate 25 mg., Vit. B-12 300

mcg., methionine 25 mg., niacin 10 mg./cc. Vial 10 cc.

**ADENOCREST.** (Nutrition) Adenosine-5-monophosphate sod. 25 mg., Vit. B-12 100 mcg., nicotinic acid as sod. salt 25 mg., gelatin 100 mg. w/benzyl alcohol 1.5%/cc. Vials 10 cc.

Use: Capillary and arterial vasodilation.

**ADENOLIN FORTE.** (Lincoln) Adenosine-5-monophosphate, 25 mg., Vit. B-12 300 mcg., methionine 25 mg., niacin 10 mg./cc. Vial 15 cc.

Use: Arthritis, bursitis, tendinitis & other degenerative diseases.

• **ADENOSINE.** F.D.A. 6-Amino-9-beta-D-Ribofuranosyl-9H-purine.

**ADENOSINE B-12.** (Vale) Adenosine-5-phosphoric acid 25 mg., cyanocobalamin 100 mcg./ml. Vial 10 ml.

**ADENOSINE B-12 GEL.** (Medical Chem.) Adenosine-5-monophosphate 100 mg., Vit. B-12 100 mcg./cc. Vial 10 cc.

**ADENOSINE B-12-NICOTINIC.** (Med. Chem.) Adenosine-5-monophosphate 25 mg., B-12 75 mg., nicotinic acid 20 mg./cc. Vial 10 cc.

**ADENOSINE & VIT. B-12 IN GELATIN.** (Fellows) **Forte:** Adenosine-5-monophosphate 50 mg., Vit. B-12 100 mcg./cc. **Super:** Adenosine-5-monophosphate 100 mg., Vit. B-12 100 mcg./cc.

**ADENOSINE W/VITAMIN B-12.** (Med. Chem.) Adenosine-5-monophosphoric acid 25 mg., Vit. B-12 60 mcg., water q. s./cc. Vials 10 cc.

**ADENOSINE COMBINATION SOLUTION.** (Ries) Adenosine-5-monophosphoric acid 25 mg., Vit. B-12 100 mcg. Vial 10 cc.

**ADENOSINE-5-MONOPHOSPHATE AS THE SODIUM SALT.**

See: Adenic, Prods. (Myers-Carter)

Adenocrest, Amp. (Nutrition)

W/Vit. B-12

See: Adenosine B-12 Gel. (Medical Chem.)

**ADENOSINE-5-MONOPHOSPHORIC ACID W/CYANOCOBALAMINE.**

See: Adenosine B-12. (Var. Mfr.)

Myo-B, Inj. (Sig)

Vitasine-Gel, Inj. (Burgin Arden)

**ADENOSINE GEL FORTE.** (Douglas) Adenosine-5-monophosphate 25 mg., sodium nicotinate 20 mg., Vit. B-12 25 mcg./cc. in gelatin base 4%. Bot. 10 cc.

• **ADENOSINE PHOSPHATE.** 5-Adenylic acid. B.A.N. Adenosine-5'-(dihydrogen phosphate). Adenosine monophosphate, AMP.

Use: Nutrient.

See: Adenyl

Cobalazine, Inj. (Keene Pharm.)

My-B-Den (Dome)

**ADENOTINATE.** (Medics) Adenosine-5-monophosphate 25 mg., Vit. B-12 75 mcg., nicotinic acid 20 mg., benzyl alcohol/cc. Vials 10 cc.

Use: Neuritis, bursitis, etc.

**ADENYL.** Adenosine Phosphate, B.A.N.

**ADEPSINE OIL**

See: Petrolatum Liquid (Various Mfr.)

**ADHES-OFF.** (Delta) Spray can 3 oz.

Use: Tape removal.