

# ACRIDINES

*Edited by*

R. M. Acheson

SECOND EDITION

---

---

# ACRIDINES

*Edited by*

**R. M. Acheson**

*The Department of Biochemistry*

*and*

*The Queen's College*

*University of Oxford*

**SECOND EDITION**

---

---

**INTERSCIENCE PUBLISHERS**

*a division of*

**JOHN WILEY & SONS**

**NEW YORK • LONDON • SYDNEY • TORONTO**

---

---

Copyright © 1973, by John Wiley & Sons, Inc.

All rights reserved. Published simultaneously in Canada.

No part of this book may be reproduced by any means, nor transmitted, nor translated into a machine language without the written permission of the publisher.

***Library of Congress Cataloging in Publication Data:***

Acheson, Richard Morris.

Acridines.

(The Chemistry of heterocyclic compounds, 9)

Includes bibliographies.

1. Acridine. [DNLM: 1. Acridines. W1 CH364H v. 9  
1973. XNLM: [QD 401 A187 1972]]

QD401.A23 1973      547'.593      72-5847

ISBN 0-471-37753-8

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

## Contributors

- R. M. Acheson**, *The Department of Biochemistry and The Queen's College, University of Oxford, England*
- B. A. Adcock**, *Flintshire College of Technology, Flintshire, England*
- N. R. Ayyangar**, *National Chemical Laboratory, Poona, India*
- Margaret L. Bailey**, *Chemistry Department, Victoria University of Wellington, Wellington, New Zealand*
- R. G. Bolton**, *I.C.I. Pharmaceuticals Division Research Laboratories, Cheshire, England*
- David B. Clayson**, *Department of Experimental Pathology and Cancer Research, School of Medicine, Leeds, England*
- A. C. R. Dean**, *Physical Chemistry Laboratory, University of Oxford, England*
- J. M. F. Gagan**, *Department of Chemistry and Chemical Technology, Bradford University, Bradford, England*
- David W. Henry**, *Department of Bio-Organic Chemistry, Stanford Research Institute, Menlo Park, California*
- Frank McCapra**, *The Chemical Laboratory, University of Sussex, Brighton, England*
- B. H. Nicholson**, *Department of Physiology and Biochemistry, The University of Reading, Whiteknights Reading, England*
- A. R. Peacocke**, *St. Peter's College, Oxford, England*
- N. R. Raulins**, *Department of Chemistry, University of Wyoming, Laramie, Wyoming*
- D. A. Robinson**, *Molecular Pharmacology Research Unit, Medical Research Council, Cambridge, England*
- J. E. Saxton**, *Department of Organic Chemistry, The University of Leeds, Leeds, England*
- I. A. Selby**, *Pharmaceutical Division, Reckitt and Colman, Hull, England*
- B. D. Tilak**, *Director, National Chemical Laboratory, Poona, India*

## The Chemistry of Heterocyclic Compounds

The chemistry of heterocyclic compounds is one of the most complex branches of organic chemistry. It is equally interesting for its theoretical implications, for the diversity of its synthetic procedures, and for the physiological and industrial significance of heterocyclic compounds.

A field of such importance and intrinsic difficulty should be made as readily accessible as possible, and the lack of a modern detailed and comprehensive presentation of heterocyclic chemistry is therefore keenly felt. It is the intention of the present series to fill this gap by expert presentations of the various branches of heterocyclic chemistry. The subdivisions have been designed to cover the field in its entirety by monographs which reflect the importance and the interrelations of the various compounds, and accommodate the specific interests of the authors.

In order to continue to make heterocyclic chemistry as readily accessible as possible, new editions are planned for those areas where the respective volumes in the first edition have become obsolete by overwhelming progress. If, however, the changes are not too great so that the first editions can be brought up-to-date by supplementary volumes, supplements to the respective volumes will be published in the first edition.

ARNOLD WEISSBERGER

*Research Laboratories  
Eastman Kodak Company  
Rochester, New York*

EDWARD C. TAYLOR

*Princeton University  
Princeton, New Jersey*

## Preface

In the 15 years since the publication of the first edition, there have been many developments in acridine chemistry and biochemistry. These have taken place over a broad front, too broad in fact for one person to review the field both adequately and rapidly enough to ensure up-to-date publication. I have been very fortunate in having a number of colleagues who were willing and able to find the time necessary to revise the chapters of the first edition, or to write entirely new chapters where necessary, for this new edition. Building a book of contributed chapters always presents difficulties of possible duplication and accidental omission of material that is on the borderlines of two or more chapters. A small amount of overlap has resulted, but this has been left in order to maintain continuity in the individual chapters.

All those working with acridines should be grateful that agreement has now been achieved concerning the numbering of the acridine ring. The agreed system is that employed in both editions of this book, and by *Chemical Abstracts*, and the *Ring Index*, and recommended by the International Union of Pure and Applied Chemistry.

I should like to thank most sincerely all my co-authors, all of whom have given up much leisure time in order to help. I also wish to thank The Queen's College and The University of Oxford for leave during which the manuscript was edited, and Professor C. A. Grob for the hospitality of the Institute for Organic Chemistry, The University of Basel, which was greatly appreciated.

I also thank Miss M. B. Acheson, Mrs. R. F. Flowerday, Messers P. J. Abbott, M. P. Acheson, and N. D. Wright for their help in preparing the index, and Miss P. Lloyd for typing the copy. Last, but not least, I must express my gratitude to all those working in Wiley-Interscience and associated with this volume for their advice and continuous assistance, which greatly facilitated my task.

R. M. ACHESON

March 1972  
The Department of Biochemistry  
and  
The Queen's College  
University of Oxford  
Oxford, England

## Contents

<b>Nomenclature and Numbering System</b>	<b>1</b>
R. M. ACHESON	
<b>I. Acridines</b>	<b>9</b>
N. R. RAULINS	
<b>II. Aminoacridines</b>	<b>109</b>
B. ADCOCK	
<b>III. 9-Acridanones</b>	<b>141</b>
J. M. F. GAGAN	
<b>IV. The Acridine Alkaloids</b>	<b>379</b>
J. E. SAXTON	
<b>V. Acridinium Salts and Reduced Acridines</b>	<b>433</b>
I. A. SELBY	
<b>VI. Biacridines</b>	<b>519</b>
FRANK MCCAPRA	
<b>VII. Benzacridines and Condensed Acridines</b>	<b>529</b>
D. A. ROBINSON	
<b>VIII. Acridine Dyes</b>	<b>579</b>
B. D. TILAK AND N. R. AYYANGAR	
<b>IX. Chemiluminescent Reactions of Acridines</b>	<b>615</b>
FRANK MCCAPRA	

<b>X. Ultraviolet and Visible Absorption Spectra</b>	<b>631</b>
MARGARET L. BAILEY	
<b>XI. The Infrared Spectra of Acridines</b>	<b>665</b>
R. M. ACHESON	
<b>XII. The Nuclear Magnetic Resonance Spectra of Acridines</b>	<b>687</b>
R. M. ACHESON	
<b>XIII. The Mass Spectra of Acridines</b>	<b>709</b>
R. G. BOLTON	
<b>XIV. The Interaction of Acridines with Nucleic Acids</b>	<b>723</b>
A. R. PEACOCKE	
<b>XV. Acridines and Enzymes</b>	<b>759</b>
B. H. NICHOLSON	
<b>XVI. The Antibacterial Action of Acridines</b>	<b>789</b>
A. C. R. DEAN	
<b>XVII. Carcinogenic and Anticarcinogenic Properties of Acridines</b>	<b>815</b>
DAVID B. CLAYSON	
<b>XVIII. Acridine Antimalarials</b>	<b>829</b>
DAVID W. HENRY	
<b>Index</b>	<b>851</b>



# Nomenclature and Numbering System


R. M. ACHESON

*Department of Biochemistry, and The Queen's College,  
University of Oxford, England*


The discovery of a new basic material in the anthracene fraction of coal tar was announced by Graebe and Caro<sup>1</sup> in 1870. On account of its acrid smell and irritating action on the skin and mucous membrane, this new substance was called "acridin" (acris = sharp, or pungent). Apart from the addition of a terminal "e," and the replacement of the "c" by "k" in a few older papers, this name has not subsequently been changed. The general nomenclature and numbering system employed for acridine and its derivatives in this monograph is the same as that used in *Chemical Abstracts* since 1937.

At least seven different systems of numbering have been used for the simple acridine ring system, and many more methods have been proposed for fused ring systems containing the acridine nucleus. Much difficulty, therefore, arises when a literature search for acridine derivatives becomes necessary, especially in regard to early publications.


The first numbering system (1) was suggested by Hess and Bernthsen<sup>2</sup> but found no support. Schöpf's system (2), suggested in 1892,<sup>3</sup> was almost as unpopular, although a slight modification (3) was used in 1922 and occasionally later.<sup>4</sup>



1






2





3

In 1893 Graebe,<sup>5</sup> the discoverer of acridine, suggested a numbering system (4) based on the then accepted numbering used for anthracene, xanthene, etc. This system was generally approved at the time. In 1900, however, method



(5), which was largely ignored, was propounded by von Richter in his textbook,<sup>6</sup> while in the same year M. M. Richter<sup>7</sup> used another system (6) in his *Lexikon der Kohlenstoff-Verbindungen*.



This method of numbering gained some popularity. Borsche<sup>8</sup> suggested another system (7), which did not find acceptance. In 1921 Stelzner<sup>9</sup> extended Graebe's system to 8.



Another variation (9) suggested by Patterson,<sup>10</sup> which is in conformity with the *Ring Index* rules,<sup>11</sup> was used in a standard textbook<sup>12</sup> in 1926, but the numbering was changed to that of Graebe in the 1936 edition of the book. Yet another system<sup>13</sup> (10) appeared in 1947, adding to the confusion.



At the present time only two systems of numbering, 4 and 6, are significantly used for acridine derivatives. A record was kept during the preparation of the first edition (1956) of this monograph of the number and year of publication of all original papers available and referred to (whether mentioned subsequently in the monograph or not) using these systems. Figure 1 shows the results. From the graph it is clear that (1) the great majority of publications use Graebe's system; and (2) although an increase in the popularity of Richter's system in recent years is evident, Graebe's system is still used in the majority of publications. The *Ring Index* has adopted the latter system, which is the same as that generally used for anthracene, although it is not consistent with Patterson's rules for the numbering of cyclic compounds.




Fig. 1. The number of papers published yearly using Graebe's (—) and Richter's (.....) systems.

Graebe's system of numbering is used in *Beilstein's Handbuch* and in *Chemisches Zentralblatt*. Until 1937 *Chemical Abstracts* officially used Richter's system of numbering; at that time a change was made to Graebe's system. *British Chemical Society Abstracts* uses Richter's system, which was also employed by Albert.<sup>14</sup> Although the British and American abstracts officially use particular methods of numbering, papers in which the numbering of acridine derivatives differs from that officially used in the abstracts may be abstracted in the original numberings. This calls for much care when using abstracts. The stage has even been reached at which both systems are used in the *same* chapter of a standard textbook.<sup>15</sup>

The continued use of two numbering systems for the acridine ring system added to the nomenclature difficulties, so that a decision as to which system should be used in the future became essential. This decision has recently been made (see below). The arguments put forward by Albert<sup>14</sup> in support of Richter's system (6) are:

(a) It has been used in a significant portion of the literature including almost the whole of the literature in the English language; (b) it makes no suggestion of there being more than five possible mono-derivatives whereas 4 gives rise to such names


as 9-chloroacridine which suggests that there are nine; and (c) a number of medical products have been introduced in Britain and Australia in recent years not under trade names but described systematically according to 6—e.g. 5-aminoacridine; it would be confusing to the physician if they should simultaneously be numbered in two ways, and he would be apt to think that two isomeric substances are involved.

The first statement is incorrect; the increase in popularity of Graebe's system is largely a result of its use in American publications. With regard to (b), it is necessary to point out that many accepted numbering systems involve this disadvantage, which is now unlikely to be a source of confusion. As to (c), the medical profession can be little interested in acridine nomenclature, since the systematic name "5-aminoacridine hydrochloride" (system 6) has been officially replaced by "Aminacrine Hydrochloride B.P.," and the trade names for this substance, e.g., "Acramine," "Dermacrine," and "Monacrin," are widely used. Incidentally, in spite of his support for Richter's system, Albert has, in fact, found it convenient to use Graebe's system in one section (on carbazimes) in the first edition of his own monograph.<sup>14</sup>

From the evidence discussed above, it is clear that Graebe's system (4), which is employed throughout this monograph, is used in the large majority (ca. 75.3%) of the papers published before 1955 in which a numbering system is required, and in the more important abstract journals.




Richter's system (6) offers no particular advantage to compensate for its minority position and should therefore be completely dispensed with in the future. This view has also been taken by the Union of Pure and Applied Chemistry<sup>16</sup> and now appears to have been generally accepted.<sup>17</sup>

The carbon atoms common to both rings can bear substituents only when the acridine ring is suitably reduced, and in this situation it becomes necessary to number these positions. Albert<sup>17</sup> employs Stelzner's extension (7) of Graebe's system for these carbon atoms. This use is unfortunate and should be discontinued. It is inconsistent with the generally agreed rules for the numbering of such atoms, and it is inconsistent with current *Chemical Abstracts* practice. The atoms common to both rings should be numbered as shown in structure 11.



Many arbitrary numberings, as well as generic numberings, have been used for fused ring systems containing the acridine nucleus. Generic numbering is built up from the numberings of the constituent ring systems regarded as being fused together, subscripts being used to differentiate between the figures. In this system 12 is 1',2'-1,2-benzacridine, or 1,2-benzacridine, since this

simplification leads to no ambiguity. This type of numbering is much used in the German literature. A similar system, based on Richter's method of numbering acridine, is also largely used in *British Chemical Abstracts*. It was often used in *Chemical Abstracts* until 1937, when the *Ring Index* numbering was adopted for condensed acridines. In order to minimize the use of numbers and subscripts, the sides of the parent ring system have been lettered *a, b, c, . . .* (13) starting from position 1. Thus 1,2-benzacridine (12) becomes benz[*a*]-acridine, and 14 is 3'-methylbenz[*a*]acridine.













Although generic numbering for comparatively simple, fused ring systems has the advantage of being easily worked out from the numbering of the constituent ring systems, it has the disadvantage of requiring the use of many figures and brackets. For this reason, and in order to conform to current usage in *Chemical Abstracts*, the arbitrary *Ring Index*<sup>10,18</sup> numbering of condensed acridine derivatives has been adopted here. 14 is then known as 4-methylbenz[*a*]acridine. The main disadvantage in using an arbitrary numbering system for polycyclic compounds is that similar compounds may have substantially different numberings. This, however, is a small price to pay for the convenience offered by a generally accepted, and used, numbering system. In view of the variety of current methods of numbering employed for condensed acridines, as much care is necessary when making a literature search for particular compounds of this type as in the case of simple acridine derivatives.

A variety of unsystematic and misleading names have been used in the past for most condensed acridines, an example being benz[*a*]acridine, which has also been referred to as  $\beta$ -chrysidine,  $\beta$ -phenonaphthacridine and 1,2-naphthacridine. Misleading nomenclature of the latter type is hardly ever used in current literature and should not be revived. The only remaining example of unsystematic nomenclature in the acridine series officially used by *Chemical Abstracts* until about 1957 is that of "carbazime" for 2,9-dihydro-2-iminoacridine. Obsolete names, as well as the *Chemical Abstracts* names and numberings used in this monograph for condensed acridines, are given in many cases when the individual compounds are discussed.



In the naming of a substituted acridine derivative, there are still two points to be considered. First, the lowest possible numbers should be chosen for the substituents; a full discussion of this is available in *Chemical Abstracts*<sup>19</sup>

TABLE I. Numbering Systems used for Acridine and the Benzacridines

Compound and Ring Index number <sup>18</sup>	Numbering used here and in <i>Chem.</i> <i>Abstr.</i> from 1937	Alternative (minority) used in <i>Chem.</i> <i>Abstr.</i> before 1937
Acridine R.R.I. 3523 (R.I. 1973) <sup>10</sup>		
Benz[a]acridine R.R.I. 5144 (R.I. 2735) <sup>10</sup>		
Benz[b]acridine R.R.I. 5140 (R.I. 2731) <sup>10</sup>		
Benz[c]acridine R.R.I. 5148 (R.I. 2737) <sup>10</sup>		
7(H)-Benz[k]acridine R.R.I. 5564 (R.I. not listed) <sup>10</sup>		
1, 7(H)-Pyrid[3,2,1- de]acridine R.R.I. 5104 (R.I. 2712) <sup>10</sup>		

Note: The *Chemical Abstracts* numberings for more complicated condensed compounds are given in the sections in which the compounds are discussed.

and in International Union of Pure and Applied Chemistry publications.<sup>20</sup> Since the acridine ring can be numbered in either direction, **15** may be called either 9-chloro-6-dimethylamino-7-fluoro-4,5-dimethylacridine or 9-chloro-3-dimethylamino-2-fluoro-4,5-dimethylacridine, the latter being "correct." Such a compound, however, may be found "incorrectly" numbered in *Chemical Abstracts*. Second, substituents should be placed in alphabetical order, regardless of their number or position in the molecule, and compound radical names should be treated as a unit according to their first letter. Agreement on this issue has been reached between American and British workers.<sup>21</sup> For instance **15** should be called 9-chloro-3-dimethylamino-2-fluoro-4,5-dimethylacridine, and not 9-chloro-4,5-dimethyl-3-dimethylamino-2-fluoroacridine. An order of preference for the last radical in the naming of complex organic compounds is also used by *Chemical Abstracts*: onium compound, acid, acid halide, amide, imide, amidine, aldehyde, nitrile, isocyanide, ketone, alcohol, thiol, amine, imine, ether, sulfide, sulfoxide, sulfone, etc. On this basis **16** should be known as 2-(6-chloro-2-methoxy-9-acridylamino)-ethanol.



However, the heterocyclic part of this molecule is generally considered more important than the aliphatic part, so that the compound is called 6-chloro-9-(2-hydroxyethylamino)-2-methoxyacridine in this monograph and usually in *Chemical Abstracts*, where it may also be indexed under its alternative name.

The system of numbering approved by the *Ring Index* and by the International Union of Pure and Applied Chemistry, used in this monograph and officially used in *Chemical Abstracts* for acridine and the benzacridines, and alternative numberings used in *Chemical Abstracts* before 1937, are given in Table I.

## References

1. C. Graebe and H. Caro, *Chem. Ber.*, **3**, 746 (1870).
2. W. Hess and A. Bernthsen, *Chem. Ber.*, **13**, 689 (1885).
3. M. Schöpf, *Chem. Ber.*, **25**, 1980 (1892).
4. C. H. Browning, J. B. Cohen, R. Gaunt, and R. Gulbransen, *Proc. Roy. Soc., Ser. B*, **93**, 329 (1922).

5. C. Graebe and K. Lagodzinski, *Justus Liebigs Ann. Chem.*, **276**, 35 (1893).
6. V. von Richter, *Organic Chemistry*, 3rd American ed., from the 8th German ed., 1900, p. 559.
7. M. M. Richter, *Lexikon der Kohlenstoff-Verbindungen*, Voss, Hamburg, and Leipzig, 1900, p. 24.
8. W. Borsche, *Justus Liebigs Ann. Chem.*, **377**, 70 (1910).
9. R. Stelzner, *Literatur-Register der organischen Chemie*, Vieweg and Sohn, Braunschweig, **3**, 59 (1921).
10. A. M. Patterson, *J. Amer. Chem. Soc.*, **47**, 556 (1925).
11. A. M. Patterson and L. T. Capell, *The Ring Index*, 1st ed. Reinhold, New York, 1940.
12. J. Schmidt, *Organic Chemistry*, 1st English ed., transl. by H. G. Rule, Gurney and Jackson, London, 1926, p. 619.
13. B. Pullman, *Bull. Assoc. Franc., Etude Cancer*, **34**, 245 (1947).
14. A. Albert, *The Acridines*, 1st ed., Arnold, London, 1951, p. xi; A. Albert, "The Acridines," in *Heterocyclic Compounds*, Vol. 4, Elderfield, Ed., Wiley, New York, 1952, p. 491.
15. H. Ing, in *Organic Chemistry*, Vol. 3, Chap. V., H. Gilman, Ed., Wiley, New York, 1953.
16. *International Union of Pure and Applied Chemistry, Nomenclature of Organic Chemistry*, Sections A and B, 2nd ed., Butterworths, London, 1966, p. 58.
17. A. Albert, *The Acridines*, 2nd ed., Arnold, London, 1966, p. xi.
18. A. M. Paterson, L. T. Capell, and D. F. Walker, *The Ring Index*, 2nd ed., American Chemical Society, Washington, D.C., 1960.
19. *Chem. Abstr.*, **39**, 5867 (1945).
20. *International Union of Pure and Applied Chemistry, Nomenclature of Organic Chemistry*, Section C, Butterworths, London, 1965.
21. "Editorial Report on Nomenclature," *J. Chem. Soc.*, 3699 (1950).



## CHAPTER I

# Acridines

N. R. RAULINS

*Department of Chemistry,  
University of Wyoming,  
Laramie, Wyoming*

1. Historical Introduction and the Formulation of Acridine . . . . .	10
2. Methods of Preparation of Acridines . . . . .	13
A. Preparation of Acridines from 9-Acridanones or 9-Chloroacridines . . . . .	13
(1) 9-Substituted Acridines . . . . .	17
(2) Zinc Dust Reduction of 9-Acridanones . . . . .	18
(3) Acridines from 9-Chloroacridines with Hydrazides and Hydrazine . . . . .	18
(4) Reduction of 9-Acridanones or 9-Chloroacridines to Acridans, Followed by Oxidation to Acridines . . . . .	19
(a) Reduction Procedures . . . . .	20
(b) Oxidation of Acridans to Acridines . . . . .	21
B. The Bernthsen Reaction and Its Modifications . . . . .	23
C. Preparation of Acridines from <i>m</i> -Phenylenediamines and Formic (or Oxalic) Acid . . . . .	30
(1) 3,6-Diaminoacridines . . . . .	30
(2) 3-Aminoacridines . . . . .	36
D. Acridines and Acridans from Aromatic Amines and Aldehydes and Similar Syntheses . . . . .	38
E. Cyclization of Diphenylamine-2-aldehydes and Ketones . . . . .	41
(1) Aldehyde Cyclizations . . . . .	45
(2) Ketone Cyclizations . . . . .	46
F. Other Methods of Preparing Acridines . . . . .	50
(1) Dehydrogenations Giving Acridines . . . . .	50
(2) Cyclodehydrogenations Giving Acridines . . . . .	51
(3) Acridines from 2-Aminobenzaldehyde and Phenols . . . . .	53
(4) Other Acridine Syntheses . . . . .	54
3. General Properties of Acridine . . . . .	58
4. Chemical Properties of Acridine . . . . .	63