



# Filtering Complex Turbulent Systems

Andrew J. Majda and John Harlim

CAMBRIDGE


# FILTERING COMPLEX TURBULENT SYSTEMS

ANDREW J. MAJDA

*New York University*

JOHN HARLIM

*North Carolina State University*



CAMBRIDGE  
UNIVERSITY PRESS

CAMBRIDGE UNIVERSITY PRESS

Cambridge, New York, Melbourne, Madrid, Cape Town,  
Singapore, São Paulo, Delhi, Mexico City

Cambridge University Press  
The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by Cambridge University Press, New York

[www.cambridge.org](http://www.cambridge.org)

Information on this title: [www.cambridge.org/9781107016668](http://www.cambridge.org/9781107016668)

© Andrew J. Majda and John Harlim 2012

This publication is in copyright. Subject to statutory exception  
and to the provisions of relevant collective licensing agreements,  
no reproduction of any part may take place without the written  
permission of Cambridge University Press.

First published 2012

Printed in the United Kingdom at the University Press, Cambridge

*A catalogue record for this publication is available from the British Library*

ISBN 978-1-107-01666-8 Hardback

Cambridge University Press has no responsibility for the persistence or  
accuracy of URLs for external or third-party internet websites referred to  
in this publication, and does not guarantee that any content on such  
websites is, or will remain, accurate or appropriate.

## FILTERING COMPLEX TURBULENT SYSTEMS

Many natural phenomena ranging from climate through to biology are described by complex dynamical systems. Getting information about these phenomena involves filtering noisy data and making predictions based on incomplete information, and often we need to do this in real time, e.g. for weather forecasting or pollution control. All this is further complicated by the sheer number of parameters involved, leading to further problems associated with the “curse of dimensionality” and the “curse of small ensemble size”.

The authors develop, for the first time in book form, a systematic perspective on all these issues from the standpoint of applied mathematics. Their approach follows several strands:

- blending classical stability analysis of partial differential equations and their finite difference approximations;
- extending classical Kalman filters and applying them to stochastic models of turbulence to deal with large model errors;
- developing test suites of statistically exactly solvable models and new SPEKF algorithms for filtering slow-fast systems, moist convection, turbulent tracers, and geophysical turbulent systems.

The book contains enough background material from filtering, turbulence theory, and numerical analysis to make the presentation self-contained, and is suitable for graduate courses as well as for researchers in a range of disciplines across science and engineering where applied mathematics is required to enlighten observations and models.

ANDREW J. MAJDA is the Morse Professor of Arts and Sciences at the Courant Institute of New York University.

JOHN HARLIM is an Assistant Professor in the Department of Mathematics at North Carolina State University.

# Preface

This book is an outgrowth of lectures by both authors in the graduate course of the first author at the Courant Institute during spring 2008 and 2010 on the topic of filtering turbulent dynamical systems as well as lectures by the second author at the North Carolina State University in a graduate course in fall 2009. The material is based on the authors' joint research as well as collaborations with Marcus Grote and Boris Gershgorin; the authors thank these colleagues for their explicit and implicit contributions to this material. Chapter 1 presents a detailed overview and summary of the viewpoint and material in the book. This book is designed for applied mathematicians, scientists and engineers, ranging from first- and second-year graduate students to senior researchers interested in filtering large-dimensional complex nonlinear systems.

The first author acknowledges the generous support of DARPA through Ben Mann and ONR through Reza Malek-Madani which funded the research on these topics and helped make this book a reality.

# Contents

*Preface*

*page ix*

<b>1</b>	<b>Introduction and overview: Mathematical strategies for filtering turbulent systems</b>	1
1.1	Turbulent dynamical systems and basic filtering	3
1.2	Mathematical guidelines for filtering turbulent dynamical systems	9
1.3	Filtering turbulent dynamical systems	11
<b>Part I</b>	<b>Fundamentals</b>	13
<b>2</b>	<b>Filtering a stochastic complex scalar: The prototype test problem</b>	15
2.1	Kalman filter: One-dimensional complex variable	15
2.2	Filtering stability	20
2.3	Model error	23
<b>3</b>	<b>The Kalman filter for vector systems: Reduced filters and a three-dimensional toy model</b>	30
3.1	The classical $N$ -dimensional Kalman filter	30
3.2	Filter stability	32
3.3	Example: A three-dimensional toy model with a single observation	33
3.4	Reduced filters for large systems	39
3.5	A priori covariance stability for the unstable mode filter given strong observability	43
<b>4</b>	<b>Continuous and discrete Fourier series and numerical discretization</b>	47
4.1	Continuous and discrete Fourier series	47
4.2	Aliasing	49
4.3	Differential and difference operators	52
4.4	Solving initial value problems	53
4.5	Convergence of the difference operator	55

<b>Part II</b>	<b>Mathematical guidelines for filtering turbulent signals</b>	59
<b>5</b>	<b>Stochastic models for turbulence</b>	61
5.1	The stochastic test model for turbulent signals	61
5.2	Turbulent signals for the damped forced advection–diffusion equation	65
5.3	Statistics of turbulent solutions in physical space	66
5.4	Turbulent Rossby waves	68
	Appendix A: Temporal correlation function for each Fourier mode	70
	Appendix B: Spatio-temporal correlation function	71
<b>6</b>	<b>Filtering turbulent signals: Plentiful observations</b>	72
6.1	A mathematical theory for Fourier filter reduction	73
6.2	Theoretical guidelines for filter performance under mesh refinement for turbulent signals	77
6.3	Discrete filtering for the stochastically forced dissipative advection equation	81
<b>7</b>	<b>Filtering turbulent signals: Regularly spaced sparse observations</b>	94
7.1	Theory for filtering sparse regularly spaced observations	94
7.2	Fourier domain filtering for sparse regular observations	99
7.3	Approximate filters in the Fourier domain	102
7.4	New phenomena and filter performance for sparse regular observations	107
<b>8</b>	<b>Filtering linear stochastic PDE models with instability and model error</b>	116
8.1	Two-state continuous-time Markov process	117
8.2	Idealized spatially extended turbulent systems with instability	119
8.3	The mean stochastic model for filtering	123
8.4	Numerical performance of the filters with and without model error	127
<b>Part III</b>	<b>Filtering turbulent nonlinear dynamical systems</b>	131
<b>9</b>	<b>Strategies for filtering nonlinear systems</b>	133
9.1	The extended Kalman filter	134
9.2	The ensemble Kalman filter	136
9.3	The ensemble square-root filters	139
9.4	Ensemble filters on the Lorenz-63 model	143
9.5	Ensemble square-root filters on stochastically forced linear systems	149
9.6	Advantages and disadvantages with finite ensemble strategies	151
<b>10</b>	<b>Filtering prototype nonlinear slow–fast systems</b>	153
10.1	The nonlinear test model for filtering slow–fast systems with strong fast forcing: An overview	153
10.2	Exact solutions and exactly solvable statistics in the nonlinear test model	159
10.3	Nonlinear extended Kalman filter (NEKF)	171
10.4	Experimental designs	174

10.5	Filter performance	177
10.6	Summary	190
<b>11</b>	<b>Filtering turbulent nonlinear dynamical systems by finite ensemble methods</b>	192
11.1	The L-96 model	192
11.2	Ensemble square-root filters on the L-96 model	195
11.3	Catastrophic filter divergence	200
11.4	The two-layer quasi-geostrophic model	204
11.5	Local least-square EAKF on the QG model	210
<b>12</b>	<b>Filtering turbulent nonlinear dynamical systems by linear stochastic models</b>	214
12.1	Linear stochastic models for the L-96 model	215
12.2	Filter performance with plentiful observation	220
12.3	Filter performance with regularly spaced sparse observations	223
<b>13</b>	<b>Stochastic parametrized extended Kalman filter for filtering turbulent signals with model error</b>	236
13.1	Nonlinear filtering with additive and multiplicative biases: One-mode prototype test model	238
13.2	Filtering spatially extended turbulent systems with SPEKF	251
13.3	Application of SPEKF to the two-layer QG model	263
	Appendix	269
<b>14</b>	<b>Filtering turbulent tracers from partial observations: An exactly solvable test model</b>	276
14.1	Model description	278
14.2	System statistics	279
14.3	Nonlinear extended Kalman filter	292
14.4	Filter performance	297
<b>15</b>	<b>The search for efficient skillful particle filters for high-dimensional turbulent dynamical systems</b>	316
15.1	The basic idea of a particle filter	317
15.2	Innovative particle filter algorithms	319
15.3	Filter performance on the L-63 model	326
15.4	Filter performance on the L-96 model	339
15.5	Discussion	346
	<i>References</i>	349
	<i>Index</i>	356



# 1

## Introduction and overview: Mathematical strategies for filtering turbulent systems

Filtering is the process of obtaining the best statistical estimate of a natural system from partial observations of the true signal from nature. In many contemporary applications in science and engineering, real-time filtering of a turbulent signal from nature involving many degrees of freedom is needed to make accurate predictions of the future state. This is obviously a problem with significant practical impact. Important contemporary examples involve the real-time filtering and prediction of weather and climate as well as the spread of hazardous plumes or pollutants. Thus, an important emerging scientific issue is the real-time filtering through observations of noisy signals for turbulent nonlinear dynamical systems as well as the statistical accuracy of spatio-temporal discretizations for filtering such systems. From the practical standpoint, the demand for operationally practical filtering methods escalates as the model resolution is significantly increased. In the coupled atmosphere–ocean system, the current practical models for prediction of both weather and climate involve general circulation models where the physical equations for these extremely complex flows are discretized in space and time and the effects of unresolved processes are parametrized according to various recipes; the result of this process involves a model for the prediction of weather and climate from partial observations of an extremely unstable, chaotic dynamical system with several billion degrees of freedom. These problems typically have many spatio-temporal scales, rough turbulent energy spectra in the solutions near the mesh scale, and a very large-dimensional state space, yet real-time predictions are needed.

Particle filtering of low-dimensional dynamical systems is an established discipline (Bain and Crisan, 2009). When the system is low dimensional or when it has a low-dimensional attractor, Monte Carlo approaches such as the particle filter (Chorin and Krause, 2004) with its various up-to-date resampling strategies (Del Moral, 1996; Del Moral and Jacod, 2001; Rossi and Vila, 2006) provide better estimates in the presence of strong nonlinearity and highly non-Gaussian distributions. However, with the above practical computational constraint in mind, these accurate nonlinear particle filtering strategies are not feasible since sampling a high-dimensional variable is computationally impossible for the foreseeable future. Recent mathematical theory strongly supports this curse of dimensionality for particle filters (Bengtsson *et al.*, 2008; Bickel *et al.*, 2008). Nevertheless

some progress in developing particle filtering with small ensemble size for non-Gaussian turbulent dynamical systems is discussed in Chapter 15. These approaches, including the new maximum entropy particle filter (MEPF) due to the authors, all make judicious use of partial marginal distributions to avoid particle collapse. In the second direction, Bayesian hierarchical modeling (Berliner *et al.*, 2003) and reduced-order filtering strategies (Miller *et al.*, 1999; Ghil and Malanotte-Rizolli, 1991; Todling and Ghil, 1994; Anderson, 2001, 2003; Chorin and Krause, 2004; Farrell and Ioannou, 2001, 2005; Ott *et al.*, 2004; Hunt *et al.*, 2007; Harlim and Hunt, 2007b) based on the Kalman filter (Anderson and Moore, 1979; Chui and Chen, 1999; Kaipio and Somersalo, 2005) have been developed with some success in these extremely complex high-dimensional nonlinear systems. There is an inherently difficult practical issue of small ensemble size in filtering statistical solutions of these complex problems due to the large computational overload in generating individual ensemble members through the forward dynamical operator (Haven *et al.*, 2005). Numerous ensemble-based Kalman filters (Evensen, 2003; Bishop *et al.*, 2001; Anderson, 2001; Szunyogh *et al.*, 2005; Hunt *et al.*, 2007) show promising results in addressing this issue for synoptic-scale mid-latitude weather dynamics by imposing suitable spatial localization on the covariance updates; however, all these methods are very sensitive to model resolution, observation frequency and the nature of the turbulent signals when a practical limited ensemble size (typically less than 100) is used. They are also less skillful for more complex phenomena like gravity waves coupled with condensational heating from clouds which are important for the tropics and severe local weather.

Here is a list of fundamental new difficulties in the real-time filtering of turbulent signals that need to be addressed as mentioned briefly above.

- 1(a) **Turbulent dynamical systems to generate the true signal.** The true signal from nature arises from a turbulent nonlinear dynamical system with extremely complex noisy spatio-temporal signals which have significant amplitude over many spatial scales.
- 1(b) **Model errors.** A major difficulty in accurate filtering of noisy turbulent signals with many degrees of freedom is model error; the fact that the true signal from nature is processed for filtering and prediction through an imperfect model where by practical necessity, important physical processes are parametrized due to inadequate numerical resolution or incomplete physical understanding. The model errors of inadequate resolution often lead to rough turbulent energy spectra for the truth signal to be filtered on the order of the mesh scale for the dynamical system model used for filtering.
- 1(c) **Curse of ensemble size.** For forward models for filtering, the state space dimension is typically large, of order  $10^4$ – $10^8$ , for these turbulent dynamical systems, so generating an ensemble size with such a direct approach of order 50–100 members is typically all that is available for real-time filtering.
- 1(d) **Sparse, noisy, spatio-temporal observations for only a partial subset of the variables.** In systems with multiple spatio-temporal scales, the sparse observations of the truth signal might automatically couple many spatial scales, as shown below in Chapter 7 or in Harlim and Majda (2008b), while the observation of a partial

subset of variables might mix together temporal slow and fast components of the system (Gershgorin and Majda, 2008, 2010) as discussed in Chapter 10. For example, observations of pressure or temperature in the atmosphere mix slow vortical and fast gravity wave processes.

This book is an introduction to filtering with an emphasis on the central new issues in 1(a)–(d) for filtering turbulent dynamical systems through the “modus operandi” of the modern applied mathematics paradigm (Majda, 2000a) where rigorous mathematical theory, asymptotic and qualitative models, and novel numerical algorithms are all blended together interactively to give insight into central “cutting edge” practical science problems. In the last several years, the authors have utilized the synergy of modern applied mathematics to address the following:

- 2(a) How to develop simple off-line mathematical test criteria as guidelines for filtering extremely stiff multiple space–time scale problems that often arise in filtering turbulent signals through plentiful and sparse observations? (Majda and Grote, 2007; Castronovo *et al.*, 2008; Grote and Majda, 2006; Harlim and Majda, 2008b)
- 2(b) For turbulent signals from nature with many scales, even with mesh refinement, the model has inaccuracies from parametrization, under-resolution, etc. Can judicious model errors help filtering and simultaneously overcome the curse of dimensionality? (Castronovo *et al.*, 2008; Harlim and Majda, 2008a,b, 2010a)
- 2(c) Can new computational strategies based on stochastic parametrization algorithms be developed to overcome the curse of dimensionality, to reduce model error and improve the filtering as well as the prediction skill? (Gershgorin *et al.*, 2010a,b; Harlim and Majda, 2010b)
- 2(d) Can exactly solvable models be developed to elucidate the central issues in 1(d) for turbulent signals, to develop unambiguous insight into model errors and to lead to efficient new computational algorithms? (Gershgorin and Majda, 2008, 2010)

The main goals of this book are the following: first, to introduce the reader to filtering from this viewpoint in an elementary fashion where no prior background on these topics is assumed (Chapters 2–4); secondly, to describe in detail the recent and ongoing developments, emphasizing the remarkable new mathematical and physical phenomena that emerge from the modern applied mathematics modus operandi applied to filtering turbulent dynamical systems. Next, in this introductory chapter, we provide an overview of turbulent dynamical systems and basic filtering followed by an overview of the basic applied mathematics motivation which leads to the new developments and viewpoint emphasized in this book.

## 1.1 Turbulent dynamical systems and basic filtering

The large-dimensional turbulent dynamical systems which define the true signal from nature to be filtered in the class of problems studied here have a fundamentally different statistical character than in more familiar low-dimensional chaotic dynamical systems. The

most well-known low-dimensional chaotic dynamical system is Lorenz's famous three-equation model (Lorenz, 1963) which is weakly mixing with one unstable direction on an attractor with high symmetry. In contrast, realistic turbulent dynamical systems have a large phase space dimension, a large dimensional unstable manifold on the attractor, and are strongly mixing with exponential decay of correlations. The simplest prototype example of a turbulent dynamical system is also due to Lorenz and is called the L-96 model (Lorenz, 1996; Lorenz and Emanuel, 1998). It is widely used as a test model for algorithms for prediction, filtering and low-frequency climate response (Majda *et al.*, 2005; Majda and Wang, 2006). The L-96 model is a discrete periodic model given by the following system

$$\frac{du_j}{dt} = (u_{j+1} - u_{j-2})u_{j-1} - u_j + F, \quad j = 0, \dots, J - 1, \quad (1.1)$$

with  $J = 40$  and with  $F$  the forcing parameter. The model is designed to mimic baroclinic turbulence in the mid-latitude atmosphere with the effects of energy-conserving nonlinear advection and dissipation represented by the first two terms in (1.1). For sufficiently strong forcing values such as  $F = 6, 8, 16$ , the L-96 model is a prototype turbulent dynamical system which exhibits features of weakly chaotic turbulence ( $F = 6$ ), strongly chaotic turbulence ( $F = 8$ ), and strong turbulence ( $F = 16$ ) (Majda *et al.*, 2005). In order to quantify and compare the different types of turbulent chaotic dynamics in the L-96 model as  $F$  is varied, it is convenient to rescale the system to have unit energy for statistical fluctuations around the constant mean statistical state,  $\bar{u}$  (Majda *et al.*, 2005); thus, the transformation  $u_j = \bar{u} + E_p^{1/2} \tilde{u}_j$ ,  $t = \tilde{t} E_p^{-1/2}$  is utilized where  $E_p$  represents the energy fluctuations (Majda *et al.*, 2005). After this normalization, the mean state becomes zero and the energy fluctuations are unity for all values of  $F$ . The dynamical equation in terms of the new variables,  $\tilde{u}_j$ , becomes

$$\frac{d\tilde{u}_j}{d\tilde{t}} = (\tilde{u}_{j+1} - \tilde{u}_{j-2})\tilde{u}_{j-1} + E_p^{-1/2}((\tilde{u}_{j+1} - \tilde{u}_{j-2})\bar{u} - \tilde{u}_j) + E_p^{-1}(F - \bar{u}). \quad (1.2)$$

Table 1.1 lists, in the non-dimensional coordinates, the leading Lyapunov exponent,  $\lambda_1$ , the dimension of the unstable manifold,  $N^+$ , the sum of the positive Lyapunov exponents (the KS entropy) and the correlation time,  $T_{\text{corr}}$ , of any  $\tilde{u}_j$  variable with itself as  $F$  is varied

Table 1.1 Dynamical properties of the L-96 model for regimes with  $F = 6, 8, 16$ .  $\lambda_1$  denotes the largest Lyapunov exponent,  $N^+$  denotes the dimension of the expanding subspace of the attractor,  $KS$  denotes the Kolmogorov–Sinai entropy and  $T_{\text{corr}}$  denotes the decorrelation time of the energy-rescaled time correlation function.

	$F$	$\lambda_1$	$N^+$	KS	$T_{\text{corr}}$
Weakly chaotic	6	1.02	12	5.547	8.23
Strongly chaotic	8	1.74	13	10.94	6.704
Fully turbulent	16	3.945	16	27.94	5.594




Figure 1.1 Space-time diagrams of numerical solutions of the L-96 model for the weakly chaotic ( $F = 6$ ), strongly chaotic ( $F = 8$ ) and fully turbulent ( $F = 16$ ) regimes.

through  $F = 6, 8, 16$ . Note that  $\lambda_1$ ,  $N^+$  and KS increase significantly as  $F$  increases while  $T_{\text{corr}}$  decreases in these non-dimensional units; furthermore, the weakly turbulent case with  $F = 6$  already has a 12-dimensional unstable manifold in the 40-dimensional phase space. Snapshots of the time series for (1.1) with  $F = 6, 8, 16$ , as depicted in Fig. 1.1, qualitatively confirm the above quantitative intuition with weakly turbulent patterns for  $F = 6$ , strongly chaotic wave turbulence for  $F = 8$ , and fully developed wave turbulence for  $F = 16$ . It is worth remarking here that smaller values of  $F$  around  $F = 4$  exhibit the more familiar low-dimensional weakly chaotic behavior associated with the transition to turbulence.

In regimes to realistically mimic properties of nature, virtually all atmosphere, ocean and climate models with sufficiently high resolution are turbulent dynamical systems with features as described above. The simplest paradigm model of this type is the two-layer quasi-geostrophic (QG) model in doubly periodic geometry that is externally forced by a mean vertical shear (Smith *et al.*, 2002), which has baroclinic instability (Salmon, 1998); the properties of the turbulent cascade have been extensively discussed in this setting, e.g. see Salmon (1998) and citations in Smith *et al.* (2002). The governing equations for the two-layer QG model with a flat bottom, rigid lid and equal-depth layers  $H$  can be written as

$$\begin{aligned} \frac{\partial q_1}{\partial t} + J(\psi_1, q_1) + U \frac{\partial q_1}{\partial x} + (\beta + k_d^2 U) \frac{\partial \psi_1}{\partial x} + \nu \nabla^8 q_1 &= 0, \\ \frac{\partial q_2}{\partial t} + J(\psi_2, q_2) - U \frac{\partial q_2}{\partial x} + (\beta - k_d^2 U) \frac{\partial \psi_2}{\partial x} + \kappa \nabla^2 \psi_2 + \nu \nabla^8 q_2 &= 0, \end{aligned} \quad (1.3)$$

where subscript 1 denotes the top layer and 2 the bottom layer;  $\psi$  is the perturbed stream function;  $J(\psi, q) = \psi_x q_y - \psi_y q_x$  is the Jacobian term representing nonlinear advection;  $U$  is the zonal mean shear;  $\beta$  is the meridional gradient of the Coriolis parameter;  $q$  is the perturbed quasi-geostrophic potential vorticity, defined as follows

$$q_i = \beta y + \nabla^2 \psi_i + \frac{k_d^2}{2} (\psi_{3-i} - \psi_i), \quad i = 1, 2, \quad (1.4)$$

where  $k_d = \sqrt{8}/L_d$  is the wavenumber corresponding to the Rossby radius  $L_d$ ;  $\kappa$  is the Ekman bottom drag coefficient; and  $\nu$  is the hyperviscosity coefficient. Note that Eqns (1.3) are the prognostic equations for perturbations around a uniform shear with stream function  $\Psi_1 = -Uy$ ,  $\Psi_2 = Uy$  as the background state, and the hyperviscosity term,  $\nu \nabla^8 q$ , is added to filter out the energy buildup on the smaller scales.

This is the simplest climate model for the poleward transport of heat in the atmosphere or ocean and with a modest resolution of  $128 \times 128 \times 2$  grid points has a phase space of more than 30,000 variables. Again for modeling the atmosphere and ocean, this model in the appropriate parameter regimes is a strongly turbulent dynamical system with strong cascades of energy (Salmon, 1998; Smith *et al.*, 2002; Kleeman and Majda, 2005); it has been utilized recently as a test model for algorithms for filtering sparsely observed turbulent signals in the atmosphere and ocean (Harlim and Majda, 2010b).

### 1.1.1 Basic filtering

We assume that observations are made at uniform discrete times,  $m\Delta t$ , with  $m = 1, 2, 3, \dots$ . For example, in global weather prediction models, the observations are given as inputs in the model every six hours and for large-dimensional turbulent dynamical systems, it is a challenge to implement continuous observations, practically. As depicted in Fig. 1.2, filtering is a two-step process involving statistical prediction of a probability distribution for the state variable  $u$  through a forward operator on the time interval between observations followed by an analysis step at the next observation time which corrects this probability distribution on the basis of the statistical input of noisy observations of the system. In the present applications, the forward operator is a large-dimensional dynamical system perhaps with noise written in the Itô sense as

$$\frac{du}{dt} = F(u, t) + \sigma(u, t)\dot{W}(t) \quad (1.5)$$

for  $u \in \mathbb{R}^N$ , where  $\sigma$  is an  $N \times K$  noise matrix and  $\dot{W} \in \mathbb{R}^K$  is  $K$ -dimensional white noise. The Fokker–Planck equation for the probability density,  $p(u, t)$ , associated with (1.5) is




Figure 1.2 Filtering: Two-step predictor–corrector method.

$$p_t = -\nabla_u \cdot (F(u, t)p) + \frac{1}{2} \nabla_u \cdot \nabla_u (Qp) \equiv L_{\text{FP}} p \quad (1.6)$$

$$p_t|_{t=t_0} = p_0(u)$$

with  $Q(t) = \sigma\sigma^T$ . For simplicity in exposition, here and throughout the remainder of the book we assume  $M$  linear observations,  $\vec{v}_m \in \mathbb{R}^M$ , of the true signal from nature given by

$$\vec{v}_m = Gu(m\Delta t) + \vec{\sigma}_m^o, \quad m = 1, 2, \dots \quad (1.7)$$

where  $G$  maps  $\mathbb{R}^N$  into  $\mathbb{R}^M$  while the observational noise,  $\vec{\sigma}_m^o \in \mathbb{R}^M$ , is assumed to be a zero-mean Gaussian random variable with  $M \times M$  covariance matrix,

$$R^o = \langle \vec{\sigma}_m^o \otimes (\vec{\sigma}_m^o)^T \rangle. \quad (1.8)$$

Gaussian random variables are uniquely determined by their mean and covariance; here and below, we utilize the standard notation  $\mathcal{N}(\vec{X}, R)$  to denote a vector Gaussian random variable with mean  $\vec{X}$  and covariance matrix  $R$ . With these preliminaries, we describe the two-step filtering algorithm with the dynamics in (1.5), (1.6) and the noisy observations in (1.7), (1.8). Start at time step  $m\Delta t$  with a posterior probability distribution,  $p_{m,+}(u)$ , which takes into account the observations in (1.7) at time  $m\Delta t$ . Calculate a prediction or forecast probability distribution,  $p_{m+1,-}(u)$ , by using (1.6), in other words, let  $p$  be the solution of the Fokker–Planck equation,

$$p_t = L_{\text{FP}} p, \quad m\Delta t < t \leq (m+1)\Delta t \quad (1.9)$$

$$p|_{t=m\Delta t} = p_{m,+}(u).$$

Define  $p_{m+1,-}(u)$ , the prior probability distribution before taking observations at time  $m+1$  into account, by

$$p_{m+1,-}(u) \equiv p(u, (m+1)\Delta t) \quad (1.10)$$

with  $p$  determined by the forward dynamics in (1.9). Next, the analysis step at time  $(m+1)\Delta t$  which corrects this forecast and takes the observations into account is implemented by using Bayes' theorem

$$p_{m+1,+}(u)p(v_{m+1}) = p_{m+1}(u|v_{m+1})p(v_{m+1})$$

$$= p_{m+1}(u, v) = p_{m+1}(v_{m+1}|u)p_{m+1,-}(u). \quad (1.11)$$

With Bayes' formula in (1.11), we calculate the posterior distribution

$$p_{m+1,+}(u) \equiv p_{m+1}(u|v_{m+1}) = \frac{p_{m+1}(v_{m+1}|u)p_{m+1,-}(u)}{\int p_{m+1}(v_{m+1}|u)p_{m+1,-}(u)du}. \quad (1.12)$$

The two steps described in (1.9), (1.10), (1.12) define the basic nonlinear filtering algorithm which forms the theoretical basis for practical design of algorithms for filtering turbulent dynamical systems (Jazwinski, 1970; Bain and Crisan, 2009). While this is conceptually clear, practical implementation of (1.9), (1.10), (1.12), directly in turbulent dynamical systems, is impossible due to large state space,  $N \gg 1$ , as well as the fundamental difficulties elucidated in 1(a)–(d) in the introduction.

The most important and famous example of filtering is the Kalman filter where the analysis step in (1.5) is associated with linear dynamics which can be integrated between observation time steps  $m\Delta t$  and  $(m+1)\Delta t$  to yield the forward operator

$$u_{m+1} = Fu_m + \bar{f}_{m+1} + \sigma_{m+1}. \quad (1.13)$$

Here  $F$  is the  $N \times N$  system operator matrix and  $\sigma_m$  is the system noise assumed to be zero-mean and Gaussian with  $N \times N$  covariance matrix

$$R = \langle \sigma_m \otimes \sigma_m^T \rangle, \forall m, \quad (1.14)$$

while  $\bar{f}_m$  is a deterministic forcing. Next, we present the simplified Kalman filter equations for the linear case. First assume the initial probability density  $p_0(u)$  is Gaussian, i.e.  $p_0(u) = \mathcal{N}(\bar{u}_0, R_o)$  and assume by recursion that the posterior probability distribution,  $p_{m,+}(u) = \mathcal{N}(\bar{u}_{m,+}, R_{m,+})$ , is also Gaussian. By using the linear dynamics in (1.13), the forecast or prediction distribution at time  $(m+1)\Delta t$  is also Gaussian,

$$\begin{aligned} p_{m+1,-}(u) &= \mathcal{N}(\bar{u}_{m+1,-}, R_{m+1,-}) \\ \bar{u}_{m+1,-} &= F\bar{u}_{m,+} + \bar{f}_{m+1} \\ R_{m+1,-} &= FR_{m,+}F^T + R. \end{aligned} \quad (1.15)$$

With the assumptions in (1.7), (1.8) and (1.13), (1.15), the analysis step in (1.12) becomes an explicit regression procedure for Gaussian random variables (Chui and Chen, 1999; Anderson and Moore, 1979) so that the posterior distribution,  $p_{m+1,+}(u)$ , is also Gaussian yielding the **Kalman filter**

$$\begin{aligned} p_{m+1,+}(u) &= \mathcal{N}(\bar{u}_{m+1,+}, R_{m+1,+}) \\ \bar{u}_{m+1,+} &= (\mathcal{I} - K_{m+1}G)\bar{u}_{m+1,-} + K_{m+1}v_{m+1} \\ R_{m+1,+} &= (\mathcal{I} - K_{m+1}G)R_{m+1,-} \\ K_{m+1} &= R_{m+1,-}G^T(GR_{m+1,-}G^T + R^o)^{-1}. \end{aligned} \quad (1.16)$$

The  $N \times M$  matrix,  $K_{m+1}$ , is the Kalman gain matrix. Note that the posterior mean after processing the observations is a weighted sum of the forecast and analysis contributions through the Kalman gain matrix and also that the observations reduce the covariance,  $R_{m+1,+} \leq R_{m+1,-}$ . In this Gaussian case with linear observations, the analysis step going from (1.15) to (1.16) is a standard linear least-squares regression. An excellent treatment of this can be found in chapter 3 of Kaipio and Somersalo (2005). There is a huge literature on Kalman filtering; two excellent basic texts are Chui and Chen (1999) and Anderson and Moore (1979) where more details and references can be found. Our intention in the introductory parts in this book in Chapters 2 and 3 is not to repeat the well-known material in (1.15), (1.16) in detail; instead we introduce this elementary material in a fashion to set the stage for the mathematical guidelines developed in Part II (Chapters 5–8) and the applications to filtering turbulent nonlinear dynamical systems presented in Part III (Chapters 9–15).



Naively, the reader might expect that everything is known about filtering linear systems; however, when the linear system is high dimensional, i.e.  $N \gg 1$ , the same issues elucidated in 1(a)–(d) occur for linear systems in a more transparent fashion. This is the viewpoint emphasized and developed in Part II of the book (Chapters 5–8) which is motivated next. For linear systems without model errors, the recursive Kalman filter is an optimal estimator but the recursive nonlinear filter in (1.7)–(1.12) may not be an optimal estimator for the nonlinear stochastic dynamical system without model error in (1.5).

## 1.2 Mathematical guidelines for filtering turbulent dynamical systems

How can useful mathematical guidelines be developed in order to elucidate and ameliorate the central new issues in 1(a)–(d) from the introduction for turbulent dynamical systems? This is the topic of this section. Of course, to be useful, such mathematical guidelines have to be general yet still involve simplified models with analytical tractability. Such criteria have been developed recently by Majda and Grote (2007); Castronovo *et al.* (2008) and Harlim and Majda (2008b) through the modern applied mathematics paradigm and the goal here is to outline this development and discuss some of the remarkable phenomena which occur. The starting point for this development for filtering turbulent dynamical systems involves the symbiotic interaction of three different disciplines in applied mathematics/physics, as depicted in Fig. 1.3: stochastic modeling of turbulent signals, numerical analysis of PDEs and classical filtering theory outlined in (1.13)–(1.16) of Section 1.1. Here is the motivation from the three legs of the triangle.

First, the simplest stochastic models for modeling turbulent fluctuations consist of replacing the nonlinear interaction at these modes by additional dissipation and white noise




Figure 1.3 Modern applied mathematics paradigm for filtering.