Ovidiu Cârjă Ioan I Vrabie editors

APPLIED ANALYSIS
AND DIF AL
EQUATIONS

editors Ovidiu Cârjă Ioan I Vrabie

Al I Cuza University, România & O Mayer Mathematics Institute of the Romanian Academy, România

APPLIED ANALYSIS AND DIFFERENTIAL EQUATIONS

las' namber 2006





Published by

World Scientific Publishing Co. Pte. Ltd.

5 Toh Tuck Link, Singapore 596224

USA office: 27 Warren Street, Suite 401-402, Hackensack, NJ 07601 UK office: 57 Shelton Street, Covent Garden, London WC2H 9HE

British Library Cataloguing-in-Publication Data

A catalogue record for this book is available from the British Library.

APPLIED ANALYSIS AND DIFFERENTIAL EQUATIONS

Copyright © 2007 by World Scientific Publishing Co. Pte. Ltd.

All rights reserved. This book, or parts thereof, may not be reproduced in any form or by any means, electronic or mechanical, including photocopying, recording or any information storage and retrieval system now known or to be invented, without written permission from the Publisher.

For photocopying of material in this volume, please pay a copying fee through the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, USA. In this case permission to photocopy is not required from the publisher.

ISBN-13 978-981-270-594-5 ISBN-10 981-270-594-5

PREFACE

The International Conference of Applied Analysis and Differential Equations was held September 4-9, 2006 at the "Al. I. Cuza" University of Iaşi and the "Octav Mayer" Mathematical Institute of the Romanian Academy of Iaşi, Romania.

It was supported by the Project CERES 4-194/2004.

The conference was devoted to recent advances in *Nonsmooth Analysis* and *Optimizations, ODE, PDE, Control Theory, Stochastic Analysis* and was well attended by mathematicians all over the world. There were 40 minutes plenary invited lectures, 20 minutes talks and a poster session.

This volume includes 29 selected articles, many of them written by leading specialists in their fields and covering the main topics of the conference.

We take this opportunity to express our gratitude to all who have chosen to publish their contributions in this volume as well as to the referees of the submitted papers for their promptitude and high exigence. Furthermore, we thank Ms. Carmen Savin for her excellent secretarial work, as well as Ms. Elena Mocanu for her extremely efficient and skilful assistance in preparation of the camera-ready copy.

Finally, we express our warmest thanks to World Scientific for the very pleasant and fruitful cooperation during the publication of this volume.

Iași, December 19, 2006

Ovidiu Cârjă and Ioan I. Vrabie

ORGANIZING COMMITTEE SERIES EDITORIAL BOARD

ORGANIZING COMMITTEE

of the International Conference on Applied Analysis and Differential Equations

O. Cârjă (Chairman)	-	University "Al.I.Cuza", Iaşi, RO
V. Barbu		University "Al.I.Cuza", Iaşi, RO
M. Durea (Secretary)		University "Al.I.Cuza", Iaşi, RO
M. Necula		University "Al.I.Cuza", Iaşi, RO
A. Răşcanu		University "Al.I.Cuza", Iaşi, RO
I.I. Vrabie		University "Al.I.Cuza", Iaşi, RO
C. Zălinescu		University "Al.I.Cuza", Iaşi, RO

CONTENTS

Preface	7
Organizing Committees	vi
A Stability Criterion for Delay Differential Equations with Impulse Effects $ J.O.\ Alzabut$	1
On the Use of Korn's Type Inequalities in the Existence Theory for Cosserat Elastic Surfaces with Voids $M.\ B\hat{\imath}rsan$	11
Time Periodic Viscosity Solutions of Hamilton–Jacobi Equations M. Bostan and G. Namah	21
A Viability Result for Semilinear Reaction-diffusion Systems M. Burlică and D. Roşu	31
Necessary Optimality Conditions for Hyperbolic Discrete Inclusions $A.\ Cernea$	45
Controllability of Damped Second-Order Initial Value Problem for a Class of Differential Inclusions with Nonlocal Conditions on Noncompact Intervals D.N. Chalishajar	55
Averaging of Evolution Inclusions in Banach Spaces $T.\ Donchev$	69
Influence of Variable Permeability on Vortex Instability of a Horizontal Combined Free and Mixed Convection Flow in a Saturated Porous Medium A.M. Elaiw, F.S. Ibrahim and A.A. Bakr	79

Analyticity of Stable Invariant Manifolds for Ginzburg-Landau Equation A.V. Fursikov	93
Approximate Solutions for Non-linear Autonomous ODEs on the Basis of PWL Approximation Theory A.G. Garcia, S.I. Biagiola, J.L. Figueroa, O.E. Agamennoni and L.S. Castro	113
On the Influence of a Subquadratic Convection Term in Singular Elliptic Problems M. Ghergu and V. Rădulescu	127
Existence and Uniqueness Results in the Micropolar Mixture Theory of Porous Media ID. Ghiba	139
Approximate Controllability for Linear Stochastic Differential Equations with Control Acting on the Noise D. Goreac	153
Numerical Solutions of Two-point Boundary Value Problems for Ordinary Differential Equations Using Particular Newton Interpolating Series G. Groza and N. Pop	165
On the Integral and Asymptotic Representation of Singular Solutions of Elliptic Boundary Value Problems Near Components of Small Dimensions of Boundary N. Jitaraşu	177
An Existence Result for a Class of Nonlinear Differential Systems R. Luca-Tudorache	185
Well-posedness for a Non-autonomous Model of Fast Diffusion G. Marinoschi	199

C	fontents	xi
Backward Stochastic Generalized Variational Inequality L. Maticiuc and A. Răşcanu	2	217
Trichotomy for Linear Skew-product Semiflows M. Megan, C. Stoica and L. Buliga	2	227
Feedback Control of Constrained Parabolic Systems in Uncertainty Conditions via Asymmetric Games B.S. Mordukhovich and T.I. Seidman	2	237
Implicit Parallel Solvers in Computational Electrocardiology M. Munteanu and L.F. Pavarino	2	255
Fan's Inequality in the Context of M_P -Convexity $C.P.$ Niculescu and I. Rovenţa	2	267
Range Condition in Optimization and Optimal Control $N.H.\ Pavel$	2	275
Limits of Solutions to the Initial Boundary Dirichlet Problem for Semilinear Hyperbolic Equation with Small Parameter A. Perjan	2	281
Well-posedness of the Fixed Point Problem for Multivalued Operators A. Petruşel and I.A. Rus	2	295
Some Remarks on a Nonlinear Semigroup Acting on Positive Measures E. Popa	3	307
Integral Inclusions in Banach Spaces Using Henstock-Type Integrals	3	319
B. Satco		
Good Intermediate-rank Lattice Rules Based on the Weighted Star Discrepancy V. Sinescu and S. Joe	3	329

xii Contents

On the Minimum Energy Problem for Linear Systems in	
Hilbert Spaces	343
$A.I.\ Vieru$	
Author Index	351

A STABILITY CRITERION FOR DELAY DIFFERENTIAL EQUATIONS WITH IMPULSE EFFECTS

J.O. ALZABUT

Department of Mathematics and Computer Science Çankaya University, 06530 Ankara, Turkey E-mail: jehad@cankaya.edu.tr

In this paper, we prove that if a delay differential equation with impulse effects of the form

$$x'(t) = A(t)x(t) + B(t)x(t-\tau)$$
, $t \neq \theta_i$,

$$\Delta x(\theta_i) = C_i x(\theta_i) + D_i x(\theta_{i-j}), \quad i \in \mathbb{N},$$

verifies a Perron condition then its trivial solution is uniformly asymptotically stable.

Keywords: Impulse; Delay; Adjoint; Perron; Uniform asymptotic stability.

1. Introduction and preliminaries

Delay differential equations with impulse effects can suitably model various evolutionary processes that exhibit both delay and impulse characteristics. In particular, they provide a natural description of the motion of several real world processes which, on one hand, depends on the processes history that often turns out to be the cause of phenomena substantially affecting the motion and, on other hand, is subject to short time perturbations whose duration is almost negligible. Such processes are often investigated in various fields of science and technology, such as physics, population dynamics, ecology, biological systems, optimal control, etc., see Refs. 1–11 and reference quoted therein.

It is well known in the theory of ordinary differential equations (see eg. Ref. 12 [p. 120]) that if for every continuous function f(t) bounded on $[0,\infty)$, the solution of the equation

$$x'(t) = A(t)x(t) + f(t),$$

satisfying x(0) = 0 is bounded on $[0, \infty)$, then the trivial solution of the corresponding homogeneous equation

$$x'(t) = A(t)x(t)$$

is uniformly asymptotically stable. This result is referred as Perron theorem Ref. 13. Later, Perron theorem has been extended to delay differential equations Ref. 12 [p. 371]. Indeed, it was shown that if for every continuous function f(t) bounded on $[0, \infty)$, the solution of the equation

$$x'(t) = A(t)x(t) + B(t)x(t - \tau) + f(t), \ t > 0$$

satisfying x(t)=0 for $t\in [-\tau,0]$ is bounded on $[0,\infty)$, then the trivial solution of the equation

$$x'(t) = A(t)x(t) + B(t)x(t - \tau),$$

is uniformly asymptotically stable. For more related materials, see the papers Refs. 14,15.

In this paper, we carry out the above result to a type of linear delay differential equations with impulse effects. Indeed, we consider equation of the form

$$x'(t) = A(t)x(t) + B(t)x(t - \tau), t \neq \theta_i,$$

$$\Delta x(\theta_i) := x(\theta_i^+) - x(\theta_i) = C_i x(\theta_i) + D_i x(\theta_{i-j}), i \in \mathbb{N},$$
(1)

and show that its trivial solution is uniformly asymptotically stable under a Perron condition. Our equation differs from the previous ones, see also Refs. 16–19, not only it is more general but also it allows delay terms in the impulse conditions. Such impulse conditions are more natural for delay differential equations.

With regard to equation (4) it is assumed that

- (i) A and B are $n \times n$ continuous bounded matrices, $\tau > 0$ is a positive real number;
- (ii) C_i and D_i are $n \times n$ bounded matrices, $j \in \mathbb{N}$ is fixed;
- (iii) $\{\theta_i\}$ is an increasing sequence of real numbers with $\lim_{i\to\infty}\theta_i=\infty$.

We also assume that $\det(I+C_i) \neq 0$ and that there exist a positive real numbers ρ and ν such that $||D_i|| \leq \rho$ and $||(I+C_i)^{-1}|| \leq \rho$ and $\theta_i - \theta_{i-j} \leq \nu$ for all $i \in \mathbb{N}$. $||\cdot||$ denotes any matrix norm.

Definition 1.1. Equation (4) is said to verify Perron condition if for every continuous bounded on $[0, \infty)$ function f(t) and every bounded sequence

 β_i the solution of

$$x'(t) = A(t)x(t) + B(t)x(t-\tau) + f(t), \quad t \neq \theta_i,$$

$$\Delta x(\theta_i) = C_i x(\theta_i) + D_i x(\theta_{i-j}) + \beta_i, \quad i \in \mathbb{N},$$
(2)

satisfying x(t) = 0 for $t \in [-\tau, 0]$ is bounded on $t \in [0, \infty)$.

By a solution of (6) on an interval J, we mean a function x defined on J such that x is continuous on J except possibly at $\theta_i \in J$ for $i \in \mathbb{N}$, where $x(\theta_i^+): \lim_{t \to \theta_i^+} x(t)$ and $x(\theta_i^-):= \lim_{t \to \theta_i^-} x(t)$ exist, $x(\theta_i^-):= x(\theta_i)$, and that $x(\theta_i^+): \lim_{t \to \theta_i^+} x(t)$ satisfies (6) on J. Clearly, if $f \equiv 0$ and $\beta_i = 0$ for all $i \in \mathbb{N}$ then (6) reduces to (4).

Let $PLC([-\tau,0],\mathbb{R}^n)$ denote the set of piecewise left continuous functions $\phi: [-\tau, 0] \to \mathbb{R}^n$ having a finite number of discontinuity points of the first kind. Under the above conditions, one can easily show that for given $\sigma \geq 0$ and $\phi \in PLC([-\tau,0],\mathbb{R}^n)$ there is a unique solution x(t) of (6) such that

$$x(t+\sigma) = \phi(t), \quad t \in [-\tau, 0]. \tag{3}$$

2. Preparatory lemmas

The following lemmas, see Ref. 12 for delay differential equations without impulse effects, are essential in proving the main result of this paper. Lemma 4.1 is needed to define an adjoint equation of (4), Lemma 2.2 provides representation of solutions, and Lemma 2.3 is concerned with the boundedness of fundamental matrices of (4).

Consider the equation

$$y'(t) = -A^{T}(t)y(t) - B^{T}(t+\tau)y(t+\tau), t \neq \theta_{i},$$

$$\Delta y(\theta_{i}) - (I + C_{i}^{T})^{-1}C_{i}^{T}y(\theta_{i}) - (I + C_{i}^{T})^{-1}D_{i+j}^{T}y(\theta_{i+j}^{+}), i \in \mathbb{N}.$$
(4)

We claim that equation (9) is an adjoint of (4) with respect to a function resembles the one used by Halanay in Ref. 12 [p. 371]. It turns out that this function has the form

$$< y(t), x(t) >= y^{T}(t)x(t) + \int_{t}^{t+\tau} y^{T}(s)B(s)x(s-\tau)ds + \sum_{n(t) \le k < n(t)+j} y^{T}(\theta_{k}^{+})D_{k}x(\theta_{k-j}),$$
 (5)

where

$$n(t) = \min\{i \in \mathbb{N} : \theta_i \ge t\}.$$

Lemma 2.1. If x(t) is a solution of (4) and y(t) is a solution of (9) then

$$\langle y(t), x(t) \rangle = c = constant,$$

where \langle , \rangle is defined by (10).

Proof. Let $t \in (\theta_i, \theta_{i+1})$. Then

$$\frac{\mathrm{d}}{\mathrm{d}t} \langle y(t), x(t) \rangle = -y^T(t)A(t)x(t) - y^T(t+\tau)B(t+\tau)x(t)$$
$$+ y^T(t+\tau)B(t+\tau)x(t) - y^T(t)B(t)x(t-\tau)$$
$$+ y^T(t)A(t)x(t) + y^T(t)B(t)x(t-\tau) = 0,$$

and hence $\langle y(t), x(t) \rangle = c_i = \text{constant for } t \in (\theta_i, \theta_{i+1})$. We may claim that $c_i = c$ for $i \in \mathbb{N}$. Indeed, since

$$c_{i+1} - c_i = \Delta \langle y(t), x(t) \rangle |_{t=\theta_i}$$

by (10) we have

$$c_{i+1} - c_i = y^T(\theta_i^+) x(\theta_i^+) - y^T(\theta_i) x(\theta_i)$$

$$+ \sum_{n(\theta_i^+) \le k < n(\theta_i^+) + j} y^T(\theta_k^+) D_k x(\theta_{k-j})$$

$$- \sum_{n(\theta_i) \le k < n(\theta_i^+) + j} y^T(\theta_k^+) D_k x(\theta_{k-j}).$$

Since $n(\theta_i^+) = i + 1$ and $n(\theta_i) = i$, we have

$$c_{i+1} - c_i = y^T(\theta_i^+) x(\theta_i^+) - y^T(\theta_i) x(\theta_i) - y^T(\theta_i^+) D_i x(\theta_{i-j}) + y^T(\theta_{i+j}^+) D_{i+j} x(\theta_i).$$

Using the impulse conditions in (4)

$$x(\theta_i^+) = (I + C_i)x(\theta_i) + D_ix(\theta_{i-1})$$

and the impulse conditions in (9)

$$y^{T}(\theta_{i+j}^{+})D_{i+j} = y^{T}(\theta_{i}) - y^{T}(\theta_{i}^{+}) - y^{T}(\theta_{i}^{+})C_{i}$$

we deduce that $c_{i+1} - c_i = 0$ for all $i \in \mathbb{N}$ and thus $\langle y(t), x(t) \rangle = c$.

Remark 2.1. It is easy to verify also that the adjoint of (9) is (4), i.e they are mutually adjoint of each other.

Definition 2.1. A matrix solution $X(t,\alpha)$ of (4) satisfying $X(\alpha,\alpha)=I$ and $X(t, \alpha) = 0$ for $t < \alpha$ is called a fundamental matrix of (4).

Definition 2.2. A matrix solution $Y(t,\alpha)$ of (9) satisfying $Y(\alpha,\alpha) = I$ and $Y(t, \alpha) = 0$ for $t > \alpha$ is said to be a fundamental matrix of (9).

Lemma 2.2. Let $X(t,\alpha)$ be a fundamental matrix of (4) and $\sigma \geq 0$ a real number. If x(t) is a solution of (6), then

$$x(t) = X(t,\sigma)x(\sigma)$$

$$+ \int_{\sigma-\tau}^{\sigma} X(t,\alpha+\tau)B(\alpha+\tau)x(\alpha)d\alpha + \int_{\sigma}^{t} X(t,\alpha)f(\alpha)d\alpha$$

$$+ \sum_{n(\sigma)-j\leq k < n(\sigma)} X(t,\theta_{k+j}^{+})D_{k+j}x(\theta_{k}) + \sum_{n(\sigma)\leq k < n(t)} X(t,\theta_{k}^{+})\beta_{k}.$$
(6)

Proof. Multiplying the differential equation in (6) by the matrix $Y^{T}(\alpha, t)$ and integrating with respect to α from σ to t, we obtain

$$x(t) = Y^{T}(\sigma, t)x(\sigma) - \int_{\sigma}^{t} Y^{T}(\alpha + \tau, t)B(\alpha + \tau)x(\alpha)d\alpha$$
$$+ \int_{\sigma}^{t} Y^{T}(\alpha, t)B(\alpha)x(\alpha - \tau)d\alpha + \int_{\sigma}^{t} Y^{T}(\alpha, t)f(\alpha)d\alpha$$
$$+ \sum_{n(\sigma) \leq k < n(t)} \left[Y^{T}(\theta_{k}^{+}, t)x(\theta_{k}^{+}) - Y^{T}(\theta_{k}, t)x(\theta_{k}) \right].$$

Replacing α by $\alpha + \tau$ in the second integral and using the impulse conditions in (6) and (9), we have

$$x(t) = Y^{T}(\sigma, t)x(\sigma) + \int_{\sigma - \tau}^{\sigma} Y^{T}(\alpha + \tau, t)B(\alpha + \tau)x(\alpha)d\alpha$$

$$+ \sum_{n(\sigma) - j \le k < n(\sigma)} Y^{T}(\theta_{k+j}^{+}, t)D_{k+j}x(\theta_{k})$$

$$+ \int_{\sigma}^{t} Y^{T}(\alpha, t)f(\alpha)d\alpha + \sum_{n(\sigma) \le k < n(t)} Y^{T}(\theta_{k}^{+}, t)\beta_{k}. \tag{7}$$

Since $X(t,\sigma) = Y^T(\sigma,t)$, which can be seen by replacing x(t) by the fundamental matrix $X(t,\sigma)$ in (7) with $f \equiv 0$ and $\beta_i = 0$ for all $i \in \mathbb{N}$, (7) is the same as (6). Corollary 2.1. Let $X(t, \alpha)$ be a fundamental matrix of (4) and $Y(t, \alpha)$ be a fundamental matrix of (9). Then

$$X(t, \alpha) = Y^T(\alpha, t).$$

Lemma 2.3. If (4) verifies Perron condition, then

$$||X(t,\alpha)|| < M \text{ for } t \ge \alpha \ge 0.$$

Proof. We first claim that there exists a constant d such that

$$\int_{0}^{t} \|X(t,\alpha)\| d\alpha + \sum_{0 \le m < n(t)} \|X(t,\theta_{m}^{+})\| < d \text{ for } t \ge 0.$$
 (8)

Define the space $\Pi = CB \times S$, where CB is the set of bounded functions $f \in C([0,\infty),\mathbb{R}^n)$ and S is the set of bounded sequences $\beta = \{\beta_m\}$, $\beta_m \in \mathbb{R}^n$, $m \in \mathbb{N}$. The elements are represented by the pair (f,β) supplied by the norm $\|(f,\beta)\| = \sup_{t \in [0,\infty)} \|f(t)\| + \sup_{m \in \mathbb{N}} \|\beta_m\|$. Consider the operator U defined on the Banach space Π by

$$U(f, \beta_m) = \int_0^t X(t, \alpha) f(\alpha) d\alpha + \sum_{0 \le m < n(t)} X(t, \theta_m^+) \beta_m.$$

We may use the Banach Steinhaus theorem Ref. 20 by employing similar arguments developed in Ref. 12 to arrive at (8).

Now let us consider (9) satisfied by $Y(\alpha, t)$. Integrating both sides from σ to t leads to

$$Y^{T}(\sigma,t) = I + \int_{\sigma}^{t} Y^{T}(\alpha,t)A(\alpha)d\alpha + \int_{\sigma}^{t} Y^{T}(\alpha+\tau,t)B(\alpha+\tau)d\alpha$$
$$-\sum_{n(\sigma)\leq i < n(t)} \Delta Y^{T}(\theta_{i},t).$$

Observing that $Y^{T}(\theta_{i}, t) = Y^{T}(\theta_{i}^{+}, t)(I + C_{i}) + Y^{T}(\theta_{i+j}^{+}, t)D_{i}$, we obtain

$$\sum_{n(\sigma) \le i < n(t)} \|\Delta Y^{T}(\theta_{i}, t)\| \le 2\rho \sum_{0 \le i < n(t)} \|Y^{T}(\theta_{i}^{+}, t)\|.$$

It follows that

$$||Y^{T}(\sigma,t)|| \le 1 + 2\gamma \int_{0}^{t} ||Y^{T}(\alpha,t)|| d\alpha + 2\rho \sum_{0 \le i < n(t)} ||Y^{T}(\theta_{i}^{+},t)||,$$

where

$$\gamma = \max \left\{ \sup_{t \ge 0} \|A(t)\|, \sup_{t \ge 0} \|B(t)\| \right\}.$$

Replacing $Y^{T}(\sigma,t)$ by $X(t,\sigma)$ and using inequality (8) result in the desired conclusion.

3. The main result

Theorem 3.1. If equation (4) verifies Perron condition then its trivial solution is uniformly asymptotically stable.

Proof. Let $x(t; \sigma, \phi)$ denote the solution of (4) satisfying (3). From Lemma 2.2,

$$x(t; \sigma, \phi) = X(t, \sigma)\phi(0) + \int_{-\tau}^{0} X(t, \alpha + \sigma + \tau)B(\alpha + \sigma + \tau)\phi(\alpha)d\alpha$$
$$+ \sum_{-j \le k \le 0} X(t, \theta_{k+n(\sigma)+j}^{+})D_{k+n(\sigma)+j}\phi(\theta_{k}).$$

By Lemma 2.3, there exists M > 0 such that ||X(t,r)|| < M. Hence

$$||x(t;\sigma,\phi)|| \le M(1+\tau\gamma+j\rho)||\phi||_0 = M_1||\phi||_0$$

where

$$M_1 = M(1 + \tau \gamma + j\rho)$$

and

$$\|\phi\|_0 = \sup_{r \in [-\tau, 0]} \|\phi(r)\|.$$

Thus, the zero solution of (4) is uniformly stable. To complete the proof we need to show that

$$\lim_{t \to \infty} x(t; \sigma, \phi) = 0 \quad \text{uniformly with respect to } \sigma \text{ and } \phi. \tag{9}$$

For our purpose, let $\mu \geq \sigma$. It is clear that $x(t) = x(t; \sigma, \phi)$ satisfies

$$x(t; \sigma, \phi) = X(t, \mu)x(\mu; \sigma, \phi) + \int_{\mu - \tau}^{\mu} X(t, \alpha + \tau)B(\alpha + \tau)x(\alpha; \sigma, \phi)d\alpha$$
$$+ \sum_{n(\mu) - j \le k < n(\mu)} X(t, \theta_{k+j}^+)D_{k+j}x(\theta_k; \sigma, \phi).$$

Integrating both sides from σ to t and then changing the order of integrations and the order of summation and the integral, we have

$$(t - \sigma)x(t; \sigma, \phi) = \int_{\sigma - \tau}^{\sigma} \int_{\sigma}^{\alpha + \tau} X(t, \alpha + \tau)B(\alpha + \tau)x(\alpha; \sigma, \phi) d\mu d\alpha$$

$$+ \int_{\sigma}^{t - \tau} \int_{\alpha}^{\alpha + \tau} X(t, \alpha + \tau)B(\alpha + \tau)x(\alpha; \sigma, \phi) d\mu d\alpha$$

$$+ \int_{\sigma}^{t} X(t, \mu)x(\mu; \sigma, \phi) d\mu$$

$$+ \sum_{n(\sigma) - j \le k < n(\sigma)} \int_{\sigma}^{\theta_{k+j}} X(t, \theta_{k+j}^{+}) D_{k+j}x(\theta_{k}; \sigma, \phi) d\mu$$

$$+ \sum_{n(\sigma) \le k < n(t) - j} \int_{\theta_{k}}^{\theta_{k+j}} X(t, \theta_{k+j}^{+}) D_{k+j}x(\theta_{k}; \sigma, \phi) d\mu.$$

We easily see from above that

$$(t - \sigma) \|x(t; \sigma, \phi)\| \le \gamma \tau^{2} M M_{1} \|\phi\|_{0} + j \rho \nu M M_{1} \|\phi\|_{0}$$

$$+ M_{1} \max\{\tau \gamma, \nu \rho, 1\} \|\phi\|_{0} \Big[\int_{0}^{t} \|X(t, s)\| ds$$

$$+ \sum_{0 \le r < n(t)} \|X(t, \theta_{r})\| \Big].$$

$$(10)$$

In view of (8), the right side of (10) is bounded. Hence

$$||x(t;\sigma,\phi)|| \le \frac{M_2}{t-\sigma} ||\phi||_0, \tag{11}$$

where M_2 is chosen so that

$$M_2 < MM_1(\gamma \tau^2 + j\rho \nu) + M_1 \max\{\tau \gamma_2, \nu \rho, 1\}d$$

Obviously, (9) follows from (11).

此为试读,需要完整PDF请访问: www.ertongbook.com