

# ORDINARY DIFFERENTIAL EQUATIONS

Second Edition

PHILIP HARTMAN



# ORDINARY DIFFERENTIAL EQUATIONS

Second Edition

PHILIP HARTMAN



1982

**Birkhäuser**

**Boston • Basel • Stuttgart**

Professor Philip Hartman\*  
Department of Mathematics  
The Johns Hopkins University  
Baltimore, Maryland 21218

*\*This work is a reprint of the second edition.*

**CIP - Kurztitelaufnahme der Deutschen Bibliothek**

Hartman, Philip:  
Ordinary differential equations / Philip Hartman.  
- Reprint of the 2. ed. -  
Boston ; Basel ; Stuttgart : Birkhäuser, 1982.  
ISBN 3-7643-3068-6

**Library of Congress Cataloging in Publication Data**

Hartman, Philip, 1915-  
Ordinary differential equations.  
Reprint. Originally published: Baltimore, Md., 1973.  
Bibliography: p.  
Includes index.  
1. Differential equations. I. Title.  
QA372.H33 1982 515.3'52 81-21738  
ISBN 3-7643-3068-6 AACR2

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the copyright owner.

©Birkhäuser Boston, 1982

ISBN 3-7643-3068-6

Printed in USA

# Preface to the First Edition

This book is based on lecture notes of courses on ordinary differential equations which I have given from time to time for advanced undergraduates and graduate students in mathematics, physics, and engineering. It assumes a knowledge of matrix theory and, if not a thorough knowledge of, at least a certain maturity in the handling of functions of real variables.

I was never tempted to scatter asterisks liberally throughout this book and claim that it could serve as a sophomore-junior-senior textbook, for I believe that a course of this type should give way to basic courses in analysis, algebra, and topology.

This book contains more material than I ever covered in one year but not all of the topics which I treated in the many courses. The contents of these courses always included the subject matter basic to the theory of differential equations and its many applications to other disciplines (as, for example, differential geometry). A "basic course" is covered in Chapter I; §§ 1-3 of Chapter II; §§ 1-6 and 8 of Chapter III; Chapter IV except for the "Application" in § 3 and part (ix) in § 8; §§ 1-4 of Chapter V; §§ 1-7 of Chapter VII; §§ 1-3 of Chapter VIII; §§ 1-12 of Chapter X; §§ 1-4 of Chapter XI; and §§ 1-4 of Chapter XII.

Many topics are developed in depth beyond that found in standard textbooks. The subject matter in a chapter is arranged so that more difficult, less basic, material is usually put at the end of the chapter (and/or in an appendix). In general, the content of any chapter depends only on the material in that chapter and the portion of the "basic course" preceding it. For example, after completing the basic course, an instructor can discuss Chapter IX, or the remainder of the contents of Chapter XII, or Chapter XIV, etc. There are two exceptions: Chapter VI, Part I, as written, depends on Chapter V, §§ 5-12; Part III of Chapter XII is not essential but is a good introduction to Chapter XIII.

Exercises have been roughly graded into three types according to difficulty. Many of the exercises are of a routine nature to give the student an opportunity to review or test his understanding of the techniques just explained. For more difficult exercises, there are hints in the back of the book (in some cases, these hints simplify available proofs). Finally, references are given for the most difficult exercises; these serve

to show extensions and further developments, and to introduce the student to the literature.

The theory of differential equations depends heavily on the "integration of differential inequalities" and this has been emphasized by collecting some of the main results on this topic in Chapter III and § 4 of Chapter IV. Much of the material treated in this book was selected to illustrate important techniques as well as results: the reduction of problems on differential equations to problems on "maps" (cf. Chapter VII, Appendix, and Chapter IX); the use of simple topological arguments (cf. Chapters VIII, § 1; X, §§ 2-7; and XIV, § 6); and the use of fixed point theorems and other basic facts in functional analysis (cf. Chapters XII and XIII).

I should like to acknowledge my deep indebtedness to the late Professor Aurel Wintner from whom and with whom I learned about differential equations, first as a student and later as a collaborator. My debt to him is at once personal, in view of my close collaboration with him, and impersonal, in view of his contributions to the resurgence of the theory of ordinary differential equations since the Second World War.

I wish to thank several students at Hopkins, in particular, N. Max, C. C. Pugh, and J. Wavrik, for checking parts of the manuscript. I also wish to express my appreciation to Miss Anna Lea Russell for the excellent typescript created from nearly illegible copy, numerous revisions, and changes in the revisions.

My work on this book was partially supported by the Air Force Office of Scientific Research.

PHILIP HARTMAN

Baltimore, Maryland

August, 1964

## *Preface to the Second Edition*

This edition is essentially that of 1964 (John Wiley and Sons, Inc.) with some corrections and additions. The only major changes are in Chapter IX. A few pertinent additions have been made to the bibliography as a Supplement, but no effort has been made to bring it up to date.

I should like to thank Professors T. Butler, K.T. Chen, W.A. Coppel, L. Lorch, M.E. Muldoon, L. Nicolson and C. Olech for corrections and/or suggestions, and Mrs. Margaret A. Einstein for able secretarial assistance.

P.H. (10/73)



# Contents

<b>I. Preliminaries</b>	<b>1</b>
1. Preliminaries, 1	
2. Basic theorems, 2	
3. Smooth approximations, 6	
4. Change of integration variables, 7	
Notes, 7	
<b>II. Existence</b>	<b>8</b>
1. The Picard-Lindelöf theorem, 8	
2. Peano's existence theorem, 10	
3. Extension theorem, 12	
4. H. Kneser's theorem, 15	
5. Example of nonuniqueness, 18	
Notes, 23	
<b>III. Differential inequalities and uniqueness</b>	<b>24</b>
1. Gronwall's inequality, 24	
2. Maximal and minimal solutions, 25	
3. Right derivatives, 26	
4. Differential inequalities, 26	
5. A theorem of Wintner, 29	
6. Uniqueness theorems, 31	
7. van Kampen's uniqueness theorem, 35	
8. Egress points and Lyapunov functions, 37	
9. Successive approximations, 40	
Notes, 44	
<b>IV. Linear differential equations</b>	<b>45</b>
1. Linear systems, 45	
2. Variation of constants, 48	
3. Reductions to smaller systems, 49	
4. Basic inequalities, 54	
5. Constant coefficients, 57	
6. Floquet theory, 60	

7. Adjoint systems, 62
8. Higher order linear equations, 63
9. Remarks on changes of variables, 68

#### APPENDIX. ANALYTIC LINEAR EQUATIONS, 70

10. Fundamental matrices, 70
11. Simple singularities, 73
12. Higher order equations, 84
13. A nonsimple singularity, 87
- Notes, 91

#### V. Dependence on initial conditions and parameters . . . . 93

1. Preliminaries, 93
2. Continuity, 94
3. Differentiability, 95
4. Higher order differentiability, 100
5. Exterior derivatives, 101
6. Another differentiability theorem, 104
7.  $S$ - and  $L$ -Lipschitz continuity, 107
8. Uniqueness theorem, 109
9. A lemma, 110
10. Proof of Theorem 8.1, 111
11. Proof of Theorem 6.1, 113
12. First integrals, 114
- Notes, 116

#### VI. Total and partial differential equations . . . . 117

##### PART I. A THEOREM OF FROBENIUS, 117

1. Total differential equations, 117
2. Algebra of exterior forms, 120
3. A theorem of Frobenius, 122
4. Proof of Theorem 3.1, 124
5. Proof of Lemma 3.1, 127
6. The system (1.1), 128

##### PART II. CAUCHY'S METHOD OF CHARACTERISTICS, 131

7. A nonlinear partial differential equation, 131
8. Characteristics, 135
9. Existence and uniqueness theorem, 137
10. Haar's lemma and uniqueness, 139
- Notes, 142

<b>VII. The Poincaré-Bendixson theory</b>	<b>144</b>
1. Autonomous systems, 144	
2. Umlaufsatz, 146	
3. Index of a stationary point, 149	
4. The Poincaré-Bendixson theorem, 151	
5. Stability of periodic solutions, 156	
6. Rotation points, 158	
7. Foci, nodes, and saddle points, 160	
8. Sectors, 161	
9. The general stationary point, 166	
10. A second order equation, 174	
<b>APPENDIX. POINCARÉ-BENDIXSON THEORY ON 2-MANIFOLDS, 182</b>	
11. Preliminaries, 182	
12. Analogue of the Poincaré-Bendixson theorem, 185	
13. Flow on a closed curve, 190	
14. Flow on a torus, 195	
Notes, 201	
<b>VIII. Plane stationary points</b>	<b>202</b>
1. Existence theorems, 202	
2. Characteristic directions, 209	
3. Perturbed linear systems, 212	
4. More general stationary point, 220	
Notes, 227	
<b>IX. Invariant manifolds and linearizations</b>	<b>228</b>
1. Invariant manifolds, 228	
2. The maps $T'$ , 231	
3. Modification of $F(\xi)$ , 232	
4. Normalizations, 233	
5. Invariant manifolds of a map, 234	
6. Existence of invariant manifolds, 242	
7. Linearizations, 244	
8. Linearization of a map, 245	
9. Proof of Theorem 7.1, 250	
10. Periodic solution, 251	
11. Limit cycles, 253	
<b>APPENDIX. SMOOTH EQUIVALENCE MAPS, 256</b>	
12. Smooth linearizations, 256	
13. Proof of Lemma 12.1, 259	
14. Proof of Theorem 12.2, 261	
<b>APPENDIX. SMOOTHNESS OF STABLE MANIFOLDS, 271</b>	
Notes, 271 L	



<b>X. Perturbed linear systems</b>	<b>273</b>
1. The case $E = 0$ , 273	
2. A topological principle, 278	
3. A theorem of Wazewski, 280	
4. Preliminary lemmas, 283	
5. Proof of Lemma 4.1, 290	
6. Proof of Lemma 4.2, 291	
7. Proof of Lemma 4.3, 292	
8. Asymptotic integrations. Logarithmic scale, 294	
9. Proof of Theorem 8.2, 297	
10. Proof of Theorem 8.3, 299	
11. Logarithmic scale (continued), 300	
12. Proof of Theorem 11.2, 303	
13. Asymptotic integration, 304	
14. Proof of Theorem 13.1, 307	
15. Proof of Theorem 13.2, 310	
16. Corollaries and refinements, 311	
17. Linear higher order equations, 314	
Notes, 320	
<b>XI. Linear second order equations</b>	<b>322</b>
1. Preliminaries, 322	
2. Basic facts, 325	
3. Theorems of Sturm, 333	
4. Sturm-Liouville boundary value problems, 337	
5. Number of zeros, 344	
6. Nonoscillatory equations and principal solutions, 350	
7. Nonoscillation theorems, 362	
8. Asymptotic integrations. Elliptic cases, 369	
9. Asymptotic integrations. Nonelliptic cases, 375	
<b>APPENDIX. DISCONJUGATE SYSTEMS, 384</b>	
10. Disconjugate systems, 384	
11. Generalizations, 396	
Notes, 401	
<b>XII. Use of implicit function and fixed point theorems</b>	<b>404</b>
<b>PART I. PERIODIC SOLUTIONS, 407</b>	
1. Linear equations, 407	
2. Nonlinear problems, 412	

**PART II. SECOND ORDER BOUNDARY VALUE PROBLEMS, 418**

3. Linear problems, 418
4. Nonlinear problems, 422
5. A priori bounds, 428

**PART III. GENERAL THEORY, 435**

6. Basic facts, 435
7. Green's functions, 439
8. Nonlinear equations, 441
9. Asymptotic integration, 445
- Notes, 447

**XIII. Dichotomies for solutions of linear equations . . . . . 450****PART I. GENERAL THEORY, 451**

1. Notations and definitions, 451
2. Preliminary lemmas, 455
3. The operator  $T$ , 461
4. Slices of  $\|Py(t)\|$ , 465
5. Estimates for  $\|y(t)\|$ , 470
6. Applications to first order systems, 474
7. Applications to higher order systems, 478
8.  $P(B, D)$ -manifolds, 483

**PART II. ADJOINT EQUATIONS, 484**

9. Associate spaces, 484
10. The operator  $T'$ , 486
11. Individual dichotomies, 486
12.  $P'$ -admissible spaces for  $T'$ , 490
13. Applications to differential equations, 493
14. Existence of  $PD$ -solutions, 497
- Notes, 498

**XIV. Miscellany on monotony . . . . . 500****PART I. MONOTONE SOLUTIONS, 500**

1. Small and large solutions, 500
2. Monotone solutions, 506
3. Second order linear equations, 510
4. Second order linear equations (continuation), 515

## PART II. A PROBLEM IN BOUNDARY LAYER THEORY, 519

5. The problem, 519
6. The case  $\lambda > 0$ , 520
7. The case  $\lambda < 0$ , 525
8. The case  $\lambda = 0$ , 531
9. Asymptotic behavior, 534

## PART III. GLOBAL ASYMPTOTIC STABILITY, 537

10. Global asymptotic stability, 537
  11. Lyapunov functions, 539
  12. Nonconstant  $G$ , 540
  13. On Corollary 11.2, 545
  14. On " $J(y)x \cdot x \leq 0$  if  $x \cdot f(y) = 0$ ", 548
  15. Proof of Theorem 14.2, 550
  16. Proof of Theorem 14.1, 554
- Notes, 554

## HINTS FOR EXERCISES, 557

## REFERENCES, 581

## INDEX, 607

# Chapter I

## Preliminaries

### 1. Preliminaries

Consider a system of  $d$  first order differential equations and an initial condition

$$(1.1) \quad y' = f(t, y), \quad y(t_0) = y_0,$$

where  $y' = dy/dt$ ,  $y = (y^1, \dots, y^d)$  and  $f = (f^1, \dots, f^d)$  are  $d$ -dimensional vectors, and  $f(t, y)$  is defined on a  $(d+1)$ -dimensional  $(t, y)$ -set  $E$ . For the most part, it will be assumed that  $f$  is continuous. In this case,  $y = y(t)$  defined on a  $t$ -interval  $J$  containing  $t = t_0$  is called a solution of the initial value problem (1.1) if  $y(t_0) = y_0$ ,  $(t, y(t)) \in E$ ,  $y(t)$  is differentiable, and  $y'(t) = f(t, y(t))$  for  $t \in J$ . It is clear that  $y(t)$  then has a continuous derivative. These requirements on  $y$  are equivalent to the following:  $y(t_0) = y_0$ ,  $(t, y(t)) \in E$ ,  $y(t)$  is continuous and

$$(1.2) \quad y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds$$

for  $t \in J$ .

An initial value problem involving a system of equations of  $m$ th order,

$$(1.3) \quad z^{(m)} = F(t, z, z^{(1)}, \dots, z^{(m-1)}), \quad z^{(j)}(x_0) = z_0^{(j)} \text{ for } j = 0, \dots, m-1,$$

where  $z^{(j)} = d^j z / dt^j$ ,  $z$  and  $F$  are  $e$ -dimensional vectors, and  $F$  is defined on an  $(me+1)$ -dimensional set  $E$ , can be considered as a special case of (1.1), where  $y$  is a  $d = me$ -dimensional vector, symbolically,  $y = (z, z^{(1)}, \dots, z^{(m-1)})$  (or more exactly,  $y = (z^1, \dots, z^e, z^{1'}, \dots, z^{e'}, \dots, z^{e(m-1)'})$ ); correspondingly,  $f(t, y) = (z^{(1)}, \dots, z^{(m-1)}, F(t, y))$  and  $y_0 = (z_0, z_0^{(1)}, \dots, z_0^{(m-1)})$ . For example, if  $e = 1$  so that  $z$  is a scalar, (1.3) becomes

$$y^{1'} = y^2, \dots, y^{m-1'} = y^m, \quad y^{m'} = F(t, y^1, \dots, y^m),$$

$$y^j(t_0) = z_0^{(j-1)} \quad \text{for } j = 1, \dots, m,$$

where  $y^1 = z$ ,  $y^2 = z'$ ,  $\dots$ ,  $y^m = z^{(m-1)}$ .

## 2 Ordinary Differential Equations

The first set of questions to be considered will be (1) local existence (does (1.1) have a solution  $y(t)$  defined for  $t$  near  $t_0$ ?); (2) existence in the large (on what  $t$ -ranges does a solution of (1.1) exist?); and (3) uniqueness of solutions.

The significance of question (2) is clear from the following situation: Let  $y, f$  be scalars,  $f(t, y)$  defined for  $0 \leq t \leq 1, |y| \leq 1$ . A solution  $y = y(t)$  of (1.1), with  $(t_0, y_0) = (0, 0)$ , may exist for  $0 \leq t \leq \frac{1}{2}$  and increase from 0 to 1 as  $t$  goes from 0 to  $\frac{1}{2}$ , then one cannot expect to have an extension of  $y(t)$  for any  $t > \frac{1}{2}$ . Or consider the following scalar case where  $f(t, y)$  is defined for all  $(t, y)$ :

$$(1.4) \quad y' = y^2, \quad y(0) = c (> 0).$$

It is easy to see that  $y = c/(1 - ct)$  is a solution of (1.4), but this solution exists only on the range  $-\infty < t < 1/c$ , which depends on the initial condition.

In order to illustrate the significance of the question of uniqueness, let  $y$  be a scalar and consider the initial value problem

$$(1.5) \quad y' = |y|^{1/2}, \quad y(0) = 0.$$

This has more than one solution, in fact, it has, e.g., the solution  $y(t) \equiv 0$




Figure 1.

and the 1-parameter family of solutions defined by  $y(t) = 0$  for  $t \leq c$ ,  $y(t) = (t - c)^2/4$  for  $t \geq c$ , where  $c \geq 0$ ; see Figure 1. This situation is typical in that if (1.1) has more than one solution, then it has a "continuum" of solutions; cf. Theorem II 4.1.

### 2. Basic Theorems

This section introduces some conventions, notions and theorems to be used later. The proofs of most of the theorems will be omitted.



The symbols  $O$ ,  $o$  will be used from time to time where, e.g.,  $f(t) = O(g(t))$  as  $t \rightarrow \infty$  means that there exists a constant  $C$  such that  $|f(t)| \leq C|g(t)|$  for large  $t$ , while  $f(t) = o(g(t))$  as  $t \rightarrow \infty$  means that  $C > 0$  can be chosen arbitrarily small (so that if  $g(t) \neq 0$ ,  $f(t)/g(t) \rightarrow 0$  as  $t \rightarrow \infty$ ).

"Function" below generally means a map from some specified set of a vector space  $R^e$  into a space  $R^d$ , not always of the same dimension.  $R^d$  denotes a normed, real  $d$ -dimensional vector space of elements  $y = (y^1, \dots, y^d)$  with norm  $|y|$ . Unless otherwise specified,  $|y|$  will be the norm

$$(2.1) \quad |y| = \max(|y^1|, \dots, |y^d|),$$

and  $\|y\|$  the Euclidean norm.

If  $y_0$  is a point and  $E$  a subset of  $R^d$ , then  $\text{dist}(y_0, E)$ , the distance from  $y_0$  to  $E$ , is defined to be  $\inf |y_0 - y|$  for  $y \in E$ . If  $E_1, E_2$  are two subsets of  $R^d$ , then  $\text{dist}(E_1, E_2)$  is defined to be  $\inf |y_1 - y_2|$  for  $y_1 \in E_1, y_2 \in E_2$ , and is called the distance between  $E_1$  and  $E_2$ . If  $E_1$  (or  $E_2$ ) is compact and  $E_1, E_2$  are closed and disjoint, then  $\text{dist}(E_1, E_2) > 0$ .

If  $E$  is an open set or a closed parallelepiped in  $R^d$ ,  $f \in C^n(E)$ ,  $0 \leq n < \infty$ , means that  $f(y)$  is continuous on  $E$  and that the components of  $f$  have continuous partial derivatives of all orders  $k \leq n$  with respect to  $y^1, \dots, y^d$ .

A function  $f(y, z) = f(y^1, \dots, y^d, z^1, \dots, z^e)$  defined on a  $(y, z)$ -set  $E$ , where  $y \in R^d$ , is said to be *uniformly Lipschitz continuous on  $E$  with respect to  $y$*  if there exists a constant  $K$  satisfying

$$(2.2) \quad |f(y_1, z) - f(y_2, z)| \leq K|y_1 - y_2| \quad \text{for all } (y_j, z) \in E$$

with  $j = 1, 2$ . Any constant  $K$  satisfying (2.1) is called a *Lipschitz constant* (for  $f$  on  $E$ ). (The admissible values of  $K$  depend, of course, on the norms in the  $f$ - and  $y$ -spaces.)

A family  $F$  of functions  $f(y)$  defined on some  $y$ -set  $E \subset R^d$  is said to be *equicontinuous* if, for every  $\epsilon > 0$ , there exists a  $\delta = \delta_\epsilon > 0$  such that  $|f(y_1) - f(y_2)| \leq \epsilon$  whenever  $y_1, y_2 \in E$ ,  $|y_1 - y_2| \leq \delta$  and  $f \in F$ . The point of this definition is that  $\delta_\epsilon$  does not depend on  $f$  but is admissible for all  $f \in F$ . The most frequently encountered equicontinuous families  $F$  below will occur when all  $f \in F$  are uniformly Lipschitz continuous on  $E$  and there exists a  $K > 0$  which is a Lipschitz constant for all  $f \in F$ ; in which case,  $\delta$  can be chosen to be  $\delta = \epsilon/K$ .

**Lemma 2.1.** *If a sequence of continuous functions on a compact set  $E$  is uniformly convergent on  $E$ , then it is uniformly bounded and equicontinuous.*

**Cantor Selection Theorem 2.1.** *Let  $f_1(y), f_2(y), \dots$  be a uniformly bounded sequence of functions on a  $y$ -set  $E$ . Then for any countable set  $D \subset E$ , there exists a subsequence  $f_{n(1)}(y), f_{n(2)}(y), \dots$  convergent on  $D$ .*

#### 4 Ordinary Differential Equations

In order to prove Cantor's theorem, let  $D$  consist of the points  $y_1, y_2, \dots$ . Also assume that  $f_n(y)$  is real-valued; the proof for the case that  $f_n(y) = (f_n^1(y), \dots, f_n^d(y))$  is a  $d$ -dimensional vector is similar. The sequence of numbers  $f_1(y), f_2(y), \dots$  is bounded, thus, by the theorem of Bolzano-Weierstrass, there is a sequence of integers  $n_1(1) < n_1(2) < \dots$  such that  $\lim f_{n_1(k)}(y_1)$  exists as  $k \rightarrow \infty$ , where  $n = n_1(k)$ . Similarly there is a subsequence  $n_2(1) < n_2(2) < \dots$  of  $n_1(1), n_1(2), \dots$  such that  $\lim f_{n_2(k)}(y_2)$  exists as  $k \rightarrow \infty$  for  $n = n_2(k)$ . Continuing in this fashion one obtains successive subsequences of positive integers, such that if  $n_j(1) < n_j(2) < \dots$  is the  $j$ th one, then  $\lim f_{n_j(k)}(y_i)$  exists on  $k \rightarrow \infty$ , where  $n = n_j(k)$  and  $i = 1, \dots, j$ . The desired subsequence is the "diagonal sequence"  $n_1(1) < n_2(2) < n_3(3) < \dots$ . Variants of this proof will be referred to as the "standard diagonal process."

The next two assertions usually have the names Ascoli or Arzela attached to them.

**Propagation Theorem 2.2.** *On a compact  $y$ -set  $E$ , let  $f_1(y), f_2(y), \dots$  be a sequence of functions which is equicontinuous and convergent on a dense subset of  $E$ . Then  $f_1(y), f_2(y), \dots$  converges uniformly on  $E$ .*

**Selection Theorem 2.3.** *On a compact  $y$ -set  $E \subset \mathbb{R}^d$ , let  $f_1(y), f_2(y), \dots$  be a sequence of functions which is uniformly bounded and equicontinuous. Then there exists a subsequence  $f_{n(1)}(y), f_{n(2)}(y), \dots$  which is uniformly convergent on  $E$ .*

This last theorem can be obtained as a consequence of the preceding two. By applying Theorem 2.3 to a suitable subsequence, we obtain the following:

**Remark 1.** If, in the last theorem,  $y_0 \in E$  and  $f_0$  is a cluster point of the sequence  $f_1(y_0), f_2(y_0), \dots$ , then the subsequence  $f_{n(1)}(y), f_{n(2)}(y), \dots$  in the assertion can be chosen so that the limit function  $f(y)$  satisfies  $f(y_0) = f_0$ .

**Remark 2.** If, in Theorem 2.3, it is known that all (uniformly) convergent subsequences of  $f_1(y), f_2(y), \dots$  have the same limit, say  $f(y)$ , then a selection is unnecessary and  $f(y)$  is the uniform limit of  $f_1(y), f_2(y), \dots$ . This follows from Remark 1.

Theorem 2.3 and the following consequences of it will be used repeatedly.

**Theorem 2.4.** *Let  $y, f \in \mathbb{R}^d$  and  $f_0(t, y), f_1(t, y), f_2(t, y), \dots$  be a sequence of continuous functions on the parallelepiped  $R: t_0 \leq t \leq t_0 + a, |y - y_0| \leq b$  such that*

$$(2.3) \quad f_0(t, y) = \lim_{n \rightarrow \infty} f_n(t, y) \quad \text{uniformly on } R.$$

Let  $y_n(t)$  be a solution of

$$(2.4_n) \quad y' = f_n(t, y), \quad y(t_n) = y_n,$$

on  $[t_0, t_0 + a]$ , where  $n = 1, 2, \dots$ , and

$$(2.5) \quad t_n \rightarrow t_0, \quad y_n \rightarrow y_0 \quad \text{as} \quad n \rightarrow \infty.$$

Then there exists a subsequence  $y_{n(1)}(t), y_{n(2)}(t), \dots$  which is uniformly convergent on  $[t_0, t_0 + a]$ . For any such subsequence, the limit

$$(2.6) \quad y_0(t) = \lim_{k \rightarrow \infty} y_{n(k)}(t)$$

is a solution of (2.4<sub>0</sub>) on  $[t_0, t_0 + a]$ . In particular, if (2.4<sub>0</sub>) possesses a unique solution  $y = y_0(t)$  on  $[t_0, t_0 + a]$ , then

$$(2.7) \quad y_0(t) = \lim_{n \rightarrow \infty} y_n(t) \quad \text{uniformly on } [t_0, t_0 + a].$$

**Proof.** Since  $f_1, f_2, \dots$  are continuous and (2.3) holds uniformly on  $R$ , there is a constant  $K$  such that  $|f_n(t, y)| \leq K$  for  $n = 0, 1, \dots$  and  $(t, y) \in R$ ; Lemma 2.1. Since  $|y_n'(t)| \leq K$ , it is clear that  $K$  is a Lipschitz constant for  $y_1, y_2, \dots$ , so that this sequence is equicontinuous. It is also uniformly bounded since  $|y_n(t) - y_0| \leq b$ . Thus the existence of uniformly convergent subsequences follows from Theorem 2.3. By (2.3), Lemma 2.1, and the uniformity of (2.6), it is easy to see that

$$f_{n(k)}(t, y_{n(k)}(t)) \rightarrow f_0(t, y_0(t))$$

uniformly on  $[t_0, t_0 + a]$  as  $k \rightarrow \infty$ . Thus term-by-term integration is applicable to

$$y_n(t) = y_n + \int_{t_n}^t f_n(s, y_n(s)) \, ds$$

where  $n = n(k)$  and  $k \rightarrow \infty$ . It follows that the limit (2.6) is a solution of (2.4<sub>0</sub>).

As to the last assertion, note that the assumed uniqueness of the solution  $y_0(t)$  of (2.4<sub>0</sub>) shows that the limit of every (uniformly) convergent subsequence of  $y_1(t), y_2(t), \dots$  is the solution  $y_0(t)$ . Hence a selection is unnecessary and (2.7) holds by Remark 2 above.

**Implicit Function Theorem 2.5.** Let  $x, y, f, g$  be  $d$ -dimensional vectors and  $z$  an  $e$ -dimensional vector. Let  $f(y, z)$  be continuous for  $(y, z)$  near a point  $(y_0, z_0)$  and have continuous partial derivatives with respect to the components of  $y$ . Let the Jacobian  $\det(\partial f^i / \partial y^k) \neq 0$  at  $(y, z) = (y_0, z_0)$ . Let  $x_0 = f(y_0, z_0)$ . Then there exist positive numbers,  $\epsilon$  and  $\delta$ , such that if  $x$  and  $z$  are fixed,  $|x - x_0| < \delta$  and  $|z - z_0| < \delta$ , then the equation  $x = f(y, z)$  has a unique solution  $y = g(x, z)$  satisfying  $|y - y_0| < \epsilon$ . Furthermore,  $g(x, z)$  is continuous for  $|x - x_0| < \delta, |z - z_0| < \delta$  and has continuous partial derivatives with respect to the components of  $x$ .

For a sharper form of this theorem, see Exercise II 2.3.

## 3. Smooth Approximations

In some situations, it will be convenient to extend the definition of a function  $f$ , say, given continuous on a closed parallelepiped, or to approximate it uniformly by functions which are smooth ( $C^1$  or  $C^\infty$ ) with respect to certain variables. The following devices can be used to obtain such extensions or approximations (which have the same bounds as  $f$ ).

Let  $f(t, y)$  be defined on  $R: t_0 \leq t \leq t_1, |y| \leq b$  and let  $|f(t, y)| \leq M$ . Let  $f^*(t, y)$  be defined for  $t_0 \leq t \leq t_1$  and all  $y$  by placing  $f^*(t, y) = f(t, y)$  if  $|y| \leq b$  and  $f^*(t, y) = f(t, by/|y|)$  if  $|y| > b$ . It is clear that  $f^*(t, y)$  is continuous for  $t_0 \leq t \leq t_1, y$  arbitrary, and that  $|f^*(t, y)| \leq M$ . In some cases, it is more convenient to replace  $f^*$  by an extension of  $f$  which is 0 for large  $|y|$ . Such an extension is given by  $f^0(t, y) = f^*(t, y)\varphi^0(|y|)$ , where  $\varphi^0(s)$  is a continuous function for  $t \geq 0$  satisfying  $0 \leq \varphi^0(s) \leq 1$  for  $s \geq 0$ ,  $\varphi^0(s) = 1$  for  $0 \leq s \leq b$ , and  $\varphi^0(s) = 0$  for  $s \geq b + 1$ .

In order to approximate  $f(t, y)$  uniformly on  $R$  by functions  $f^\epsilon(t, y)$  which are, say, smooth with respect to the components of  $y$ , let  $\varphi(s)$  be a function of class  $C^\infty$  for  $s \geq 0$  satisfying  $\varphi(s) > 0$  for  $0 \leq s < 1$  and  $\varphi(s) = 0$  for  $s \geq 1$ . Then there is a constant  $c > 0$  depending only on  $\varphi(s)$  and the dimension  $d$ , such that for every  $\epsilon > 0$ ,

$$(3.1) \quad c\epsilon^{-d} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} \varphi(\epsilon^{-2} \|y\|^2) dy^1 \dots dy^d = 1,$$

where  $\|y\| = (\sum |y^k|^2)^{1/2}$  is the Euclidean length of  $y$ . Put

$$(3.2) \quad f^\epsilon(t, y) = c\epsilon^{-d} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} f^0(t, \eta) \varphi(\epsilon^{-2} \|y - \eta\|^2) d\eta^1 \dots d\eta^d,$$

where  $\eta = (\eta^1, \dots, \eta^d)$ , so that

$$(3.3) \quad f^\epsilon(t, y) = c\epsilon^{-d} \int_{-\infty}^{+\infty} \dots \int_{-\infty}^{+\infty} f^0(t, y - \eta) \varphi(\epsilon^{-2} \|\eta\|^2) d\eta^1 \dots d\eta^d.$$

Since  $f^\epsilon(t, y)$  is an "average" of the values of  $f^0$  in a sphere  $\|\eta - y\| \leq \epsilon$  for a fixed  $t$ , it is clear that  $f^\epsilon \rightarrow f^0$  as  $\epsilon \rightarrow 0$  uniformly on  $t_0 \leq t \leq t_1, y$  arbitrary. Note that  $|f^\epsilon| \leq M$  for all  $\epsilon > 0$  and that  $f^\epsilon(t, y) = 0$  for  $|y| \geq b + 1 + \epsilon$ . Furthermore,  $f^\epsilon(t, y)$  has continuous partial derivatives of all orders with respect to  $y^1, \dots, y^d$ .

The last formula can be used to show that if  $f^0(t, y)$  has continuous partial derivatives of order  $k$  with respect to  $y^1, \dots, y^d$ , then the corresponding partial derivatives of  $f^\epsilon(t, y)$  tend uniformly to those of  $f^0(t, y)$  as  $\epsilon \rightarrow 0$ .