



IRMA Lectures in Mathematics
and Theoretical Physics 6

Athanase Papadopoulos

Metric Spaces, Convexity and Nonpositive Curvature



European Mathematical Society

Athanase Papadopoulos

Metric Spaces, Convexity and Nonpositive Curvature



European Mathematical Society

Author:

Athanase Papadopoulos
CNRS and Université Louis Pasteur
Institut de Recherche Mathématiques Avancée
7 rue René Descartes
F-67084 Strasbourg Cedex
France

2000 Mathematics Subject Classification 26-01; 30F25; 30F45; 30F60; 32G15; 32Q45; 51-01; 51K05; 51K10; 51M09; 51M10; 51F99; 52-01; 52A07; 52A41; 53-01; 53C70; 54-01; 54E35.

Bibliographic information published by Die Deutsche Bibliothek

Die Deutsche Bibliothek lists this publication in the Deutsche Nationalbibliografie;
detailed bibliographic data are available in the Internet at <http://dnb.ddb.de>.

ISBN 3-03719-010-8

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, re-use of illustrations, recitation, broadcasting, reproduction on microfilms or in other ways, and storage in data banks. For any kind of use permission of the copyright owner must be obtained.

© 2005 European Mathematical Society

Contact address:

European Mathematical Society Publishing House
Seminar for Applied Mathematics
ETH-Zentrum FLI C4
CH-8092 Zürich
Switzerland

Phone: +41 (0)1 632 34 36
Email: info@ems-ph.org
Homepage: www.ems-ph.org

Typeset using the author's \TeX files: I. Zimmermann, Freiburg
Printed in Germany

9 8 7 6 5 4 3 2 1

IRMA Lectures in Mathematics and Theoretical Physics 6

Edited by Vladimir G. Turaev

Institut de Recherche Mathématique Avancée
Université Louis Pasteur et CNRS
7 rue René Descartes
67084 Strasbourg Cedex
France

IRMA Lectures in Mathematics and Theoretical Physics

Edited by Vladimir G. Turaev

This series is devoted to the publication of research monographs, lecture notes, and other materials arising from programs of the Institut de Recherche Mathématique Avancée (Strasbourg, France). The goal is to promote recent advances in mathematics and theoretical physics and to make them accessible to wide circles of mathematicians, physicists, and students of these disciplines.

Previously published in this series:

- 1 Deformation Quantization, *Gilles Halbout* (Ed.)
- 2 Locally Compact Quantum Groups and Groupoids, *Leonid Vainerman* (Ed.)
- 3 From Combinatorics to Dynamical Systems, *Frédéric Fauvet and Claude Mitschi* (Eds.)
- 4 Three courses on Partial Differential Equations, *Eric Sonnendrücker* (Ed.)
- 5 Infinite Dimensional Groups and Manifolds, *Tilman Wurzbacher* (Ed.)

Volumes 1–5 are available from Walter de Gruyter (www.degruyter.de)

Dedicated to the memory of my parents

Preface

This book is about metric spaces of nonpositive curvature in the sense of Busemann, that is, metric spaces whose distance function is convex. We have also included a systematic introduction to the theory of geodesics and related matters in metric spaces, as well as a detailed presentation of a few facets of convexity theory that are useful in the study of nonpositive curvature.

We have tried to start from first principles and to give full proofs, but there are two exceptions, which occur several times in examples that are spread through the book. These exceptions are the following:

- we give without proof some classical well-known facts about hyperbolic spaces \mathbb{H}^n ;
- we refer to some elements of the theory of Teichmüller space.

The material that we present in connection with these spaces is used as an illustration for the general concepts that are developed. It can be asserted without exaggeration that Teichmüller space is the most beautiful space ever, and the metric theory of that space, in particular its convexity properties, is most interesting.

This book can be divided into three parts. Part I (Chapters 1 to 4) contains basic material on metric spaces, Part II (Chapters 5 to 7) concerns convexity in vector spaces and Part III (Chapters 8 to 12) concerns convexity in metric spaces and related matters.

Each chapter contains an introduction in which we describe its content. At the end of each chapter we have included a few notes, some of them historical and others indicating some further developments. In both cases, there is no pretention for completeness. The only reason for which a note is included is that the author of this book thinks that it is interesting information.

Acknowledgements. This text grew up from notes I wrote for graduate courses I gave at the Universities of Strasbourg and of Florence. The reader will easily realize that its content owes a lot to the work of Herbert Busemann. The course in Italy was supported by the Istituto Nazionale di Alta Matematica (Roma). I would like to thank Raffaella Lefkou and Vincenzo Ancona who invited me to give that course.

Contents

Preface	vii
Introduction: Some historical markers	1
The work of Hadamard	1
The works of Menger and Wald	4
The works of Busemann and Alexandrov	6
Convexity	8
1 Lengths of paths in metric spaces	10
1.1 The length of a path	11
1.2 Arclength as parameter	19
1.3 Differentiable paths in Euclidean space	22
1.4 The space of paths	24
Notes on Chapter 1	31
2 Length spaces and geodesic spaces	34
2.1 Length spaces	35
2.2 Geodesics	50
2.3 Limits of geodesics	56
2.4 Geodesic spaces	58
2.5 Geodesic convexity	67
2.6 Menger convexity	69
Notes on Chapter 2	77
3 Maps between metric spaces	79
3.1 K -Lipschitz maps and K -length-non-increasing maps	80
3.2 Non-expanding maps	82
3.3 Distance non-decreasing maps	87
3.4 Local isometries	89
3.5 Covering spaces	96
Notes on Chapter 3	101
4 Distances	103
4.1 The Hausdorff distance	105
4.2 The Busemann–Hausdorff distance	110
4.3 Closed limits of subsets	112
4.4 Metrics on the isometry group	119
Notes on Chapter 4	124

5	Convexity in vector spaces	127
5.1	Affinely convex subsets	127
5.2	Convex hull	133
5.3	Convexity in normed vector spaces	137
5.4	Limits of convex sets	143
5.5	Minkowski's construction	144
5.6	The Hilbert geometry	148
	Notes on Chapter 5	156
6	Convex functions	159
6.1	Convex functions	160
6.2	Convex functions of one variable	170
	Notes on Chapter 6	175
7	Strictly convex normed vector spaces	178
7.1	Strictly convex normed vector spaces	178
7.2	Uniquely geodesic spaces	180
7.3	Inner products and ℓ^p norms	185
	Notes on Chapter 7	186
8	Busemann spaces	187
8.1	Busemann spaces	187
8.2	Local geodesics in Busemann spaces	196
8.3	Geodesic convexity in Busemann spaces	198
8.4	Convex functions on Busemann spaces	199
	Notes on Chapter 8	204
9	Locally convex spaces	210
9.1	Locally convex spaces	211
9.2	Variation of local geodesics	218
9.3	The universal covering of a locally convex metric space	224
	Notes on Chapter 9	227
10	Asymptotic rays and the visual boundary	229
10.1	Asymptotic rays	230
10.2	The visual boundary	235
	Notes on Chapter 10	239
11	Isometries	241
11.1	Minimal displacement, minimal sets and the isometry types	242
11.2	Axial isometries	249
11.3	Periodic geodesics	253
11.4	Axial isometries of Busemann spaces	256

11.5 Parallel lines	258
Notes on Chapter 11	260
12 Busemann functions, co-rays and horospheres	261
12.1 Busemann functions	261
12.2 Co-rays	265
12.3 Horospheres	269
Notes on Chapter 12	273
References	275
Index	283

Introduction: Some historical markers

The aim of this introduction is threefold. First, we give a brief account of a paper by Jacques Hadamard which is at the genesis of many works on global properties of surfaces of nonpositive Gaussian curvature. Then, we make a brief account of several different notions of nonpositive curvature in metric spaces, each of which generalizes in its own manner the notion of nonpositive Gaussian curvature for surfaces or of nonpositive sectional curvature for Riemannian manifolds. In particular, we mention important works by Menger, Busemann and Alexandrov in this domain. Finally, we review some of the connections between convexity theory and the theory of nonpositive curvature, which constitute the main theme that we develop in this book.

The work of Hadamard

The theory of spaces of nonpositive curvature has a long and interesting history, and it is good to look at its sources. We start with a brief review of the pioneering paper by Hadamard, “Les surfaces à courbures opposées et leurs lignes géodésiques” [62]. This paper, which was written at the end of the nineteenth century, can be considered as the foundational paper for the study of global properties of nonpositively curved spaces.

Hadamard was one of the first mathematicians (and may have been the first) who strongly emphasized the importance of topological methods in the study of spaces of nonpositive curvature. In the introduction to the paper cited above, he writes : “The only theory which has to be studied profoundly, as a basis to the current work, is the *Analysis situs*, which, as one can expect after reading the work of Poincaré, plays an essential role in everything that will follow.”

Let us look more closely at the content of that paper.

Hadamard considers a surface S equipped with a Riemannian metric of nonpositive curvature. The surface is smoothly embedded in \mathbb{R}^3 , the Riemannian metric is induced from that inclusion and the set of points at which the curvature is zero is finite.

Hadamard starts by noting that this surface is necessarily unbounded since if one of the coordinates assumes a maximum or a minimum, then, in a neighborhood of that point, the surface would be situated at one side of its tangent plane and therefore the curvature would be positive. He then proves that such a surface can always be decomposed into the union of a compact region and of a collection of *infinite sheets* (we are translating the term “nappes infinies” used by Hadamard). Each infinite sheet is homeomorphic to a cylinder, and is connected to the compact region along a boundary curve. Hadamard constructs examples of surfaces with nonpositive curvature having

an arbitrary number of infinite sheets. He classifies the infinite sheets into two types: the *flared* infinite sheets (“nappes infinies évasées”), which we call “funnels”, and the *unflared* infinite sheets. An unflared infinite sheet is characterized by the fact that one can continuously push to infinity a homotopically non-trivial closed curve on such a sheet while keeping its length bounded from above. To each unflared infinite sheet, Hadamard associates an *asymptotic direction*. The simple closed curves that connect the funnels to the compact region can be taken to be closed geodesics that are pairwise disjoint.

Let us now suppose that the fundamental group of S is finitely generated (Hadamard says that S has “finite connectivity”). For such a surface, Hadamard describes a finite collection of homotopically non-trivial and pairwise non-homotopic simple closed curves C_1, \dots, C_n such that any non-trivial homotopy class of closed curves on the surface can be uniquely represented by a finite word written in the letters $C_1, \dots, C_n, C_1^{-1}, \dots, C_n^{-1}$. Hadamard pushes forward this analysis to make it include not only representations of closed curves, but also representations of long segments of open infinite curves. In fact, with these elements, Hadamard initiates the approximation of long segments of bi-infinite geodesic lines by closed geodesics, and therefore he also initiates the theory of symbolic representation of bi-infinite geodesics on the surface, that is, the theory of representation of elements of the geodesic flow associated to S by bi-infinite words in the finite alphabet $\{C_1, \dots, C_n, C_1^{-1}, \dots, C_n^{-1}\}$. We note in passing that the theory of symbolic dynamics for the geodesic flow associated to a nonpositively curved surface was thoroughly developed about twenty years later by Marston Morse, and that Morse, in his paper [113], refers to Hadamard as his source of inspiration. Hadamard calls a closed curve a “contour”, and the curves C_1, \dots, C_n the “elementary contours”. There are two sorts of closed curves in the collection C_1, \dots, C_n which Hadamard considers: the curves that are the boundaries of the infinite sheets, and those that correspond to the genus of the surface. (These curves come in pairs and they correspond to the “holes” of the compact part of the surface S , as Hadamard calls them.) He notices that there is (up to circular permutation) one and only one relation in the representation he obtains that allows us to consider any of these contours as a product of the elementary contours, and therefore, by eliminating one of the contours, he obtains a symbolic representation that is unique up to circular permutation.

In the same paper, Hadamard proves that in each homotopy class of paths on S with fixed endpoints, there is a unique geodesic.¹ He also obtains an analogous result for the free homotopy classes of closed curves on S that are neither homotopic to a point nor to the core curve of a flared infinite sheet.

Hadamard then makes the fundamental observation that the distance function from a fixed point in S to a point that moves along a global geodesic in this surface is convex. This observation is at the basis of the definition of nonpositive curvature in the sense

¹Here, the word geodesic is used, as in Riemannian geometry, in the sense of local geodesic. We warn the reader that this is not the definition that we are adopting in the rest of these notes, where a geodesic is a distance-minimizing map between its endpoints (and therefore it is a *global* geodesic).

of Busemann, which is the main topic of this book.

Next, Hadamard introduces the notion of *asymptotic geodesic rays*, and he constructs such rays as follows. Given a geodesic ray, he considers a sequence of geodesic segments joining a fixed point on the surface to a sequence of points on that ray that tends to infinity. He proves that this sequence of geodesic segments converges to a geodesic ray, which he calls *asymptotic* to the initial ray. He studies the asymptoticity relation and introduces through an analogous construction the notion of (local) geodesics that are asymptotic to a given closed geodesic.

Using an argument that is based on the convexity of the distance function from a point in a funnel to the closed geodesic that connects this funnel to the compact region in S , Hadamard proves that a geodesic that penetrates a funnel cannot get out of it.²

Finally, in the case where all the infinite sheets of S are funnels, Hadamard investigates the distribution of geodesics that stay in the compact part of the surface. These geodesics are the flowlines of the geodesic flow associated with the compact nonpositively curved surface with geodesic boundary, obtained from the surface S by deleting the funnels. For a given point x in S , Hadamard considers the set of initial directions of geodesic rays that start at x and stay in that compact surface. He proves that this is a perfect set with empty interior, whereas the set of directions of geodesic rays that tend to infinity is open.

We already mentioned that the setting of the paper [62] by Hadamard is that of a differentiable surface S embedded in \mathbb{R}^3 , such that at each point of S , the Gaussian curvature is negative except for a finite set of points where it is zero. A theory of *metric spaces with nonpositive curvature*, that is, a theory that does not make any differentiability assumption and whose methods use the distance function alone, without the local coordinates provided by an embedding in Euclidean space or by another Riemannian metric structure, was developed several decades after the paper [62]. It is good to remember, in this respect, that the theory of metric spaces itself was developed long after the theory of spaces equipped with differentiable structures. For instance, Gauss's treatise on the differential geometry of surfaces [51], in which he defines (Gaussian) curvature and proves that this curvature depends only on the intrinsic geometry of the surface and not on its embedding in \mathbb{R}^3 , was published in 1827,³ whereas the axioms for metric spaces were set down by Fréchet, some 90 years later.

Of course, the setting of surfaces embedded in \mathbb{R}^3 has the advantage of providing visual characteristics for the sign of the curvature. For instance, it is well-known that for such a surface, we have the following:

- if the Gaussian curvature at some point is > 0 , then the surface, in the neighborhood of that point, is situated on one side of the tangent plane;
- if the Gaussian curvature at some point is < 0 , then the surface, at that point, crosses its tangent plane.

²Hadamard already obtained a similar result in [60].

³We recall that the surfaces considered by Gauss were always embedded in \mathbb{R}^3 . It is only 30 years after Gauss's paper was written that Riemann introduced the concept of spaces equipped with (Riemannian) metrics with no embedding in \mathbb{R}^3 involved in the definition.

We also recall that if the Gaussian curvature at some point is 0, then any one of the above configurations can occur.

Hadamard indicated in the note [61] how to extend some of the results he proved for surfaces to higher dimensions. The development of these ideas in the general setting of Riemannian manifolds of nonpositive curvature has been carried out by Elie Cartan, in particular in his famous “Leçons sur la géométrie des espaces de Riemann” [36].⁴

The works of Menger and Wald

A few years after the introduction by Fréchet of the axioms for metric spaces, Karl Menger initiated a theory of geodesics in these spaces. A *geodesic* in a metric space is a path whose length is equal to the distance between its endpoints. Menger generalized several results of classical geometry to this new setting and he introduced new methods that did not make any use of local coordinates or of differentials, but only of equalities involving the distance function, and of the triangle inequality. Menger wrote several important papers involving this new notion of geodesic, and he also introduced a notion of “discrete geodesic”, which is based on his definition of *betweenness*: a point z in a metric space is said to lie between two distinct points x and y if z is distinct from x and from y and if we have

$$d(x, y) = d(x, z) + d(z, y).$$

In a complete metric space, the existence of a geodesic joining any two points is equivalent to the existence, for any distinct points, of a point that lies between them. We refer the reader to the commented edition of Menger’s papers in the volume [108], published on the one hundredth anniversary of Menger’s birth. We shall have several opportunities to mention Menger’s work in the following chapters, but here, we mention an important notion that he introduced, which we shall not consider further in this book. This notion contains an idea that is at the basis of the various definitions of curvature that make sense in general metric spaces. The idea is to construct “comparison configurations” for sets of points (say of finite cardinality) in a given metric space X . The comparison configurations are built in a model space, which is generally one of the complete simply connected surfaces M_κ of constant Gaussian curvature κ , that is, either the Euclidean plane (of curvature $\kappa = 0$), or a sphere of curvature $\kappa > 0$, or a hyperbolic plane of curvature $\kappa < 0$. The comparison configuration for a given subset $F \subset X$ is a subset F^* of M_κ , equipped with a map from F to F^* , which is generally taken to be distance-preserving and which is called a “comparison map”. Then, one can define notions like curvature at a given point x in X by requiring the *ad hoc* property for the comparison configurations associated to certain classes of subsets contained in a neighborhood of x . Of course, a comparison

⁴Note that the later editions of Cartan’s book [36] contains additional material, in particular the development of the work of Hadamard to higher dimensions.

configuration does not always exist, but in the case where F is of cardinality 3, one can always construct a comparison configuration F^* of F in any one of the surfaces M_κ with $\kappa \leq 0$. In fact, the axioms for a metric space X are equivalent to the fact that one can construct a comparison configuration in the Euclidean plane for any triple of points in X .⁵ Let us now consider an example.

Given three pairwise distinct points a, b and c in the Euclidean plane M_0 , either they lie on a unique circle (the circumscribed circle), or they lie on a Euclidean straight line. It is useful to consider here such a line as being a circle of radius ∞ , in order to avoid taking subcases. Now for any triple of pairwise distinct points in a metric space X , Menger defined its “curvature” as being equal to $1/R$, where R is the radius of a circle in the Euclidean plane which is circumscribed to a comparison configuration associated to that triple. With this definition, the three points in X are aligned (that is, they satisfy a degenerate triangle inequality) if and only if their curvature is zero. Given an arc (or, say, a one-dimensional object) A contained in X , Menger says that the curvature of A at a point $a \in A$ is equal to κ if for any triple of pairwise distinct points in A that are sufficiently close to a , the curvature of this triple is close to κ .

With this definition, Menger introduced the notion of *curvature* for one-dimensional objects in an arbitrary metric space, and he posed the problem of the definition of curvature for higher-dimensional objects.

Now, we must mention the work of Abraham Wald, a student of Menger, who introduced a notion of two-dimensional curvature in an arbitrary metric space.⁶ The definition again uses a limiting process, but now it involves quadruples of points in that space. The problem is that in general, a quadruple of points in a metric space does not necessarily possess a comparison configuration in the Euclidean plane. However, Wald starts by proving that if the metric space X is a differentiable surface, then, for every point x in X , there exists a real number $\kappa(x)$ satisfying the following property:

(\star) for each $\epsilon > 0$, there exists a neighborhood $V(x)$ of x such that for every quadruple of points Q in $V(x)$, there is an associated real number $\kappa(Q)$ satisfying $|\kappa(x) - \kappa(Q)| < \epsilon$ such that the quadruple of points Q possesses a comparison configuration in the model surface $M_{\kappa(Q)}$.

Wald then proves that the quantity $\kappa(x)$ is equal to the Gaussian curvature of the surface at the given point x .

Now let X be a metric space that is “Menger convex”, that is, a metric space X in which for every pair of distinct points x and y , there exists a point z that lies between them. Suppose furthermore that the Wald two-dimensional curvature exists at each point of X . In other words, suppose that one can associate to each point x in X a real number $\kappa(x)$ satisfying property (\star) that is stated above. Under these assumptions, Wald shows that the space X has the structure of a differentiable surface embedded in \mathbb{R}^3 that induces the same *length structure* as that of the original metric on X . In

⁵We also mention that an early version of the idea of a comparison map is already contained in the very definition of the Gauss map.

⁶Wald gave this 2-dimensional curvature the name “surface curvature”, and Menger refers to it as “Wald curvature”.

other words, the lengths of an arbitrary curve in X , on the one hand measured using the original metric and on the other hand measured using this differentiable surface structure, coincide. Furthermore, at each point a in X , the quantity $\kappa(a)$ is equal to the Gaussian curvature induced by the differentiable embedding of the surface in \mathbb{R}^3 . With this result, Wald solved a problem that had been posed by Menger in [105], which asked for a metric characterization of Gauss surfaces among Menger convex metric spaces. We refer the reader to the Comptes Rendus Note [141] by Wald, presented by Elie Cartan, which describes this work.

We note in passing that for extra-mathematical reasons, Wald stopped working on this subject soon after he published the solution to Menger's problem, and his research interests switched to statistics and econometry. The story is interesting and it is told by Menger in [109]. It seems that there was no direct continuation to Wald's work.

Now, after the notion of curvature in metric spaces, we pass to the notion of nonpositive curvature.

The works of Busemann and Alexandrov

For the development of the theory of nonpositively curved metric spaces, we shall consider works that have been carried out in two different directions: the works of H. Busemann and the works of A. D. Alexandrov and his collaborators. Both Busemann and Alexandrov started their works in the 1940s, and the two approaches gave rise to rich and fruitful developments, with no real interaction between the two. The ramifications of these two theories continue to grow today, especially since the rekindling of interest that was given to nonpositive curvature by M. Gromov in the 1970s.

Let us briefly describe the basic underlying ideas of these works. First we need to recall a few definitions. Consider a metric space X in which each point x possesses a neighborhood U such that any two points in U can be joined by a geodesic path in U . A metric space X with such a property is said to be a *locally geodesic* space. We say that such a neighborhood U is a *geodesically convex* neighborhood of x . A *geodesic segment* $[a, b]$ in X is, by definition, the image of a geodesic path in X joining a and b .⁷ A *triangle* in U is the union of three geodesic segments $[a, b]$, $[a, c]$ and $[b, c]$ contained in U . The segments $[a, b]$, $[a, c]$ and $[b, c]$ are called the *sides* of this triangle.

We start by presenting Busemann's definition of nonpositive curvature, which has the advantage of being the simplest to describe, and on which we shall focus in these notes.

We say that the space X has *nonpositive curvature in the sense of Busemann* if every point x in X possesses a geodesically convex neighborhood U such that for any

⁷From now on, we use the term "geodesic" in the sense of "global geodesic", that is, a path whose length is equal to the distance between its endpoints.

geodesic triangle with sides $[a, b]$, $[a, c]$ and $[b, c]$ contained in U , we have

$$\text{dist}(m, m') \leq (1/2)\text{dist}(b, c),$$

where m and m' are respectively the midpoints of $[a, b]$ and $[a, c]$. This property can be stated in terms of a convexity property of the distance function, defined on the product of any two geodesic segments $[a, b] \times [a, c]$ in X equipped with their natural (barycentric) coordinates. We shall develop this point of view in later chapters.

A nonpositively curved space in the sense of Busemann is sometimes referred to as a “locally convex metric space”, or “local Busemann metric space”. The terminology “nonpositively curved space”, in this sense, is due to Busemann.⁸

A complete Riemannian manifold of nonpositive sectional curvature is an example of a metric space of nonpositive curvature in the sense of Busemann.

The most important writings of Busemann on this subject are certainly the paper [26] and the book [28].

Now we consider the point of view of A. D. Alexandrov.

Before giving the definition of nonpositive curvature in the sense of Alexandrov, we must recall the notion of angle which two geodesic segments (or more generally two paths) in a metric space, that start at a common point, make at that point.

The notion of angle in a metric space has been introduced by Alexandrov as a generalization of the notion of angle in a surface. The first papers by A. D. Alexandrov deal with the intrinsic geometry of surfaces, and the notion of angle is certainly the most important notion in that theory, after the notion of distance and that of length of a path.⁹ In fact, Alexandrov introduced several notions of angle, and these notions coincide provided some reasonable hypotheses on the ambient metric spaces are satisfied. For our needs, it suffices to consider the notion of *upper angle* that two geodesic segments with a common initial point make at that point.

Let X be a locally geodesic metric space, let U be a geodesically convex open subset of X and let us consider a geodesic triangle Δ in U , with sides $[a, b]$, $[a, c]$ and $[b, c]$. Let us define the *upper angle* α of the triangle Δ at the vertex a . For every point x on the segment $[a, b]$ and for every point y on the segment $[a, c]$, we consider a triangle $\Delta_{x,y}$ in X with sides $[a, x]$, $[a, y]$ and $[x, y]$, where $[a, x]$ and $[a, y]$ are subsegments of $[a, b]$ and $[a, c]$ respectively and where $[x, y]$ is a geodesic segment joining the points x and y . Let $\Delta_{x,y}^*$ be an associated comparison triangle in the Euclidean plane and let $\alpha_{x,y}$ be the angle of $\Delta_{x,y}^*$ at the vertex that corresponds to the vertex a of $\Delta_{x,y}$. Then, the *upper angle* of the triangle Δ at the vertex a is defined as

$$\alpha = \limsup_{x,y \rightarrow a} \alpha_{x,y}.$$

⁸Busemann has some additional hypotheses on the metric spaces that he considers. One such hypothesis is the uniqueness of the prolongation of geodesics. There are interesting examples of spaces that do not satisfy this hypothesis and for which the results that we are interested in are valid, and for that reason we have tried to avoid this hypothesis in this book. We recall the precise definition of the spaces considered by Busemann (which he calls G -spaces) in the Notes on Chapter 2 below.

⁹One of the basic papers of Alexandrov on that theory is [2]. We also refer the reader to [3] and [4].