


Progress in Mathematics

Edited by
J. Coates and
S. Helgason



Victor G. Kac

Infinite Dimensional Lie Algebras



Birkhäuser

Victor G. Kac

Infinite Dimensional Lie Algebras

An Introduction

1983

Birkhäuser
Boston • Basel • Stuttgart

Author:

Victor G. Kac
Mathematics Department
Massachusetts Institute of Technology
77 Massachusetts Avenue
Cambridge, MA 02139

Library of Congress Cataloging in Publication Data

Kac, Victor G., 1943—

Infinite dimensional Lie algebras.

(Progress in mathematics ; vol. 44)

Bibliography: p.

Includes index.

1. Lie algebras. I. Title. II. Series: Progress in mathematics (Boston, Mass.) ; vol. 44.

QA252.3.K33 1983 512'.55 83-25809

ISBN 0-8176-3118-6

CIP-Kurztitelaufnahme der Deutschen Bibliothek

Kac, Victor G.:

Infinite dimensional Lie algebras / an introd. /

Victor G. Kac. - Basel ; Boston ; Stuttgart :

Birkhäuser, 1984.

(Progress in mathematics ; Vol. 44)

ISBN 3-7643-3118-6

sfr 62.00

NE: GT

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the copyright owner.

© Birkhäuser Boston, Inc., 1983

ISBN 0-8176-3118-6

ISBN 3-7643-3118-6

Printed in USA

9 8 7 6 5 4 3 2 1

Progress in Mathematics
Vol. 44

Edited by
J. Coates and
S. Helgason

Birkhäuser
Boston · Basel · Stuttgart

Dedicated to my teacher,
Ernest Borisovich Vinberg
with gratitude and admiration

Grau, teurer Freund, ist alle Theorie,
Und grün des Lebens goldner Baum.
J. W. Goethe, *Faust I*

И я выхожу из пространства
В запущенный сад величин
O. Mandelstam

Introduction.

0.1. The creators of the Lie theory viewed a Lie group as a group of symmetries of an algebraic or a geometric object; the corresponding Lie algebra, from their point of view, was the set of infinitesimal transformations. Since the group of symmetries of the object is not necessarily finite-dimensional, S. Lie considered not only the problem of classification of subgroups of GL_n , but also the problem of classification of infinite-dimensional groups of transformations.

The problem of classification of simple finite-dimensional Lie algebras over the field of complex numbers was solved by the end of the 19th century by W. Killing and E. Cartan. (A vivid description of the history of this discovery, one of the most remarkable in all of mathematics, can be found in Hawkins [1982].) And just over a decade later, Cartan classified simple infinite-dimensional Lie algebras of vector fields on a finite-dimensional space.

Starting with the works of Lie, Killing, and Cartan, the theory of finite-dimensional Lie groups and Lie algebras has developed systematically in depth and scope. On the other hand, Cartan's works on simple infinite-dimensional Lie algebras had been virtually forgotten until the mid-sixties. A resurgence of interest in this area began with the work of Guillemin-Sternberg [1964] and Singer-Sternberg [1965], which developed an adequate algebraic language and the machinery of filtered and graded Lie algebras. They were, however, unable to find an algebraic proof of Cartan's classification theorem (see Guillemin-Quillen-Sternberg [1966]). This was done by Weisfeiler [1968], who reduced the problem to the classification of simple \mathbb{Z} -graded Lie algebras of finite "depth" $\mathfrak{g} = \bigoplus_{j \geq -d} \mathfrak{g}_j$, where $\dim \mathfrak{g}_j < \infty$ and the \mathfrak{g}_0 -module \mathfrak{g}_{-1} is irreducible.

0.2. At the present time there is no general theory of infinite-dimensional Lie groups and algebras and their representations. There are, however, four classes of infinite-dimensional Lie groups and algebras that underwent a more or less intensive study. These are, first of all, the above-mentioned Lie algebras of vector fields and the corresponding groups of diffeomorphisms of a manifold. Starting with the works of Gelfand-Fuchs [1969], [1970A,B], there emerged an important direction having many geometric applications, which is the cohomology theory of infinite-dimensional Lie algebras of vector fields on a finite-dimensional manifold. There is also a rather large number of works which study and classify various

classes of representations of the groups of diffeomorphisms of a manifold. One should probably include in the first class the groups of biregular automorphisms of an algebraic variety (see Shafarevich [1982]).

The second class consists of Lie groups (resp. Lie algebras) of smooth mappings of a given manifold into a finite-dimensional Lie group (resp. Lie algebra). In other words, this is a group (resp. Lie algebra) of matrices over some function algebra but viewed over the base field. (The physicists refer to certain central extensions of these Lie algebras as current algebras.) The main subject of study in this case has been certain special families of representations.

The third class consists of the classical Lie groups and algebras of operators in a Hilbert or Banach space. There is a rather large number of scattered results in this area, which study the structure of these Lie groups and algebras and their representations. A representation which plays an important role in quantum field theory is the Segal-Shale-Weil (or metaplectic) representation of an infinite-dimensional symplectic group.

I shall not discuss in this book the three classes of infinite-dimensional Lie algebras listed above, with the exception of those closely related to the Lie algebras of the fourth class, which we consider below. The reader interested in these three classes should consult the literature cited at the end of the book.

Finally, the fourth class of infinite-dimensional Lie algebras is the class of the so-called Kac-Moody algebras, the subject of the present book.

0.3. Let us briefly discuss the main concepts of the structural theory of Kac-Moody algebras. Let $A = (a_{ij})_{i,j=1}^n$ be a *generalized Cartan matrix*, i.e. an integral $n \times n$ matrix such that $a_{ii} = 2$, $a_{ij} \leq 0$, and $a_{ij} = 0$ implies $a_{ji} = 0$. The associated *Kac-Moody algebra* $\mathfrak{g}(A)$ is a complex Lie algebra on $3n$ generators e_i, f_i, h_i ($i = 1, \dots, n$) and the following defining relations ($i, j = 1, \dots, n$):

$$(0.3.1) \quad \begin{cases} [h_i, h_j] = 0, & [e_i, f_i] = h_i, & [e_i, f_j] = 0 \text{ if } i \neq j, \\ [h_i, e_j] = a_{ij}e_j, & [h_i, f_j] = -a_{ij}f_j, \\ (\text{ad } e_i)^{1-a_{ij}}e_j = 0, & (\text{ad } f_i)^{1-a_{ij}}f_j = 0 \text{ if } i \neq j. \end{cases}$$

(The definition given in the main text of the book (see Chapter 1) is different from the above; it is more convenient for a number of reasons. The proof of the fact that the derived algebra of the Lie algebra $\mathfrak{g}(A)$ defined in Chapter 1 coincides with the Lie algebra $\mathfrak{g}'(A)$ defined by relations (0.3.1) has been obtained quite recently; this proof appears in Chapter 9.)

I came to consider these Lie algebras while trying to understand and generalize the works of Guillemin-Quillen-Singer-Sternberg-Weisfeiler on Cartan's classification. The key idea was to consider arbitrary simple \mathbb{Z} -graded Lie algebras $\mathfrak{g} = \bigoplus_j \mathfrak{g}_j$; but since there are too many such Lie algebras, the point was to require the dimension of \mathfrak{g}_j to grow no faster than some polynomial in j . (One can show that Lie algebras of finite depth do satisfy this condition, and that this condition does not depend on the gradation.) Such Lie algebras were classified under some

technical hypotheses (see Kac [1968 B]). It turned out that in addition to Cartan's four series of Lie algebras of polynomial vector fields, there is another class of infinite-dimensional Lie algebras of polynomial growth, which are called affine Lie algebras (more precisely, they are the quotients of affine Lie algebras by the 1-dimensional center). At the same time, Moody [1968] independently undertook the study of the Lie algebras $\mathfrak{g}'(A)$.

The class of Kac-Moody algebras breaks up into three subclasses. To describe them, it is convenient to assume that the matrix A is *indecomposable* (i.e. there is no partition of the set $\{1, \dots, n\}$ into two non-empty subsets so that $a_{ij} = 0$ whenever i belongs to the first subset, while j belongs to the second; this is done without loss of generality since the direct sum of matrices corresponds to the direct sum of Kac-Moody algebras). Then there are the following three mutually exclusive possibilities:

- a) There is a vector θ of positive integers such that all the coordinates of the vector $A\theta$ are positive. In such case all the principal minors of the matrix A are positive and the Lie algebra $\mathfrak{g}'(A)$ is finite-dimensional.
- b) There is a vector δ of positive integers such that $A\delta = 0$. In such case all the principal minors of the matrix A are non-negative and $\det A = 0$; the algebra $\mathfrak{g}'(A)$ is infinite-dimensional, but is of polynomial growth (moreover, it admits a \mathbb{Z} -gradation by subspaces of uniformly bounded dimension). The Lie algebras of this subclass are called *affine Lie algebras*.
- c) There is a vector α of positive integers such that all the coordinates of the vector $A\alpha$ are negative. In such case the Lie algebra $\mathfrak{g}'(A)$ is of exponential growth.

The main achievement of the Killing-Cartan theory may be formulated as follows: a simple finite-dimensional complex Lie algebra is isomorphic to one of the Lie algebras of the subclass a). (Note that the classification of matrices of type a) and b) is a rather simple problem.) The existence of the generators satisfying relations (0.3.1) was pointed out by Chevalley [1948]. (Much later Serre [1966] showed that these are defining relations.)

It turned out that most of the classical concepts of the Killing-Cartan-Weyl theory can be carried over to the entire class of Kac-Moody algebras, such as the Cartan subalgebra, the root system, the Weyl group, etc. In doing so one discovers a series of new phenomena, which the book treats in detail (see Chapters 1–6). I shall only point out here that $\mathfrak{g}'(A)$ does not always possess a nonzero invariant bilinear form. This is the case if and only if the matrix A is *symmetrizable*, i.e. the matrix DA is symmetric for some non-degenerate diagonal matrix D (see Chapter 2).

0.4. It is an important property of affine Lie algebras that they possess a simple realization (see Chapters 7 and 8). Here I shall explain this realization for the example of the Kac-Moody algebra associated to the extended Cartan matrix A of a simple finite-dimensional complex Lie algebra \mathfrak{g} . (All such matrices are “affine” generalized Cartan matrices; the corresponding algebra $\mathfrak{g}'(A)$ is called a non-twisted affine Lie algebra.) Namely, the affine Lie algebra $\mathfrak{g}'(A)$ is a central extension by the

1-dimensional center of the Lie algebra of polynomial maps of the circle into the simple finite-dimensional complex Lie algebra \mathfrak{g} (so that it is the simplest example of a Lie algebra of the second class mentioned in 0.2).

More precisely, let us consider the Lie algebra \mathfrak{g} in some faithful finite-dimensional representation. Then the Lie algebra $\mathfrak{g}'(A)$ is isomorphic to the Lie algebra on the complex space $(\mathbb{C}[t, t^{-1}] \otimes_{\mathbb{C}} \mathfrak{g}) \oplus \mathbb{C}c$ with the bracket

$$[a(t) \oplus \lambda c, b(t) \oplus \mu c] = (a(t)b(t) - b(t)a(t)) \oplus \left(\text{Res}_0 \text{tr} \frac{da(t)}{dt} b(t) \right) c,$$

so that $\mathbb{C}c$ is the (1-dimensional) center. This realization allows us to study affine Lie algebras from another point of view. In particular, the algebra of vector fields on the circle (the simplest algebra of the first class) plays an important role in the theory of affine Lie algebras.

Note also that the Lie algebras of the fourth class are closely related to the affine Lie algebras of infinite rank, considered in Chapters 7 and 14.

Unfortunately, no simple realization has been found up to now for any non-affine infinite-dimensional Kac-Moody algebra. This question appears to be one of the most important open problems of the theory.

0.5. An important concept missing from the first works in Kac-Moody algebras was the concept of an integrable highest weight representation (introduced in Kac [1974]). Given a sequence of non-negative integers $\Lambda = (\lambda_1, \dots, \lambda_n)$, the *integrable highest weight representation* of a Kac-Moody algebra $\mathfrak{g}'(A)$ is an irreducible representation π_Λ of $\mathfrak{g}'(A)$ on a complex vector space $L(\Lambda)$, which is determined by the property that there is a non-zero vector $v_\Lambda \in L(\Lambda)$ such that

$$\pi_\Lambda(e_i)v_\Lambda = 0 \text{ and } \pi_\Lambda(h_i)v_\Lambda = \lambda_i v_\Lambda \quad (i = 1, \dots, n).$$

(This terminology is explained by the fact that Λ is called the highest weight, and the conditions on Λ are necessary and sufficient for being able to integrate π_Λ and obtain a representation of the group.)

Cartan's theorem on the highest weight asserts that all the representations π_Λ of a complex simple finite-dimensional Lie algebra are finite-dimensional, and that every finite-dimensional irreducible representation is equivalent to one of the π_Λ .

That the representations π_Λ are finite-dimensional (the most nontrivial part of Cartan's theorem) was proved by Cartan by examining the cases, one by one. A purely algebraic proof was found much later by C. Chevalley [1948] (a "transcendental" proof had been found earlier by H. Weyl). This brief note by Chevalley appears in retrospect as the precursor of the algebraization of the representation theory of Lie groups. This note also contains, in an embryonic form, many of the basic concepts of the theory of Kac-Moody algebras.

The algebraization of the representation theory of Lie groups, which has undergone such an explosive development during the last decade, started with the work Bernstein-Gelfand-Gelfand [1971] on Verma modules (the first nontrivial results about these modules were obtained by Verma [1968]). In particular, using the

Verma modules, Bernstein-Gelfand-Gelfand gave a transparent algebraic proof of Weyl's formula for the characters of finite-dimensional irreducible representations of finite-dimensional simple Lie algebras.

At about the same time Macdonald [1972] obtained his remarkable identities. In this work he undertook to generalize the Weyl denominator identity to the case of affine root systems. He remarked that a straightforward generalization is actually false. To salvage the situation he had to add some "mysterious" factors, which he was able to determine as a result of lengthy calculations. The simplest example of Macdonald's identities is the famous Jacobi triple product identity:

$$\prod_{n \geq 1} (1 - u^n v^n)(1 - u^{n+1} v^n)(1 - u^n v^{n+1}) = \sum_{n \in \mathbb{Z}} (-1)^n u^{\frac{1}{2}n(n+1)} v^{\frac{1}{2}n(n-1)}.$$

The "mysterious" factors which do not correspond to affine roots are the factors $(1 - u^n v^n)$.

After the appearance of the two works mentioned above very little remained to be done: one had to place them on the desk next to one another to understand that Macdonald's result is only the tip of the iceberg—the representation theory of Kac-Moody algebras. Namely, it turned out that a simplified version of Bernstein-Gelfand-Gelfand's proof may be applied to the proof of a formula generalizing Weyl's formula, for the formal character of the representation π_A of an arbitrary Kac-Moody algebra $\mathfrak{g}(A)$ corresponding to a symmetrizable matrix A . In the case of the simplest 1-dimensional representation π_0 , this formula becomes the generalization of Weyl's denominator identity. In the case of an affine Lie algebra, the generalized Weyl denominator identity turns out to be equivalent to the Macdonald identities. In the process, the "mysterious" factors receive a simple interpretation: they correspond to the so-called imaginary roots (i.e. roots that one should add to the affine roots to obtain all the roots of the affine Lie algebra). Note that the simplest example of the Jacobi triple product identity turns out to be just the generalized denominator identity for the affine Lie algebra corresponding to the matrix $\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$.

The exposition of these results (obtained in the note Kac [1974]) may be found in Chapter 10. Chapters 9–14 are devoted to the general theory of highest weight representations and their applications.

The main tool of the theory of representations with highest weight is the generalized Casimir operator (see Chapter 2). Unfortunately, the construction of this operator depends on whether the matrix A is symmetrizable. The question whether one can lift the hypothesis of symmetrizability of the matrix A remains open. Another, more important question is the following: what is an integrable highest weight representation of the Lie algebra of maps of an arbitrary manifold into a simple finite-dimensional Lie algebra?

Once the integrable highest weight representations had been introduced, the theory of Kac-Moody algebras got off the ground and has been developing since at an accelerating speed. In the past decade this theory has emerged as a field that has close connections with many areas of mathematics and mathematical physics,

such as invariant theory, combinatorics, topology, the theory of modular forms and theta functions, the theory of singularities, Hamiltonian mechanics, and quantum field theory.

This book contains a detailed exposition of the foundations of the theory of Kac-Moody algebras and their integrable representations. Besides the application to the Macdonald identities mentioned above (Chapter 12), the book discusses the application to the classification of finite order automorphisms of simple finite-dimensional Lie algebras (Chapter 8), and the connection with the theory of modular forms and theta functions (Chapter 13). The last chapter discusses the remarkable connection between the representation theory of affine Lie algebras and the Korteweg-de Vries type equations, discovered recently by the Kyoto school.

A theory of Lie algebras is usually interesting, insofar as it is related to group theory, and Kac-Moody algebras are no exception. Recently there appeared a series of deep results on groups associated with Kac-Moody algebras. A discussion of these results would require writing another book. I chose to make only a few comments regarding this subject at the end of some chapters.

0.7. Throughout the book the base field is the field of complex numbers. However, all the results of the book, except, of course, for the ones concerning Hermitian forms and convergence problems, can be extended without difficulty to the case of an arbitrary field of characteristic zero.

0.8. Motivations are provided at the beginning of each chapter, which ends with a discussion of related results. The main text of each chapter is followed by exercises (whose total number exceeds 250). Some of them are elementary, others constitute a brief exposition of original works. I hope that these expositions are sufficiently detailed for the diligent reader to reconstruct all the proofs. The square brackets at the end of some exercises contain hints for their solution.

The exposition in the book is practically self-contained. Although I had in mind a reader familiar with the theory of finite-dimensional semisimple Lie algebras, what would suffice for the most part is a knowledge of the elements of Lie algebras, their enveloping algebras and representations. For example, the book of Humphreys [1972] is more than sufficient.

One finds a rather extensive bibliography at the end of the book. I hope that the collection of references to mathematical works in the theory of Kac-Moody algebras is at least everywhere dense. This is not at all so in the case of the works in physics. The choice of references in this case was rather arbitrary and often depended on whether I had a copy of the paper or discussed it with the author. The same should be said as regards the references to the works on the other classes of infinite-dimensional Lie algebras.

0.9. This book is based on lectures given at MIT in 1978, 1980, and 1982, and at the Collège de France in 1981. I would like to thank those who attended for helpful comments and corrections of the notes, in particular F. Arnold, R. Coley, R. Gross, Z. Haddad, M. Haiman, G. Heckman, F. Levstein, A. Rocha, and T. Vongiouklis.

I am grateful to M. Duflo, G. Heckman, B. Kupershmidt and B. Weisfeiler for reading some parts of the manuscript and pointing out errors. I apologize for those errors that remain. My thanks go to F. Rose, B. Katz and M. Katz without whose help and support this book would never have come out. I also owe thanks to K. Manning and C. Macpherson for help with the language. Finally I would like on this occasion to express my deep gratitude to D. Peterson, whose collaboration had a great influence not only on this book, but also on most of my mathematical work in the past few years.

The author was supported in part by a Sloan foundation grant and by grants from the National Science Foundation.

\mathbb{Z}	Notational conventions the set of integers
\mathbb{Z}_+	the set of non-negative integers
\mathbb{Q}	the set of rational numbers
\mathbb{R}	the set of real numbers
\mathbb{R}_+	the set of non-negative real numbers
\mathbb{C}	the set of complex numbers
\mathbb{C}^\times	the set of non-zero complex numbers
$ z $	modulus of $z \in \mathbb{C}$
$\operatorname{Re} z$ and $\operatorname{Im} z$	real and imaginary parts of $z \in \mathbb{C}$
$\log z$	for $z \in \mathbb{C}^\times$: $e^{\log z} = z$ and $-\pi \leq \operatorname{Im} \log z < \pi$
z^α	$= e^{\alpha \log z}$ for $\alpha \in \mathbb{C}$, $z \in \mathbb{C}^\times$
$U \oplus V$ or $\bigoplus_\alpha U_\alpha$	direct sum of vector spaces
$\sum_\alpha U_\alpha$	sum of subspaces of a vector space
$\prod_\alpha U_\alpha$	direct product of vector spaces
kS	the linear k -span of S ($k = \mathbb{Z}, \mathbb{Z}_+, \mathbb{Q}, \mathbb{R}$, or \mathbb{C})
$U \otimes V$	tensor product of vector k -spaces over k ($k = \mathbb{Q}, \mathbb{R}$, or \mathbb{C})
U^*	the dual of a vector k -space over k ($k = \mathbb{Q}, \mathbb{R}$, or \mathbb{C})
\mathbb{C}^n	direct sum of n copies of the vector space \mathbb{C} ($n \in \mathbb{Z}_+ \cup \{\infty\}$)
I_V or I_n or I	the identity operator on the n -dimensional vector space V
$\langle \cdot, \cdot \rangle$	pairing between a vector space and its dual
$ u ^2 = (u u)$	square length of a vector u
$ S $	cardinality of a set S
$P \bmod Q$	a set of representatives of cosets of an abelian group P with respect to a subgroup Q
$a \equiv b \bmod C$	means that $a - b \in C$.
$U(\mathfrak{g})$	universal enveloping algebra of a Lie algebra \mathfrak{g}
$U_0(\mathfrak{g}) = U(\mathfrak{g})\mathfrak{g}$	the augmentation ideal of $U(\mathfrak{g})$
$g(v)$ or $g \cdot v$	action of an element g of a Lie algebra or a group on an element v of a module; all modules are assumed to be left modules unless otherwise specified
$G \cdot v$ or $G(v)$	$= \{g \cdot v \mid g \in G\}$ the orbit of v under the action of a group G
$G \cdot V$ or $G(V)$	union of orbits of elements from a set V
$\mathfrak{g} \cdot V$ or $\mathfrak{g}(V)$	$= \{g(v) \mid g \in \mathfrak{g}, v \in V\}$.
UV	linear span of the set $\{uv \mid u \in U, v \in V\}$, where U and V are subspaces of an algebra

Progress in Mathematics

Edited by J. Coates and S. Helgason

Progress in Physics

Edited by A. Jaffe and D. Ruelle

- A collection of research-oriented monographs, reports, notes arising from lectures or seminars
- Quickly published concurrent with research
- Easily accessible through international distribution facilities
- Reasonably priced
- Reporting research developments combining original results with an expository treatment of the particular subject area
- A contribution to the international scientific community: for colleagues and for graduate students who are seeking current information and directions in their graduate and post-graduate work.

Manuscripts

Manuscripts should be no less than 100 and preferably no more than 500 pages in length.

They are reproduced by a photographic process and therefore must be typed with extreme care. Symbols not on the typewriter should be inserted by hand in indelible black ink. Corrections to the typescript should be made by pasting in the new text or painting out errors with white correction fluid.

The typescript is reduced slightly (75%) in size during reproduction; best results will not be obtained unless the text on any one page is kept within the overall limit of $6 \times 9\frac{1}{2}$ in (16×24 cm). On request, the publisher will supply special paper with the typing area outlined.

Manuscripts should be sent to the editors or directly to:
Birkhäuser Boston, Inc., P.O. Box 2007, Cambridge,
Massachusetts 02139

PROGRESS IN MATHEMATICS

Already published

- PM 1 Quadratic Forms in Infinite-Dimensional Vector Spaces
Herbert Gross
ISBN 3-7643-1111-8, 432 pages, paperback
- PM 2 Singularités des systèmes différentiels de Gauss-Manin
Frédéric Pham
ISBN 3-7643-3002-3, 346 pages, paperback
- PM 3 Vector Bundles on Complex Projective Spaces
C. Okonek, M. Schneider, H. Spindler
ISBN 3-7643-3000-7, 396 pages, paperback
- PM 4 Complex Approximation, Proceedings, Quebec, Canada,
July 3-8, 1978
Edited by Bernard Aupetit
ISBN 3-7643-3004-X, 128 pages, paperback
- PM 5 The Radon Transform
Sigurdur Helgason
ISBN 3-7643-3006-6, 207 pages, hardcover
- PM 6 The Weil Representation, Maslov Index and Theta Series
Gérard Lion, Michèle Vergne
ISBN 3-7643-3007-4, 348 pages, paperback
- PM 7 Vector Bundles and Differential Equations
Proceedings, Nice, France, June 12-17, 1979
Edited by André Hirschowitz
ISBN 3-7643-3022-8, 256 pages, paperback
- PM 8 Dynamical Systems, C.I.M.E. Lectures, Bressanone, Italy,
June 1978
John Guckenheimer, Jürgen Moser, Sheldon E. Newhouse
ISBN 3-7643-3024-4, 305 pages, hardcover
- PM 9 Linear Algebraic Groups
T. A. Springer
ISBN 3-7643-3029-5, 314 pages, hardcover

- PM 10 Ergodic Theory and Dynamical Systems I
A. Katok
ISBN 3-7643-3036-8, 346 pages, hardcover
- PM 11 18th Scandinavian Congress of Mathematicians, Aarhus,
Denmark, 1980
Edited by Erik Balslev
ISBN 3-7643-3040-6, 526 pages, hardcover
- PM 12 Séminaire de Théorie des Nombres, Paris 1979-80
Edited by Marie-José Bertin
ISBN 3-7643-3035-X, 404 pages, hardcover
- PM 13 Topics in Harmonic Analysis on Homogeneous Spaces
Sigurdur Helgason
ISBN 3-7643-3051-1, 152 pages, hardcover
- PM 14 Manifolds and Lie Groups, Papers in Honor of Yozô Matsushima
*Edited by J. Hano, A. Marimoto, S. Murakami, K. Okamoto,
and H. Ozeki*
ISBN 3-7643-3053-8, 476 pages, hardcover
- PM 15 Representations of Real Reductive Lie Groups
David A. Vogan, Jr.
ISBN 3-7643-3037-6, 776 pages, hardcover
- PM 16 Rational Homotopy Theory and Differential Forms
Phillip A. Griffiths, John W. Morgan
ISBN 3-7643-3041-4, 258 pages, hardcover
- PM 17 Triangular Products of Group Representations and
their Applications
S. M. Vovsi
ISBN 3-7643-3062-7, 142 pages, hardcover
- PM 18 Géométrie Analytique Rigide et Applications
Jean Fresnel, Marius van der Put
ISBN 3-7643-3069-4, 232 pages, hardcover
- PM 19 Periods of Hilbert Modular Surfaces
Takayuki Oda
ISBN 3-7643-3084-8, 144 pages, hardcover
- PM 20 Arithmetic on Modular Curves
Glenn Stevens
ISBN 3-7643-3088-0, 236 pages, hardcover

- PM 21 Ergodic Theory and Dynamical Systems II
A. Katok, editor
ISBN 3-7643-3096-1, 226 pages, hardcover
- PM 22 Séminaire de Théorie des Nombres, Paris 1980-81
Marie-José Bertin, editor
ISBN 3-7643-3066-X, 374 pages, hardcover
- PM 23 Adeles and Algebraic Groups
A. Weil
ISBN 3-7643-3092-9, 138 pages, hardcover
- PM 24 Enumerative Geometry and Classical Algebraic Geometry
Patrick Le Barz, Yves Hervier, editors
ISBN 3-7643-3106-2, 260 pages, hardcover
- PM 25 Exterior Differential Systems and the Calculus of Variations
Phillip A. Griffiths
ISBN 3-7643-3103-8, 349 pages, hardcover
- PM 26 Number Theory Related to Fermat's Last Theorem
Neal Koblitz, editor
ISBN 3-7643-3104-6, 376 pages, hardcover
- PM 27 Differential Geometric Control Theory
Roger W. Brockett, Richard S. Millman, Hector J. Sussmann, editors
ISBN 3-7643-3091-0, 349 pages, hardcover
- PM 28 Tata Lectures on Theta I
David Mumford
ISBN 3-7643-3109-7, 254 pages, hardcover
- PM 29 Birational Geometry of Degenerations
Robert Friedman and David R. Morrison, editors
ISBN 3-7643-3111-9, 410 pages, hardcover
- PM 30 CR Submanifolds of Kaehlerian and Sasakian Manifolds
Kentaro Yano, Masahiro Kon
ISBN 3-7643-3119-4, 223 pages, hardcover
- PM 31 Approximations Diophantiennes et Nombres Transcendants
D. Bertrand and M. Waldschmidt, editors
ISBN 3-7643-3120-8, 349 pages, hardcover