



# Pharmacogenetics and Individualized Therapy

Edited by  
*Anke-Hilse Maitland-van der Zee*  
and *Ann K. Daly*



---

# PHARMACOGENETICS AND INDIVIDUALIZED THERAPY

---

Edited by

Anke-Hilse Maitland-van der Zee

Utrecht University

Utrecht, The Netherlands

Ann K. Daly

Newcastle University,

Newcastle upon Tyne, UK



 **WILEY**

A JOHN WILEY & SONS, INC., PUBLICATION



Copyright © 2012 by John Wiley & Sons, Inc. All rights reserved.

Published by John Wiley & Sons, Inc., Hoboken, New Jersey  
Published simultaneously in Canada

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400, fax 978-750-4470, or on the web at [www.copyright.com](http://www.copyright.com). Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, 201-748-6011, fax 201-748-6008, or online at <http://www.wiley.com/go/permission>.

**Limit of Liability/Disclaimer of Warranty:** While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at 877-762-2974, outside the United States at 317-572-3993 or fax 317-572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at [www.wiley.com](http://www.wiley.com).

*Library of Congress Cataloging-in-Publication Data:*

Pharmacogenetics and individualized therapy / edited by Anke-Hilse Mailand-van der Zee, Ann K. Daly.  
p. ; cm.

Includes bibliographical references and index.

ISBN 978-0-470-43354-6 (cloth)

1. Pharmacogenetics. I. Maitland-van der Zee, Anke-Hilse. II. Daly, Ann K.

[DNLM: 1. Pharmacogenetics—methods. 2. Drug Therapy—methods. 3. Individualized Medicine—methods. QV 38]

RM301.3.G45P396 2011

615'.19—dc23

2011015209

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

## PREFACE

---

Pharmacogenetics and individualized therapy is a rapidly evolving field that is likely to have important consequences for clinical practice in the coming decades. This book is aimed at a general audience including advanced undergraduate and graduate students in medicine, pharmacy, pharmacology, and other related disciplines as well as researchers based in either academia or the pharmaceutical industry. Some familiarity with basic pharmacology and genetics is assumed.

This book is organized in five parts. Part I describes the basic principles of pharmacogenetics including factors relevant to drug disposition (phase I and phase II metabolizing enzymes, and drug transporters) and the role of pharmacodynamics (drug targets).

Part II includes discussions of state-of-the art pharmacogenetics in many important therapeutic areas [cardiovascular, psychiatry, cancer, asthma/chronic obstructive pulmonary disease (COPD), adverse drug reactions, transplantation, inflammatory bowel disease, pain medication].

Part III describes ethical and related issues in implementing pharmacogenetics into clinical practice.

In Part IV important developments in the technology supporting pharmacogenetics research are discussed. More recent developments in genotyping techniques provide opportunities for genotyping over 1 million single-nucleotide polymorphisms in many patients at affordable prices. Further developments in analysis techniques provide investigators with the opportunity to consider gene–environment and epistatic interactions as well as the possibility of whole-genome sequencing.

Part V discusses the impact of pharmacogenetics in the pharmaceutical industry and also the role that pharmacogenetics currently plays in the registration process.

It has been a privilege to interact with the distinguished expert authors who have provided chapters for this book, and we would like to express our sincere gratitude to them for their excellent contributions. We also wish to thank Lisa Gilhuijs-Pederson, PhD for assistance in editing this book.

ANN K. DALY, PhD

ANKE-HILSE MAITLAND-VAN DER ZEE, PharmD PhD

## CONTRIBUTORS

---

**Katherine J. Aitchison**, Institute of Psychiatry, King's College London, London, UK

**Martin Armstrong**, Clinical Development and Medical Affairs, Shire AG, Geneva, Switzerland

**Maria Arranz**, Institute of Psychiatry, King's College London, London, UK

**Anthonius de Boer**, Faculty of Science, Utrecht Institute for Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands

**Daniel K. Burns**, Deane Drug Discovery Institute, Duke University, Durham, North Carolina, USA

**Angel Carracedo**, Galician Foundation of Genomic Medicine (SERGAS), University of Santiago de Compostela, Santiago de Compostela, Spain

**Ingolf Cascorbi**, Institute of Experimental and Clinical Pharmacology, University Hospital Schleswig-Holstein, Kiel, Germany

**Martina Cornel**, Department of Human Genetics/EMGO Institute for Health Care and Research, VU University Medical Center, Amsterdam, The Netherlands and Center for Society and Genomics, Radboud University, Nijmegen, The Netherlands

**Sarah Curran**, Institute of Psychiatry, King's College London, London, UK

**Ann K. Daly**, Institute of Cellular Medicine, Newcastle University Medical School, Newcastle upon Tyne, UK

**Vita Dolžan**, Institute of Biochemistry, Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia

**Markus Grube**, Department of Pharmacology, Ernst Moritz Arndt-University of Greifswald, Greifswald, Germany

**Shiew-Mei Huang**, Office of Clinical Pharmacology, Center for Drug Evaluation and Research, Food and Drug Administration, Silver Spring, Maryland, USA

**Susanne Karner**, Dr. Margarete Fischer-Bosch Institute of Clinical Pharmacology, Stuttgart, Germany

**Myong-Jin Kim**, Office of Clinical Pharmacology, Center for Drug Evaluation and Research, Food and Drug Administration, Silver Spring, Maryland, USA

**Olaf H. Klungel**, Faculty of Science, Utrecht Institute of Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands

**Gerard H. Koppelman**, Pediatric Pulmonology and Pediatric Allergology, Beatrix Childrens Hospital, University Medical Center Groningen, Groningen, The Netherlands

**Ellen S. Koster**, Faculty of Science, Utrecht Institute of Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands

**Heyo K. Kroemer**, Department of Pharmacology, Ernst Moritz Arndt-University of Greifswald, Greifswald, Germany

**Lawrence J. Lesko**, Office of Clinical Pharmacology, Center for Drug Evaluation and Research, Food and Drug Administration, Silver Spring, Maryland, USA

**Jörn Lötsch**, Pharmazentrum Frankfurt and Department of Clinical Pharmacology, Johann Wolfgang Goethe-University Hospital, Frankfurt, Germany

**Anke-Hilse Maitland-van der Zee**, Faculty of Science, Utrecht Institute of Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands

**Sharon Marsh**, Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta, Edmonton, Alberta, Canada

**Henriette E. Meyer zu Schwabedissen**, Department of Pharmacology, Ernst Moritz Arndt-University of Greifswald, Greifswald, Germany

**Alison A. Motsinger-Reif**, Bioinformatics Research Center, Department of Statistics, North Carolina State University, Raleigh, North Carolina, USA

**Ruth I. Ohlsen**, Institute of Psychiatry, King's College London, London, UK

**Bas J. M. Peters**, Faculty of Science, Utrecht Institute for Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands

**Toine Pieters**, Faculty of Science, Utrecht Institute for Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands; and Department of Medical Humanities, VU University Medical Center, Amsterdam, The Netherlands

**Munir Pirmohamed**, Institute of Translational Medicine, University of Liverpool, Liverpool, UK

**Jan A. M. Raaijmakers**, Faculty of Science, Utrecht Institute of Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands

**Elke Schaeffeler**, Dr. Margarete Fischer-Bosch Institute of Clinical Pharmacology, Stuttgart, Germany

**Tom Schalekamp**, Faculty of Science, Utrecht Institute for Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands

**Matthias Schwab**, Dr. Margarete Fischer-Bosch Institute of Clinical Pharmacology, Stuttgart, Germany and Institute of Experimental and Clinical Pharmacology and Toxicology, University Hospital, Tübingen, Germany

**Beatriz Sobrino**, Galician Foundation of Genomic Medicine (SERGAS), University of Santiago de Compostela, Santiago de Compostela, Spain

**Scott S. Sundseth**, Cabernet Pharmaceuticals, Durham, North Carolina, USA

**Alexander Teml**, Dr. Margarete Fischer-Bosch Institute of Clinical Pharmacology, Stuttgart, Germany

**Evangelia M. Tsapakis**, Institute of Psychiatry, King's College London, London, UK

**Susanne Vijverberg**, Faculty of Science, Utrecht Institute for Pharmaceutical Sciences (UIPS), Division of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, The Netherlands

**Nora S. Vyas**, Institute of Psychiatry, King's College London, London, UK

**Issam Zineh**, Office of Clinical Pharmacology, Center for Drug Evaluation and Research, Food and Drug Administration, Silver Spring, Maryland, USA

## **CONTENTS**

---

<b>PREFACE</b>	<b>vii</b>
<b>CONTRIBUTORS</b>	<b>ix</b>
<b>1. Pharmacogenetics: A Historical Perspective</b> <i>Ann K. Daly</i>	<b>1</b>
<b>PART I PHARMACOGENETICS: RELATIONSHIP TO PHARMACOKINETICS AND PHARMACODYNAMICS</b>	
<b>2. Pharmacogenetics in Drug Metabolism: Role of Phase I Enzymes</b> <i>Vita Dolžan</i>	<b>15</b>
<b>3. Pharmacogenetics of Phase II Drug Metabolizing Enzymes</b> <i>Ingolf Cascorbi</i>	<b>81</b>
<b>4. Pharmacogenetics of Drug Transporters</b> <i>Henriette E. Meyer zu Schwabedissen, Markus Grube, and Heyo K. Kroemer</i>	<b>101</b>
<b>5. Pharmacogenetics of Drug Targets</b> <i>Ann K. Daly and Maria Arranz</i>	<b>149</b>
<b>PART II PHARMACOGENETICS: THERAPEUTIC AREAS</b>	
<b>6. Cardiovascular Pharmacogenetics</b> <i>Bas J. M. Peters, Anthonius de Boer, Tom Schalekamp, Olaf H. Klungel, and Anke-Hilse Maitland-van der Zee</i>	<b>185</b>
<b>7. Pharmacogenetics in Psychiatry</b> <i>Evangelia M. Tsapakis, Sarah Curran, Ruth I. Ohlsen, Nora S. Vyas, Katherine J. Aitchison, and Ann K. Daly</i>	<b>215</b>
<b>8. Pharmacogenetics in Cancer</b> <i>Sharon Marsh</i>	<b>251</b>



<b>9. Pharmacogenetics of Asthma and COPD</b>	<b>271</b>
<i>Ellen S. Koster, Jan A. M. Raaijmakers, Anke-Hilse Maitland-van der Zee, and Gerard H. Koppelman</i>	
<b>10. Pharmacogenetics of Adverse Drug Reactions</b>	<b>295</b>
<i>Ann K. Daly, Martin Armstrong, and Munir Pirmohamed</i>	
<b>11. Pharmacogenomics of Inflammatory Bowel Diseases</b>	<b>323</b>
<i>Alexander Teml, Susanne Karner, Elke Schaeffeler, and Matthias Schwab</i>	
<b>12. Pharmacogenetics of Pain Medication</b>	<b>353</b>
<i>Jörn Lötsch</i>	
<b>PART III PHARMACOGENETICS: IMPLEMENTATION IN CLINICAL PRACTICE</b>	
<b>13. Ethical and Social Issues in Pharmacogenomics Testing</b>	<b>377</b>
<i>Susanne Vijverberg, Toine Pieters, and Martina Cornel</i>	
<b>PART IV DEVELOPMENTS IN PHARMACOGENETIC RESEARCH</b>	
<b>14. High-Throughput Genotyping Technologies for Pharmacogenetics</b>	<b>403</b>
<i>Beatriz Sobrino and Angel Carracedo</i>	
<b>15. Developments in Analyses in Pharmacogenetic Datasets</b>	<b>415</b>
<i>Alison A. Motsinger-Reif</i>	
<b>PART V PHARMACOGENETICS: INDUSTRY AND REGULATORY AFFAIRS</b>	
<b>16. Applications of Pharmacogenetics in Pharmaceutical Research and Development</b>	<b>439</b>
<i>Daniel K. Burns and Scott S. Sundseth</i>	
<b>17. Role of Pharmacogenetics in Registration Processes</b>	<b>461</b>
<i>Myong-Jin Kim, Issam Zineh, Shiew-Mei Huang, and Lawrence J. Lesko</i>	
<b>PART VI CONCLUSIONS</b>	
<b>18. Pharmacogenetics: Possibilities and Pitfalls</b>	<b>479</b>
<i>Anke-Hilse Maitland-van der Zee and Ann K. Daly</i>	
<b>INDEX</b>	<b>485</b>

# **Pharmacogenetics: A Historical Perspective**

**ANN K. DALY**

Newcastle University, Newcastle upon Tyne, UK

## **1.1 INTRODUCTION**

It has been known for thousands of years that some individuals show toxic responses following consumption of fava beans, especially in countries bordering the Mediterranean. This is probably the earliest pharmacogenetic observation, although the biological basis for this has been established only quite recently (see Section 1.2). The foundation for much of modern pharmacogenetics came from experiments on chemical metabolism during the 19th century. These studies included the establishment that benzoic acid undergoes conjugation with glycine *in vivo* in both humans and animals, that benzene is oxidized to phenol in both dogs and humans and that some compounds can undergo conjugation with acetate (for a review, see Ref. 1).

## **1.2 EARLY PHARMACOGENETICS STUDIES (FROM 1900 TO 1970)**

The development of genetics and Mendelian inheritance together with observations by Archibald Garrod on the possibility of variation in chemical metabolism in the early 20th century has been well reviewed elsewhere see [2]. Probably the first direct pharmacogenetic study was reported in 1932 when Synder's study on the ability to taste phenylthiocarbamide within families showed that this trait was genetically determined [3]. The gene responsible for this variation and common genetic polymorphisms have only recently been identified (for a perspective, see Ref. 4).

---

*Pharmacogenetics and Individualized Therapy*, First Edition.

Edited by Anke-Hilse Maitland-van der Zee and Ann K. Daly.

© 2012 John Wiley & Sons, Inc. Published 2012 by John Wiley & Sons, Inc.

Although not a prescribed drug, phenylthiocarbamide shows homology to drugs such as propylthiouracil.

The initial drug-specific pharmacogenetics observations appeared in the literature during the 1950s. These were concerned with three widely used drugs at that time, that are all still used today: isoniazid, primaquine, and succinylcholine. The earliest observation concerned primaquine, which was found by Alf Alving to be associated with acute hemolysis in a small number of individuals [5]. Subsequent work by Alving and colleagues found that this toxicity was due to absence of the enzyme glucose-6-phosphate dehydrogenase in red blood cells of affected individuals [6]. The molecular genetic basis of this deficiency was later established by Ernest Beutler and colleagues in 1988 [7].

Isoniazid was first used against tuberculosis in the early 1950s, although it had been developed originally a number of years previously as an antidepressant. As reviewed recently, its use in tuberculosis patients represented an important advance in treatment of this disease [8]. Variation between individuals in urinary excretion profiles was described by Hettie Hughes [9], who soon afterwards also found an association between the metabolic profile and the incidence of a common adverse reaction, peripheral neuritis, with those showing slow conversion of the parent drug to acetylisoniazid more susceptible [10]. Further studies by several different workers, particularly Mitchell and Bell [11], Harris [12], and David Price Evans [13], led to the conclusion that isoniazid acetylation was subject to a genetic polymorphism and that some individuals (~10% of East Asians but 50% of Europeans) were slow acetylators. Slow acetylation was shown to be a recessive trait. As summarized in Section 1.4, the biochemical and genetic basis of slow acetylation is now well understood.

Also during the 1950s, a rare adverse response to the muscle relaxant succinylcholine was found to be due to an inherited deficiency in the enzyme cholinesterase [14]. Succinylcholine is used as a muscle relaxant during surgery, and those with the deficiency show prolonged paralysis (succinylcholine apnea). This observation was then further developed by Werner Kalow, who showed that the deficiency is inherited as an autosomal recessive trait and devised a biochemical test to screen for the deficiency, as he described in a description of his early work [15]. The gene encoding this enzyme, which is now usually referred to as *butyrylcholinesterase*, has been well studied, and a number of different mutations responsible for the deficiency have been identified. However, the original biochemical test is still the preferred method for identifying those affected by succinylcholine apnea due to the rarity of both the problem and the number of different mutations.

While these initial studies showing the clear role for genetics in determining adverse responses to primaquine, isoniazid, and succinylcholine were in progress, the general importance of the area was increasingly recognized. Arno Motulsky published a key review on the relationship between biochemical genetics and drug reactions that highlighted the adverse reactions to primaquine and succinylcholine in 1957 [16]. The term *pharmacogenetics* was first used in 1959 by Friedrich Vogel in an article on human genetics written in German [17] and was soon adopted by others working in the field.

### 1.3 PHARMACOGENETICS OF DRUG OXIDATION

As described in Section 1.1, studies in the 19th century had demonstrated oxidation of benzene to phenol *in vivo* [1]. Pioneering studies on drug metabolism, especially those in the laboratories of the Millers and of Brodie and Gillette during the 1950s, showed that many drugs undergo oxidative metabolism in the presence of NADPH and molecular oxygen in liver microsomes [18,19]. In 1962, Omura and Sato described cytochrome P450 from a rat liver microsome preparation as a hemoprotein that showed a peak at 450 nm in the presence of carbon monoxide and dithionite [20]. Shortly afterwards Ron Estabrook, David Cooper, and Otto Rosenthal showed that cytochrome P450 had steroid hydroxylase activity [21], and further studies confirmed its role in the metabolism of drugs such as codeine, aminopyrene, and acetanilide [22]. At this time, it was still assumed that cytochrome P450 was a single enzyme, but evidence for multiple forms emerged in the late 1960s [23,24] with purification of a range of rat and rabbit enzymes achieved during the 1970s [25,26].

Independent metabolism studies on two newly developed drugs sparteine and debrisoquine in Germany by Michel Eichelbaum and in the United Kingdom by Robert Smith in the mid 1970s resulted in findings indicating that some individuals were unable to oxidize these drugs, although the majority of individuals showed normal metabolism [27,28]. These studies estimated that 10% of Europeans showed absence of activity, and the term *poor metabolizer* was first used. At this time, the enzymes responsible for this absence of activity were not known, but further studies confirmed that the deficiency in metabolism of both drugs cosegregated [29] and that the trait was inherited recessively [30]. It became clear that a number of different drugs, including tricyclic antidepressants, were also metabolized by this enzyme [31]. Studies on human liver microsomes confirmed that the enzyme responsible was a cytochrome P450 [32,33], and this enzyme was then purified to homogeneity [34]. The availability of antibodies to the purified protein facilitated the cloning of the relevant cDNA by Frank Gonzalez and colleagues, who initially termed the gene CYP1D1 [35]. On the basis of emerging data for cytochrome P450 genes in humans and other animal species, it was decided subsequently that the gene encoding the debrisoquine/sparteine hydroxylase should be termed CYP2D6. Studies on human genomic DNA led to the identification of several polymorphisms in CYP2D6 associated with the poor metabolizer phenotype, including the most common splice site variant, a large deletion, and a small deletion [36–40]. A major additional contribution to the field was made in 1993 by Johansson, Ingelman-Sundberg, and colleagues, who described the phenomenon of ultrarapid metabolizers with one or more additional copies of CYP2D6 present [41]. These ultrarapid metabolizers had been previously identified on the basis of poor response to tricyclic antidepressants, and this was one of the first accounts of copy number variation in the human genome. Agreement regarding the current nomenclature for variant alleles in CYP2D6 and other cytochromes P450 was reached in 1996 [42].

In an approach similar to that used in the discovery of the CYP2D6 polymorphism, Kupfer and Preisig found that some individuals showed absence of metabolism of the anticonvulsant *S*-mephenytoin [43]. It was demonstrated that *S*-mephenytoin

metabolism did not cosegregate with that of debrisoquine and sparteine, as this polymorphism was due to a separate gene defect. Identification of the gene responsible for *S*-mephenytoin hydroxylase proved difficult initially, probably because the relevant enzyme was expressed at a low level in the liver. The gene, now termed CYP2C19, was cloned by Goldstein and Meyer and colleagues in 1994, and the two most common polymorphisms associated with absence of *S*-mephenytoin hydroxylase activity were identified [44,45].

A number of other cytochrome P450 genes are now known to be subject to functionally significant polymorphisms. In the case of one of these, CYP2C9, which metabolizes a range of drugs, including warfarin, tolbutamide, and nonsteroidal antiinflammatory drugs, some evidence for the existence of a polymorphism appeared in 1979 when a trimodal distribution in the metabolism of tolbutamide was reported [46]. Subsequently, it was shown that tolbutamide metabolism was distinct from debrisoquine metabolism [47]. The enzyme involved was purified and cloned and later named CYP2C9 [48,49]. Analysis of CYP2C9 cDNA sequences provided evidence for the presence of coding region polymorphisms resulting in amino acid substitutions, and expression studies suggested these were functionally significant [48,50,51]. Genotyping of patients undergoing warfarin treatment confirmed the functional importance of the two most common coding region CYP2C9 polymorphisms [52–54].

Using a similar approach involving comparison of cloned cDNA sequences, evidence for a nonsynonymous polymorphism in CYP2A6 was obtained [55]. Following expression studies and population screening, it was demonstrated that this polymorphism was associated with a rare absence of CYP2A6 activity, but additional polymorphisms (including a large deletion) in CYP2A6 that also lead to loss of activity have been reported [56].

Biochemical studies on human liver demonstrated that some individuals express an additional cytochrome P450 with homology but not identity to the major drug metabolizing P450 CYP3A4 [57–59]. Expression of this isoform, now termed CYP3A5, is also determined by a common genetic polymorphism affecting splicing that was first identified by Erin Schutz and colleagues in 2001 [60].

From the early studies in the 1970s, it is now clear that at least four CYPs, namely, CYP2D6, CYP2C19, CYP2A6, and CYP3A5, are subject to polymorphisms leading to absence of enzyme activity in significant numbers of individuals and that CYP2C9 activity is very low (although not completely absent) in some individuals. There are also a large number of polymorphisms leading to smaller changes in cytochrome P450 activities (see Chapter 3 for more details). Current knowledge of phenotype–genotype relationships within the cytochrome P450 family is now more comprehensive than for the majority of human genes, although a better understanding of some aspects such as regulation of gene expression is still needed.

## 1.4 PHARMACOGENETICS OF DRUG CONJUGATION

As discussed in Section 1.2, a polymorphism affecting conjugation of drugs such as isoniazid with acetyl CoA had been known to exist since the 1950s. Other

conjugation polymorphisms were subsequently described from phenotyping studies. In particular, Richard Weinshilboum identified several polymorphisms affecting methylation of xenobiotics and endogenous compounds by measurement of enzymatic activities in blood cells. He described the most pharmacologically important of these polymorphisms, in thiopurine methyltransferase (TPMT), in 1980 [61]. Approximately 1 in every 300 Europeans lacks this enzyme with lower activity observed in heterozygotes. Other conjugation polymorphisms identified by phenotypic approaches included a deficiency in the glutathione *S*-transferase M1 (GSTM1), which affects 50% of Europeans and was originally detected by measurement of *trans*-stilbene oxide conjugation in lymphocytes [62]. The classic paper by Motulsky on genetic variability in metabolism [16] mentioned the mild hyperbilirubinemia described previously by Gilbert in 1901 and usually referred to as *Gilbert's syndrome* [63]. This was later shown to relate to impaired activity in glucuronidation by a form of the enzyme UDP-glucuronosyltransferase, and there were suggestions that glucuronidation of prescribed drugs might also be affected in this syndrome [64].

With the development of molecular cloning techniques, the basis of the various conjugation polymorphisms known previously became clear during the late 1980s and early 1990s, and evidence also emerged for additional functionally significant polymorphisms by sequence comparisons. The molecular basis of the GSTM1 deficiency was established quite early in 1988, probably because it is due to a large gene deletion that was readily detectable by a number of different approaches [65]. Cloning of the NAT2 cDNA, encoding the enzyme responsible for isoniazid metabolism, was achieved in 1991 by Blum and Meyer with two common variant alleles with several base substitutions in their coding regions found to be associated with absence of activity [66]. Other inactive variants were identified elsewhere [67,68], and, as in the case of the cytochrome P450 alleles, a standardized nomenclature system was developed [69]. In the case of TPMT, gene cloning and identification of two common alleles associated with absence of activity was achieved in 1996 [70,71]. The most common variant allele giving rise to Gilbert's syndrome was identified in the same year and found to be a TA insertion in the promoter region of the UGT1A1 gene, which encodes the major UDP-glucuronosyltransferase responsible for bilirubin conjugation [72].

Genotyping for the TPMT polymorphisms in patients being prescribed 6-mercaptopurine or azathioprine and the UGT1A1 variant associated with Gilbert's disease in patients receiving irinotecan are now recommended but not mandated by the US Food and Drug Administration (FDA). Knowledge of genotype can enable either dose adjustment or an alternative drug to be prescribed.

## 1.5 PHARMACOGENETIC STUDIES ON RECEPTORS AND TRANSPORTERS

Progress on pharmacogenetics of drug receptors and other targets has been slower mainly because phenotypic evidence for the existence of functionally significant polymorphisms was generally not available. However, as discussed in



Chapter 5, data from the human genome sequencing project have provided new insights into this area. Studies on polymorphisms in both the adrenergic receptor and dopamine receptor genes appeared in the early 1990s with Stephen Liggett leading in the area of adrenergic receptors [73]. As discussed in Chapter 6, polymorphisms in the various adrenergic receptors have been demonstrated to be of considerable relevance to drug response, especially for the  $\beta_2$ -receptor, but the overall pharmacological importance of polymorphisms in dopamine receptors is still less well established.

Among other drug targets, vitamin K epoxide reductase, the target for coumarin anticoagulants, which is also discussed in detail in Chapter 6, is another example of a gene with well-established pharmacogenetics. In particular, limited phenotypic data from the 1970s suggested that the target for warfarin was subject to interindividual variation in some individuals with resistance to the drug occurring in some families [74]. The gene encoding vitamin K epoxide reductase in humans was finally identified only in 2004 [75,76], but this advance quickly led to identification of isolated mutations associated with warfarin resistance and also to common genetic polymorphisms affecting response to anticoagulants [77–79].

## 1.6 PHARMACOGENOMICS, GENOMEWIDE STUDIES, AND PERSONALIZED MEDICINE

As reviewed by Meyer [2], the term *pharmacogenomics* first appeared in the literature in 1997. One of the first articles using this term [80] described its relevance to personalized medicine. Pharmacogenomics is often described as the whole-genome application of *pharmacogenetics*. There is clearly a large overlap between the two disciplines, but pharmacogenomics is broader and may involve the development of new drugs to target specific genes as well as more effective use of existing medicines. Prior to the 1990s, pharmacogenetic studies were concerned with the effects of single genes, but in the era of pharmacogenomics, the combined effects of a number of genes on a particular phenotype is typically investigated.

Probably the best example of an area in which there has been some implementation of pharmacogenomics is in cancer chemotherapy. Although pharmacogenetic polymorphisms such as TPMT (see Section 1.4) are important in determining the metabolism of selected drugs used in chemotherapy and their possible toxicity, it was realized that tumor genotype and phenotype in addition to host genotype will be predictors of response. The licensing of trastuzumab (Herceptin) as a targeted therapy for breast tumors in 1998 is the earliest example of a drug used as a personalized medicine on the basis of tumor phenotype (for review, see Ref. 81). A test to determine estrogen receptor status is needed before the drug is prescribed as only tumors that are estrogen receptor-positive respond. Other similar drugs followed, most notably imatinib (Gleevec) in 2001. Imatinib is a tyrosine kinase inhibitor effective only in tumors with a particular chromosomal translocation [81]. In a separate development, it is now possible to classify tumors by signature for expression of a number of different genes and to use this signature to predict the

most appropriate cancer chemotherapy regimen. As discussed by Bonnefoi and colleagues [82], clinical trials are now in progress in breast cancer patients to confirm previous retrospective trials showing that determining mRNA expression levels for a set of either 21 or 70 genes in tumor tissue was of value in predicting whether patients with early-stage breast cancer should undergo chemotherapy or be treated only by hormone therapy.

Another area of pharmacogenomics that is currently developing is the use of genomewide association studies to identify genotypes associated with either drug response or drug toxicity. Such studies have been widely reported for complex polygenic diseases with some interesting novel genes affecting disease susceptibility already identified [83]. A number of genomewide association studies on drug response or adverse reactions have appeared since 2007 [84–86], but these have generally pointed to only one or two genes having a major effect rather than the larger number of genes each with a small effect typically seen in the complex polygenic disease studies. Further similar studies, especially on serious adverse drug reactions, are already in progress.

## 1.7 CONCLUSION

During 1957–1997, pharmacogenetics evolved to pharmacogenomics. There has been considerable further progress in the subsequent 12 years. Our understanding of single gene effects, especially in relation to drug metabolism, is now comprehensive, but our understanding of effects by multiple genes is still limited. In addition, we still need to translate the range of well-validated and clinically relevant pharmacogenetic discoveries that have been made over the years into more widespread use in patient care. Despite the predictions that we are entering an era of personalized medicine [80], except for the few examples discussed in Sections 1.5 and 1.6 in relation to cancer treatment, this has not yet occurred to any great extent.

## REFERENCES

1. Conti A, Bickel MH. History of drug-metabolism—discoveries of major pathways in 19th-century. *Drug Metab. Rev.* 1977;**6**:1–50.
2. Meyer UA. Pharmacogenetics—five decades of therapeutic lessons from genetic diversity. *Nat. Rev. Genet.* 2004;**5**:669–676.
3. Snyder LH. Studies in human inheritance IX. The inheritance of taste deficiency in man. *Ohio J. Sci.* 1932;**32**:436–468.
4. Wooding S. Phenylthiocarbamide: A 75-year adventure in genetics and natural selection. *Genetics* 2006;**172**:2015–2023.
5. Clayman CB, Arnold J, Hockwald RS, Yount EH Jr, Edgcomb JH, Alving AS. Toxicity of primaquine in Caucasians. *J. Am. Med. Assoc. (JAMA)* 1952;**149**:1563–1568.
6. Alving AS, Carson PE, Flanagan CL, Ickes CE. Enzymatic deficiency in primaquine-sensitive erythrocytes. *Science* 1956;**124**:484–485.



7. Hirono A, Beutler E. Molecular cloning and nucleotide sequence of cDNA for human glucose-6-phosphate dehydrogenase variant A(-). *Proc. Natl. Acad. Sci. USA* 1988;**85**:3951-3954.
8. Rieder HL. Fourth-generation fluoroquinolones in tuberculosis. *Lancet* 2009;**373**:1148-1149.
9. Hughes HB. On the metabolic fate of isoniazid. *J. Pharmacol. Exp. Ther.* 1953;**109**:444-452.
10. Hughes HB, Biehl JP, Jones AP, Schmidt LH. Metabolism of isoniazid in man as related to the occurrence of peripheral neuritis. *Am. Rev. Tuberc.* 1954;**70**:266-273.
11. Mitchell RS, Bell JC. Clinical implications of isoniazid, PAS and streptomycin blood levels in pulmonary tuberculosis. *Trans. Am. Clin. Climatol. Assoc.* 1957;**69**:98-102; discussion. 103-105.
12. Harris HW, Knight RA, Selin MJ. Comparison of isoniazid concentrations in the blood of people of Japanese and European descent; therapeutic and genetic implications. *Am. Rev. Tuberc.* 1958;**78**:944-948.
13. Evans DAP, Manley KA, McKusick VA. Genetic control of isoniazid metabolism in man. *Br. Med. J.* 1960;**2**:485-491.
14. Lehmann H, Ryan E. The familial incidence of low pseudocholinesterase level. *Lancet* 1956;**271**:124.
15. Kalow W. The Pennsylvania State University College of Medicine 1990 Bernard B. Brodie Lecture. Pharmacogenetics: Past and future. *Life Sci.* 1990;**47**:1385-1397.
16. Motulsky AG. Drug reactions enzymes, and biochemical genetics. *J. Am. Med. Assoc.* 1957;**165**:835-837.
17. Vogel F. Moderne probleme der humangenetik. *Ergebnisse der Innere Medizinische und Kinderheilkunde* 1959;**12**:52-62.
18. Conney AH, Miller EC, Miller JA. Substrate-induced synthesis and other properties of benzpyrene hydroxylase in rat liver. *J. Biol. Chem.* 1957;**228**:753-766.
19. Brodie BB, Gillette JR, La Du BN. Enzymatic metabolism of drugs and other foreign compounds. *Annu. Rev. Biochem.* 1958;**27**:427-454.
20. Omura T, Sato R. A new cytochrome in liver microsomes. *J. Biol. Chem.* 1962;**237**:1375-1376.
21. Cooper DY, Estabrook RW, Rosenthal O. The stoichiometry of C21 hydroxylation of steroids by adrenocortical microsomes. *J. Biol. Chem.* 1963;**238**:1320-1323.
22. Cooper DY, Levin S, Narasimhulu S, Rosenthal O. Photochemical action spectrum of the terminal oxidase of mixed function oxidase systems. *Science* 1965;**147**:400-402.
23. Sladek NE, Mannering GJ. Induction of drug metabolism. I. Differences in the mechanisms by which polycyclic hydrocarbons and phenobarbital produce their inductive effects on microsomal N-demethylating systems. *Mol. Pharmacol.* 1969;**5**:174-185.
24. Alvares AP, Schilling G, Levin W, Kuntzman R. Studies on the induction of CO-binding pigments in liver microsomes by phenobarbital and 3-methylcholanthrene. *Biochem. Biophys. Res. Commun.* 1967;**29**:521-526.
25. Cheng KC, Schenkman JB. Purification and characterization of two constitutive forms of rat liver microsomal cytochrome P-450. *J. Biol. Chem.* 1982;**257**:2378-2385.