

**SCHAUM'S OUTLINE SERIES**

**THEORY AND PROBLEMS OF**

# **MODERN PHYSICS**

**RONALD GAUTREAU**

**WILLIAM SAVIN**

**INCLUDING 486 SOLVED PROBLEMS**

**SCHAUM'S OUTLINE SERIES IN SCIENCE**

**McGRAW-HILL BOOK COMPANY**

**SCHAUM'S OUTLINE OF**

**THEORY AND PROBLEMS**

of

**MODERN PHYSICS**

by

**RONALD GAUTREAU, Ph.D.**

*Associate Professor of Physics  
New Jersey Institute of Technology*

and

**WILLIAM SAVIN, Ph.D.**

*Associate Professor of Physics  
New Jersey Institute of Technology*

**SCHAUM'S OUTLINE SERIES**

McGRAW-HILL BOOK COMPANY

|                 |                  |                      |                 |                  |                   |                     |
|-----------------|------------------|----------------------|-----------------|------------------|-------------------|---------------------|
| <i>New York</i> | <i>St. Louis</i> | <i>San Francisco</i> | <i>Auckland</i> | <i>Bogotá</i>    | <i>Düsseldorf</i> | <i>Johannesburg</i> |
| <i>London</i>   | <i>Madrid</i>    | <i>Mexico</i>        | <i>Montreal</i> | <i>New Delhi</i> | <i>Panama</i>     | <i>Paris</i>        |
|                 | <i>São Paulo</i> | <i>Singapore</i>     | <i>Sydney</i>   | <i>Tokyo</i>     | <i>Toronto</i>    |                     |

*Dedicated to the memory of  
Professor Marcus M. Mainardi*

Copyright © 1978 by McGraw-Hill, Inc. All rights reserved. Printed in the United States of America. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher.

0-07-023062-5

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 SH SH 7 8 3 2 1 0 9 8

**Library of Congress Cataloging in Publication Data**

Gautreau, Ronald.

Schaum's outline of theory and problems of modern physics.

(Schaum's outline series)

Includes index.

I. Physics—Problems, exercises, etc. I. Savin, William, joint author. II. Title.

QC32.G34 530'.076 78-8752

ISBN 0-07-023062-5

## Preface

The area of Modern Physics embraces topics that have evolved since roughly the turn of this century. These developments can be mind-boggling, as with the effects on time predicted by Einstein's Special Theory of Relativity, or quite practical, like the many devices based upon semiconductors, whose explanation lies in the band theory of solids.

The scope of the present book may be gauged from the Table of Contents. Each chapter consists of a succinct presentation of the principles and "meat" of a particular subject, followed by a large number of completely solved problems that naturally develop the subject and illustrate the principles. It is the authors' conviction that these solved problems are a valuable learning tool. The solved problems have been made short and to the point, and have been ordered in terms of difficulty. They are followed by unsolved supplementary problems, with answers, which allow the reader to check his grasp of the material.

It has been assumed that the reader has had the standard introductory courses in general physics, and the book is geared primarily at the sophomore or junior level, although we have also included problems of a more advanced nature. While it will certainly serve as a supplement to any standard Modern Physics text, this book is sufficiently comprehensive and self-contained to be used by itself to learn the principles of Modern Physics.

We extend special thanks to David Beckwith for meticulous editing and for input that improved the final version of the book. Any mistakes are ours, of course, and we would appreciate having these pointed out to us. Finally, we are indebted to our families for their enormous patience with us throughout the long preparation of this work.

RONALD GAUTREAU  
WILLIAM SAVIN

New Jersey Institute of Technology

### **Catalog**

**If you are interested in a list of SCHAUM'S  
OUTLINE SERIES in Science, Mathematics,  
Engineering and other subjects, send your name  
and address, requesting your free catalog, to:**

**SCHAUM'S OUTLINE SERIES, Dept. C  
McGRAW-HILL BOOK COMPANY  
1221 Avenue of Americas  
New York, N.Y. 10020**

# Contents

|                  |  |           |
|------------------|--|-----------|
| <b>PART I</b>    | <b>THE SPECIAL THEORY OF RELATIVITY</b>                                      | <b>1</b>  |
| <b>Chapter 1</b> | <b>Galilean Transformations</b>  | <b>1</b>  |
| 1.1              | Events and Coordinates   | 1         |
| 1.2              | Galilean Coordinate Transformations  | 1         |
| 1.3              | Galilean Velocity Transformations  | 1         |
| 1.4              | Galilean Acceleration Transformations  | 2         |
| 1.5              | Invariance of an Equation  | 2         |
| <b>Chapter 2</b> | <b>The Postulates of Einstein</b>  | <b>7</b>  |
| 2.1              | Absolute Space and the Ether   | 7         |
| 2.2              | The Michelson-Morley Experiment  | 7         |
| 2.3              | Length and Time Measurements—A Question of Principle                         | 7         |
| 2.4              | The Postulates of Einstein   | 7         |
| <b>Chapter 3</b> | <b>The Lorentz Coordinate Transformations</b>                                | <b>12</b> |
| 3.1              | The Constancy of the Speed of Light  | 12        |
| 3.2              | The Invariance of Maxwell's Equations  | 12        |
| 3.3              | General Considerations in Solving Problems Involving Lorentz Transformations | 13        |
| 3.4              | Simultaneity   | 13        |
| <b>Chapter 4</b> | <b>Relativistic Length Measurements</b>                                      | <b>17</b> |
| 4.1              | The Definition of Length   | 17        |
| <b>Chapter 5</b> | <b>Relativistic Time Measurements</b>  | <b>20</b> |
| 5.1              | Proper Time  | 20        |
| 5.2              | Time Dilation  | 20        |
| <b>Chapter 6</b> | <b>Relativistic Space-Time Measurements</b>                                  | <b>24</b> |
| <b>Chapter 7</b> | <b>Relativistic Velocity Transformations</b>                                 | <b>34</b> |
| 7.1              | The Lorentz Velocity Transformations and the Speed of Light                  | 34        |
| 7.2              | General Considerations in Solving Velocity Problems                          | 34        |
| <b>Chapter 8</b> | <b>Mass, Energy and Momentum in Relativity</b>                               | <b>39</b> |
| 8.1              | The Need to Redefine Classical Momentum                                      | 39        |
| 8.2              | The Variation of Mass with Velocity  | 39        |
| 8.3              | Newton's Second Law in Relativity  | 39        |

## CONTENTS

|            |   |    |
|------------|---|----|
| 8.4        | Mass and Energy Relationship: $E = mc^2$                    | 40 |
| 8.5        | Momentum and Energy Relationship                            | 40 |
| 8.6        | Units for Energy and Momentum                               | 40 |
| 8.7        | General Considerations in Solving Mass-Energy Problems      | 41 |
| Chapter 9  | <b>The Relativistic Doppler Effect</b>                      | 50 |
| PART II    | <b>THE QUANTUM THEORY OF ELECTROMAGNETIC RADIATION</b>      | 53 |
| Chapter 10 | <b>The Theory of Photons</b>                                | 53 |
| Chapter 11 | <b>The Photoelectric Effect</b>                             | 56 |
| 11.1       | Experimental Results  | 56 |
| 11.2       | Theory of the Photoelectric Effect                          | 56 |
| Chapter 12 | <b>The Compton Effect</b>                                   | 62 |
| Chapter 13 | <b>Pair Production and Annihilation</b>                     | 68 |
| 13.1       | Pair Production   | 68 |
| 13.2       | Pair Annihilation   | 68 |
| Chapter 14 | <b>Absorption of Photons</b>                                | 74 |
| PART III   | <b>MATTER WAVES</b>   | 77 |
| Chapter 15 | <b>De Broglie Waves</b>                                     | 77 |
| 15.1       | The Wave-Particle Duality of Electromagnetic Radiation      | 77 |
| 15.2       | The Wave-Particle Duality of Matter                         | 77 |
| Chapter 16 | <b>Experimental Verification of De Broglie's Hypothesis</b> | 82 |
| 16.1       | The Bragg Law of Diffraction                                | 82 |
| 16.2       | Electron Diffraction Experiments                            | 83 |
| Chapter 17 | <b>The Probability Interpretation of De Broglie Waves</b>   | 88 |
| 17.1       | A Probability Interpretation for Electromagnetic Radiation  | 88 |
| 17.2       | A Probability Interpretation of Matter                      | 88 |
| Chapter 18 | <b>The Heisenberg Uncertainty Principle</b>                 | 91 |
| 18.1       | Measurements and Uncertainties                              | 91 |
| 18.2       | The Uncertainty Relation for Position and Momentum          | 92 |
| 18.3       | The Uncertainty Relation for Energy and Time                | 92 |
| 18.4       | The Principle of Complementarity                            | 92 |

## CONTENTS

|                |  |            |
|----------------|--|------------|
| <b>PART IV</b> | <b>HYDROGENLIKE ATOMS</b>  | <b>99</b>  |
| Chapter 19     | <b>The Bohr Atom</b>   | <b>99</b>  |
| 19.1           | The Hydrogen Spectrum  | 99         |
| 19.2           | The Bohr Theory of the Hydrogen Atom                                       | 99         |
| 19.3           | Emission of Radiation in Bohr's Theory                                     | 100        |
| 19.4           | Energy Level Diagrams  | 101        |
| 19.5           | Hydrogenic Atoms   | 103        |
| 19.6           | $\mu$ -Mesic and $\pi$ -Mesic Atoms  | 103        |
| Chapter 20     | <b>Electron Orbital Motion and the Zeeman Effect</b>                       | <b>112</b> |
| 20.1           | Orbital Angular Momentum from a Classical Viewpoint                        | 112        |
| 20.2           | Classical Magnetic Dipole Moment   | 112        |
| 20.3           | Classical Energy of a Magnetic Dipole Moment in an External Magnetic Field | 113        |
| 20.4           | The Zeeman Experiment  | 113        |
| 20.5           | Quantization of the Magnitude of the Orbital Angular Momentum              | 114        |
| 20.6           | Quantization of the Direction of the Orbital Angular Momentum              | 114        |
| 20.7           | Explanation of the Zeeman Effect   | 114        |
| Chapter 21     | <b>The Stern–Gerlach Experiment and Electron Spin</b>                      | <b>120</b> |
| 21.1           | The Stern–Gerlach Experiment   | 120        |
| 21.2           | Electron Spin  | 120        |
| Chapter 22     | <b>Electron Spin and Fine Structure</b>                                    | <b>124</b> |
| 22.1           | Spin-Orbit Coupling  | 124        |
| 22.2           | Fine Structure   | 124        |
| 22.3           | Total Angular Momentum (the Vector Model)                                  | 124        |
| <b>PART V</b>  | <b>MANY-ELECTRON ATOMS</b>   | <b>128</b> |
| Chapter 23     | <b>The Pauli Exclusion Principle</b>                                       | <b>128</b> |
| 23.1           | Introduction   | 128        |
| 23.2           | The Pauli Exclusion Principle  | 128        |
| 23.3           | A Single Particle in a One-Dimensional Box                                 | 128        |
| 23.4           | Many Particles in a One-Dimensional Box                                    | 128        |
| Chapter 24     | <b>Many-Electron Atoms and the Periodic Table</b>                          | <b>133</b> |
| 24.1           | Spectroscopic Notation for Electron Configurations in Atoms                | 133        |
| 24.2           | The Periodic Table and an Atomic Shell Model                               | 133        |
| 24.3           | Spectroscopic Notation for Atomic States                                   | 134        |
| 24.4           | Atomic Excited States and <i>LS</i> Coupling                               | 135        |
| 24.5           | The Anomalous Zeeman Effect  | 135        |
| Chapter 25     | <b>Inner-Electron Transitions: X-Rays</b>                                  | <b>149</b> |
| 25.1           | X-Ray Apparatus  | 149        |
| 25.2           | Production of Bremsstrahlung   | 149        |



## CONTENTS

|                |  |            |
|----------------|--|------------|
| 25.3           | Production of Characteristic X-Ray Spectra | 150        |
| 25.4           | The Moseley Relation                       | 150        |
| 25.5           | X-Ray Absorption Edges                     | 152        |
| 25.6           | Auger Effect                               | 152        |
| 25.7           | X-Ray Fluorescence                         | 153        |
| <b>PART VI</b> | <b>NUCLEAR PHYSICS</b>                     | <b>163</b> |
| Chapter 26     | <b>Nucleon and Deuteron Properties</b>     | <b>163</b> |
| 26.1           | The Nucleons                               | 163        |
| 26.2           | Nucleon Forces                             | 163        |
| 26.3           | The Deuteron                               | 164        |
| Chapter 27     | <b>Properties of Nuclei</b>                | <b>168</b> |
| 27.1           | Designation of Nuclei                      | 168        |
| 27.2           | Relative Number of Protons and Neutrons    | 168        |
| 27.3           | The Nucleus as a Sphere                    | 169        |
| 27.4           | Nuclear Binding Energy                     | 169        |
| Chapter 28     | <b>Nuclear Models</b>                      | <b>173</b> |
| 28.1           | Liquid Drop Model                          | 173        |
| 28.2           | Shell Model                                | 175        |
| Chapter 29     | <b>The Decay of Unstable Nuclei</b>        | <b>184</b> |
| 29.1           | Introduction                               | 184        |
| 29.2           | The Statistical Radioactive Decay Law      | 184        |
| 29.3           | Gamma Decay                                | 185        |
| 29.4           | Alpha Decay                                | 185        |
| 29.5           | Beta Decay and the Neutrino                | 186        |
| Chapter 30     | <b>Nuclear Reactions</b>                   | <b>199</b> |
| 30.1           | Introduction                               | 199        |
| 30.2           | Classification of Nuclear Reactions        | 199        |
| 30.3           | Laboratory and Center-of-Mass Systems      | 200        |
| 30.4           | Energetics of Nuclear Reactions            | 201        |
| 30.5           | Nuclear Cross Sections                     | 201        |
| Chapter 31     | <b>Fission and Fusion</b>                  | <b>209</b> |
| 31.1           | Nuclear Fission                            | 209        |
| 31.2           | Nuclear Fusion                             | 210        |
| Chapter 32     | <b>Elementary Particles</b>                | <b>215</b> |
| 32.1           | Elementary Particle Genealogy              | 215        |
| 32.2           | Particle Interactions                      | 216        |

## CONTENTS

|  |                |
|--|----------------|
| 32.3 Conservation Laws   | 216            |
| 32.4 Conservation of Leptons   | 217            |
| 32.5 Conservation of Baryons   | 217            |
| 32.6 Conservation of Isotopic Spin   | 217            |
| 32.7 Conservation of Strangeness   | 218            |
| 32.8 Conservation of Parity  | 219            |
| 32.9 Short-lived Particles and the Resonances                                  | 219            |
| <br><b>PART VII ATOMIC SYSTEMS</b>   | <br><b>227</b> |
| <br>Chapter 33 <b>Molecular Bonding</b>  | <br>227        |
| 33.1 Ionic Bonding   | 227            |
| 33.2 Covalent Bonding  | 227            |
| 33.3 Other Types of Bonding  | 228            |
| <br>Chapter 34 <b>Excitations of Diatomic Molecules</b>                        | <br>232        |
| 34.1 Molecular Rotations   | 232            |
| 34.2 Molecular Vibrations  | 232            |
| 34.3 Combined Excitations  | 233            |
| <br>Chapter 35 <b>Kinetic Theory</b>   | <br>241        |
| 35.1 The Ideal Gas Law   | 241            |
| <br>Chapter 36 <b>Distribution Functions</b>                                   | <br>249        |
| 36.1 Discrete Distribution Functions   | 249            |
| 36.2 Continuous Distribution Functions   | 250            |
| 36.3 Fundamental Distribution Functions and Density of States                  | 250            |
| <br>Chapter 37 <b>Classical Statistics: The Maxwell-Boltzmann Distribution</b> | <br>256        |
| <br>Chapter 38 <b>Quantum Statistics</b>                                       | <br>266        |
| 38.1 Fermi-Dirac Statistics  | 266            |
| 38.2 Bose-Einstein Statistics  | 266            |
| 38.3 High-Temperature Limit  | 267            |
| 38.4 Two Useful Integrals  | 267            |
| <br>Chapter 39 <b>The Band Theory of Solids</b>                                | <br>287        |
| <br><b>Appendix</b>  | <br>296        |
| <br><b>Index</b>   | <br>305        |

## Chapter 1

### Galilean Transformations

#### 1.1 EVENTS AND COORDINATES

We begin by considering the concept of a physical event. The event might be the striking of a tree by a lightning bolt or the collision of two particles, and happens at a point in space and at an instant in time. The particular event is specified by an observer by assigning to it four coordinates: the three position coordinates  $x, y, z$  that measure the distance from the origin of a coordinate system where the observer is located, and the time coordinate  $t$  that the observer records with his clock.

Consider now two observers,  $O$  and  $O'$ , where  $O'$  travels with a constant velocity  $v$  with respect to  $O$  along their common  $x$ - $x'$  axis (Fig. 1-1). Both observers are equipped with metersticks and clocks so that they can measure coordinates of events. Further, suppose both observers adjust their clocks so that when they pass each other at  $x = x' = 0$ , the clocks read  $t = t' = 0$ . Any given event  $P$  will have eight numbers associated with it, the four coordinates  $(x, y, z, t)$  assigned by  $O$  and the four coordinates  $(x', y', z', t')$  assigned (to the same event) by  $O'$ .




Fig. 1-1

#### 1.2 GALILEAN COORDINATE TRANSFORMATIONS

The relationship between the measurements  $(x, y, z, t)$  of  $O$  and the measurements  $(x', y', z', t')$  of  $O'$  for a particular event is obtained by examining Fig. 1-1:

$$x' = x - vt \quad y' = y \quad z' = z$$

In addition, in classical physics it is implicitly assumed that

$$t' = t$$

These four equations are called the *Galilean coordinate transformations*.

#### 1.3 GALILEAN VELOCITY TRANSFORMATIONS

In addition to the coordinates of an event, the velocity of a particle is of interest. Observers  $O$  and  $O'$  will describe the particle's velocity by assigning three components to it, with  $(u_x, u_y, u_z)$  being the velocity components as measured by  $O$ , and  $(u'_x, u'_y, u'_z)$  being the velocity components as measured by  $O'$ .

The relationship between  $(u_x, u_y, u_z)$  and  $(u'_x, u'_y, u'_z)$  is obtained from the time differentiation of the Galilean coordinate transformations. Thus, from  $x' = x - vt$ ,

$$u'_x = \frac{dx'}{dt'} = \frac{d}{dt}(x - vt) \frac{dt}{dt'} = \left( \frac{dx}{dt} - v \right) (1) = u_x - v$$

Altogether, the *Galilean velocity transformations* are

$$u'_x = u_x - v \quad u'_y = u_y \quad u'_z = u_z$$

#### 1.4 GALILEAN ACCELERATION TRANSFORMATIONS

The acceleration of a particle is the time derivative of its velocity, i.e.  $a_x = du_x/dt$ , etc. To find the *Galilean acceleration transformations* we differentiate the velocity transformations and use the facts that  $t' = t$  and  $v = \text{constant}$  to obtain

$$a'_x = a_x \quad a'_y = a_y \quad a'_z = a_z$$

Thus the measured acceleration components are the same for all observers moving with uniform relative velocity.

#### 1.5 INVARIANCE OF AN EQUATION

By *invariance* of an equation it is meant that the equation will have the same form when determined by two observers. In classical theory it is assumed that space and time measurements of two observers are related by the Galilean transformations. Thus, when a particular form of an equation is determined by one observer, the Galilean transformations can be applied to this form to determine the form for the other observer. If both forms are the same, the equation is invariant under the Galilean transformations. See Problems 1.11 and 1.12.

### Solved Problems

- 1.1. A passenger in a train moving at 30 m/s passes a man standing on a station platform at  $t = t' = 0$ . Twenty seconds after the train passes him, the man on the platform determines that a bird flying along the tracks in the same direction as the train is 800 m away. What are the coordinates of the bird as determined by the passenger?

The coordinates assigned to the bird by the man on the station platform are

$$(x, y, z, t) = (800 \text{ m}, 0, 0, 20 \text{ s})$$

The passenger measures the distance  $x'$  to the bird as

$$x' = x - vt = 800 \text{ m} - (30 \text{ m/s})(20 \text{ s}) = 200 \text{ m}$$

Therefore the bird's coordinates as determined by the passenger are

$$(x', y', z', t') = (200 \text{ m}, 0, 0, 20 \text{ s})$$

- 1.2. Refer to Problem 1.1. Five seconds after making the first coordinate measurement, the man on the platform determines that the bird is 850 m away. From these data find the velocity of the bird (assumed constant) as determined by the man on the platform and by the passenger on the train.

The coordinates assigned to the bird at the second position by the man on the platform are

$$(x_2, y_2, z_2, t_2) = (850 \text{ m}, 0, 0, 25 \text{ s})$$

Hence, the velocity  $u_x$  of the bird as measured by the man on the platform is

$$u_x = \frac{x_2 - x_1}{t_2 - t_1} = \frac{850 \text{ m} - 800 \text{ m}}{25 \text{ s} - 20 \text{ s}} = +10 \text{ m/s}$$

The positive sign indicates the bird is flying in the positive  $x$ -direction. The passenger finds that at the second position the distance  $x'_2$  to the bird is

$$x'_2 = x_2 - vt_2 = 850 \text{ m} - (30 \text{ m/s})(25 \text{ s}) = 100 \text{ m}$$

Thus,  $(x'_2, y'_2, z'_2, t'_2) = (100 \text{ m}, 0, 0, 25 \text{ s})$ , and the velocity  $u'_x$  of the bird as measured by the passenger is

$$u'_x = \frac{x'_2 - x'_1}{t'_2 - t'_1} = \frac{100 \text{ m} - 200 \text{ m}}{25 \text{ s} - 20 \text{ s}} = -20 \text{ m/s}$$

so that, as measured by the passenger, the bird is moving in the negative  $x'$ -direction. Note that this result is consistent with that obtained from the Galilean velocity transformation:

$$u'_x = u_x - v = 10 \text{ m/s} - 30 \text{ m/s} = -20 \text{ m/s}$$

- 1.3.** A sample of radioactive material, at rest in the laboratory, ejects two electrons in opposite directions. One of the electrons has a speed of  $0.6c$  and the other has a speed of  $0.7c$ , as measured by a laboratory observer. According to classical velocity transformations, what will be the speed of one electron as measured from the other?

Let observer  $O$  be at rest with respect to the laboratory and let observer  $O'$  be at rest with respect to the particle moving with speed  $0.6c$  (taken in the positive direction). Then, from the Galilean velocity transformation,

$$u'_x = u_x - v = -0.7c - 0.6c = -1.3c$$

This problem demonstrates that velocities greater than the speed of light are possible with the Galilean transformations, a result that is inconsistent with Special Relativity.

- 1.4.** A train moving with a velocity of  $60 \text{ mi/hr}$  passes through a railroad station at 12:00. Twenty seconds later a bolt of lightning strikes the railroad tracks one mile from the station in the same direction that the train is moving. Find the coordinates of the lightning flash as measured by an observer at the station and by the engineer of the train.

Both observers measure the time coordinate as

$$t = t' = (20 \text{ s}) \left( \frac{1 \text{ hr}}{3600 \text{ s}} \right) = \frac{1}{180} \text{ hr}$$

The observer at the station measures the spatial coordinate to be  $x = 1 \text{ mi}$ . The spatial coordinate as determined by the engineer of the train is

$$x' = x - vt = 1 \text{ mi} - (60 \text{ mi/hr}) \left( \frac{1}{180} \text{ hr} \right) = \frac{2}{3} \text{ mi}$$

- 1.5.** A hunter on the ground fires a bullet in the northeast direction which strikes a deer  $0.25 \text{ miles}$  from the hunter. The bullet travels with a speed of  $1800 \text{ mi/hr}$ . At the instant when the bullet is fired, an airplane is directly over the hunter at an altitude of one mile and is traveling due east with a velocity of  $600 \text{ mi/hr}$ . When the bullet strikes the deer, what are the coordinates as determined by an observer in the airplane?

Using the Galilean transformations,

$$t' = t = \frac{0.25 \text{ mi}}{1800 \text{ mi/hr}} = 1.39 \times 10^{-4} \text{ hr}$$

$$x' = x - vt = (0.25 \text{ mi}) \cos 45^\circ - (600 \text{ mi/hr})(1.39 \times 10^{-4} \text{ hr}) = 0.094 \text{ mi}$$

$$y' = y = (0.25 \text{ mi}) \sin 45^\circ = 0.177 \text{ mi}$$

$$z' = z - h = 0 - 1 \text{ mi} = -1 \text{ mi}$$

- 1.6. An observer, at rest with respect to the ground, observes the following collision. A particle of mass  $m_1 = 3$  kg moving with velocity  $u_1 = 4$  m/s along the  $x$ -axis approaches a second particle of mass  $m_2 = 1$  kg moving with velocity  $u_2 = -3$  m/s along the  $x$ -axis. After a head-on collision the ground observer finds that  $m_2$  has velocity  $u_2^* = 3$  m/s along the  $x$ -axis. Find the velocity  $u_1^*$  of  $m_1$  after the collision.

initial momentum = final momentum

$$\begin{aligned} m_1 u_1 + m_2 u_2 &= m_1 u_1^* + m_2 u_2^* \\ (3 \text{ kg})(4 \text{ m/s}) + (1 \text{ kg})(-3 \text{ m/s}) &= (3 \text{ kg})u_1^* + (1 \text{ kg})(3 \text{ m/s}) \\ 9 \text{ kg} \cdot \text{m/s} &= (3 \text{ kg})u_1^* + 3 \text{ kg} \cdot \text{m/s} \end{aligned}$$

Solving,  $u_1^* = 2$  m/s.

- 1.7. A second observer,  $O'$ , who is walking with a velocity of 2 m/s relative to the ground along the  $x$ -axis observes the collision described in Problem 1.6. What are the system momenta before and after the collision as determined by him?

Using the Galilean velocity transformations,

$$\begin{aligned} u_1' &= u_1 - v = 4 \text{ m/s} - 2 \text{ m/s} = 2 \text{ m/s} \\ u_2' &= u_2 - v = -3 \text{ m/s} - 2 \text{ m/s} = -5 \text{ m/s} \\ u_1^{*'} &= u_1^* - v = 2 \text{ m/s} - 2 \text{ m/s} = 0 \\ u_2^{*'} &= u_2^* - v = 3 \text{ m/s} - 2 \text{ m/s} = 1 \text{ m/s} \\ (\text{initial momentum})' &= m_1 u_1' + m_2 u_2' = (3 \text{ kg})(2 \text{ m/s}) + (1 \text{ kg})(-5 \text{ m/s}) = 1 \text{ kg} \cdot \text{m/s} \\ (\text{final momentum})' &= m_1 u_1^{*'} + m_2 u_2^{*'} = (3 \text{ kg})(0) + (1 \text{ kg})(1 \text{ m/s}) = 1 \text{ kg} \cdot \text{m/s} \end{aligned}$$

Thus, as a result of the Galilean transformations,  $O'$  also determines that momentum is conserved (but at a different value from that found by  $O$ ).

- 1.8. An open car traveling at 100 ft/s has a boy in it who throws a ball upward with a velocity of 20 ft/s. Write the equation of motion (giving position as a function of time) for the ball as seen by (a) the boy, (b) an observer stationary on the road.

(a) For the boy in the car the ball travels straight up and down, so

$$\begin{aligned} y' &= v_0 t' + \frac{1}{2} a t'^2 = (20 \text{ ft/s})t' + \frac{1}{2}(-32 \text{ ft/s}^2)t'^2 = 20t' - 16t'^2 \\ x' &= z' = 0 \end{aligned}$$

(b) For the stationary observer, one obtains from the Galilean transformations

$$\begin{aligned} t &= t' \\ x &= x' + vt = 0 + 100t & y &= y' = 20t - 16t^2 & z &= z' = 0 \end{aligned}$$

- 1.9. Consider a mass attached to a spring and moving on a horizontal, frictionless surface. Show, from the classical transformation laws, that the equations of motion of the mass are the same as determined by an observer at rest with respect to the surface and by a second observer moving with constant velocity along the direction of the spring.

The equation of motion of the mass, as determined by an observer at rest with respect to the surface, is  $F = ma$ , or

$$-k(x - x_0) = m \frac{d^2 x}{dt^2} \quad (1)$$

To determine the equation of motion as found by the second observer we use the Galilean transformations to obtain

$$x = x' + vt' \quad x_0 = x'_0 + vt' \quad \frac{d^2 x}{dt^2} = \frac{d^2 x'}{dt'^2}$$

Substituting these values in (1) gives

$$-k(x' - x'_0) = m \frac{d^2 x'}{dt'^2} \quad (2)$$

Because (1) and (2) have the same form, the equation of motion is invariant under the Galilean transformations.

**1.10.** Show that the electromagnetic wave equation,

$$\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} + \frac{\partial^2 \phi}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} = 0$$

is not invariant under the Galilean transformations.

The equation will be invariant if it retains the same form when expressed in terms of the new variables  $x', y', z', t'$ . We first find from the Galilean transformations that

$$\begin{aligned} \frac{\partial x'}{\partial x} &= 1 & \frac{\partial x'}{\partial t} &= -v & \frac{\partial t'}{\partial t} &= \frac{\partial y'}{\partial y} = \frac{\partial z'}{\partial z} = 1 \\ \frac{\partial x'}{\partial y} &= \frac{\partial x'}{\partial z} = \frac{\partial y'}{\partial x} = \frac{\partial t'}{\partial x} = \dots = 0 \end{aligned}$$

From the chain rule and using the above results we have

$$\frac{\partial \phi}{\partial x} = \frac{\partial \phi}{\partial x'} \frac{\partial x'}{\partial x} + \frac{\partial \phi}{\partial y'} \frac{\partial y'}{\partial x} + \frac{\partial \phi}{\partial z'} \frac{\partial z'}{\partial x} + \frac{\partial \phi}{\partial t'} \frac{\partial t'}{\partial x} = \frac{\partial \phi}{\partial x'} \quad \text{and} \quad \frac{\partial^2 \phi}{\partial x^2} = \frac{\partial^2 \phi}{\partial x'^2}$$

Similarly,

$$\frac{\partial^2 \phi}{\partial y^2} = \frac{\partial^2 \phi}{\partial y'^2} \quad \frac{\partial^2 \phi}{\partial z^2} = \frac{\partial^2 \phi}{\partial z'^2}$$

Moreover,

$$\frac{\partial \phi}{\partial t} = -v \frac{\partial \phi}{\partial x'} + \frac{\partial \phi}{\partial t'} \quad \frac{\partial^2 \phi}{\partial t^2} = \frac{\partial^2 \phi}{\partial t'^2} - 2v \frac{\partial^2 \phi}{\partial x' \partial t'} + v^2 \frac{\partial^2 \phi}{\partial x'^2}$$

Substituting these expressions in the wave equation gives

$$\frac{\partial^2 \phi}{\partial x'^2} + \frac{\partial^2 \phi}{\partial y'^2} + \frac{\partial^2 \phi}{\partial z'^2} - \frac{1}{c^2} \frac{\partial^2 \phi}{\partial t'^2} + \frac{1}{c^2} \left( 2v \frac{\partial^2 \phi}{\partial x' \partial t'} - v^2 \frac{\partial^2 \phi}{\partial x'^2} \right) = 0$$

Therefore the wave equation is *not* invariant under the Galilean transformations, for the form of the equation has changed.

The electromagnetic wave equation follows from Maxwell's equations of electromagnetic theory. By applying the procedure described here to Maxwell's equations, one finds that Maxwell's equations also are *not* invariant under Galilean transformations. Compare with Problem 6.23.

## Supplementary Problems

- 1.11.** A man ( $O'$ ) in the back of a 20-ft flatcar moving at 30 ft/s records that a flashbulb is fired in the front of the flatcar two seconds after he has passed a man ( $O$ ) on the ground. Find the coordinates of the event as determined by each observer. *Ans.*  $(x', t') = (20 \text{ ft}, 2 \text{ s}); (x, t) = (80 \text{ ft}, 2 \text{ s})$
- 1.12.** A boy sees a deer run directly away from him. The deer is running with a speed of 20 mi/hr. The boy gives chase and runs with a speed of 8 mi/hr. What is the speed of the deer relative to the boy? *Ans.* 12 mi/hr
- 1.13.** A boy in a train throws a ball in the forward direction with a speed of 20 mi/hr. If the train is moving with a speed of 80 mi/hr, what is the speed of the ball as measured by a man on the ground? *Ans.* 100 mi/hr

- 1.14.** A passenger walks backward along the aisle of a train with a speed of 2 mi/hr as the train moves along a straight track at a constant speed of 60 mi/hr with respect to the ground. What is the passenger's speed as measured by an observer standing on the ground? *Ans.* 58 mi/hr
- 1.15.** A conductor standing on a railroad platform synchronizes his watch with the engineer in the front of a train traveling at 60 mi/hr. The train is  $\frac{1}{4}$  mile long. Two minutes after the train leaves the platform a brakeman in the caboose lights a cigarette. What are the coordinates of the brakeman, as determined by the engineer and by the conductor, when the cigarette is lit?  
*Ans.*  $(x', t') = (-\frac{1}{4} \text{ mi}, 2 \text{ min})$ ;  $(x, t) = (1\frac{3}{4} \text{ mi}, 2 \text{ min})$
- 1.16.** A man sitting in a train lights two cigarettes, one ten minutes after the other. The train is moving in a straight line with a velocity of 20 m/s. What is the distance separation as measured by a man on the ground? *Ans.* 12,000 m
- 1.17.** A one-kilogram ball is constrained to move to the north at 3 m/s. It makes a perfectly elastic collision with an identical second ball which is at rest, and both balls move on a north-south axis after the collision. Compute, in the laboratory system, the total momentum before and after the collision.  
*Ans.* 3 kg · m/s
- 1.18.** For Problem 1.17 calculate the total energy before and after the collision. *Ans.* 4.5 J
- 1.19.** Refer to Problem 1.17. Calculate the total momentum before and after the collision as measured by an observer moving northwards at 1.5 m/s. *Ans.* 0
- 1.20.** For the observer in Problem 1.19 calculate the total energy before and after the collision.  
*Ans.* 2.25 J
- 1.21.** Repeat Problems 1.19 and 1.20 for an observer moving eastwards at 2 m/s.  
*Ans.* 5 kg · m/s 37° north of west; 8.5 J
- 1.22.** A person is in a boat moving eastwards with a speed of 15 ft/s. At the instant that the boat passes a dock a person on the dock throws a rock northwards. The rock strikes the water 6 s later at a distance of 150 ft from the dock. Find the coordinates of the splash as measured by the person in the boat.  
*Ans.*  $(x, y, t) = (-90 \text{ ft}, 150 \text{ ft}, 6 \text{ s})$
- 1.23.** Consider a one-dimensional, elastic collision that takes place along the  $x$ -axis of  $O$ . Show, from the classical transformation equations, that kinetic energy will also be conserved as determined by a second observer,  $O'$ , who moves with constant velocity  $u$  along the  $x$ -axis of  $O$ .



# Chapter 2

## The Postulates of Einstein

### 2.1 ABSOLUTE SPACE AND THE ETHER

A consequence of the Galilean velocity transformations is that if a certain observer measures a light signal to travel with the velocity  $c = 3 \times 10^8$  m/s, then any other observer moving relative to him will measure the same light signal to travel with a velocity different from  $c$ . What determines the particular reference frame such that if an observer is at rest relative to this frame, this privileged observer will measure the value  $c$  for the velocity of light signals?

Before Einstein it was generally believed that this privileged observer was the same observer for whom Maxwell's equations were valid. Maxwell's equations describe electromagnetic theory and predict that electromagnetic waves will travel with the speed  $c = 1/\sqrt{\epsilon_0 \mu_0} = 3 \times 10^8$  m/s. The space that was at rest with respect to this privileged observer was called "absolute space." Any other observer moving with respect to this absolute space would find the speed of light to be different from  $c$ . Since light is an electromagnetic wave, it was felt by 19th century physicists that a medium must exist through which the light propagated. Thus it was postulated that the "ether" permeated all of absolute space.

### 2.2 THE MICHELSON-MORLEY EXPERIMENT

If an ether exists, then an observer on the earth moving through the ether should notice an "ether wind." An apparatus with the sensitivity to measure the earth's motion through the hypothesized ether was developed by Michelson in 1881, and refined by Michelson and Morley in 1887. The outcome of the experiment was that *no motion through the ether was detected*. See Problems 2.4, 2.5 and 2.6.

### 2.3 LENGTH AND TIME MEASUREMENTS— A QUESTION OF PRINCIPLE

The one element common to both the null result of the Michelson-Morley experiment and the fact that Maxwell's equations hold only for a privileged observer is the Galilean transformations. These "obvious" transformations were reexamined by Einstein from what might be termed an "operational" point of view. Einstein took the approach that any quantity relevant to physical theories should, at least in principle, have a well-defined procedure by which it is measured. If such a procedure cannot be formulated, then the quantity should not be employed in physics.

Einstein could find no way to justify operationally the Galilean transformation  $t' = t$ , i.e. the statement that two observers *can* measure the time of an event to be the same. Consequently, the transformation  $t' = t$ , and with it the rest of the Galilean transformations, was rejected by Einstein.

### 2.4 THE POSTULATES OF EINSTEIN

Einstein's guiding idea, which he called the *Principle of Relativity*, was that *all* nonaccelerating observers should be treated equally in all respects, even if they are moving (at constant velocity) relative to each other. This principle can be formalized as follows:

**Postulate 1:** The laws of physics are the same (invariant) for all inertial (nonaccelerating) observers.